

## TENSILE STRENGTH OF COMPRESSED SNOW FORMED ON ROADS

By T. NAKAMURA, H. NAKAMURA, M. HIGASHIURA and O. ABE

(Shinjo Branch, National Research Center for Disaster Prevention, Shinjo, Yamagata-ken, Japan)

**ABSTRACT.** Tensile fracture strengths of fine-grained compressed high-density snow, of compressed and metamorphosed high-density snow-ice, of fine-grained naturally settled snow, and of snow-ice artificially made from the settled snow by freezing with absorbed water, were obtained at constant deformation speeds (constant strain-rates) in a temperature range of 264 to 270K. No remarkable temperature dependence of fracture stresses was observed in this temperature range. A critical deformation speed in a tensile test above which all the snow samples, except settled snow, fractured, was  $4.2 \times 10^{-7} \text{ m s}^{-1}$ . The fracture strength  $\sigma$  (in  $\text{N m}^{-2}$ ) varied with snow density  $\rho$  ( $\text{kg m}^{-3}$ ) as  $\sigma = 2.5 \times 10^4 \times 1.004 6^{\rho}$ . In a power-law relationship between strain-rate and maximum stress,  $\dot{\epsilon} \propto \sigma^n$ , the constant  $n$  obtained was 5.3 for all the unfractured snow samples.

## USING LINEAR DISCRIMINANT ANALYSIS TO CLASSIFY SNOWFALL SITUATIONS INTO AVALANCHING AND NON-AVALANCHING ONES

By N. F. DROZDOVSKAYA

(Central-Asian Regional Research Hydrometeorological Institute, Observatorskaya 72, Tashkent, GSP 700000, U.S.S.R.)

**ABSTRACT.** The existing methods of predicting avalanche danger often do not meet users' demands because of the empiric character of the insufficient volume of information used. In such forecasts the contribution of each individual parameter into the prognostic information is unknown, and this is very important when studying such an event as avalanche formation, which is conditioned by a complex interaction of numerous factors, including snow accumulation, the state of snow thickness, and the conditions of its development.

It is obvious that such problems can be successfully solved by statistical methods, and that explains the growing interest in numerical methods of avalanche forecasting. Problems of multi-dimensional observations arises in many scientific fields. The method suited for this problem is discriminant analysis, the purpose of which is to divide a multi-dimensional observation vector into predetermined classes.

This study considers the prognostic (diagnostic) problems of fresh-snow avalanches released during snowfall or in the two days after it has ceased. The theoretical basis is a complex of statistical methods: correlation and dispersion analysis, "sifting" for the choice of predictors' informative groups, construction of linear parametric discriminant functions, predictions based on training sample, and verification of discriminant functions based on independent material.

The archive used in the study consisted of 500 avalanching cases and 1 300 non-avalanching ones. All situations were grouped according to geomorphological characteristics. Each