

Nonlocal anisotropic interactions of Coulomb type

Maria Giovanna Mora 

Dipartimento di Matematica, Università di Pavia, via Ferrata 5, 27100 Pavia, Italy (mariagiovanna.mora@unipv.it)

(Received 22 May 2023; accepted 29 January 2024)

In this paper, we review some recent results on nonlocal interaction problems. The focus is on interaction kernels that are anisotropic variants of the classical Coulomb kernel. In other words, while preserving the same singularity at zero of the Coulomb kernel, they present preferred directions of interaction. For kernels of this kind and general confinement we will prove existence and uniqueness of minimizers of the corresponding energy. In the case of a quadratic confinement we will review a recent result by Carrillo and Shu about the explicit characterization of minimizers, and present a new proof, which has the advantage of being extendable to higher dimensions. In light of this result, we will re-examine some previous works motivated by applications to dislocation theory in materials science. Finally, we will discuss some related results and open questions.

Keywords: nonlocal energy; potential theory; anisotropic interaction; Coulomb potential; equilibrium measure

2020 *Mathematics Subject Classification* 31A15 (primary); 49K20 (secondary)

1. Introduction

The general goal of this review paper is the study of the minimization problem for an energy of the form:

$$I(\mu) = \iint_{\mathbb{R}^N \times \mathbb{R}^N} W(x - y) \, d\mu(y) \, d\mu(x) + \int_{\mathbb{R}^N} V(x) \, d\mu(x) \quad (1.1)$$

defined for $\mu \in \mathcal{P}(\mathbb{R}^N)$. Here, $\mathcal{P}(\mathbb{R}^N)$ stands for the space of probability measures in \mathbb{R}^N . In this formulation a measure $\mu \in \mathcal{P}(\mathbb{R}^N)$ represents the distribution of a family of particles in \mathbb{R}^N , the first integral in $I(\mu)$ is called the *interaction energy*, whereas the last integral usually plays the role of a *confinement energy*.

Energies as I arise as *mean-field limits* of discrete energies. More precisely, let us consider n particles in \mathbb{R}^N located at points x^1, x^2, \dots, x^n in \mathbb{R}^N and let us define as their interaction energy the quantity:

$$\frac{1}{n^2} \sum_{j=1}^n \sum_{k \neq j}^n W(x^j - x^k).$$

Since the order in which particles are considered is irrelevant, it is natural to assume the interaction kernel W to be an even function. This discrete energy describes *nonlocal* interactions in the sense that each particle interacts with any other particle in the system and not only with those in its immediate neighbourhood. If one identifies the distribution of particles with the so-called *empirical measure*:

$$\frac{1}{n} \sum_{j=1}^n \delta_{x^j},$$

one can show, under suitable assumptions of W and V , that the discrete energies:

$$\frac{1}{n^2} \sum_{j=1}^n \sum_{k \neq j} W(x^j - x^k) + \frac{1}{n} \sum_{j=1}^n V(x^j)$$

Γ -converge to I , as $n \rightarrow \infty$, with respect to the narrow convergence in $\mathcal{P}(\mathbb{R}^N)$ (see, e.g. [21]). In other words, minimizers of I describe the asymptotic behaviour of optimal distributions at the discrete level in the many-particle limit.

In many applications the typical interaction among particles is short-range repulsive and long-range attractive. This behaviour can be reproduced in the energy I by assuming:

- $W(x) \rightarrow +\infty$, as $x \rightarrow 0$, so that the interaction energy blows up when particles get too close to one another;
- $V(x) \rightarrow +\infty$ fast enough, as $|x| \rightarrow +\infty$, so that the confinement energy blows up when particles escape at infinity.

As a model example, we can consider as W the Coulomb kernel:

$$W(x) = \begin{cases} -\log|x| & \text{if } N = 2, \\ \frac{1}{|x|^{N-2}} & \text{if } N \geq 3, \end{cases}$$

and, as confinement potential, a power law $V(x) = |x|^p$ with $p > 0$ or the indicator of a given compact set $K \subset \mathbb{R}^N$:

$$V(x) = \begin{cases} 0 & \text{if } x \in K, \\ +\infty & \text{if } x \notin K. \end{cases}$$

In this last case (which we call *physical confinement*) minimizing I is equivalent to minimizing the sole interaction energy on the class of probability measures supported in K .

Continuum energies as I , as well as their discrete counterparts, are relevant in a variety of applications, ranging from physics (electrostatics, Coulomb gases, Ginzburg–Landau theory) to biology (population dynamics) and materials science. In particular, the Coulomb kernel is probably the most studied interaction kernel in physics and in mathematics.

Besides existence and uniqueness, one of the main questions in minimizing I is whether minimizers can be identified or at least some of their qualitative properties

can be established. For instance, can we determine the dimension of their supports and their shape? Is the distribution ‘regular’ on the support? One of the key difficulties in addressing these questions is the nonlocal nature of the problem: given a distribution μ , any local perturbation of μ , however small, will have a global impact on the interaction energy. Moreover, numerical simulations show that, according to the different choice of W and V , minimizers may present a rich variety of geometries and shapes (see, e.g. [12]).

In this paper, we will focus on the two-dimensional (2D) case $N = 2$ and on interaction kernels of the form:

$$W(x) = -\log|x| + \kappa(x), \tag{1.2}$$

where κ is an even 0-homogeneous function, smooth enough outside 0. The kernel W can be seen as a perturbation (not small though) of the 2D Coulomb kernel. Since κ is 0-homogeneous, $\kappa(x)$ depends only on the angle that x forms with respect to a given reference axis. In this sense we call κ an *anisotropic kernel*, meaning that it introduces some preferred directions of interaction.

The goal of this paper is to review the most recent results about existence, uniqueness, and characterization of minimizers for kernels W of the form (1.2) and general confinements V . The common thread of these results is the following key idea. A clever way to look at this class of problems is via Fourier analysis. In fact, if we denote by \widehat{f} the Fourier transform of f given by

$$\widehat{f}(\xi) = \frac{1}{2\pi} \int_{\mathbb{R}^2} f(x)e^{-i\xi \cdot x} dx \quad \text{for } \xi \in \mathbb{R}^2,$$

formally we have

$$\widehat{W * \mu} = 2\pi \widehat{W} \widehat{\mu}$$

and by Plancherel theorem:

$$\begin{aligned} \iint_{\mathbb{R}^2 \times \mathbb{R}^2} W(x - y) d\mu(y) d\mu(x) &= \int_{\mathbb{R}^2} (W * \mu) d\mu = \int_{\mathbb{R}^2} \widehat{W * \mu}(\xi) \overline{\widehat{\mu}(\xi)} d\xi \\ &= 2\pi \int_{\mathbb{R}^2} \widehat{W} |\widehat{\mu}|^2 d\xi. \end{aligned} \tag{1.3}$$

In other words, the nonlocal interaction can be expressed in a local form in the Fourier space. Note, however, that (1.3) holds true only under specific assumptions for \widehat{W} and μ (see proposition 3.2). Using (1.3) we will show that a sign condition on \widehat{W} guarantees strict convexity of the energy and, therefore, uniqueness of minimizers (see § 3). Moreover, the inversion formula for the Fourier transform will be a crucial ingredient in the characterization results of § 4.

1.1. Motivation

The study of interaction kernels of the form (1.2) is motivated by materials science, more precisely, by *dislocation theory*. Dislocations are defects in the crystalline lattice of a metal, whose presence and concerted movement favour plastic





Figure 1. Cubic crystal with a dislocation.

Figure 2. Burgers circuit and the Burgers vector b .

slips, that is, relative slips of atomic layers, that macroscopically result into a shearing plastic deformation.

Let us consider an idealized three-dimensional (3D) cubic lattice, where all 2D sections along a certain direction are assumed to be identical. In this simplified 2D setting a dislocation of edge type looks as shown in figure 1.

The presence of a dislocation can be detected as follows. One draws a so-called *Burgers circuit*, that is, a closed circuit enclosing the defect. If we draw the same circuit in a perfect reference crystal, the circuit does not close up, see figure 2. The vector that needs to be added to close the circuit is defined as the Burgers vector. The Burgers vector is thus a measure of the discrepancy between the distorted lattice and a perfect lattice.

Assume now that every dislocation has the same Burgers vector (which for simplicity we set to be e_1). We would like to compute the interaction force between two dislocations, the so-called *Peach–Köhler force*. To make this computation we

consider a semi-discrete setting, where the crystal is described as a continuum medium (as if the discrete lattice structure were averaged out), but dislocations are still modelled as point singularities. In the framework of linear elasticity the fundamental strain generated by a dislocation of Burgers vector e_1 located at 0 is a solution $\beta : \mathbb{R}^2 \rightarrow \mathbb{M}^{2 \times 2}$ of the problem:

$$\begin{cases} \operatorname{div} \mathbb{C}\beta = 0 & \text{in } \mathbb{R}^2, \\ \operatorname{curl} \beta = \delta_0 e_1 & \text{in } \mathbb{R}^2, \end{cases}$$

where \mathbb{C} is the tensor of linear elasticity and δ_0 denotes the Dirac delta at 0. The Peach–Köhler force can then be computed by the formula:

$$F = \mathbb{C}\beta e_1 \times e_3,$$

where $\mathbb{C}\beta$ is the stress associated with the fundamental strain, e_1 is the Burgers vector, and e_3 represents the dislocation line (in our simplified 2D setting the dislocation line is orthogonal to the plane of the 2D section). For $x \in \mathbb{R}^2$ the Peach–Köhler force $F(x)$ is the configurational force that a dislocation at x experiences because of the dislocation at 0. It turns out (see [11, Chapter 13-4]) that

$$F = -(c\nabla W, 0)$$

for some positive material constant $c > 0$, where W is of the form (1.2) with

$$\kappa(x) = -\frac{1}{4} \frac{a+b}{a} \log \left(\frac{x_1^2 + (a+b)^2 x_2^2}{|x|^2} \right) + \frac{1}{4} \frac{b-a}{a} \log \left(\frac{x_1^2 + (b-a)^2 x_2^2}{|x|^2} \right) \quad (1.4)$$

for $x = (x_1, x_2) \in \mathbb{R}^2$. Here, $b > a > 0$ are material constants. If the medium is isotropic (which corresponds to $a \rightarrow 0^+$, $b = 1$), the above expression reduces, up to additive constants, to

$$\kappa(x) = \frac{x_1^2}{|x|^2}. \quad (1.5)$$

We would like to predict the optimal distribution of dislocations at equilibrium in this setting. For simplicity, let us focus on the isotropic case (1.5) and assume to have exactly two dislocations located at x and y . Their interaction energy is given by

$$-\log|x-y| + \frac{(x_1-y_1)^2}{|x-y|^2}.$$

The Coulomb term forces x and y to be as far as possible (this repulsive behaviour is counterbalanced by the presence of some confinement), whereas the anisotropic term is minimized when $x_1 = y_1$, that is, when the two dislocations are aligned vertically. Does the same phenomenon occur in the mesoscopic description (1.1)?

In other words, let I be the energy defined by

$$I(\mu) = \iint_{\mathbb{R}^2 \times \mathbb{R}^2} \left(-\log|x - y| + \frac{(x_1 - y_1)^2}{|x - y|^2} \right) d\mu(y) d\mu(x) + \int_{\mathbb{R}^2} V(x) d\mu(x), \tag{1.6}$$

where $\mu \in \mathcal{P}(\mathbb{R}^2)$ represents now the distribution of a family of dislocations of Burgers vector e_1 . Is it true that minimizers of I have a one-dimensional (1D) vertical support? And if so, what is their distribution? This last question is actually not difficult to answer. In fact, let $\mu \in \mathcal{P}(\mathbb{R}^2)$ be a measure of the form $\mu = \delta_{\bar{x}_1} \otimes \nu$ with $\bar{x}_1 \in \mathbb{R}$ and $\nu \in \mathcal{P}(\mathbb{R})$. When restricted to measures of this form the energy I reduces to

$$J(\bar{x}_1, \nu) := - \iint_{\mathbb{R} \times \mathbb{R}} \log|s - t| d\nu(t) d\nu(s) + \int_{\mathbb{R}} V(\bar{x}_1, t) d\nu(t).$$

Thus, if the minimizer of I has a vertical support, its vertical projection minimizes the 1D functional $J(\bar{x}_1, \cdot)$ among all measures in $\mathcal{P}(\mathbb{R})$. The functional J is known in the literature as the log-gas energy and its minimizers can be explicitly computed for several confinements. For instance, for $V(x) = |x|^2$ the unique minimizer of $J(\bar{x}_1, \cdot)$ is the so-called semicircle law, that is, the measure:

$$\frac{1}{\pi} \sqrt{2 - t^2} \mathcal{L}^1 \llcorner [-\sqrt{2}, \sqrt{2}](t)$$

(and the optimal choice for \bar{x}_1 is clearly $\bar{x}_1 = 0$). Therefore, in the case $V(x) = |x|^2$, if the energy I in (1.6) has a minimizer μ with a vertical support, then necessarily μ is given by the semicircle law on the vertical axis

$$\frac{1}{\pi} \delta_0(x_1) \otimes \sqrt{2 - x_2^2} \mathcal{H}^1 \llcorner [-\sqrt{2}, \sqrt{2}](x_2). \tag{1.7}$$

1.2. Overview of the results

We now briefly review the main results about explicit characterization of minimizers for kernels of the form (1.2). We consider an energy of the form (1.6) with confinement $V(x) = |x|^2$ and we introduce a parameter $\alpha \in \mathbb{R}$ in front of the anisotropic term. The role of α is that of tuning the strength of the anisotropic interaction. The energy

$$I_\alpha(\mu) = \iint_{\mathbb{R}^2 \times \mathbb{R}^2} \left(-\log|x - y| + \alpha \frac{(x_1 - y_1)^2}{|x - y|^2} \right) d\mu(y) d\mu(x) + \int_{\mathbb{R}^2} |x|^2 d\mu(x) \tag{1.8}$$

has a unique minimizer, which can be characterized as follows:

- if $\alpha = 0$ (purely Coulomb case), the minimizer is the so-called *circle law*

$$\frac{1}{\pi} \chi_{B_1(0)};$$

the derivation of this classical result is attributed to [8, 10, 17];

- if $\alpha = 1$ (dislocation case), the minimizer is the semicircle law on the vertical axis, that is, the measure (1.7); this was proved in [18];
- if $\alpha \in (0, 1)$, the minimizer is given by the *ellipse law*:

$$\frac{1}{|E_\alpha|} \chi_{E_\alpha}, \tag{1.9}$$

where

$$E_\alpha = \left\{ x \in \mathbb{R}^2 : \frac{x_1^2}{1 - \alpha} + \frac{x_2^2}{1 + \alpha} \leq 1 \right\};$$

this result is contained in [2].

We mention that the minimizer of I_α can be actually identified for any value of $\alpha \in \mathbb{R}$, see theorem 5.1.

The above results show that at $\alpha = 1$ the anisotropy has a dramatic effect on the structure of the minimizer and, in particular, on its dimensionality. For any $\alpha \in [0, 1)$ the minimizer is given by a uniform distribution on a 2D set and it is only at $\alpha = 1$ that an abrupt loss of dimensionality occurs. A natural question is whether this phenomenon can be explained in mathematical terms and how much of this analysis is bound to the specific choice of the anisotropy.

In [15], we conjectured that the loss of dimensionality could be related to a change of sign of the Fourier transform of the interaction kernel. Indeed, we have that

$$\widehat{W}_\alpha(\xi) = c_\alpha \delta_0 + \frac{(1 - \alpha)\xi_1^2 + (1 + \alpha)\xi_2^2}{|\xi|^4},$$

see (5.2). Thus, for $\alpha \in [0, 1)$ the Fourier transform of W_α is strictly positive outside 0, whereas it is only non-negative for $\alpha = 1$.

Let us consider now an interaction kernel of the form (1.2) with a general even and 0-homogeneous anisotropy κ , and confinement $V(x) = |x|^2$. In [14], it was proved that, if κ is small enough in $C^3(\mathbb{S}^1)$, then the minimizer is unique and given by the normalized characteristic function of the domain enclosed by an ellipse centred at the origin. Note that the smallness assumption on κ implies in particular that the Fourier transform of W outside 0 is strictly positive.

Recently, Carrillo and Shu proved that the smallness assumption in [14] can be lifted, as long as the condition on the non-negativity of the Fourier transform of W outside 0 is preserved. More precisely, in [3] the following remarkable result was proved: let W be an interaction kernel of the form (1.2) with κ even, 0-homogeneous, and smooth enough on \mathbb{S}^1 , and let the confinement be $V(x) = |x|^2$; then,

- if $\widehat{W} > 0$ outside 0, the unique minimizer is the normalized characteristic function of the domain enclosed by an ellipse centred at the origin;
- if $\widehat{W} \geq 0$ outside 0, the unique minimizer is either as above or is a semicircle law on a line passing through the origin.

This result sheds some light on the relation between loss of dimensionality and change of sign of the Fourier transform: for this class of kernels loss of dimensionality cannot occur as long as the Fourier transform is ‘not degenerate’. However, explicit examples show that for a degenerate Fourier transform both cases (ellipse or semicircle law) can indeed occur, see remark 5.2.

In this article, we will first review the existence and uniqueness results for interaction kernels of the form (1.2) and general confinements. In the case of the quadratic confinement $V(x) = |x|^2$ we will present a new proof of the result [3] by Carrillo and Shu, which follows the approach proposed in [16] for the analogous problem in 3D. This different approach has the advantage of being extendable to higher dimensions. We will then deduce the characterization of minimizers in the dislocation case (1.5) from this general result. Finally, we will examine an example with physical confinement given by the domain enclosed by an ellipse and discuss some open questions.

2. The existence result and first properties of minimizers

Throughout the paper, we will consider an interaction kernel of the form:

$$W(x) = -\log|x| + \kappa(x) \quad \text{for } x \neq 0, \quad W(0) = +\infty, \quad (2.1)$$

where κ is an even 0-homogeneous function of class H^s on \mathbb{S}^1 with $s > 3/2$. We will denote the 2D Coulomb kernel by W_0 , that is,

$$W_0(x) = -\log|x| \quad \text{for } x \neq 0, \quad W_0(0) = +\infty.$$

Since κ is bounded on $\mathbb{R}^2 \setminus \{0\}$, there exist two constants $C_1, C_2 \in \mathbb{R}$ such that

$$W_0(x) + C_1 \leq W(x) \leq W_0(x) + C_2 \quad \text{for every } x \in \mathbb{R}^2. \quad (2.2)$$

2.1. Logarithmic capacity

For any compact set $K \subset \mathbb{R}^2$ we define the logarithmic capacity of K as

$$\text{cap}(K) = \Phi \left(\inf_{\mu \in \mathcal{P}(K)} \iint_{K \times K} W_0(x-y) \, d\mu(x) \, d\mu(y) \right),$$

where $\Phi(t) = e^{-t}$ for $t \in \mathbb{R}$ and $\Phi(+\infty) = 0$. Note that the integral above is well defined (possibly equal to $+\infty$), since the integrand is bounded from below on $K \times K$.

If $B \subset \mathbb{R}^2$ is a Borel set, we define its capacity as

$$\text{cap}(B) = \sup \{ \text{cap}(K) : K \text{ compact, } K \subset B \}.$$

We will say that a property holds *quasi everywhere* (q.e.) if it holds up to sets of zero capacity.

The key property of the capacity is the following: if $\mu \in \mathcal{P}(\mathbb{R}^2)$ has compact support and satisfies:

$$\iint_{\mathbb{R}^2 \times \mathbb{R}^2} W(x - y) \, d\mu(x) \, d\mu(y) < +\infty, \tag{2.3}$$

then $\mu(B) = 0$ for every Borel set B with $\text{cap}(B) = 0$. In other words, if a property holds q.e., then it holds μ -a.e. for any $\mu \in \mathcal{P}(\mathbb{R}^2)$ with compact support and satisfying (2.3) (as we will see, these are the relevant measures for the minimization problem under study). This can be shown as follows: assume by contradiction that $\mu(B) > 0$, where $B \subset \mathbb{R}^2$ is a Borel set with $\text{cap}(B) = 0$. Then there exists a compact set $K \subset B$ such that $\mu(K) > 0$. Since μ is compactly supported, there exists $C_0 > 0$ such that

$$W(x - y) \geq -C_0 \quad \text{for every } (x, y) \in (\text{supp } \mu)^2.$$

Therefore, setting

$$\nu := \frac{1}{\mu(K)} \mu \llcorner K,$$

by (2.3) we have

$$\begin{aligned} \iint_{K \times K} W(x - y) \, d\nu(x) \, d\nu(y) &\leq \frac{1}{\mu(K)^2} \iint_{\mathbb{R}^2 \times \mathbb{R}^2} W(x - y) \, d\mu(x) \, d\mu(y) \\ &\quad + \frac{C_0}{\mu(K)^2} - C_0 < +\infty. \end{aligned}$$

On the other hand, from (2.2) it follows that

$$\iint_{K \times K} W(x - y) \, d\nu(x) \, d\nu(y) \geq \iint_{K \times K} W_0(x - y) \, d\nu(x) \, d\nu(y) + C_1.$$

This would imply that $\text{cap}(K) > 0$, contradicting the assumption $\text{cap}(B) = 0$.

In a similar way, one can show that a countable union of sets with zero capacity has zero capacity.

2.2. The confinement potential

In this section, we assume the confinement potential $V : \mathbb{R}^2 \rightarrow \mathbb{R} \cup \{+\infty\}$ to be a lower semicontinuous function, which is bounded from below and satisfies the following conditions:

$$\lim_{|x| \rightarrow +\infty} \left(\frac{1}{2} V(x) - \log |x| \right) = +\infty \tag{2.4}$$

and

$$\text{cap}(\{x \in \mathbb{R}^2 : V(x) < +\infty\}) > 0. \tag{2.5}$$

Examples of admissible confinements are the power laws $V(x) = |x|^p$ with $p > 0$ or the indicator function of a compact set of positive capacity.

2.3. Main results

For every $\mu \in \mathcal{P}(\mathbb{R}^2)$ we define:

$$I(\mu) = \iint_{\mathbb{R}^2 \times \mathbb{R}^2} \left(W(x - y) + \frac{1}{2}V(x) + \frac{1}{2}V(y) \right) d\mu(y) d\mu(x), \tag{2.6}$$

where the interaction kernel W is as in (2.1) and the confinement potential V as in § 2.2.

THEOREM 2.1 [7, 20]. *The energy I is well defined and has a minimizer in $\mathcal{P}(\mathbb{R}^2)$. If μ is a minimizer of I , then $I(\mu) < +\infty$, the support of μ is a compact set, and μ satisfies the following Euler–Lagrange conditions: there exists a constant $c = c(\mu)$ such that*

$$(W * \mu)(x) + \frac{1}{2}V(x) = c \quad \text{for } \mu\text{-a.e. } x \in \text{supp } \mu, \tag{2.7}$$

and

$$(W * \mu)(x) + \frac{1}{2}V(x) \geq c \quad \text{for q.e. } x \in \mathbb{R}^2. \tag{2.8}$$

Proof. The function $(x, y) \mapsto W(x - y) + \frac{1}{2}V(x) + \frac{1}{2}V(y)$ is lower semicontinuous and blows up at infinity by (2.4), hence it is bounded from below by a constant $-c_1$ with $c_1 > 0$. Therefore, the energy I in (2.6) is well defined (possibly equal to $+\infty$) and $\inf I > -\infty$. Note also that the two representations (1.1) and (2.4) coincide whenever the interaction energy is well defined and not equal to $-\infty$.

Assumption (2.5) guarantees that $\inf I < +\infty$. Indeed, writing $\{V < +\infty\}$ as the union over $n \in \mathbb{N}$ of the compact sets $\{V \leq n\}$, we must have $\text{cap}(\{V \leq n_0\}) > 0$ for some $n_0 \in \mathbb{N}$, that is, there exists a probability measure μ_0 with support in $\{V \leq n_0\}$ such that

$$\iint_{\mathbb{R}^2 \times \mathbb{R}^2} W_0(x - y) d\mu_0(x) d\mu_0(y) < +\infty.$$

By (2.2) this implies that

$$\iint_{\mathbb{R}^2 \times \mathbb{R}^2} W(x - y) d\mu_0(x) d\mu_0(y) < +\infty.$$

On the other hand,

$$\int_{\mathbb{R}^2} V(x) d\mu_0(x) \leq n_0,$$

hence $I(\mu_0) < +\infty$.

Existence of a minimizer follows by the Direct Method of the Calculus of Variations. Let $(\mu_n)_n$ be a minimizing sequence. Since $\inf I < +\infty$, there exists a constant $C > 0$ such that $I(\mu_n) \leq C$ for every $n \in \mathbb{N}$. By (2.4) for every $M > 0$ there exists

a compact set $K \subset \mathbb{R}^2$ such that

$$W(x - y) + \frac{1}{2}V(x) + \frac{1}{2}V(y) \geq M \quad \text{for } (x, y) \notin K \times K.$$

Therefore, for every $n \in N$:

$$\begin{aligned} C &\geq I(\mu_n) \geq M(\mu_n \otimes \mu_n)((K \times K)^c) - c_1(\mu_n(K))^2 \\ &\geq M(1 - (\mu_n(K))^2) - c_1 \\ &\geq M\mu_n(K^c) - c_1. \end{aligned}$$

This inequality implies that the sequence $(\mu_n)_n$ is tight, hence, up to subsequences, $(\mu_n)_n$ converges narrowly to some $\mu \in \mathcal{P}(\mathbb{R}^2)$ (we refer to [1, Chapter 5] for the definition of tightness, narrow convergence, and their properties). Since the integrand in I is lower semicontinuous and bounded from below, we have:

$$I(\mu) \leq \liminf_{n \rightarrow \infty} I(\mu_n),$$

hence μ is a minimizer.

Let now μ be a minimizer. In particular, $I(\mu) < +\infty$. By (2.4) there exists a compact set $K \subset \mathbb{R}^2$ such that

$$W(x - y) + \frac{1}{2}V(x) + \frac{1}{2}V(y) \geq I(\mu) + 1 \quad \text{for } (x, y) \notin K \times K. \tag{2.9}$$

By taking K larger if needed, we can assume that $\mu(K) > 0$. We claim that $\text{supp } \mu \subset K$. Assume by contradiction that $\mu(K) < 1$ and define:

$$\tilde{\mu} := \frac{1}{\mu(K)}\mu \llcorner K \in \mathcal{P}(\mathbb{R}^2).$$

By (2.9) we deduce that

$$\begin{aligned} I(\tilde{\mu}) &= \frac{1}{(\mu(K))^2} \left(I(\mu) - \iint_{(K \times K)^c} \left(W(x - y) + \frac{1}{2}V(x) + \frac{1}{2}V(y) \right) d\mu(y) d\mu(x) \right) \\ &\leq \frac{1}{(\mu(K))^2} (I(\mu) - (I(\mu) + 1)(1 - (\mu(K))^2)) \\ &= I(\mu) + 1 - \frac{1}{(\mu(K))^2} < I(\mu), \end{aligned}$$

where the last inequality follows from the fact that $\mu(K) < 1$. This contradicts the minimality of μ .

We conclude by showing that μ satisfies (2.7)–(2.8). Let $\nu \in \mathcal{P}(\mathbb{R}^2)$ be a competitor such that its support is compact and $I(\nu) < +\infty$. For $\varepsilon \in (0, 1)$ we have that

$(1 - \varepsilon)\mu + \varepsilon\nu \in \mathcal{P}(\mathbb{R}^2)$. Therefore, by minimality:

$$I(\mu) \leq I((1 - \varepsilon)\mu + \varepsilon\nu).$$

Expanding the energy on the right-hand side yields:

$$0 \leq -2\varepsilon \int_{\mathbb{R}^2} W * \mu \, d\mu + 2\varepsilon \int_{\mathbb{R}^2} W * \mu \, d\nu + \varepsilon \int_{\mathbb{R}^2} V \, d(\nu - \mu) + O(\varepsilon^2).$$

Dividing by 2ε and sending $\varepsilon \rightarrow 0^+$ lead to the following inequality:

$$\int_{\mathbb{R}^2} \left(W * \mu + \frac{1}{2}V \right) d\nu \geq \int_{\mathbb{R}^2} \left(W * \mu + \frac{1}{2}V \right) d\mu =: c \tag{2.10}$$

for every $\nu \in \mathcal{P}(\mathbb{R}^2)$ with compact support and $I(\nu) < +\infty$.

Set $P := W * \mu + \frac{1}{2}V$ and assume by contradiction that

$$\text{cap}(\{x \in \mathbb{R}^2 : P(x) < c\}) > 0.$$

Note that by Fatou’s lemma P is lower semicontinuous, thus $\{P < c\}$ is a Borel set. By definition of capacity there exists a compact set $K \subset \mathbb{R}^2$ of positive capacity such that $P(x) < c$ for every $x \in K$. Therefore, there exists $\nu \in \mathcal{P}(K)$ such that

$$\iint_{\mathbb{R}^2 \times \mathbb{R}^2} W(x - y) \, d\nu(x) \, d\nu(y) < +\infty.$$

Moreover, the confinement energy of ν is also finite, since

$$\int_{\mathbb{R}^2} \frac{1}{2}V \, d\nu = \int_K \frac{1}{2}V \, d\nu < \int_K (c - W * \mu) \, d\nu = c - \int_K W * \mu \, d\nu \tag{2.11}$$

and the right-hand side is finite by the bound from below of W on compact sets. Having finite energy and compact support, the measure ν has to satisfy condition (2.10). However, (2.11) contradicts (2.10). This proves (2.8).

To prove (2.7) we note that by (2.8) we have $W * \mu + \frac{1}{2}V \geq c$ q.e., hence μ -a.e. On the other hand, c is by definition the integral of $W * \mu + \frac{1}{2}V$ with respect to μ , so necessarily (2.7) holds true. □

REMARK 2.2. The previous result applies to much more general interaction kernels. Note in particular that the only properties of κ we used are boundedness and lower semicontinuity.

3. The uniqueness result

As mentioned in the Introduction, the key assumption to guarantee uniqueness of the minimizer is the sign condition $\widehat{W} \geq 0$ outside 0. In fact, this assumption implies the strict convexity of the energy (when restricted to a suitable class). This result bears some similarities to Bochner theorem, which characterizes functions of positive type as those whose Fourier transform is a positive finite measure, see,

e.g. [23]. Functions of positive type provide in fact interaction kernels whose corresponding energies are convex on discrete measures. However, the regularity of W and \widehat{W} does not allow for a direct application of this theorem in our context.

Once strict convexity is established, not only the minimizer is unique, but the two Euler–Lagrange conditions (2.7)–(2.8) are equivalent to minimality. Thus, identifying the unique minimizer reduces to finding a measure satisfying (2.7)–(2.8). In order to do so it is essential to have an expression for $W * \mu$, in particular inside the support of μ where we need to verify (2.7). As we will see, the main idea is to rely again on Fourier analysis applying the inversion formula for the Fourier transform.

3.1. The Fourier transform

We denote the Schwartz space of rapidly decreasing functions by \mathcal{S} and its dual space by \mathcal{S}' , the so-called space of tempered distributions. We recall that for every $\varphi \in \mathcal{S}$ its Fourier transform $\widehat{\varphi} \in \mathcal{S}$ is defined as

$$\widehat{\varphi}(\xi) = \frac{1}{2\pi} \int_{\mathbb{R}^2} \varphi(x) e^{-i\xi \cdot x} dx \quad \text{for } \xi \in \mathbb{R}^2.$$

The map $\varphi \mapsto \widehat{\varphi}$ is a continuous linear isomorphism (with continuous inverse) of \mathcal{S} into itself. This allows one to extend the definition of Fourier transform to elements of \mathcal{S}' simply by duality, that is, the Fourier transform \widehat{u} of an element $u \in \mathcal{S}'$ is defined as

$$\langle \widehat{u}, \varphi \rangle = \langle u, \widehat{\varphi} \rangle \quad \text{for every } \varphi \in \mathcal{S}.$$

We recall that the Fourier transform of the tempered distribution δ_0 (the Dirac delta at 0) is the constant $1/(2\pi)$. Moreover, for every $\varphi \in \mathcal{S}$ and $j = 1, 2$ we have

$$\widehat{\partial_{x_j} \varphi}(\xi) = i\xi_j \widehat{\varphi}(\xi) \quad \text{for } \xi \in \mathbb{R}^2.$$

3.2. The Fourier transform of the kernel W

Both W and W_0 are locally integrable functions with sublinear growth at infinity, hence they are tempered distributions. Thus, their Fourier transforms are well defined as tempered distributions. We start by computing the Fourier transform of the Coulomb kernel W_0 . Since $\Delta W_0 = -2\pi\delta_0$, formally we have

$$-|\xi|^2 \widehat{W_0}(\xi) = \widehat{\Delta W_0}(\xi) = -2\pi \widehat{\delta_0}(\xi) = -1.$$

Hence, we would expect $\widehat{W_0}(\xi) = 1/|\xi|^2$. However, this function is not integrable at 0 and so, it does not define a tempered distribution. In fact, the correct expression of $\widehat{W_0}$ is as follows:

$$\langle \widehat{W_0}, \varphi \rangle = c_0 \varphi(0) + \int_{|\xi| \leq 1} \frac{1}{|\xi|^2} (\varphi(\xi) - \varphi(0)) d\xi + \int_{|\xi| > 1} \frac{1}{|\xi|^2} \varphi(\xi) d\xi \quad (3.1)$$

for every $\varphi \in \mathcal{S}$. Here, $c_0 = (1/2\pi)(\gamma + \log \pi)$, where γ is the Euler constant. For the proof of this formula we refer to [6] or [18].

To compute the Fourier transform of the anisotropic kernel κ , it is convenient to pass to complex variables to simplify notation. We replace x by $z \in \mathbb{C}$ and write

$z = |z|e^{i\theta}$ with $\theta \in [0, 2\pi]$. Since κ is 0-homogeneous, we have that $\kappa(z) = \kappa(e^{i\theta})$ and thus, it can be written as a Fourier series for $\theta \in [0, 2\pi]$. Moreover, since κ is even, its Fourier series contains only the even terms, that is:

$$\kappa(e^{i\theta}) = a_0 + \sum_{n=1}^{\infty} (a_{2n} \cos(2n\theta) + b_{2n} \sin(2n\theta))$$

with $(a_{2n})_{n \in \mathbb{N}}, (b_{2n})_{n \in \mathbb{N}} \in \ell^2$. Note that for $n \in \mathbb{N}$:

$$\cos(n\theta) = \operatorname{Re} \frac{z^n}{|z|^n} =: \frac{\phi_n(z)}{|z|^n} \quad \text{and} \quad \sin(n\theta) = \operatorname{Im} \frac{z^n}{|z|^n} =: \frac{\psi_n(z)}{|z|^n}.$$

The functions ϕ_n, ψ_n are harmonic homogeneous polynomials of order n (in fact, they correspond to the so-called spherical harmonics in dimension 2). We can thus rewrite:

$$\kappa(z) = a_0 + \sum_{n=1}^{\infty} \left(a_{2n} \frac{\phi_{2n}(z)}{|z|^{2n}} + b_{2n} \frac{\psi_{2n}(z)}{|z|^{2n}} \right). \tag{3.2}$$

Without loss of generality, we can assume that $a_0 = 0$. In fact, adding a constant to κ does not affect the minimization problem under study. Expression (3.2) is particularly convenient to compute the Fourier transform of κ , owing to the following result, whose proof can be found in [22].

LEMMA 3.1. *Let ϕ be a harmonic homogeneous polynomial of degree $m \geq 1$ in \mathbb{R}^2 . Then the Fourier transform of $\phi(x)/|x|^m$ is given by*

$$\gamma_m \frac{\phi(\xi)}{|\xi|^{m+2}} \tag{3.3}$$

for a suitable constant γ_m . For m even, $m = 2n$, one has $\gamma_{2n} = (-1)^n 2n$.

Note that $\phi(\xi)/|\xi|^{m+2}$ behaves as $1/|\xi|^2$ for ξ close to zero, therefore it is not integrable at 0. Formula (3.3) has to be interpreted as in (3.1), that is:

$$\left\langle \gamma_m \frac{\phi}{|\cdot|^{m+2}}, \varphi \right\rangle = \gamma_m \int_{|\xi| \leq 1} \frac{\phi(\xi)}{|\xi|^{m+2}} (\varphi(\xi) - \varphi(0)) \, d\xi + \gamma_m \int_{|\xi| > 1} \frac{\phi(\xi)}{|\xi|^{m+2}} \varphi(\xi) \, d\xi$$

for every $\varphi \in \mathcal{S}$.

By lemma 3.1 and (3.2) with $a_0 = 0$ we deduce that

$$\widehat{\kappa}(\xi) = \sum_{n=1}^{\infty} \left((-1)^n 2n a_{2n} \frac{\phi_{2n}(\xi)}{|\xi|^{2n+2}} + (-1)^n 2n b_{2n} \frac{\psi_{2n}(\xi)}{|\xi|^{2n+2}} \right). \tag{3.4}$$

The argument can be made rigorous if $(2na_{2n})_{n \in \mathbb{N}}, (2nb_{2n})_{n \in \mathbb{N}} \in \ell^2$, that is, if $\kappa \in H^1(\mathbb{S}^1)$.

By (3.1) and (3.4) we can write:

$$\widehat{W}(\xi) = c_0 \delta_0 + \frac{1}{|\xi|^2} + \frac{\widehat{\kappa}(\xi/|\xi|)}{|\xi|^2} =: c_0 \delta_0 + \frac{\widehat{\Psi}(\xi)}{|\xi|^2}, \tag{3.5}$$

where, with an abuse of notation, $\widehat{\Psi}$ denotes the ‘angular part’ of \widehat{W} . Note that $\widehat{\Psi}$ is even and 0-homogeneous, and that the above formula has to be interpreted as

in (3.1). In the following, we will need $\widehat{\Psi} \in C^0(\mathbb{S}^1)$. By Sobolev embedding this is true if $\widehat{\kappa} \in H^p(\mathbb{S}^1)$ with $p > 1/2$. This is guaranteed by our assumption $\kappa \in H^s(\mathbb{S}^1)$ with $s > 3/2$, owing to (3.4).

In the following, we will refer to $\widehat{\Psi}$ as the even and 0-homogeneous function given by formula (3.5). Note that the purely Coulomb case corresponds to $\widehat{\Psi} \equiv 1$. Moreover, by (3.4) and (3.5) we have

$$\frac{1}{2\pi} \int_{\mathbb{S}^1} \widehat{\Psi}(\xi) \, d\mathcal{H}^1(\xi) = \frac{1}{2\pi} \int_{\mathbb{S}^1} (1 + \widehat{\kappa}(\xi)) \, d\mathcal{H}^1(\xi) = 1. \tag{3.6}$$

PROPOSITION 3.2. *Assume $\widehat{\Psi} \geq 0$ on \mathbb{S}^1 . Let $\mu_0, \mu_1 \in \mathcal{P}(\mathbb{R}^2)$ be two measures with compact support and finite interaction energy. Let $\nu := \mu_0 - \mu_1$. Then*

$$\int_{\mathbb{R}^2} W * \nu \, d\nu = 2\pi \int_{\mathbb{R}^2} \frac{\widehat{\Psi}(\xi)}{|\xi|^2} |\widehat{\nu}(\xi)|^2 \, d\xi. \tag{3.7}$$

In particular, the left-hand side is non-negative and is equal to zero if and only if $\nu = 0$, that is, if $\mu_0 = \mu_1$.

An immediate consequence of proposition 3.2 is the following: if $\widehat{\Psi} \geq 0$ on \mathbb{S}^1 , then I is strictly convex on the class of measures with compact support and finite interaction energy. Indeed, let $\mu_0, \mu_1 \in \mathcal{P}(\mathbb{R}^2)$ be two such measures with $\mu_0 \neq \mu_1$. By proposition 3.2 we have that

$$\int_{\mathbb{R}^2} W * (\mu_0 - \mu_1) \, d(\mu_0 - \mu_1) > 0,$$

hence

$$\int_{\mathbb{R}^2} W * \mu_0 \, d\mu_0 + \int_{\mathbb{R}^2} W * \mu_1 \, d\mu_1 > 2 \int_{\mathbb{R}^2} W * \mu_0 \, d\mu_1.$$

Let now $\mu_t := (1 - t)\mu_0 + t\mu_1$ with $t \in (0, 1)$. From the above inequality we conclude that

$$\begin{aligned} \int_{\mathbb{R}^2} W * \mu_t \, d\mu_t &< (1 - t)^2 \int_{\mathbb{R}^2} W * \mu_0 \, d\mu_0 + t^2 \int_{\mathbb{R}^2} W * \mu_1 \, d\mu_1 \\ &\quad + t(1 - t) \left(\int_{\mathbb{R}^2} W * \mu_0 \, d\mu_0 + \int_{\mathbb{R}^2} W * \mu_1 \, d\mu_1 \right) \\ &= (1 - t) \int_{\mathbb{R}^2} W * \mu_0 \, d\mu_0 + t \int_{\mathbb{R}^2} W * \mu_1 \, d\mu_1. \end{aligned}$$

Since minimizers have compact support and finite interaction energy by theorem 2.1, this convexity property is enough to guarantee uniqueness and the equivalence of (2.7)–(2.8) with minimality.

Proof of proposition 3.2. The heuristic idea to prove (3.7) is to apply Plancherel theorem, as we did formally in (1.3), and write:

$$\int_{\mathbb{R}^2} W * \nu \, d\nu = 2\pi \int_{\mathbb{R}^2} \widehat{W}(\xi) |\widehat{\nu}(\xi)|^2 \, d\xi.$$

Taking into account (3.5) and the fact that

$$\widehat{\nu}(0) = \frac{1}{2\pi} \int_{\mathbb{R}^2} d\nu = 0,$$

the right-hand side reduces to

$$2\pi \int_{\mathbb{R}^2} \widehat{W}(\xi) |\widehat{\nu}(\xi)|^2 \, d\xi = 2\pi \int_{\mathbb{R}^2} \frac{\widehat{\Psi}(\xi)}{|\xi|^2} |\widehat{\nu}(\xi)|^2 \, d\xi.$$

However, this argument is only formal, since we do not have enough regularity to apply Plancherel theorem. Our strategy is to prove (3.7) by approximation. This is a rather delicate argument, since both sides of (3.7) may *a priori* be not finite and neither W nor ν have a sign.

For $\varepsilon > 0$, let φ_ε be a radial mollifier supported on $B_\varepsilon(0)$. Let $\nu_\varepsilon := \nu * \varphi_\varepsilon \in C_c^\infty(\mathbb{R}^2) \subset \mathcal{S}$. Thus, $\widehat{\nu}_\varepsilon$ belongs to \mathcal{S} and, in particular, to $L^\infty(\mathbb{R}^2)$. We note that $W * \nu_\varepsilon \in C^\infty(\mathbb{R}^2)$ and, since $\widehat{\nu}_\varepsilon$ is smooth and $\widehat{\nu}_\varepsilon(0) = 0$, we have that

$$\widehat{W * \nu_\varepsilon} = 2\pi \widehat{W} \widehat{\nu}_\varepsilon \in L^1(\mathbb{R}^2).$$

Using these properties one can show that Plancherel theorem holds for $W * \nu_\varepsilon$ and ν_ε , that is,

$$\int_{\mathbb{R}^2} (W * \nu_\varepsilon)(x) \nu_\varepsilon(x) \, dx = 2\pi \int_{\mathbb{R}^2} \widehat{W}(\xi) |\widehat{\nu}_\varepsilon(\xi)|^2 \, d\xi = 2\pi \int_{\mathbb{R}^2} \frac{\widehat{\Psi}(\xi)}{|\xi|^2} |\widehat{\nu}_\varepsilon(\xi)|^2 \, d\xi \quad (3.8)$$

for every $\varepsilon > 0$.

We now want to pass to the limit in (3.8), as $\varepsilon \rightarrow 0^+$. For every $\xi \in \mathbb{R}^2$ we have

$$\widehat{\nu}_\varepsilon(\xi) = 2\pi \widehat{\nu}(\xi) \widehat{\varphi}(\varepsilon\xi) \rightarrow 2\pi \widehat{\nu}(\xi) \widehat{\varphi}(0) = \widehat{\nu}(\xi),$$

as $\varepsilon \rightarrow 0^+$. Therefore, $\widehat{\Psi}(\xi) |\widehat{\nu}_\varepsilon(\xi)|^2 / |\xi|^2$ converges to $\widehat{\Psi}(\xi) |\widehat{\nu}(\xi)|^2 / |\xi|^2$ for a.e. $\xi \in \mathbb{R}^2$, as $\varepsilon \rightarrow 0^+$. Moreover, $|\widehat{\varphi}(\varepsilon \cdot)| \leq C \|\varphi\|_{L^1}$. Either by dominated convergence or by Fatou's lemma we can thus pass to the limit in the last integral in (3.8).

To pass to the limit on the left-hand side of (3.8), we observe that

$$\int_{\mathbb{R}^2} (W * \nu_\varepsilon)(x) \nu_\varepsilon(x) \, dx = \int_{\mathbb{R}^2} (W * \varphi_\varepsilon * \varphi_\varepsilon) * \nu \, d\nu,$$

where we used that φ_ε is a radial, hence even, function. Set $\psi_\varepsilon := \varphi_\varepsilon * \varphi_\varepsilon$ and note that ψ_ε has the same properties as φ_ε (it is radial, compactly supported, non-negative, and integrates to one). Since W is continuous as a function with values

in $\mathbb{R} \cup \{+\infty\}$, we have that $(W * \psi_\varepsilon)(x) \rightarrow W(x)$, as $\varepsilon \rightarrow 0^+$, for every $x \in \mathbb{R}^2$. Moreover, by (2.2):

$$(W * \psi_\varepsilon)(x) \leq (W_0 * \psi_\varepsilon)(x) + C_2 \quad \text{for every } x \in \mathbb{R}^2. \tag{3.9}$$

Since W_0 is superharmonic and ψ_ε is radial, we have

$$(W_0 * \psi_\varepsilon)(x) \leq W_0(x) \leq W(x) - C_1 \quad \text{for every } x \in \mathbb{R}^2, \tag{3.10}$$

where the last inequality follows again from (2.2).

Let $M > 0$ be such that $\text{supp } \nu \subset B_M(0)$ and let c_M be the minimum of W on $B_{4M}(0)$. Combining (3.9) and (3.10), we deduce that

$$0 \leq (W * \psi_\varepsilon)(x) - c_M \leq W(x) + C_2 - C_1 - c_M \quad \text{for every } x \in B_{2M}(0) \tag{3.11}$$

and for every $\varepsilon > 0$ small enough.

We now write:

$$\begin{aligned} \int_{\mathbb{R}^2} (W * \psi_\varepsilon) * \nu \, d\nu &= \iint_{\mathbb{R}^2 \times \mathbb{R}^2} (W * \psi_\varepsilon)(x - y) \, d\mu_0(x) \, d\mu_0(y) \\ &\quad + \iint_{\mathbb{R}^2 \times \mathbb{R}^2} (W * \psi_\varepsilon)(x - y) \, d\mu_1(x) \, d\mu_1(y) \\ &\quad - 2 \iint_{\mathbb{R}^2 \times \mathbb{R}^2} (W * \psi_\varepsilon)(x - y) \, d\mu_0(x) \, d\mu_1(y). \end{aligned}$$

Since μ_0 and μ_1 have finite interaction energy, we can pass to the limit in the first two integrals on the right-hand side by (3.11) and dominated convergence. As for the last integral, it goes to the limit either by dominated convergence or by Fatou's lemma. This completes the proof of (3.7).

To conclude, assume that

$$0 = \int_{\mathbb{R}^2} W * \nu \, d\nu = 2\pi \int_{\mathbb{R}^2} \frac{\widehat{\Psi}(\xi)}{|\xi|^2} |\widehat{\nu}(\xi)|^2 \, d\xi. \tag{3.12}$$

Since $\widehat{\Psi}$ cannot be identically zero (otherwise, W would be constant) and is continuous outside 0, there exist $\delta > 0$, $\eta > 0$, and $\xi_0 \neq 0$ such that $\widehat{\Psi}(\xi) > \eta$ for all $\xi \in B_\delta(\xi_0)$. Thus, (3.12) implies that $\widehat{\nu} = 0$ on $B_\delta(\xi_0)$. On the other hand, ν is a distribution with compact support, so by Paley–Wiener theorem $\widehat{\nu}$ is the restriction to \mathbb{R}^2 of an entire function. Therefore, $\widehat{\nu}$ has to be identically zero on \mathbb{R}^2 , that is, $\nu = 0$. □

4. Characterization of minimizers

Throughout this section, we consider the confinement $V(x) = |x|^2$ and focus on the characterization of the minimizer of I for this specific choice of the confinement.

4.1. The Coulomb case

In the purely Coulomb case with confinement $V(x) = |x|^2$ the energy:

$$I_0(\mu) = - \iint_{\mathbb{R}^2 \times \mathbb{R}^2} \log|x - y| \, d\mu(y) \, d\mu(x) + \int_{\mathbb{R}^2} |x|^2 \, d\mu(x), \quad \mu \in \mathcal{P}(\mathbb{R}^2), \quad (4.1)$$

is invariant by rotations. Therefore, by uniqueness the minimizer μ_0 has to be invariant by rotations, too. If we formally take the Laplacian of both sides of (2.7) (with W_0 in place of W), we obtain:

$$0 = \Delta W_0 * \mu_0 + 2 = -2\pi\mu_0 + 2 \quad \text{on } \text{supp } \mu_0,$$

where the last equality follows from the relation $\Delta W_0 = -2\pi\delta_0$. Hence, $\mu_0 = 1/\pi$ on its support. Since μ_0 is a probability measure, one can conjecture that μ_0 has to be the normalized characteristic function of $B_1(0)$. This is indeed the case, as shown in the next theorem.

THEOREM 4.1 [8, 10, 17]. *The unique minimizer of the energy I_0 in (4.1) is the so-called circle law, that is, the measure:*

$$\mu_0 = \frac{1}{\pi} \chi_{B_1(0)}.$$

The proof of this result is based on the Gauss averaging principle: for any $r > 0$:

$$-\frac{1}{2\pi} \int_{-\pi}^{\pi} \log|z - re^{i\theta}| \, d\theta = \begin{cases} -\log r & \text{if } |z| < r, \\ -\log|z| & \text{if } |z| \geq r, \end{cases} \quad (4.2)$$

where z is a complex variable. A notable consequence of this principle is the well-known fact that the Coulomb potential due to a homogeneous spherical body is the same, outside the body, as if all the mass were at its centre.

Proving (4.2) is straightforward: if $|z| > r$, the map $w \mapsto -\log|z - w|$ is harmonic for $|w| \leq r$, so (4.2) follows by the mean value property. If $|z| < r$, one can write:

$$-\log|z - re^{i\theta}| = -\log|ze^{-i\theta} - r| = -\log|\bar{z}e^{i\theta} - r|.$$

The map $w \mapsto -\log|w - r|$ is harmonic for $|w| \leq |z|$, so one can conclude again by applying the mean value property. Finally, for $|z| = r$ one can argue by approximation with radii $\rho \rightarrow r^-$ and dominated convergence.

Proof of theorem 4.1. By applying the Gauss averaging principle we can explicitly compute the Coulomb potential of μ_0 : indeed, using polar coordinates we have:

$$(W_0 * \mu_0)(x) = -\frac{1}{\pi} \int_{B_1(0)} \log|x - y| \, dy = -\frac{1}{\pi} \int_0^1 \int_{-\pi}^{\pi} \log|x - re^{i\theta}| \, d\theta \, r \, dr.$$

By (4.2) we deduce that

$$(W_0 * \mu_0)(x) = \begin{cases} \frac{1}{2} - \frac{1}{2}|x|^2 & \text{for } |x| \leq 1, \\ -\log|x| & \text{for } |x| > 1. \end{cases}$$

Using this formula, one can immediately check that $(W_0 * \mu_0)(x) + \frac{1}{2}|x|^2 = \frac{1}{2}$ for $x \in B_1(0)$ and $(W_0 * \mu_0)(x) + \frac{1}{2}|x|^2 \geq \frac{1}{2}$ for $x \notin B_1(0)$, that is, μ_0 satisfies the Euler–Lagrange conditions. By the results of the previous section this is enough to conclude that μ_0 is the unique minimizer of I_0 . \square

4.2. The anisotropic case

In this section, we discuss the characterization of the minimizer for the energy:

$$I(\mu) = \iint_{\mathbb{R}^2 \times \mathbb{R}^2} (-\log|x - y| + \kappa(x - y)) \, d\mu(y) \, d\mu(x) + \int_{\mathbb{R}^2} |x|^2 \, d\mu(x), \quad (4.3)$$

where κ is even, 0-homogeneous, and of regularity $H^s(\mathbb{S}^1)$ with $s > 3/2$. We start by considering the case where the Fourier transform of the interaction kernel (computed in (3.5)) is strictly positive outside 0.

THEOREM 4.2. *Assume $\widehat{\Psi} > 0$ on \mathbb{S}^1 , where $\widehat{\Psi}$ is the function introduced in (3.5). Then the unique minimizer of the energy I in (4.3) is given by the normalized characteristic function of the domain enclosed by an ellipse centred at the origin, whose semi-axes a_1, a_2 satisfy the relation $a_1^2 + a_2^2 = 2$.*

Theorem 4.2 was originally proved by Carrillo and Shu in [3]. Here, we present an alternative proof inspired by [16].

Proof of theorem 4.2. We write a general domain enclosed by an ellipse as $E = RE_0$, where $R \in SO(2)$ and

$$E_0 = \left\{ x \in \mathbb{R}^2 : \frac{x_1^2}{a_1^2} + \frac{x_2^2}{a_2^2} \leq 1 \right\}$$

and we set $\chi := \chi_E/|E|$. The theorem is proved if we show that there exist $a_1, a_2 > 0$ and $R \in SO(2)$ such that

$$(W * \chi)(x) + \frac{1}{2}|x|^2 = c \quad \text{for every } x \in E, \quad (4.4)$$

and

$$(W * \chi)(x) + \frac{1}{2}|x|^2 \geq c \quad \text{for every } x \in \mathbb{R}^2 \setminus E \quad (4.5)$$

for some constant $c \in \mathbb{R}$.

Note that $W * \chi$ is a C^1 function in \mathbb{R}^2 . To compute its expression we would like to make use of the inversion formula, that is, write $W * \chi$ as an integral of its Fourier transform. However, \widehat{W} is not a function, it is only a tempered distribution. To circumvent this difficulty we apply this strategy to the gradient of $W * \chi$. In fact, the Fourier transform of $\nabla(W * \chi)$ is given by

$$i\xi \widehat{W * \chi}(\xi) = 2\pi i \xi \widehat{W}(\xi) \widehat{\chi}(\xi),$$

so the presence of the factor ξ annihilates the singular part of \widehat{W} . Note that $\nabla(W * \chi)$ is a continuous function that behaves as $1/|x|$ at infinity, so it is a tempered distribution.

Step 1: The inversion formula for $\nabla(W * \chi)$. We recall that

$$\widehat{\chi_{B_1(0)}}(\xi) = \frac{J_1(|\xi|)}{|\xi|},$$

where J_1 denotes the Bessel function of the first kind of order 1. There is no explicit formula for J_1 ; however, it is well known that J_1 has the following behaviour:

$$J_1(|\xi|) \simeq \frac{1}{2}|\xi| \quad \text{as } \xi \rightarrow 0, \tag{4.6}$$

$$|J_1(|\xi|)| \leq \frac{C}{|\xi|^{1/2}} \quad \text{as } |\xi| \rightarrow \infty, \tag{4.7}$$

see, e.g. [13, Section 5.16].

For $a = (a_1, a_2) \in \mathbb{R}^2$ we denote the diagonal matrix $\text{diag}(a_1, a_2)$ by $D(a)$. By writing $x \in E$ as $x = RD(a)y$ with $y \in B_1(0)$, we obtain:

$$\widehat{\chi}(\xi) = \frac{1}{\pi} \widehat{\chi_{B_1(0)}}(D(a)R^T\xi) = \frac{1}{\pi} \frac{J_1(|D(a)R^T\xi|)}{|D(a)R^T\xi|}.$$

By the bounds (4.6)–(4.7) we deduce that the function $2\pi i\xi \widehat{W} \widehat{\chi}$ belongs to $L^1(\mathbb{R}^2)$. Therefore, since $\nabla(W * \chi)$ is a continuous function, the inversion formula holds, that is:

$$\nabla(W * \chi)(x) = \int_{\mathbb{R}^2} i\xi \frac{\widehat{\Psi}(\xi)}{|\xi|^2} \widehat{\chi}(\xi) e^{ix \cdot \xi} d\xi = - \int_{\mathbb{R}^2} \xi \frac{\widehat{\Psi}(\xi)}{|\xi|^2} \widehat{\chi}(\xi) \sin(x \cdot \xi) d\xi$$

for every $x \in \mathbb{R}^2$, where we used that $\widehat{\Psi}$ and $\widehat{\chi}$ are even functions. Passing to polar coordinates we obtain:

$$\begin{aligned} \nabla(W * \chi)(x) &= - \int_{\mathbb{S}^1} \int_0^\infty y \widehat{\Psi}(y) \widehat{\chi}(\rho y) \sin(\rho x \cdot y) d\rho d\mathcal{H}^1(y) \\ &= - \frac{1}{\pi} \int_{\mathbb{S}^1} \int_0^\infty y \widehat{\Psi}(y) \frac{J_1(\rho |D(a)R^T y|)}{\rho |D(a)R^T y|} \sin(\rho x \cdot y) d\rho d\mathcal{H}^1(y). \end{aligned}$$

Setting $r := \rho |D(a)R^T y|$, we deduce that

$$\nabla(W * \chi)(x) = - \frac{1}{\pi} \int_{\mathbb{S}^1} \frac{y \widehat{\Psi}(y)}{|D(a)R^T y|} \int_0^\infty \frac{J_1(r)}{r} \sin(r\alpha(x, y)) dr d\mathcal{H}^1(y) \tag{4.8}$$

for every $x \in \mathbb{R}^2$, where we set:

$$\alpha(x, y) := \frac{x \cdot y}{|D(a)R^T y|}. \tag{4.9}$$

Using formula (5), p. 99, in [5], one can compute the improper integral:

$$\int_0^\infty \frac{J_1(r)}{r} \sin(r\alpha) dr = \begin{cases} \alpha & \text{if } 0 \leq \alpha \leq 1, \\ \frac{1}{\alpha + \sqrt{\alpha^2 - 1}} & \text{if } \alpha > 1. \end{cases} \tag{4.10}$$

Moreover, we have that, if $x \in E$, then

$$|x \cdot y| = |D(a)^{-1}R^T x \cdot D(a)R^T y| \leq |D(a)R^T y|,$$

that is, $|\alpha(x, y)| \leq 1$ for every $x \in E$. We conclude that

$$\begin{aligned} \nabla(W * \chi)(x) &= -\frac{1}{\pi} \int_{\mathbb{S}^1} \frac{\widehat{\Psi}(y)}{|D(a)R^T y|} \alpha(x, y) y \, d\mathcal{H}^1(y) \\ &= -\frac{1}{\pi} \int_{\mathbb{S}^1} \frac{\widehat{\Psi}(y)}{|D(a)R^T y|^2} (x \cdot y) y \, d\mathcal{H}^1(y) \end{aligned} \tag{4.11}$$

for every $x \in E$. Formula (4.11) shows that $\nabla(W * \chi)$ is a homogeneous polynomial of degree 1 inside E . In other words, up to an additive constant, $W * \chi$ is a homogeneous polynomial of degree 2 inside E .

Step 2: Solving the first Euler–Lagrange condition. Solving (4.4) corresponds to finding $a_1, a_2 > 0$ and $R \in SO(2)$ such that

$$\nabla(W * \chi)(x) + x = 0 \quad \text{for every } x \in E.$$

By formula (4.11) this translates into the following system of three equations:

$$\frac{1}{\pi} \int_{\mathbb{S}^1} \frac{\widehat{\Psi}(y)}{|D(a)R^T y|^2} y_j y_k \, d\mathcal{H}^1(y) = \delta_{jk} \quad \text{for every } j, k = 1, 2, \tag{4.12}$$

where δ_{jk} is the Kronecker delta.

Let us denote by $\mathbb{M}_{\text{sym},+}^{2 \times 2}$ the set of positive definite 2×2 symmetric matrices. Note that $M := RD(a)^2 R^T \in \mathbb{M}_{\text{sym},+}^{2 \times 2}$ and $|D(a)R^T y|^2 = My \cdot y$. Solving system (4.12) is therefore equivalent to finding $M \in \mathbb{M}_{\text{sym},+}^{2 \times 2}$ such that

$$\frac{1}{\pi} \int_{\mathbb{S}^1} \frac{\widehat{\Psi}(y)}{My \cdot y} y_j y_k \, d\mathcal{H}^1(y) = \delta_{jk} \quad \text{for every } j, k = 1, 2. \tag{4.13}$$

Let us denote the entries of M by M_{jk} for $j, k = 1, 2$. Note that, if we multiply by M_{jk} the jk equation in (4.13) and we sum over j, k , we obtain:

$$\text{tr}(M) = M_{11} + M_{22} = \frac{1}{\pi} \int_{\mathbb{S}^1} \widehat{\Psi}(y) \, d\mathcal{H}^1(y) = 2,$$

where we used (3.6). Since $\text{tr}(M) = a_1^2 + a_2^2$, where a_1, a_2 are the semi-axes of the candidate ellipse, we deduce that, if a minimizing ellipse exists, then necessarily its semi-axes satisfy the relation:

$$a_1^2 + a_2^2 = 2.$$

To solve (4.13) it is convenient to introduce the function:

$$f(M) := -\frac{1}{\pi} \int_{\mathbb{S}^1} \widehat{\Psi}(y) \log(My \cdot y) \, d\mathcal{H}^1(y) + \text{tr}(M)$$

defined for every $M \in \mathbb{M}_{\text{sym},+}^{2 \times 2}$. It is immediate to see that conditions (4.13) are satisfied if we find $M_0 \in \mathbb{M}_{\text{sym},+}^{2 \times 2}$ such that $\nabla_M f(M_0) = 0$, where with a slight abuse

of notation we denoted by ∇_M the operator:

$$\nabla_M = \left(\frac{\partial}{\partial M_{11}}, \frac{\partial}{\partial M_{22}}, \frac{\partial}{\partial M_{12}} \right).$$

We claim that f has a minimizer in $\mathbb{M}_{\text{sym},+}^{2 \times 2}$. Since $\mathbb{M}_{\text{sym},+}^{2 \times 2}$ is an open set, the claim will imply the existence of a critical point of f in this set and thus, of a solution to (4.13).

Given $M \in \mathbb{M}_{\text{sym},+}^{2 \times 2}$, we first look at the behaviour of the function f along the line tM for $t > 0$, that is, we consider the function:

$$g(t) := f(tM) = -\frac{1}{\pi} \int_{\mathbb{S}^1} \widehat{\Psi}(y) \log(My \cdot y) \, d\mathcal{H}^1(y) - 2 \log t + t \operatorname{tr}(M)$$

for $t > 0$, where the last equality follows from (3.6). It is immediate to see that g is minimized at $t_* = 2/\operatorname{tr}(M)$. Therefore, minimizing f over $\mathbb{M}_{\text{sym},+}^{2 \times 2}$ is equivalent to minimizing f on the set:

$$\mathcal{M} := \{M \in \mathbb{M}_{\text{sym},+}^{2 \times 2} : \operatorname{tr}(M) = 2\}.$$

By diagonalization, any $M \in \mathcal{M}$ can be written as $M = QD(b)Q^T$ with $Q \in SO(2)$ and $b = (\beta, 2 - \beta)$, $\beta \in (0, 2)$. Using this representation and a change of variables we have that

$$f(M) = f(QD(b)Q^T) = -\frac{1}{\pi} \int_{\mathbb{S}^1} \widehat{\Psi}(Qy) \log(\beta y_1^2 + (2 - \beta)y_2^2) \, d\mathcal{H}^1(y) + 2.$$

Therefore, setting

$$\gamma(\beta, Q) := -\frac{1}{\pi} \int_{\mathbb{S}^1} \widehat{\Psi}(Qy) \log(\beta y_1^2 + (2 - \beta)y_2^2) \, d\mathcal{H}^1(y)$$

for every $\beta \in [0, 2]$ and $Q \in SO(2)$, it is enough to show that γ has a minimizer in $(0, 2) \times SO(2)$ to conclude that f has a minimizer in \mathcal{M} and thus in $\mathbb{M}_{\text{sym},+}^{2 \times 2}$.

The function γ is finite and continuous on the compact set $[0, 2] \times SO(2)$. Therefore, it has a minimizer (β_0, Q_0) in this set. It remains to prove that β_0 is neither 0 nor 2.

By assumption there exists a constant $C_0 > 0$ such that

$$\widehat{\Psi}(\xi) \geq C_0 \quad \text{for every } \xi \in \mathbb{S}^1.$$

Using this inequality and (3.6), we obtain that

$$\begin{aligned} \frac{\partial}{\partial \beta} \gamma(\beta, Q_0) &= \frac{1}{\pi} \int_{\mathbb{S}^1} \widehat{\Psi}(Q_0 y) \frac{2y_2^2 - 1}{\beta y_1^2 + (2 - \beta)y_2^2} \, d\mathcal{H}^1(y) \\ &\leq \frac{4}{2 - \beta} - \frac{1}{\pi} \int_{\mathbb{S}^1} \frac{C_0}{\beta y_1^2 + (2 - \beta)y_2^2} \, d\mathcal{H}^1(y) \end{aligned}$$

for every $\beta \in (0, 2)$. Since the right-hand side goes to $-\infty$ as $\beta \rightarrow 0^+$, we deduce that there exists some $\delta > 0$ such that

$$\frac{\partial}{\partial \beta} \gamma(\beta, Q_0) < 0$$

for every $\beta \in (0, \delta)$. Hence, β_0 cannot be equal to zero. One can show in a similar way that β_0 cannot be 2 either. This concludes the proof of the step.

Step 3: The first Euler–Lagrange condition implies the second one. To complete the proof, we show that, if $\chi = \chi_E/|E|$ satisfies (4.4), then it satisfies (4.5), as well. Assume χ satisfies (4.4). By (4.4) it is enough to prove that the potential $(W * \chi)(x) + \frac{1}{2}|x|^2$ grows in the radial direction, that is:

$$\nabla(W * \chi)(x) \cdot x + |x|^2 \geq 0 \quad \text{for every } x \in \mathbb{R}^2 \setminus E.$$

Let $x \in \mathbb{R}^2 \setminus E$. We can write $x = tx^0$ with $x^0 \in E$. By (4.4) we have that

$$\nabla(W * \chi)(x^0) \cdot x^0 + |x^0|^2 = 0,$$

which can be written by (4.11) as

$$-\frac{1}{\pi} \int_{\mathbb{S}^1} \widehat{\Psi}(y) \alpha^2(x^0, y) \, d\mathcal{H}^1(y) + |x^0|^2 = 0,$$

where α is defined in (4.9). Multiplying this equation by t^2 yields:

$$-\frac{1}{\pi} \int_{\mathbb{S}^1} \widehat{\Psi}(y) \alpha^2(x, y) \, d\mathcal{H}^1(y) + |x|^2 = 0. \tag{4.14}$$

By the inversion formula (4.8) and (4.10) we can write:

$$\begin{aligned} \nabla(W * \chi)(x) \cdot x + |x|^2 &= |x|^2 - \frac{1}{\pi} \int_{\mathbb{S}^1} \widehat{\Psi}(y) \alpha^2(x, y) \chi_{[-1,1]}(\alpha(x, y)) \, d\mathcal{H}^1(y) \\ &\quad - \frac{1}{\pi} \int_{\mathbb{S}^1} \widehat{\Psi}(y) \frac{|\alpha(x, y)|}{|\alpha(x, y)| + \sqrt{\alpha^2(x, y) - 1}} \chi_{\mathbb{R} \setminus [-1,1]}(\alpha(x, y)) \, d\mathcal{H}^1(y). \end{aligned}$$

Owing to (4.14), the expression above reduces to

$$\begin{aligned} &\nabla(W * \chi)(x) \cdot x + |x|^2 \\ &= \frac{1}{\pi} \int_{\mathbb{S}^1} \widehat{\Psi}(y) |\alpha(x, y)| \sqrt{\alpha^2(x, y) - 1} \chi_{\mathbb{R} \setminus [-1,1]}(\alpha(x, y)) \, d\mathcal{H}^1(y) \geq 0. \end{aligned}$$

This concludes the proof. □

The next result shows how to combine the previous theorem and an approximation argument to characterize the minimizer in the ‘degenerate’ case where the Fourier transform of the interaction kernel is only non-negative outside 0. In particular, we prove that loss of dimensionality may occur and in this case the support of the minimizer is contained in a straight line orthogonal to a direction $\xi \in \mathbb{S}^1$ such that $\widehat{\Psi}(\xi) = 0$.

COROLLARY 4.3. *Assume $\widehat{\Psi} \geq 0$ on \mathbb{S}^1 , where $\widehat{\Psi}$ is the function introduced in (3.5). Then the unique minimizer of the energy I in (4.3) is either the normalized characteristic function of the domain enclosed by an ellipse centred at the origin and with semiaxes a_1, a_2 satisfying $a_1^2 + a_2^2 = 2$, or is the push-forward of the semicircle*

law (1.7) through a rotation map $g(x) = Rx$ for $x \in \mathbb{R}^2$ with $R \in SO(2)$ satisfying $\widehat{\Psi}(Re_1) = 0$.

Proof. For $\varepsilon > 0$ we consider the kernels:

$$W_\varepsilon(x) = -(1 + \varepsilon) \log |x| + \kappa(x)$$

and we denote by I_ε the corresponding energies with confinement $|x|^2$. Arguing as in (3.4), it is immediate to see that

$$\widehat{W}_\varepsilon(\xi) = c_\varepsilon \delta_0 + \frac{\widehat{\Psi}(\xi) + \varepsilon}{|\xi|^2}$$

with $\widehat{\Psi} + \varepsilon > 0$ on \mathbb{S}^1 . By theorem 4.2 for every $\varepsilon > 0$ the unique minimizer μ_ε of I_ε is the normalized characteristic function of a set E_ε of the form:

$$R_\varepsilon^T E_\varepsilon = \left\{ x \in \mathbb{R}^2 : \frac{x_1^2}{(a_{1,\varepsilon})^2} + \frac{x_2^2}{(a_{2,\varepsilon})^2} \leq 1 \right\}$$

with $R_\varepsilon \in SO(2)$ and $(a_{1,\varepsilon})^2 + (a_{2,\varepsilon})^2 = 2$. The sequence $(\mu_\varepsilon)_\varepsilon$ is tight, since the support of μ_ε is contained in the closed ball of centre 0 and radius $\sqrt{2}$ for every $\varepsilon > 0$. Therefore, up to subsequences, $(\mu_\varepsilon)_\varepsilon$ converges narrowly to a measure $\mu_0 \in \mathcal{P}(\mathbb{R}^2)$. We claim that μ_0 is the minimizer of I . Indeed, since the supports of μ_ε and μ_0 are uniformly bounded, on these sets W_0 is bounded from below by some constant $-c_0$, with $c_0 > 0$. Therefore:

$$I_\varepsilon(\mu_\varepsilon) \geq I(\mu_\varepsilon) - c_0\varepsilon$$

and by lower semicontinuity:

$$\liminf_{\varepsilon \rightarrow 0^+} I_\varepsilon(\mu_\varepsilon) \geq \liminf_{\varepsilon \rightarrow 0^+} I(\mu_\varepsilon) \geq I(\mu_0).$$

On the other hand, if μ is any measure in $\mathcal{P}(\mathbb{R}^2)$ with compact support and such that

$$\iint_{\mathbb{R}^2 \times \mathbb{R}^2} (-\log |x - y| + \kappa(x - y)) \, d\mu(y) \, d\mu(x) < +\infty, \tag{4.15}$$

then by minimality:

$$\limsup_{\varepsilon \rightarrow 0^+} I_\varepsilon(\mu_\varepsilon) \leq \lim_{\varepsilon \rightarrow 0^+} I_\varepsilon(\mu) = I(\mu).$$

Therefore, μ_0 minimizes I over all measures with compact support and satisfying (4.15). By theorem 2.1 and proposition 3.2 we conclude that μ_0 is the minimizer of I on the whole $\mathcal{P}(\mathbb{R}^2)$.

Up to subsequences, we can assume that the rotations and the semi-axes converge, that is, $R_\varepsilon \rightarrow R$, $a_{1,\varepsilon} \rightarrow a_1$, and $a_{2,\varepsilon} \rightarrow a_2$, as $\varepsilon \rightarrow 0^+$, with $R \in SO(2)$, $a_1, a_2 \geq 0$, and $a_1^2 + a_2^2 = 2$. Therefore, we have two cases: either both a_1 and a_2 are strictly

positive, or one of them is 0 and the other one is $\sqrt{2}$. In the first case μ_0 is the normalized characteristic function of the set E given by

$$R^T E = \left\{ x \in \mathbb{R}^2 : \frac{x_1^2}{a_1^2} + \frac{x_2^2}{a_2^2} \leq 1 \right\}.$$

In the second case, μ_0 is the push-forward of measure (1.7) through the rotation map $g(x) = Rx$ for $x \in \mathbb{R}^2$ if $a_1 = 0$, or $g(x) = RJx$ for $x \in \mathbb{R}^2$ if $a_2 = 0$. Here, J denotes a $\pi/2$ -rotation.

To conclude the proof assume that $a_1 = 0$ and $a_2 = \sqrt{2}$. We want to prove that $\widehat{\Psi}(Re_1) = 0$. By (4.12) and the change of variable $z = R_\varepsilon^T y$ we have that

$$\frac{1}{\pi} \int_{\mathbb{S}^1} \frac{\widehat{\Psi}(R_\varepsilon z) + \varepsilon}{(a_{1,\varepsilon})^2 z_1^2 + (a_{2,\varepsilon})^2 z_2^2} (R_\varepsilon z \cdot e_j)^2 d\mathcal{H}^1(z) = 1 \quad \text{for } j = 1, 2$$

for every $\varepsilon > 0$. Summing over j and applying Fatou lemma yield:

$$\frac{1}{2\pi} \int_{\mathbb{S}^1} \frac{\widehat{\Psi}(Rz)}{z^2} d\mathcal{H}^1(z) \leq \liminf_{\varepsilon \rightarrow 0} \frac{1}{\pi} \int_{\mathbb{S}^1} \frac{\widehat{\Psi}(R_\varepsilon z) + \varepsilon}{(a_{1,\varepsilon})^2 z_1^2 + (a_{2,\varepsilon})^2 z_2^2} d\mathcal{H}^1(z) = 2.$$

This implies that the function $z \mapsto z^{-2} \widehat{\Psi}(Rz)$ is integrable on \mathbb{S}^1 , hence $\widehat{\Psi}(Re_1) = 0$.

The case $a_1 = \sqrt{2}$ and $a_2 = 0$ can be treated analogously. □

REMARK 4.4. If the quadratic confinement $V(x) = |x|^2$ is replaced by a positive definite quadratic form $Q(x)$, theorem 4.2 and corollary 4.3 remain true, up to a scaling of the semiaxes of the ellipse and of the semicircle law. Indeed, by rotating the whole system if needed, one can always assume that $Q(x) = D(\lambda)x \cdot x$, where $\lambda = (\lambda_1, \lambda_2)$ and $\lambda_1, \lambda_2 > 0$. Arguing as in the proof of theorem 4.2, one can show that solving the first Euler–Lagrange condition for the normalized characteristic function of the domain enclosed by an ellipse is equivalent to finding $M \in \mathbb{M}_{\text{sym},+}^{2 \times 2}$ such that

$$\frac{1}{\pi} \int_{\mathbb{S}^1} \frac{\widehat{\Psi}(y)}{My \cdot y} y_j y_k d\mathcal{H}^1(y) = \lambda_j \delta_{jk} \quad \text{for every } j, k = 1, 2. \tag{4.16}$$

The existence of a solution to this system can be proved by showing that the function

$$\tilde{f}(M) := -\frac{1}{\pi} \int_{\mathbb{S}^1} \widehat{\Psi}(y) \log(D(\lambda)^{-1} My \cdot y) d\mathcal{H}^1(y) + \text{tr}(M) \quad \text{for } M \in \mathbb{M}_{\text{sym},+}^{2 \times 2}$$

has a minimizer M_0 in $\mathbb{M}_{\text{sym},+}^{2 \times 2}$. Indeed, it is immediate to see that $D(\lambda)^{-1} M_0$ satisfies (4.16). The existence of a minimizer of \tilde{f} in $\mathbb{M}_{\text{sym},+}^{2 \times 2}$ follows by the same argument in step 2 of the proof of theorem 4.2. The proofs of step 3 and of corollary 4.3 can then be easily adapted.

REMARK 4.5. The results of theorem 4.2 and corollary 4.3 extend to \mathbb{R}^3 , see [16], where the kernel W is given by the following anisotropic variant of the 3D Coulomb kernel:

$$W(x) = \frac{\Psi(x)}{|x|} \quad \text{for } x \in \mathbb{R}^3, x \neq 0, \quad W(0) = +\infty,$$

with Ψ strictly positive, even, 0-homogeneous, and smooth enough on \mathbb{S}^2 . More precisely, we proved that, if $\widehat{W} > 0$, then the unique minimizer of the energy

$$I_{3D}(\mu) = \iint_{\mathbb{R}^3 \times \mathbb{R}^3} W(x - y) \, d\mu(y) \, d\mu(x) + \int_{\mathbb{R}^3} |x|^2 \, d\mu(x), \quad \mu \in \mathcal{P}(\mathbb{R}^3),$$

is the normalized characteristic function of the domain enclosed by an ellipsoid centred at the origin. If $\widehat{W} \geq 0$, the minimizer is either of this form or it may collapse on a 2D measure, whose support is given by the domain enclosed by an ellipse.

REMARK 4.6. The original proof by Carrillo and Shu in [3] is based on a different technique. Their starting point is a formula expressing the interaction kernel as an integral of 1D logarithmic kernels on projections. More precisely, they show that

$$-\log |x| + \kappa(x) = -\frac{1}{2\pi} \int_{-\pi}^{\pi} \widehat{\Psi}(e^{i\theta}) \log |x \cdot e^{i\theta}| \, d\theta + \text{constant},$$

where $\widehat{\Psi}$ is the same function as in (3.5). They observe that the projection of $\chi_E/|E|$ (with E the domain enclosed by an ellipse centred at the origin) on any line passing through the origin is a semicircle law. Using this remark and the representation formula above, they argue by projection and conclude by applying the minimality of the semicircle law for the 1D logarithmic kernel. The same approach can be used to treat 2D anisotropic Riesz kernels of the form:

$$\frac{\Psi(x)}{|x|^s}, \quad 0 < s < 2,$$

with Ψ strictly positive, even, 0-homogeneous, and smooth enough on \mathbb{S}^1 . However, their strategy of proof extends to \mathbb{R}^3 only under some rather strong symmetry assumptions on the anisotropy, that essentially reduce the problem to a 2D setting (see [4]).

5. Related results and open questions

In this last section, we show how to explicitly determine the optimal distributions in some concrete cases and we discuss some open problems.

5.1. The dislocation case

As a first result, we show how to deduce the characterization of the minimizer of I_α in (1.8) from the results of the previous sections. The original proofs in [2, 18] are based on completely different arguments.

THEOREM 5.1. *Let I_α be the functional defined in (1.8). Then for $|\alpha| < 1$ the unique minimizer of I_α is given by the measure (1.9). For $\alpha \geq 1$ the unique minimizer of I_α is the semicircle law (1.7), whereas for $\alpha \leq -1$ it is the semicircle law on the horizontal axis:*

$$\frac{1}{\pi} \sqrt{2 - x_1^2} \mathcal{H}^1 \llcorner [-\sqrt{2}, \sqrt{2}](x_1) \otimes \delta_0(x_2). \tag{5.1}$$

Proof. We set $W_\alpha(x) = -\log|x| + \kappa_\alpha(x)$, where

$$\kappa_\alpha(x) := \alpha \frac{x_1^2}{|x|^2}.$$

Since we can write:

$$\kappa_\alpha(x) = \frac{\alpha}{2} \frac{x_1^2 - x_2^2}{|x|^2} + \frac{\alpha}{2},$$

lemma 3.1 ensures that

$$\widehat{W}_\alpha(\xi) = c_\alpha \delta_0 + \frac{\widehat{\Psi}_\alpha(\xi)}{|\xi|^2} \tag{5.2}$$

with

$$\widehat{\Psi}_\alpha(\xi) = (1 - \alpha) \frac{\xi_1^2}{|\xi|^2} + (1 + \alpha) \frac{\xi_2^2}{|\xi|^2}, \quad \xi \neq 0.$$

Assume $|\alpha| < 1$. Since $\widehat{\Psi}_\alpha > 0$ on \mathbb{S}^1 in this case, by theorem 4.2 the unique minimizer is the normalized characteristic function of the domain enclosed by an ellipse of semi-axes $a_{1,\alpha}$, $a_{2,\alpha}$. We note that W_α is symmetric with respect to the coordinate axes, that is:

$$W_\alpha(-x_1, x_2) = W_\alpha(x_1, -x_2) = W_\alpha(x) \quad \text{for every } x \in \mathbb{R}^2.$$

By uniqueness the minimizer must have the same symmetry, that is, the ellipse is symmetric with respect to the coordinate axes. Therefore, in system (4.12) the rotation R is necessarily the identity matrix and the equation for $j = 1$, $k = 2$ is trivially satisfied by symmetry. In other words, using the expression of $\widehat{\Psi}_\alpha$, the semi-axes $a_{1,\alpha}$, $a_{2,\alpha}$ satisfy:

$$\frac{1}{\pi} \int_{\mathbb{S}^1} \frac{(1 - \alpha)y_1^2 + (1 + \alpha)y_2^2}{(a_{1,\alpha})^2 y_1^2 + (a_{2,\alpha})^2 y_2^2} y_j^2 d\mathcal{H}^1(y) = 1 \quad \text{for every } j = 1, 2.$$

It is immediate to see that $a_{1,\alpha} = \sqrt{1 - \alpha}$, $a_{2,\alpha} = \sqrt{1 + \alpha}$ is a solution of this system. It is indeed the unique solution, since we proved in step 3 of the proof of theorem 4.2 that any solution of (4.12) is automatically a minimizer and the minimizer is unique. We conclude that for $|\alpha| < 1$ the minimizer is given by the measure (1.9).

Let now $\alpha = 1$. Arguing as in the proof of corollary 4.3, one can show that the minimizer of I_1 has to be the limit of the minimizer of I_α , as $\alpha \rightarrow 1^-$. Therefore, the minimizer is the semicircle law (1.7). The same argument applies to $\alpha = -1$.

If $\alpha > 1$, let us denote by μ_{sc} the semicircle law in (1.7) and let μ be any other measure in $\mathcal{P}(\mathbb{R}^2)$. Then

$$I_\alpha(\mu) \geq I_1(\mu) > I_1(\mu_{sc}) = I_\alpha(\mu_{sc}),$$

where the last equality follows from the fact that the anisotropic interaction is 0 on the semicircle law. A similar argument shows that (5.1) is the unique minimizer of I_α for $\alpha < -1$. □

REMARK 5.2. For anisotropy (1.4) and quadratic confinement one can prove that the unique minimizer is the normalized characteristic function of the domain enclosed by an ellipse (with explicit semi-axes) if $b^2 < 1 + a^2$, whereas is the semicircle law (1.7) if $b^2 \geq 1 + a^2$, see [9]. Since the Fourier transform of the kernel W in this case is ‘degenerate’ outside 0 for any $b > a > 0$, this example shows that both options predicted by corollary 4.3 can indeed occur. From a mechanical viewpoint the quantity $H = b^2 - a^2 - 1$ is the so-called anisotropy factor, which measures the degree of anisotropy in a cubic crystal, see [11, eq. (13-27)].

5.2. Elliptic physical confinement

In this section, we provide a full characterization of minimizers for a general anisotropy in the case of the physical confinement:

$$V(x) = \begin{cases} 0 & \text{if } x \in E, \\ +\infty & \text{if } x \notin E, \end{cases}$$

where E is the domain enclosed by an ellipse centred at the origin. More precisely, we have the following result.

THEOREM 5.3. *Let $E = RE_0$ with $R \in SO(2)$ and*

$$E_0 = \left\{ x \in \mathbb{R}^2 : \frac{x_1^2}{a_1^2} + \frac{x_2^2}{a_2^2} \leq 1 \right\}.$$

Let J be the functional defined by

$$J(\mu) = \iint_{E \times E} (-\log|x - y| + \kappa(x - y)) \, d\mu(y) \, d\mu(x)$$

for every $\mu \in \mathcal{P}(E)$, where the anisotropy κ is even, 0-homogeneous, and of class H^s on \mathbb{S}^1 with $s > 3/2$. Assume $\widehat{\Psi} \geq 0$ on \mathbb{S}^1 , where $\widehat{\Psi}$ is the function introduced in (3.5). Then the unique minimizer of J is given by the push-forward $g\# \mu_{E_0}$ of the measure

$$\mu_{\partial E_0} := \frac{1}{2\pi a_1 a_2} \left(\frac{x_1^2}{a_1^4} + \frac{x_2^2}{a_2^4} \right)^{-1/2} \mathcal{H}^1 \llcorner \partial E_0$$

through the rotation map $g(x) = Rx$ for $x \in \mathbb{R}^2$.

REMARK 5.4. This result was proved in [19] for a special class of anisotropies. Note that when E is the closed ball B_r of radius $r > 0$ and centre 0 the minimizer is given by the uniform distribution on the boundary of the ball, that is, by the measure:

$$\mu_{\partial B_r} := \frac{1}{2\pi r} \mathcal{H}^1 \llcorner \partial B_r.$$

Proof of theorem 5.3. Up to rotating the axes and replacing $\kappa(x)$ by $\kappa(Rx)$, we can assume without loss of generality that R is the identity matrix and $E = E_0$. Theorem 2.1 and proposition 3.2 guarantee that the minimizer exists and is unique. Moreover, it is characterized by Euler–Lagrange conditions (2.7)–(2.8), which take the following form:

$$(W * \mu)(x) = c \quad \text{for } \mu\text{-a.e. } x \in \text{supp } \mu, \tag{5.3}$$

$$(W * \mu)(x) \geq c \quad \text{for q.e. } x \in E_0. \tag{5.4}$$

To prove the theorem it is enough to show that the measure $\mu_{\partial E_0}$ satisfies (5.3)–(5.4).

For $t > 0$, we consider the set:

$$E_t = \left\{ x \in \mathbb{R}^2 : \frac{x_1^2}{a_1^2} + \frac{x_2^2}{a_2^2} \leq 1 + t \right\}.$$

By formula (4.11) for the potential of a general ellipse we have that

$$\nabla(W * \chi_{E_t})(x) = -a_1 a_2 \int_{\mathbb{S}^1} \frac{\widehat{\Psi}(y)}{|D(a)y|^2} (x \cdot y)y \, d\mathcal{H}^1(y) \tag{5.5}$$

for every $x \in E_t$ and every $t \geq 0$.

Let now x be in the interior of E_0 . Since $x \in E_t$ for every $t \geq 0$, equation (5.5) holds for every $t \geq 0$. Differentiating both sides of (5.5) with respect to t and applying the coarea formula yield:

$$0 = \frac{d}{dt} \left(\int_{E_t} \nabla W(x - y) \, dy \right) = \frac{1}{2} \int_{\partial E_t} \nabla W(x - y) \left(\frac{y_1^2}{a_1^4} + \frac{y_2^2}{a_2^4} \right)^{-1/2} \, d\mathcal{H}^1(y)$$

for a.e. $t \geq 0$. For x in the interior of E_0 the right-hand side is a continuous function of t , therefore we deduce that

$$0 = \frac{1}{2} \int_{\partial E_0} \nabla W(x - y) \left(\frac{y_1^2}{a_1^4} + \frac{y_2^2}{a_2^4} \right)^{-1/2} \, d\mathcal{H}^1(y) = \pi a_1 a_2 \nabla(W * \mu_{\partial E_0})(x)$$

for every x in the interior of E_0 . Since $W * \mu_{\partial E_0}$ is a continuous function in \mathbb{R}^2 , this implies that $W * \mu_{\partial E_0}$ is constant in E_0 , that is, (5.3)–(5.4) are satisfied for $\mu = \mu_{\partial E_0}$. □

5.3. Some open questions and further comments

As a consequence of theorem 4.2 and corollary 4.3, energies of the form (4.3) may have minimizers of non-full dimensionality only if the Fourier transform of their kernel is degenerate. However, degeneracy of the Fourier transform is not a sufficient condition for loss of dimensionality, see remark 5.2. The arguments of proof in theorem 4.2 show that, in the convexity range, loss of dimensionality occurs if and only if system (4.13) does not have a solution M in $\mathbb{M}_{\text{sym},+}^{2 \times 2}$. Yet it would be desirable to devise a criterion for the occurrence of a lower dimensional optimal distribution, without resorting to explicit computations. Similarly, in the case of fully dimensional minimizers, there is no characterization available for the rotation of the optimal ellipse with respect to the coordinate axes.

However, some simple considerations can be made in some specific cases. Indeed, if μ is a measure with no atoms and support on a straight line passing through the origin, then by 0-homogeneity:

$$\iint_{\mathbb{R}^2 \times \mathbb{R}^2} \kappa(x - y) \, d\mu(x) \, d\mu(y) = \kappa(v)$$

where $v \in \mathbb{S}^1$ is a vector parallel to the support of μ . Since the logarithmic interaction and the confinement are radially symmetric, this implies that, among all measures with support on a straight line through the origin, the minimal energy is attained in the directions where κ is minimal. Therefore, if κ has more than one minimizer in the set $\{x \in \mathbb{S}^1 : x_1 \in (-1, 1]\}$, then loss of dimensionality cannot occur, as long as the Fourier transform of the kernel is non-negative, otherwise uniqueness would be violated.

Theorem 5.3 shows that the choice of confinement may also have a strong impact on the shape of minimizers and on their dimensionality. Preliminary computations indicate that strict positivity of the Fourier transform should guarantee full dimensionality of minimizers for smooth confinements, such as, e.g. $V(x) = |x|^p$ with $p \geq 2$.

Another interesting question is the analysis of optimal distributions outside the convexity range, that is, for kernels whose Fourier transform is negative along some directions. Numerical simulations seem to suggest the occurrence of rather complex patterns, see [3].

Acknowledgements

Part of the material of this paper is based on the content of the course ‘Nonlocal interaction problems in dislocation theory’, that the author taught at the Summer School on Analysis and Applied Mathematics in Münster in September 2022. The author would like to express her gratitude to the organizers and to the University of Münster for the support and the hospitality. The author thanks Rémy Rodiac for bringing Bochner theorem to her attention. Support from MIUR-PRIN 2017 is also acknowledged. The author is a member of GNAMPA-INdAM.

References

- 1 L. Ambrosio, N. Gigli and G. Savaré, *Gradient flows in metric spaces and in the space of probability measures*, 2nd ed., Lectures in Mathematics ETH Zürich (Birkhäuser Verlag, Basel, 2008).

- 2 J. A. Carrillo, J. Mateu, M. G. Mora, L. Rondi, L. Scardia and J. Verdera. The ellipse law: Kirchhoff meets dislocations. *Commun. Math. Phys.* **373** (2020), 507–524.
- 3 J. A. Carrillo and R. Shu. Global minimizers of a large class of anisotropic attractive–repulsive interaction energies in 2D. *Commun. Pure Appl. Math.* **77** (2024), 1353–1404.
- 4 J. A. Carrillo and R. Shu, Minimizers of 3D anisotropic interaction energies. *Adv. Calc. Var.*, to appear.
- 5 A. Erdélyi, W. Magnus, F. Oberhettinger and F. G. Tricomi. *Tables of Integral Transforms. Volume I. Based, in part, on notes left by Harry Bateman* (McGraw-Hill Book Co., New York, Toronto, London, 1954).
- 6 G. B. Folland. *Introduction to partial differential equations*, 2nd ed., (Princeton University Press, Princeton, 1995).
- 7 O. Frostman. Potentiel d'équilibre et capacité des ensembles avec quelques applications à la théorie des fonctions. Thesis. *Meddel. Lunds Univ. Mat. Sem.* **3** (1935), 1–118.
- 8 J. Ginibre. Statistical ensembles of complex, quaternion, and real matrices. *J. Math. Phys.* **6** (1965), 440–449.
- 9 R. Giorgio, Explicit minimizers for a class of nonlocal anisotropic dislocation energies. *Master Thesis, Università di Pavia*, 2022.
- 10 V. L. Girko. The circular law (Russian). *Teor. Veroyatnost. Primenen* **29** (1984), 669–679.
- 11 J. P. Hirth and J. Lothe. *Theory of dislocations* (John Wiley & Sons, New York, 1982).
- 12 T. Kolokolnikov, H. Sun, D. Uminsky and A. L. Bertozzi. Stability of ring patterns arising from two-dimensional particle interactions. *Phys. Rev. E* **84** (2011), 015203.
- 13 N. N. Lebedev. *Special Functions and their Applications* (Prentice-Hall, Englewood Cliffs, 1965).
- 14 J. Mateu, M. G. Mora, L. Rondi, L. Scardia and J. Verdera. Stability of ellipsoids as the energy minimisers of perturbed Coulomb energies. *SIAM J. Math. Anal.* **55** (2023), 3650–3676.
- 15 J. Mateu, M. G. Mora, L. Rondi, L. Scardia and J. Verdera. Energy minimisers of perturbed dislocation energies. *Nonlinear Anal.* **231** (2023), 113014.
- 16 J. Mateu, M. G. Mora, L. Rondi, L. Scardia and J. Verdera. Explicit minimisers for anisotropic Coulomb energies in 3D. *Adv. Math.* **434** (2023), 109333.
- 17 M. L. Mehta. *Random matrices and the statistical theory of energy levels* (Academic Press, New York, 1967).
- 18 M. G. Mora, L. Rondi and L. Scardia. The equilibrium measure for a nonlocal dislocation energy. *Commun. Pure Appl. Math.* **72** (2019), 136–158.
- 19 M. G. Mora and A. Scagliotti. Equilibrium measure for a nonlocal dislocation energy with physical confinement. *Adv. Calc. Var.* **15** (2022), 929–938.
- 20 E. Saff and V. Totik. *Logarithmic potentials with external fields* (Springer-Verlag, Berlin Heidelberg, 1997).
- 21 A. Scagliotti, Nonlocal interaction problems in dislocation theory. *Master Thesis, Università di Pavia*, 2018.
- 22 E. M. Stein. *Singular integrals and differentiability properties of functions* (Princeton University Press, Princeton, 1970).
- 23 M. Willem. *Analyse harmonique réelle* (in French) (Collection Méthodes, Hermann, Paris, 1995).