

JOURNAL OF GLACIOLOGY

INDEX Vol. 46 No. 152–155, 2000

- Abe-Ouchi, A., *see* Payne, A. J. and others
- Ablation:
A model of ablation-dominated medial moraines and the generation of debris-mantled glacier snouts, 459–469
- Accumulation:
The accumulation pattern across Siple Dome, West Antarctica, inferred from radar-detected internal layers, 75–87
The accumulation regime of Blue Glacier, U.S.A., 1914–96, 326–334
Mass balance of the northeast sector of the Greenland ice sheet: a remote-sensing perspective, 265–273
Relating glacier mass balance to meteorological data by using a seasonal sensitivity characteristic, 1–6
Snow accumulation rates in northern Victoria Land, Antarctica, by firn-core analysis, 541–552
- Agassiz Ice Cap (Canada):
Natural and anthropogenic levels of tritium in a Canadian Arctic ice core, Agassiz Ice Cap, Ellesmere Island, and comparison with other radionuclides, 35–40
- Ageta, Y., *see* Fujita, K.
- Albedo:
Measurement and parameterization of albedo variations at Haut Glacier d'Arolla, Switzerland, 675–688
- Alean, J., (review: Post, A. and E. R. LaChapelle. 1999. *Glacier ice*) 700
- Algae:
Himalayan ice-core dating with snow algae, 335–340
- Allison, I., *see* Fricker, H. A. and others
- Amery Ice Shelf (Antarctica):
Digital elevation models for the Lambert Glacier–Amery Ice Shelf system, East Antarctica, from ERS-1 satellite radar altimetry, 553
Mass balance of the Lambert Glacier–Amery Ice Shelf system, East Antarctica: a comparison of computed balance fluxes and measured fluxes, 561–570
- Anderson, R. S., (paper: A model of ablation-dominated medial moraines and the generation of debris-mantled glacier snouts) 459–469
- Andes:
Late Pleistocene climate conditions in the north Chilean Andes drawn from a climate–glacier model, 622–632
- Antarctica:
The accumulation pattern across Siple Dome, West Antarctica, inferred from radar-detected internal layers, 75–87
Bed properties of Siple Dome and adjacent ice streams, West Antarctica, inferred from radio-echo sounding measurements, 88–94
Calculating basal thermal zones beneath the Antarctic ice sheet, 297–310
Changes in the margin of Ice Stream C, Antarctica, 102–110
Correspondence. Subglacial water at the heads of Antarctic ice-stream tributaries, 702–703
Digital elevation models for the Lambert Glacier–Amery Ice Shelf system, East Antarctica, from ERS-1 satellite radar altimetry, 553
Elevation of ice-stream margin scars after stagnation, 111–118
Formaldehyde and peroxide concentrations in Law Dome (Antarctica) firn and ice cores, 15–19
Geophysical investigations of ice-sheet internal layering and deformation in the Dome C region of central East Antarctica, 161–166
Ice discharge of eastern Dome C drainage area, Antarctica, determined from airborne radar survey and satellite image analysis, 253
Ice motion over Lake Vostok, Antarctica: constraints on inferences regarding the accreted ice, 689–694
The link between climate warming and break-up of ice shelves in the Antarctic Peninsula, 516–530
Mass balance of the Lambert Glacier–Amery Ice Shelf system, East Antarctica: a comparison of computed balance fluxes and measured fluxes, 561–570
A millennium of variable ice flow recorded by the Ross Ice Shelf, Antarctica, 652–664
Numerical modelling of ice-sheet dynamics across the Vostok subglacial lake, central East Antarctica, 197–205
Numerical modelling of the ice sheet in western Dronning Maud Land, East Antarctica: impacts of present, past and future climates, 54–66
The onset area of Ice Stream D, West Antarctica, 95–101
Short fluctuations in the density and thickness of a dry firn column, 399–411
Snow accumulation rates in northern Victoria Land, Antarctica, by firn-core analysis, 541–552
- Arcone, S. A. and N. E. Yankielun, (paper: 1.4 GHz radar penetration and evidence of drainage structures in temperate ice: Black Rapids Glacier, Alaska, U.S.A.) 477–490
- Arcone, S. A., *see* Moran, M. L. and others
- Austria:
A geometric glacier model for sea-level change calculations, 357–368
Relating glacier mass balance to meteorological data by using a seasonal sensitivity characteristic, 1–6
The use of bulk and profile methods for determining surface heat fluxes in the presence of glacier winds, 445
- Avalanche:
Texture and strength changes of buried surface-hoar layers with implications for dry snow-slab avalanche release, 151–160
- Ayers, G. P., *see* Gillett, R. W. and others
- Bach Ice Shelf (Antarctica):
The link between climate warming and break-up of ice shelves in the Antarctic Peninsula, 516–530
- Baker, I., *see* Cullen, D.
- Baker, I., *see* Trickett, Y. L. and others
- Baltic Sea On the modelling of ice-thickness redistribution, 427–437
- Bamber, J. L. and others, (paper: An analysis of balance velocities over the Greenland ice sheet and comparison with synthetic aperture radar interferometry) 67–74
- Bamber, J. L., *see* Siegert, M. J.
- Barton, J. S., *see* Hall, D. K. and others
- Basal conditions:
Bed properties of Siple Dome and adjacent ice streams, West Antarctica, inferred from radio-echo sounding measurements, 88–94
Calculating basal thermal zones beneath the Antarctic ice sheet, 297–310
Changes in the margin of Ice Stream C, Antarctica, 102–110
Coupled ice–till deformation near subglacial channels and cavities, 580–598
Delineation of a complexly dipping temperate glacier bed using short-pulse radar arrays, 274
Diffusive mixing between shearing granular layers: constraints on bed deformation from till contacts, 641–651
Energy balance of ice streams, 665–674
Evidence for extreme pressure pulses in the subglacial water system, 206–212
Glacier motion dominated by processes deep in underlying till, 213–221
Ice-core evidence of the thickness and character of clear-facies basal ice: Glacier de Transfleuron, Switzerland, 140–150
Rheology of ice at the bed of Engabreen, Norway, 611–621
Sliding of ice past an obstacle at Engabreen, Norway, 599–610
Sliding versus till deformation in the fast motion of an ice stream over a viscous till, 633–640
- Becagli, S., *see* Stenni, B. and others
- Beer, J., *see* Eichler, A. and others
- Bennett, M. R. and others, (paper: Glaciofluvial crevasse and conduit fills as indicators of supraglacial dewatering during a surge, Skeiðarárjökull, Iceland) 25–34
- Bennett, M. R., *see* Glasser, N. F. and others
- Bindschadler, R. and others, (paper: The onset area of Ice Stream D, West Antarctica), p. 352
- Bindschadler, R. A., *see* Fahnestock, M. A. and others
- Black Rapids Glacier (U.S.A.):
1.4 GHz radar penetration and evidence of drainage structures in temperate ice: Black Rapids Glacier, Alaska, U.S.A., 477–490
Comparison of SAR-interferometric and surveyed velocities on a mountain glacier: Black Rapids Glacier, Alaska, U.S.A., 119–128
Glacier motion dominated by processes deep in underlying till, 213–221

- Blatter, H., *see* Hubbard, A. and others
- Blue Glacier (U.S.A.):
The accumulation regime of Blue Glacier, U.S.A., 1914–96, 326–334
- Bohlander, J., *see* Scambos, T. A. and others
- Bolsey, R., *see* Engelhardt, H. and others
- Boyle, P., *see* Jiskoot, H. and others
- Braithwaite, R. J. and Y. Zhang, (paper: Sensitivity of mass balance of five Swiss glaciers to temperature changes assessed by tuning a degree-day model) 7–14
- Braithwaite, R. J., *see* Raper, S. C. B. and others
- Brock, B. W. and others, (paper: Measurement and parameterization of albedo variations at Haut Glacier d'Arolla, Switzerland) 675–688
- Brown, O., *see* Raper, S. C. B. and others
- Brun, E., *see* Durand, Y. and others
- Buscarlet, G., *see* Rignot, E. and others
- Byrd (Antarctica):
Short fluctuations in the density and thickness of a dry firn column, 399–411
- Byrd Glacier (Antarctica):
Correspondence. Subglacial water at the heads of Antarctic ice-stream tributaries, 702–703
- Calov, R., *see* Payne, A. J. and others
- Calving:
Glacier calving: a numerical model of forces in the calving-speed/water-depth relation, 188–196
- Camp Century (Greenland):
Short fluctuations in the density and thickness of a dry firn column, 399–411
- Canada:
Evidence for extreme pressure pulses in the subglacial water system, 206–212
Natural and anthropogenic levels of tritium in a Canadian Arctic ice core, Agassiz Ice Cap, Ellesmere Island, and comparison with other radio-nuclides, 35–40
Relating glacier mass balance to meteorological data by using a seasonal sensitivity characteristic, 1–6
Texture and strength changes of buried surface-hoar layers with implications for dry snow-slab avalanche release, 151–160
- Cantwell Glacier (U.S.A.):
A model of ablation-dominated medial moraines and the generation of debris-mantled glacier snouts, 459–469
- Carbon dioxide:
Reconstructing past atmospheric CO₂-concentration based on ice-core analyses: open questions due to in situ production of CO₂ in ice, 45–53
- Carsey, F. D., *see* Kwok, R. and others
- Cecil, L. DeW., *see* Morin, R. H. and others
- Chemistry:
Correspondence. The chemistry of grain boundaries in Greenland ice, 703–706
Formaldehyde and peroxide concentrations in Law Dome (Antarctica) firn and ice cores, 15–19
Glaciochemical dating of an ice core from upper Grenzgletscher (4200 m a.s.l.), 507–515
Natural and anthropogenic levels of tritium in a Canadian Arctic ice core, Agassiz Ice Cap, Ellesmere Island, and comparison with other radio-nuclides, 35–40
The effects of sulfuric acid on the mechanical properties of ice single crystals, 239–243
- Chen, X., *see* Bindschadler, R. and others
- Chile:
Late Pleistocene climate conditions in the north Chilean Andes drawn from a climate–glacier model, 622–632
- China:
Effect of summer accumulation on glacier mass balance on the Tibetan Plateau revealed by mass-balance model, 244–252
Monsoon and dust signals recorded in Dasuopu glacier, Tibetan Plateau, 222–226
- Choquelimpie glacier (Chile):
Late Pleistocene climate conditions in the north Chilean Andes drawn from a climate–glacier model, 622–632
- Christensen, E. L. and others, (paper: A low-cost glacier-mapping system) 531–537
- Clarke, G. K. C., *see* Kavanaugh, J. L.
- Clausen, H. B., *see* Hempel, L. and others
- Climate:
Late Pleistocene climate conditions in the north Chilean Andes drawn from a climate–glacier model, 622–632
The link between climate warming and break-up of ice shelves in the Antarctic Peninsula, 516–530
Numerical modelling of the ice sheet in western Dronning Maud Land, East Antarctica: impacts of present, past and future climates, 54–66
Reconstructing past atmospheric CO₂-concentration based on ice-core analyses: open questions due to in situ production of CO₂ in ice, 45–53
- Cohen, D. and others, (paper: Sliding of ice past an obstacle at Engabreen, Norway) 599–610
- Cohen, D., (paper: Rheology of ice at the bed of Engabreen, Norway) 611–621
- Coleman, R., *see* Fricker, H. A. and others
- Colgan, P. M., *see* Cutler, P. M. and others
- Compression:
The orientation dependence of the strength of ice single crystals, 41–44
- Conductivity:
A comparison of radio-echo sounding data and electrical conductivity of the GRIP ice core, 369–374
- Conway, H., *see* Gades, A. M. and others
- Conway, H., *see* Rasmussen, L. A. and others
- Correspondence:
Rapid sediment entrainment and englacial deposition during jökulhlaups, 349–351
Subglacial water at the heads of Antarctic ice-stream tributaries, 702–703
The chemistry of grain boundaries in Greenland ice, 703–706
- Crabtree, M. D., *see* Murray, T. and others
- Crawford, K. R., *see* Glasser, N. F. and others
- Crevasses:
Glaciofluvial crevasse and conduit fills as indicators of supraglacial dewatering during a surge, Skeiðarárjökull, Iceland, 25–34
- Csathó, B., *see* Rignot, E. and others
- Cullen, D. and I. Baker, (correspondence: The chemistry of grain boundaries in Greenland ice) 703–706
- Cutler, P. M. and others, (paper: A numerical investigation of ice-lobe–permafrost interaction around the southern Laurentide ice sheet) 311–325
- Dalton Iceberg Tongue (Antarctica):
Calculating basal thermal zones beneath the Antarctic ice sheet, 297–310
- Dasuopu glacier (Tibet):
Monsoon and dust signals recorded in Dasuopu glacier, Tibetan Plateau, 222–226
- Dating:
Glaciochemical dating of an ice core from upper Grenzgletscher (4200 m a.s.l.), 507–515
Himalayan ice-core dating with snow algae, 335–340
- David Glacier (Antarctica):
Ice discharge of eastern Dome C drainage area, Antarctica, determined from airborne radar survey and satellite image analysis, 253
Correspondence. Subglacial water at the heads of Antarctic ice-stream tributaries, 702–703
- Delaney, A. J., *see* Moran, M. L. and others
- Denby, B. and W. Greuell, (paper: The use of bulk and profile methods for determining surface heat fluxes in the presence of glacier winds) 445–452
- Descamps, G. E., *see* Morin, R. H. and others
- Dibble Glacier (Antarctica):
Calculating basal thermal zones beneath the Antarctic ice sheet, 297–310
- Dixon, T. S., *see* Rea, B. R. and others
- Dome C [Concordia] (Antarctica):
Geophysical investigations of ice-sheet internal layering and deformation in the Dome C region of central East Antarctica, 161–166
Ice discharge of eastern Dome C drainage area, Antarctica, determined from airborne radar survey and satellite image analysis, 253
- Dongkemadi glacier (Tibet):
Effect of summer accumulation on glacier mass balance on the Tibetan Plateau revealed by mass-balance model, 244–252
- Dowdeswell, J. A., *see* Hodgkins, R. and others
- Drilling:
A hot-water ice-coring drill, 341–345
Portable system for intermediate-depth ice-core drilling, 167–172

- Durand, Y. and others, (erratum: A computer-based system simulating snowpack structures as a tool for regional avalanche forecasting) 173
- East Fork Toklat Glacier (U.S.A.):
A model of ablation-dominated medial moraines and the generation of debris-mantled glacier snouts, 459–469
- Echelmeyer, K., (review: Petrenko, Victor F. and Robert W. Whitworth. 1999. *The physics of ice.*) 347–348
- Echelmeyer, K. A., *see* Truffer, M. and others
- Eichler, A. and others, (paper: Glaciochemical dating of an ice core from upper Grenzletscher (4200 m a.s.l.)) 507–515
- EISMINT:
Results from the EISMINT model intercomparison: the effects of thermo-mechanical coupling, 227–238
- El Tatio glacier (Chile):
Late Pleistocene climate conditions in the north Chilean Andes drawn from a climate–glacier model, 622–632
- Energy balance:
Analysis of a 3 year meteorological record from the ablation zone of Morteratschgletscher, Switzerland: energy and mass balance, 571–579
Sensitivity of mass balance of five Swiss glaciers to temperature changes assessed by tuning a degree-day model, 7–14
The use of bulk and profile methods for determining surface heat fluxes in the presence of glacier winds, 445
- Engabreen (Norway):
Rheology of ice at the bed of Engabreen, Norway, 611–621
Sliding of ice past an obstacle at Engabreen, Norway, 599–610
- Engelhardt, H. and others, (paper: A hot-water ice-coring drill) 341–345
- EPICA:
Geophysical investigations of ice-sheet internal layering and deformation in the Dome C region of central East Antarctica, 161–166
- Errata:
A computer-based system simulating snowpack structures as a tool for regional avalanche forecasting, 173
Delineation of a complexly dipping temperate glacier bed using short-pulse radar arrays, 707
Numerical modelling of the ice sheet in western Dronning Maud Land, East Antarctica: impacts of present, past and future climates, 353–354
The structural glaciology of Kongsvegen, Svalbard, and its role in land-form genesis, 538
- Evans, D. J. A., *see* Rea, B. R. and others
- Fahnestock, M., *see* Scambos, T. A. and others
- Fahnestock, M. A. and others, (paper: A millennium of variable ice flow recorded by the Ross Ice Shelf, Antarctica) 652–664
- Falljökull (Iceland):
Englacial water distribution in a temperate glacier from surface and bore-hole radar velocity analysis, 389–398
- Fastook, J. L., *see* Näslund, J.-O. and others
- Fastook, J. L., *see* Payne, A. J. and others
- Fatland, D. R., *see* Rabus, B. T.
- Forsberg, R., *see* Christensen, E. L. and others
- Fountain, A. G., (review: Knight, Peter G. 1999. *Glaciers.*) 346
- France:
Dynamic behaviour analysis of glacier de Saint Sorlin, France, from 40 years of observations, 1957–97, 499–506
- Franz Josef Glacier (New Zealand):
Relating glacier mass balance to meteorological data by using a seasonal sensitivity characteristic, 1–6
- Frezzotti, M. and others, (paper: Ice discharge of eastern Dome C drainage area, Antarctica, determined from airborne radar survey and satellite image analysis) 253–264
- Frezzotti, M., *see* Stenni, B. and others
- Fricker, H. A. and others, (paper: Digital elevation models for the Lambert Glacier–Amery Ice Shelf system, East Antarctica, from ERS-1 satellite radar altimetry) 553–560
- Fricker, H. A. and others, (paper: Mass balance of the Lambert Glacier–Amery Ice Shelf system, East Antarctica: a comparison of computed balance fluxes and measured fluxes) 561–570
- Fry, M., *see* Murray, T. and others
- Fujita, K. and Y. Ageta, (paper: Effect of summer accumulation on glacier mass balance on the Tibetan Plateau revealed by mass-balance model) 244–252
- Fujita, K., *see* Yoshimura, Y. and others
- Funk, M., *see* Eichler, A. and others
- Furrer, V., *see* Eichler, A. and others
- Gades, A. M. and others, (paper: Bed properties of Siple Dome and adjacent ice streams, West Antarctica, inferred from radio-echo sounding measurements) 88–94
- Gäggeler, H. W., *see* Eichler, A. and others
- Gamble, N. H., *see* Murray, T. and others
- Garvin, J. B., *see* Hall, D. K. and others
- George VI Ice Shelf (Antarctica):
The link between climate warming and break-up of ice shelves in the Antarctic Peninsula, 516–530
- Gillett, R. W. and others, (paper: Formaldehyde and peroxide concentrations in Law Dome (Antarctica) firn and ice cores) 15–19
- Giraud, G., *see* Durand, Y. and others
- GISP2:
Correspondence. The chemistry of grain boundaries in Greenland ice, 703–706
- Glacial Geology:
Contemporaneous, localized, basal ice-flow variations: implications for bedrock erosion and the origin of p-forms, 470–476
Diffusive mixing between shearing granular layers: constraints on bed deformation from till contacts, 641–651
Glacier advance, ice-marginal lakes and the routing of meltwater and sediment: Russell Glacier, Greenland, 423–426
A model of ablation-dominated medial moraines and the generation of debris-mantled glacier snouts, 459–469
A numerical investigation of ice-lobe–permafrost interaction around the southern Laurentide ice sheet, 311
- Glacier flow:
An analysis of balance velocities over the Greenland ice sheet and comparison with synthetic aperture radar interferometry, 67–74
Calculating basal thermal zones beneath the Antarctic ice sheet, 297–310
Changes in the margin of Ice Stream C, Antarctica, 102–110
Comparison of SAR-interferometric and surveyed velocities on a mountain glacier: Black Rapids Glacier, Alaska, U.S.A., 119–128
Coupled ice–till deformation near subglacial channels and cavities, 580–598
Diffusive mixing between shearing granular layers: constraints on bed deformation from till contacts, 641–651
Dynamic behaviour analysis of glacier de Saint Sorlin, France, from 40 years of observations, 1957–97, 499–506
Elevation of ice-stream margin scars after stagnation, 111–118
Energy balance of ice streams, 665–674
Glacier mass-balance determination by remote sensing and high-resolution modelling, 491–498
Geophysical investigations of ice-sheet internal layering and deformation in the Dome C region of central East Antarctica, 161–166
Glacier motion dominated by processes deep in underlying till, 213–221
Ice discharge of eastern Dome C drainage area, Antarctica, determined from airborne radar survey and satellite image analysis, 253
Ice motion over Lake Vostok, Antarctica: constraints on inferences regarding the accreted ice, 689–694
Mass balance of the northeast sector of the Greenland ice sheet: a remote-sensing perspective, 265
A millennium of variable ice flow recorded by the Ross Ice Shelf, Antarctica, 652–664
Modelling dynamics of glaciers in volcanic craters, 177–187
Non-temperate glacier flow over wavy sloping ground, 453–458
Numerical modelling of ice-sheet dynamics across the Vostok subglacial lake, central East Antarctica, 197–205
The onset area of Ice Stream D, West Antarctica, 95–101
Results from the EISMINT model intercomparison: the effects of thermo-mechanical coupling, 227–238
Rheology of ice at the bed of Engabreen, Norway, 611–621
Sliding of ice past an obstacle at Engabreen, Norway, 599–610
Sliding versus till deformation in the fast motion of an ice stream over a viscous till, 633–640
- Glacier hydrology:
1.4 GHz radar penetration and evidence of drainage structures in temperate ice: Black Rapids Glacier, Alaska, U.S.A., 477–490
Correspondence. Subglacial water at the heads of Antarctic ice-stream tributaries, 702–703
Englacial water distribution in a temperate glacier from surface and bore-hole radar velocity analysis, 389–398
Evidence for extreme pressure pulses in the subglacial water system, 206–212

- Glacier advance, ice-marginal lakes and the routing of meltwater and sediment: Russell Glacier, Greenland, 423–426
- Glaciofluvial crevasse and conduit fills as indicators of supraglacial dewatering during a surge, Skeiðarárjökull, Iceland, 25–34
- Rapid sediment entrainment and englacial deposition during jökulhlaups, 349–351
- Glacier Inventory:
Controls on the distribution of surge-type glaciers in Svalbard, 412–422
- Glacier variations:
Evaluation of remote-sensing techniques to measure decadal-scale changes of Hofsjökull ice cap, Iceland, 375–388
A geometric glacier model for sea-level change calculations, 357–368
Glacier advance, ice-marginal lakes and the routing of meltwater and sediment: Russell Glacier, Greenland, 423–426
- Glaciers:
1.4 GHz radar penetration and evidence of drainage structures in temperate ice: Black Rapids Glacier, Alaska, U.S.A., 477–490
The accumulation regime of Blue Glacier, U.S.A., 1914–96, 326–334
Acoustic televiewer logging in glacier boreholes, 695–699
Analysis of a 3 year meteorological record from the ablation zone of Morteratschgletscher, Switzerland: energy and mass balance, 571–579
Changes in the margin of Ice Stream C, Antarctica, 102–110
Comparison of SAR-interferometric and surveyed velocities on a mountain glacier: Black Rapids Glacier, Alaska, U.S.A., 119–128
Contemporaneous, localized, basal ice-flow variations: implications for bedrock erosion and the origin of p-forms, 470–476
Controls on the distribution of surge-type glaciers in Svalbard, 412–422
Coupled ice–till deformation near subglacial channels and cavities, 580–598
Delineation of a complexly dipping temperate glacier bed using short-pulse radar arrays, 274–286
Dynamic behaviour analysis of glacier de Saint Sorlin, France, from 40 years of observations, 1957–97, 499–506
Effect of summer accumulation on glacier mass balance on the Tibetan Plateau revealed by mass-balance model, 244–252
Elevation of ice-stream margin scars after stagnation, 111–118
Energy balance of ice streams, 665–674
Englacial water distribution in a temperate glacier from surface and borehole radar velocity analysis, 389–398
Evaluation of remote-sensing techniques to measure decadal-scale changes of Hofsjökull ice cap, Iceland, 375–388
Evidence for extreme pressure pulses in the subglacial water system, 206–212
A geometric glacier model for sea-level change calculations, 357–368
Glacier advance, ice-marginal lakes and the routing of meltwater and sediment: Russell Glacier, Greenland, 423–426
Glacier calving: a numerical model of forces in the calving-speed/water-depth relation, 188–196
Glacier mass-balance determination by remote sensing and high-resolution modelling, 491–498
Glacier motion dominated by processes deep in underlying till, 213–221
Glaciofluvial crevasse and conduit fills as indicators of supraglacial dewatering during a surge, Skeiðarárjökull, Iceland, 25–34
Ice-core evidence of the thickness and character of clear-facies basal ice: Glacier de Transfleuron, Switzerland, 140–150
Late Pleistocene climate conditions in the north Chilean Andes drawn from a climate–glacier model, 622–632
The link between climate warming and break-up of ice shelves in the Antarctic Peninsula, 516–530
A low-cost glacier-mapping system, 531–537
Measurement and parameterization of albedo variations at Haut Glacier d'Arolla, Switzerland, 675–688
A model of ablation-dominated medial moraines and the generation of debris-mantled glacier snouts, 459–469
Modelling dynamics of glaciers in volcanic craters, 177–187
Non-temperate glacier flow over wavy sloping ground, 453–458
The onset area of Ice Stream D, West Antarctica, 95–101
Sensitivity of mass balance of five Swiss glaciers to temperature changes assessed by tuning a degree-day model, 7–14
Sliding versus till deformation in the fast motion of an ice stream over a viscous till, 633–640
Radar glacier zones in southeast Alaska, U.S.A.: field and satellite observations, 287–296
Relating glacier mass balance to meteorological data by using a seasonal sensitivity characteristic, 1–6
The use of bulk and profile methods for determining surface heat fluxes in the presence of glacier winds, 445–452
- Glasser, N. F. and others, (erratum: The structural glaciology of Kongsvegen, Svalbard, and its role in landform genesis) 538
- Gogineni, S., *see* Rignot, E. and others
- Ground-penetration radar (GPR):
Englacial water distribution in a temperate glacier from surface and borehole radar velocity analysis, 389–398
- Greenfield, R. J., *see* Moran, M. L. and others
- Greenland:
An analysis of balance velocities over the Greenland ice sheet and comparison with synthetic aperture radar interferometry, 67–74
A comparison of radio-echo sounding data and electrical conductivity of the GRIP ice core, 369–374
Correspondence. The chemistry of grain boundaries in Greenland ice, 703–706
Glacier advance, ice-marginal lakes and the routing of meltwater and sediment: Russell Glacier, Greenland, 423–426
A low-cost glacier-mapping system, 531–537
Mass balance of the northeast sector of the Greenland ice sheet: a remote-sensing perspective, 265
Portable system for intermediate-depth ice-core drilling, 167–172
Reconstructing past atmospheric CO₂-concentration based on ice-core analyses: open questions due to in situ production of CO₂ in ice, 45–53
Short fluctuations in the density and thickness of a dry firn column, 399–411
- Grenzgletscher (Switzerland):
Glaciochemical dating of an ice core from upper Grenzgletscher (4200 m a.s.l.), 507–515
- Greuell, W., *see* Denby, B.
- Greve, R., *see* Payne, A. J. and others
- Griegsgletscher (Switzerland) Sensitivity of mass balance of five Swiss glaciers to temperature changes assessed by tuning a degree-day model, 7–14
- GRIP:
A comparison of radio-echo sounding data and electrical conductivity of the GRIP ice core, 369–374
Reconstructing past atmospheric CO₂-concentration based on ice-core analyses: open questions due to in situ production of CO₂ in ice, 45–53
- Grosjean, M., *see* Kull, C.
- Gulkana Glacier (U.S.A.):
Delineation of a complexly dipping temperate glacier bed using short-pulse radar arrays, 274
- Gundestrup, N., *see* Hempel, L. and others
- Haapala, J., (paper: On the modelling of ice-thickness redistribution) 427–437
- Hall, D. K. and others, (paper: Evaluation of remote-sensing techniques to measure decadal-scale changes of Hofsjökull ice cap, Iceland) 375–388
- Hambrey, M. J., *see* Glasser, N. F. and others
- Hanson, B. and R. LeB. Hooke, (paper: Glacier calving: a numerical model of forces in the calving-speed/water-depth relation) 188–196
- Hardy, R. J., *see* Bamber, J. L. and others
- Harrison, W. D., *see* Truffer, M. and others
- Haut Glacier d'Arolla (Switzerland):
Glacier mass-balance determination by remote sensing and high-resolution modelling, 491–498
Measurement and parameterization of albedo variations at Haut Glacier d'Arolla, Switzerland, 675–688
- Hayes, P. S., *see* Rasmussen, L. A. and others
- Hempel, L. and others, (paper: A comparison of radio-echo sounding data and electrical conductivity of the GRIP ice core) 369–374
- Himalaya:
Himalayan ice-core dating with snow algae, 335–340
Monsoon and dust signals recorded in Dasuopu glacier, Tibetan Plateau, 222–226
- Hintereisferner (Austria):
A geometric glacier model for sea-level change calculations, 357–368
Relating glacier mass balance to meteorological data by using a seasonal sensitivity characteristic, 1–6
- Hoar:
Texture and strength changes of buried surface-hoar layers with implications for dry snow-slab avalanche release, 151–160
- Hodgkins, R. and others, (paper: Geophysical investigations of ice-sheet internal layering and deformation in the Dome C region of central East Antarctica) 161–166
- Hofsjökull (Iceland):
Evaluation of remote-sensing techniques to measure decadal-scale changes of Hofsjökull ice cap, Iceland, 375–388
- Holmlund, P., *see* Näslund, J.-O. and others
- Hooke, R. LeB., *see* Cohen, D. and others
- Hooke, R. LeB., *see* Hanson, B.

- Hooyer, T. S. and N. R. Iverson, (paper: Diffusive mixing between shearing granular layers: constraints on bed deformation from till contacts) 641–651
- Hubbard, A. and others, (paper: Glacier mass-balance determination by remote sensing and high-resolution modelling) 491–498
- Hubbard, B. and others, (paper: Ice-core evidence of the thickness and character of clear-facies basal ice: Glacier de Transfleuron, Switzerland) 140–150
- Hubbard, B., *see* Hubbard, A. and others
- Huddart, D., *see* Bennett, M. R. and others
- Huddart, D., *see* Glasser, N. F. and others
- Hughes, T. J., *see* Wilch, E.
- Hulbe, C., *see* Scambos, T. A. and others
- Huybrechts, P., *see* Payne, A. J. and others
- Hyland, G., *see* Fricker, H. A. and others
- Ice cores:
- A comparison of radio-echo sounding data and electrical conductivity of the GRIP ice core, 369–374
 - Correspondence. The chemistry of grain boundaries in Greenland ice, 703–706
 - Formaldehyde and peroxide concentrations in Law Dome (Antarctica) firn and ice cores, 15–19
 - Glaciochemical dating of an ice core from upper Grenzgletscher (4200 m a.s.l.), 507–515
 - Himalayan ice-core dating with snow algae, 335–340
 - A hot-water ice-coring drill, 341–345
 - Ice-core evidence of the thickness and character of clear-facies basal ice: Glacier de Transfleuron, Switzerland, 140–150
 - Monsoon and dust signals recorded in Dasuopu glacier, Tibetan Plateau, 222–226
 - Natural and anthropogenic levels of tritium in a Canadian Arctic ice core, Agassiz Ice Cap, Ellesmere Island, and comparison with other radionuclides, 35–40
 - Portable system for intermediate-depth ice-core drilling, 167–172
 - Reconstructing past atmospheric CO₂-concentration based on ice-core analyses: open questions due to in situ production of CO₂ in ice, 45–53
 - Snow accumulation rates in northern Victoria Land, Antarctica, by firn-core analysis, 541–552
- Ice physics:
- The effects of sulfuric acid on the mechanical properties of ice single crystals, 239–243
 - A new technique for ice-fabric analysis, 129–139
 - The orientation dependence of the strength of ice single crystals, 41–44
- Ice sheets:
- The accumulation pattern across Siple Dome, West Antarctica, inferred from radar-detected internal layers, 75–87
 - An analysis of balance velocities over the Greenland ice sheet and comparison with synthetic aperture radar interferometry, 67–74
 - Bed properties of Siple Dome and adjacent ice streams, West Antarctica, inferred from radio-echo sounding measurements, 88–94
 - Calculating basal thermal zones beneath the Antarctic ice sheet, 297–310
 - A comparison of radio-echo sounding data and electrical conductivity of the GRIP ice core, 369–374
 - Correspondence. Subglacial water at the heads of Antarctic ice-stream tributaries, 702–703
 - Diffusive mixing between shearing granular layers: constraints on bed deformation from till contacts, 641–651
 - Geophysical investigations of ice-sheet internal layering and deformation in the Dome C region of central East Antarctica, 161–166
 - Ice discharge of eastern Dome C drainage area, Antarctica, determined from airborne radar survey and satellite image analysis, 253–264
 - Mass balance of the northeast sector of the Greenland ice sheet: a remote-sensing perspective, 265
 - A numerical investigation of ice-lobe–permafrost interaction around the southern Laurentide ice sheet, 311
 - Numerical modelling of ice-sheet dynamics across the Vostok subglacial lake, central East Antarctica, 197–205
 - Numerical modelling of the ice sheet in western Dronning Maud Land, East Antarctica: impacts of present, past and future climates, 54–66
 - Portable system for intermediate-depth ice-core drilling, 167–172
 - Results from the EISMINT model intercomparison: the effects of thermo-mechanical coupling, 227–238
- Ice shelves:
- The link between climate warming and break-up of ice shelves in the Antarctic Peninsula, 516–530
 - A millennium of variable ice flow recorded by the Ross Ice Shelf, Antarctica, 652–664
- Ice Stream B (Antarctica):
- Correspondence. Subglacial water at the heads of Antarctic ice-stream tributaries, 702–703
 - Energy balance of ice streams, 665–674
- Ice Stream C (Antarctica):
- The accumulation pattern across Siple Dome, West Antarctica, inferred from radar-detected internal layers, 75–87
 - Bed properties of Siple Dome and adjacent ice streams, West Antarctica, inferred from radio-echo sounding measurements, 88–94
 - Changes in the margin of Ice Stream C, Antarctica, 102–110
 - Elevation of ice-stream margin scars after stagnation, 111–118
- Ice Stream D (Antarctica):
- The accumulation pattern across Siple Dome, West Antarctica, inferred from radar-detected internal layers, 75–87
 - Bed properties of Siple Dome and adjacent ice streams, West Antarctica, inferred from radio-echo sounding measurements, 88–94
 - Elevation of ice-stream margin scars after stagnation, 111–118
 - Energy balance of ice streams, 665–674
 - The onset area of Ice Stream D, West Antarctica, 95–101
- Ice Stream E (Antarctica):
- Energy balance of ice streams, 665–674
- Ice thickness:
- Ice-core evidence of the thickness and character of clear-facies basal ice: Glacier de Transfleuron, Switzerland, 140–150
 - The onset area of Ice Stream D, West Antarctica, 95–101
- Iceland:
- Englacial water distribution in a temperate glacier from surface and bore-hole radar velocity analysis, 389–398
 - Evaluation of remote-sensing techniques to measure decadal-scale changes of Hofsjökull ice cap, Iceland, 375–388
 - Glaciofluvial crevasse and conduit fills as indicators of supraglacial dewatering during a surge, Skeiðarárjökull, Iceland, 25–34
 - Rapid sediment entrainment and englacial deposition during jökulhlaups, 349–351
- Instruments and methods:
- Acoustic televiewer logging in glacier boreholes, 695–699
 - A hot-water ice-coring drill, 341–345
 - A low-cost glacier-mapping system, 531–537
 - A new technique for ice-fabric analysis, 129–139
 - Portable system for intermediate-depth ice-core drilling, 167–172
- Interferometry:
- An analysis of balance velocities over the Greenland ice sheet and comparison with synthetic aperture radar interferometry, 67–74
 - Comparison of SAR-interferometric and surveyed velocities on a mountain glacier: Black Rapids Glacier, Alaska, U.S.A., 119–128
- Isacks, B. L., *see* Ramage, J. M. and others
- Iverson, N. R., *see* Cohen, D. and others
- Iverson, N. R., *see* Hooyer, T. S.
- Jackson, A. V., *see* Gillett, R. W. and others
- Jacobel, R. W. and others, (paper: Changes in the margin of Ice Stream C, Antarctica) 102–110
- Jacobel, R. W., *see* Gades, A. M. and others
- Jacobel, R. W., *see* Nereson, N. A. and others
- Jamieson, J. B. and J. Schweizer, (paper: Texture and strength changes of buried surface-hoar layers with implications for dry snow-slab avalanche release) 151–160
- Janssens, L., *see* Hubbard, B. and others
- Jiskoot, H. and others, (paper: Controls on the distribution of surge-type glaciers in Svalbard) 412–422
- Jökulhlaups:
- Rapid sediment entrainment and englacial deposition during jökulhlaups, 349–351
 - Glacier advance, ice-marginal lakes and the routing of meltwater and sediment: Russell Glacier, Greenland, 423–426
- Jones, A. P., *see* Knight, P. G. and others
- Jørgensen, J. H., *see* Christensen, E. L. and others
- Joughin, I., *see* Bamber, J. L. and others
- Juneau Icefield (U.S.A./Canada):
- Radar glacier zones in southeast Alaska, U.S.A.: field and satellite observations, 287–296
- Kamb, B., *see* Engelhardt, H. and others

- Kang Shichang and others, (paper: Monsoon and dust signals recorded in Dasuopu glacier, Tibetan Plateau) 222–226
- Kavanaugh, J. L. and G. K. C. Clarke, (paper: Evidence for extreme pressure pulses in the subglacial water system) 206–212
- Kirghizstan:
Relating glacier mass balance to meteorological data by using a seasonal sensitivity characteristic, 1–6
- Knight, P. G. and others, (paper: Glacier advance, ice-marginal lakes and the routing of meltwater and sediment: Russell Glacier, Greenland) 423–426
- Knudsen, Ó., *see* Roberts, M. J. and others
- Kohler, J., *see* Cohen, D. and others
- Kohshima, S., *see* Yoshimura, Y. and others
- Konowbreen (Svalbard):
Contemporaneous, localized, basal ice-flow variations: implications for bedrock erosion and the origin of p-forms, 470–476
- Kotzer, T. G. and others, (paper: Natural and anthropogenic levels of tritium in a Canadian Arctic ice core, Agassiz Ice Cap, Ellesmere Island, and comparison with other radionuclides) 35–40
- Krabill, W., *see* Rignot, E. and others
- Kudo, A., *see* Kotzer, T. G. and others
- Kull, C. and M. Grosjean, (paper: Late Pleistocene climate conditions in the north Chilean Andes drawn from a climate–glacier model) 622–632
- Kvaran, G., *see* Fahnestock, M. A. and others
- Kwok, R. and others, (paper: Ice motion over Lake Vostok, Antarctica: constraints on inferences regarding the accreted ice) 689–694
- Lambert Glacier (Antarctica):
Digital elevation models for the Lambert Glacier–Amery Ice Shelf system, East Antarctica, from ERS-1 satellite radar altimetry, 553
Mass balance of the Lambert Glacier–Amery Ice Shelf system, East Antarctica: a comparison of computed balance fluxes and measured fluxes, 561–570
- Larsen Ice Shelf (Antarctica):
The link between climate warming and break-up of ice shelves in the Antarctic Peninsula, 516–530
- Laurentide ice sheet:
A numerical investigation of ice-lobe–permafrost interaction around the southern Laurentide ice sheet, 311
Diffusive mixing between shearing granular layers: constraints on bed deformation from till contacts, 641–651
- Law Dome (Antarctica):
Formaldehyde and peroxide concentrations in Law Dome (Antarctica) firn and ice cores, 15–19
- Le Meur, E., *see* Vincent, C. and others
- Limmerngletscher (Switzerland):
Sensitivity of mass balance of five Swiss glaciers to temperature changes assessed by tuning a degree-day model, 7–14
- MacAyeal, D. R., *see* Cutler, P. M. and others
- Mackay Glacier (Antarctica):
Ice discharge of eastern Dome C drainage area, Antarctica, determined from airborne radar survey and satellite image analysis, 253
- Maggi, V., *see* Stenni, B. and others
- Mair, D., *see* Hubbard, A. and others
- Mapping: A low-cost glacier-mapping system, 531–537
- Mars: A flow model for the polar caps of Mars, 438–444
- Marshall, S. J., *see* Payne, A. J. and others
- Marsiat, I., *see* Payne, A. J. and others
- Martin, E., *see* Durand, Y. and others
- Mass balance:
Analysis of a 3 year meteorological record from the ablation zone of Morteratschgletscher, Switzerland: energy and mass balance, 571–579
Dynamic behaviour analysis of glacier de Saint Sorlin, France, from 40 years of observations, 1957–97, 499–506
Effect of summer accumulation on glacier mass balance on the Tibetan Plateau revealed by mass-balance model, 244–252
A geometric glacier model for sea-level change calculations, 357–368
Glacier mass-balance determination by remote sensing and high-resolution modelling, 491–498
Mass balance of the Lambert Glacier–Amery Ice Shelf system, East Antarctica: a comparison of computed balance fluxes and measured fluxes, 561–570
- Mass balance of the northeast sector of the Greenland ice sheet: a remote-sensing perspective, 265
- Measurement and parameterization of albedo variations at Haut Glacier d’Arolla, Switzerland, 675–688
- Relating glacier mass balance to meteorological data by using a seasonal sensitivity characteristic, 1–6
- Sensitivity of mass balance of five Swiss glaciers to temperature changes assessed by tuning a degree-day model, 7–14
- The use of bulk and profile methods for determining surface heat fluxes in the presence of glacier winds, 445
- Matsuoka, K., *see* Salamatin, A. N. and others
- Matthes Glacier (U.S.A.):
Radar glacier zones in southeast Alaska, U.S.A.: field and satellite observations, 287–296
- Mätzler, C., (paper: A simple snowpack/cloud reflectance and transmittance model from microwave to ultraviolet: the ice-lamella pack) 20–24
- Mawson Glacier (Antarctica):
Ice discharge of eastern Dome C drainage area, Antarctica, determined from airborne radar survey and satellite image analysis, 253
- Mayer, C. and M. J. Siegert, (paper: Numerical modelling of ice-sheet dynamics across the Vostok subglacial lake, central East Antarctica) 197–205
- Mayewski, P. A., *see* Kang Shichang and others
- Mérindol, L., *see* Durand, Y. and others
- Mickelson, D. M., *see* Cutler, P. M. and others
- Microwaves:
A simple snowpack/cloud reflectance and transmittance model from microwave to ultraviolet: the ice-lamella pack, 20–24
- Miles Glacier (U.S.A.):
Radar glacier zones in southeast Alaska, U.S.A.: field and satellite observations, 287–296
- Miller, H., *see* Hempel, L. and others
- Miller, M. M., *see* Ramage, J. M. and others
- Moraines: A model of ablation-dominated medial moraines and the generation of debris-mantled glacier snouts, 459–469
- Moran, M. L. and others, (paper: Delineation of a complexly dipping temperate glacier bed using short-pulse radar arrays) 274–286
- Moran, M. L. and others, (erratum: Delineation of a complexly dipping temperate glacier bed using short-pulse radar arrays) 707
- Morin, R. H. and others, (paper: Acoustic televiewer logging in glacier boreholes) 695–699
- Morteratschgletscher (Switzerland):
Analysis of a 3 year meteorological record from the ablation zone of Morteratschgletscher, Switzerland: energy and mass balance, 571–579
- Mosley-Thompson, E., *see* Zagorodnov, V. and others
- Murav’yev, Y. D., *see* Salamatin, A. N. and others
- Murray, T. and others, (paper: Englacial water distribution in a temperate glacier from surface and borehole radar velocity analysis) 389–398
- Murray, T., *see* Jiskoot, H. and others
- Näslund, J.-O. and others, (paper: Numerical modelling of the ice sheet in western Dronning Maud Land, East Antarctica: impacts of present, past and future climates) 54–66
- Näslund, J.-O. and others, (erratum: Numerical modelling of the ice sheet in western Dronning Maud Land, East Antarctica: impacts of present, past and future climates) 353–354
- Nepal:
Himalayan ice-core dating with snow algae, 335–340
- Nereson, N. A., (paper: Elevation of ice-stream margin scars after stagnation) 111–118
- Nereson, N. A. and others, (paper: The accumulation pattern across Siple Dome, West Antarctica, inferred from radar-detected internal layers) 75–87
- Nereson, N. A., *see* Jacobel, R. W. and others
- New Zealand: Relating glacier mass balance to meteorological data by using a seasonal sensitivity characteristic, 1–6
- Ng, F. S. L., (paper: Coupled ice–till deformation near subglacial channels and cavities) 580–598
- Ng, F. S. L., (review: Van der Veen, C. J. 1999. *Fundamentals of glacier dynamics.*) 700–701
- Nienow, P., *see* Hubbard, A. and others

- Nigardsbreen (Norway):
 Relating glacier mass balance to meteorological data by using a seasonal sensitivity characteristic, 1–6
- Nioghalfjærdsbrae (Greenland):
 A low-cost glacier-mapping system, 531–537
 Mass balance of the northeast sector of the Greenland ice sheet: a remote-sensing perspective, 265
- Norway:
 Contemporaneous, localized, basal ice-flow variations: implications for bedrock erosion and the origin of p-forms, 470–476
 Relating glacier mass balance to meteorological data by using a seasonal sensitivity characteristic, 1–6
 Rheology of ice at the bed of Engabreen, Norway, 611–621
 Sliding of ice past an obstacle at Engabreen, Norway, 599–610
- Nye, J. F., (paper: A flow model for the polar caps of Mars) 438–444
- Oerlemans, J. and B. K. Reichert, (paper: Relating glacier mass balance to meteorological data by using a seasonal sensitivity characteristic) 1–6
- Oerlemans, J., (paper: Analysis of a 3 year meteorological record from the ablation zone of Morteratschgletscher, Switzerland: energy and mass balance) 571–579
- Øksfjordjøkelen (Norway):
 Contemporaneous, localized, basal ice-flow variations: implications for bedrock erosion and the origin of p-forms, 470–476
- Oxygen isotopes:
 Monsoon and dust signals recorded in Dasuopu glacier, Tibetan Plateau, 222–226
- Parizek, B. B., *see* Cutler, P. M. and others
- Pasterzenkees (Austria):
 The use of bulk and profile methods for determining surface heat fluxes in the presence of glacier winds, 445
- Patterson, C. J., *see* Knight, P. G. and others
- Payne, A. J. and others, (paper: Results from the EISMINT model intercomparison: the effects of thermomechanical coupling) 227–238
- Permafrost:
 A numerical investigation of ice-lobe–permafrost interaction around the southern Laurentide ice sheet, 311
- Peyto Glacier (Canada):
 Relating glacier mass balance to meteorological data by using a seasonal sensitivity characteristic, 1–6
- Photogrammetry:
 Glacier mass-balance determination by remote sensing and high-resolution modelling, 491–498
- Pine Island Glacier (Antarctica):
 Calculating basal thermal zones beneath the Antarctic ice sheet, 297–310
- Plattalvagletscher (Switzerland):
 Sensitivity of mass balance of five Swiss glaciers to temperature changes assessed by tuning a degree-day model, 7–14
- Plogbreen (Antarctica):
 Numerical modelling of the ice sheet in western Dronning Maud Land, East Antarctica: impacts of present, past and future climates, 54–66
- Pradhan, P. M. S., *see* Trickett, Y. L. and others
- Pradhan, P. M. S., *see* Trickett, Y. L. and others
- Priestley Glacier (Antarctica):
 Ice discharge of eastern Dome C drainage area, Antarctica, determined from airborne radar survey and satellite image analysis, 253
- Qin Dahe, *see* Kang Shichang and others
- Rabus, B. T. and D. R. Fatland, (paper: Comparison of SAR-interferometric and surveyed velocities on a mountain glacier: Black Rapids Glacier, Alaska, U.S.A.) 119–128
- Radar:
 An analysis of balance velocities over the Greenland ice sheet and comparison with synthetic aperture radar interferometry, 67–74
 1.4 GHz radar penetration and evidence of drainage structures in temperate ice: Black Rapids Glacier, Alaska, U.S.A., 477–490
 Bed properties of Siple Dome and adjacent ice streams, West Antarctica, inferred from radio-echo sounding measurements, 88–94
 Changes in the margin of Ice Stream C, Antarctica, 102–110
 Radar glacier zones in southeast Alaska, U.S.A.: field and satellite observations, 287–296
 The accumulation pattern across Siple Dome, West Antarctica, inferred from radar-detected internal layers, 75–87
- Digital elevation models for the Lambert Glacier–Amery Ice Shelf system, East Antarctica, from ERS-1 satellite radar altimetry, 553–560
 Mass balance of the Lambert Glacier–Amery Ice Shelf system, East Antarctica: a comparison of computed balance fluxes and measured fluxes, 561–570
- Radio-echo sounding (RES):
 The accumulation pattern across Siple Dome, West Antarctica, inferred from radar-detected internal layers, 75–87
 Bed properties of Siple Dome and adjacent ice streams, West Antarctica, inferred from radio-echo sounding measurements, 88–94
 Changes in the margin of Ice Stream C, Antarctica, 102–110
 A comparison of radio-echo sounding data and electrical conductivity of the GRIP ice core, 369–374
 Delineation of a complexly dipping temperate glacier bed using short-pulse radar arrays, 274
 Geophysical investigations of ice-sheet internal layering and deformation in the Dome C region of central East Antarctica, 161–166
 Ice discharge of eastern Dome C drainage area, Antarctica, determined from airborne radar survey and satellite image analysis, 253
 A low-cost glacier-mapping system, 531–537
 Mass balance of the northeast sector of the Greenland ice sheet: a remote-sensing perspective, 265
 The onset area of Ice Stream D, West Antarctica, 95–101
- Radionuclides:
 Natural and anthropogenic levels of tritium in a Canadian Arctic ice core, Agassiz Ice Cap, Ellesmere Island, and comparison with other radionuclides, 35–40
- Ramage, J. M. and others, (paper: Radar glacier zones in southeast Alaska, U.S.A.: field and satellite observations) 287–296
- Raper, S. C. B. and others, (paper: A geometric glacier model for sea-level change calculations) 357–368
- Rasmussen, L. A. and others, (paper: The accumulation regime of Blue Glacier, U.S.A., 1914–96) 326–334
- Raymond, C. F., (paper: Energy balance of ice streams) 665–674
- Raymond, C. F., *see* Gades, A. M. and others
- Raymond, C. F., *see* Jacobel, R. W. and others
- Raymond, C. F., *see* Nereson, N. A. and others
- Raymond, C. F., *see* Thorsteinsson, T.
- Rea, B. R. and others, (paper: Contemporaneous, localized, basal ice-flow variations: implications for bedrock erosion and the origin of p-forms) 470–476
- Reeh, N., *see* Christensen, E. L. and others
- Reeves Glacier (Antarctica):
 Ice discharge of eastern Dome C drainage area, Antarctica, determined from airborne radar survey and satellite image analysis, 253
- Reflectance:
 A simple snowpack/cloud reflectance and transmittance model from microwave to ultraviolet: the ice-lamella pack, 20–24
- Reichert, B. K., *see* Oerlemans, J.
- Remote Sensing:
 Evaluation of remote-sensing techniques to measure decadal-scale changes of Hofsjökull ice cap, Iceland, 375–388
 Ice discharge of eastern Dome C drainage area, Antarctica, determined from airborne radar survey and satellite image analysis, 253
 Mass balance of the northeast sector of the Greenland ice sheet: a remote-sensing perspective, 265
- Reviews:
 Knight, Peter G. 1999. *Glaciers*, 346
 Petrenko, Victor F. and Robert W. Whitworth. 1999. *The physics of ice*, 347–348
 Post, A. and E. R. LaChapelle. 1999. *Glacier ice*, 700
 Van der Veen, C. J. 1999. *Fundamentals of glacier dynamics*, 700–701
- Reynaud, L., *see* Vincent, C. and others
- Rhonegletscher (Switzerland):
 Sensitivity of mass balance of five Swiss glaciers to temperature changes assessed by tuning a degree-day model, 7–14
- Rignot, E. and others, (paper: Mass balance of the northeast sector of the Greenland ice sheet: a remote-sensing perspective) 265–273
- Ritz, C., *see* Payne, A. J. and others
- Roberts, M. J. and others, (correspondence: Rapid sediment entrainment and englacial deposition during jökulhlaups) 349–351
- Robinson, Z. P., *see* Knight, P. G. and others
- Ross Ice Shelf (Antarctica):

- Calculating basal thermal zones beneath the Antarctic ice sheet, 297–310
A millennium of variable ice flow recorded by the Ross Ice Shelf, Antarctica, 652–664
- Russell Glacier (Greenland):
Glacier advance, ice-marginal lakes and the routing of meltwater and sediment: Russell Glacier, Greenland, 423–426
- Russell, A. J., *see* Roberts, M. J. and others
- Russia:
Modelling dynamics of glaciers in volcanic craters, 177–187
Portable system for intermediate-depth ice-core drilling, 167–172
- Rutford Ice Stream (Antarctica):
Energy balance of ice streams, 665–674
- Synthetic-aperture radar (SAR):
Comparison of SAR-interferometric and surveyed velocities on a mountain glacier: Black Rapids Glacier, Alaska, U.S.A., 119–128
An analysis of balance velocities over the Greenland ice sheet and comparison with synthetic aperture radar interferometry, 67–74
Radar glacier zones in southeast Alaska, U.S.A.: field and satellite observations, 287–296
- Saint Sorlin, Glacier de (France):
Dynamic behaviour analysis of glacier de Saint Sorlin, France, from 40 years of observations, 1957–97, 499–506
- Salamatin, A. N. and others, (paper: Modelling dynamics of glaciers in volcanic craters) 177–187
- Saurer, M., *see* Eichler, A. and others
- Scambos, T. A. and others, (paper: The link between climate warming and break-up of ice shelves in the Antarctic Peninsula) 516–530
- Scambos, T. A., *see* Fahnestock, M. A. and others
- Scambos, T. A., *see* Jacobel, R. W. and others
- Schmeltz, M., *see* Rignot, E. and others
- Schweizer, J., *see* Jamieson, J. B.
- Schwikowski, M., *see* Eichler, A. and others
- Sea ice: On the modelling of ice-thickness redistribution, 427–437
- Seko, K., *see* Yoshimura, Y. and others
- Serra, F., *see* Stenni, B. and others
- Sharp, M., *see* Hubbard, A. and others
- Sharp, M. J., *see* Brock, B. W. and others
- Shiraiwa, T., *see* Salamatin, A. N. and others
- Siegert, M. J. and J. L. Bamber, (correspondence: Subglacial water at the heads of Antarctic ice-stream tributaries) 702–703
- Siegert, M. J., *see* Hodgkins, R. and others
- Siegert, M. J., *see* Kwok, R. and others
- Siegert, M. J., *see* Mayer, C.
- Sigurðsson, O., *see* Hall, D. K. and others
- Silvrettagletscher (Switzerland):
Sensitivity of mass balance of five Swiss glaciers to temperature changes assessed by tuning a degree-day model, 7–14
- Siple Dome (Antarctica):
Bed properties of Siple Dome and adjacent ice streams, West Antarctica, inferred from radio-echo sounding measurements, 88–94
Changes in the margin of Ice Stream C, Antarctica, 102–110
Elevation of ice-stream margin scars after stagnation, 111–118
The accumulation pattern across Siple Dome, West Antarctica, inferred from radar-detected internal layers, 75–87
The onset area of Ice Stream D, West Antarctica, 95–101
- Skeiðarárjökull (Iceland):
Rapid sediment entrainment and englacial deposition during jökulhlaups, 349–351
Glacioluvial crevasse and conduit fills as indicators of supraglacial dewatering during a surge, Skeiðarárjökull, Iceland, 25–34
- Skou, N., *see* Christensen, E. L. and others
- Smith, L. C., *see* Hall, D. K. and others
- Snow:
Formaldehyde and peroxide concentrations in Law Dome (Antarctica) firn and ice cores, 15–19
A simple snowpack/cloud reflectance and transmittance model from microwave to ultraviolet: the ice-lamella pack, 20–24
Himalayan ice-core dating with snow algae, 335–340
- Short fluctuations in the density and thickness of a dry firn column, 399–411
Texture and strength changes of buried surface-hoar layers with implications for dry snow-slab avalanche release, 151–160
- Spiro, B., *see* Hubbard, B. and others
- Stauffer, B., *see* Tschumi, J.
- Stenni, B. and others, (paper: Snow accumulation rates in northern Victoria Land, Antarctica, by firn-core analysis) 541–552
- Stikine Icefield (Canada):
Radar glacier zones in southeast Alaska, U.S.A.: field and satellite observations, 287–296
- Stuart, G. W., *see* Murray, T. and others
- Surging glaciers:
Controls on the distribution of surge-type glaciers in Svalbard, 412–422
Glacioluvial crevasse and conduit fills as indicators of supraglacial dewatering during a surge, Skeiðarárjökull, Iceland, 25–34
- Svalbard:
Contemporaneous, localized, basal ice-flow variations: implications for bedrock erosion and the origin of p-forms, 470–476
Controls on the distribution of surge-type glaciers in Svalbard, 412–422
- Switzerland:
Analysis of a 3 year meteorological record from the ablation zone of Morteratschgletscher, Switzerland: energy and mass balance, 571–579
Glacier mass-balance determination by remote sensing and high-resolution modelling, 491–498
Glaciochemical dating of an ice core from upper Grenzgletscher (4200 m a.s.l.), 507–515
Ice-core evidence of the thickness and character of clear-facies basal ice: Glacier de Transfleuron, Switzerland, 140–150
Measurement and parameterization of albedo variations at Haut Glacier d'Arolla, Switzerland, 675–688
Sensitivity of mass balance of five Swiss glaciers to temperature changes assessed by tuning a degree-day model, 7–14
- Synal, H.-A., *see* Eichler, A. and others
- Tabacco, I. E., *see* Frezzotti, M. and others
- Takeuchi, N., *see* Yoshimura, Y. and others
- Taku Glacier (U.S.A.):
Radar glacier zones in southeast Alaska, U.S.A.: field and satellite observations, 287–296
- Talos Dome (Antarctica):
Ice discharge of eastern Dome C drainage area, Antarctica, determined from airborne radar survey and satellite image analysis, 253
- Tarasov, L., *see* Payne, A. J. and others
- Taylor Dome (Antarctica):
Ice discharge of eastern Dome C drainage area, Antarctica, determined from airborne radar survey and satellite image analysis, 253
- Thomassen, M. P. A., *see* Payne, A. J. and others
- Thompson, L. G., *see* Zagorodnov, V. and others
- Thorsteinsson, T. and C. F. Raymond, (paper: Sliding versus till deformation in the fast motion of an ice stream over a viscous till) 633–640
- Thwaites Glacier (Antarctica):
Calculating basal thermal zones beneath the Antarctic ice sheet, 297–310
- Thyssen, F., *see* Hempel, L. and others
- Tibetan Plateau:
Effect of summer accumulation on glacier mass balance on the Tibetan Plateau revealed by mass-balance model, 244–252
Monsoon and dust signals recorded in Dasuopu glacier, Tibetan Plateau, 222–226
- Till:
Coupled ice–till deformation near subglacial channels and cavities, 580–598
Diffusive mixing between shearing granular layers: constraints on bed deformation from till contacts, 641–651
Glacier motion dominated by processes deep in underlying till, 213–221
Sliding versus till deformation in the fast motion of an ice stream over a viscous till, 633–640
- Tison, J.-L., *see* Hubbard, B. and others
- Transfleuron, Glacier de (Switzerland):
Ice-core evidence of the thickness and character of clear-facies basal ice: Glacier de Transfleuron, Switzerland, 140–150
- Transmittance:
A simple snowpack/cloud reflectance and transmittance model from microwave to ultraviolet: the ice-lamella pack, 20–24

- Trapridge Glacier (Canada):
Evidence for extreme pressure pulses in the subglacial water system, 206–212
- Traversi, R., *see* Stenni, B. and others
- Trickett, Y. L. and others, (paper: The effects of sulfuric acid on the mechanical properties of ice single crystals) 239–243
- Trickett, Y. L. and others, (paper: The orientation dependence of the strength of ice single crystals) 41–44
- Tritium:
Natural and anthropogenic levels of tritium in a Canadian Arctic ice core, Agassiz Ice Cap, Ellesmere Island, and comparison with other radio-nuclides, 35–40
- Truffer, M. and others, (paper: Glacier motion dominated by processes deep in underlying till) 213–221
- Tschumi, J. and B. Stauffer, (paper: Reconstructing past atmospheric CO₂-concentration based on ice-core analyses: open questions due to in situ production of CO₂ in ice) 45–53
- Tweed, F. S., *see* Roberts, M. J. and others
- U.S.A.:
- Comparison of SAR-interferometric and surveyed velocities on a mountain glacier: Black Rapids Glacier, Alaska, U.S.A., 119–128
 - Glacier motion dominated by processes deep in underlying till, 213–221
 - Radar glacier zones in southeast Alaska, U.S.A.: field and satellite observations, 287–296
 - The accumulation regime of Blue Glacier, U.S.A., 1914–96, 326–334
 - 1.4 GHz radar penetration and evidence of drainage structures in temperate ice: Black Rapids Glacier, Alaska, U.S.A., 477–490
 - A model of ablation-dominated medial moraines and the generation of debris-mantled glacier snouts, 459–469
 - Diffusive mixing between shearing granular layers: constraints on bed deformation from till contacts, 641–651
 - Delineation of a complexly dipping temperate glacier bed using short-pulse radar arrays, 274
 - A numerical investigation of ice-lobe–permafrost interaction around the southern Laurentide ice sheet, 311–325
- Udisti, R., *see* Stenni, B. and others
- Vallon, M., *see* Vincent, C. and others
- Van Ommen, T. D., *see* Gillett, R. W. and others
- Veststraumen (Antarctica):
Numerical modelling of the ice sheet in western Dronning Maud Land, East Antarctica: impacts of present, past and future climates, 54–66
- Vincent, C. and others, (paper: Dynamic behaviour analysis of glacier de Saint Sorlin, France, from 40 years of observations, 1957–97) 499–506
- Vornberger, P., *see* Bindshadler, R. and others
- Vostok (Antarctica):
Ice motion over Lake Vostok, Antarctica: constraints on inferences regarding the accreted ice, 689–694
Numerical modelling of ice-sheet dynamics across the Vostok subglacial lake, central East Antarctica, 197–205
- Waddington, E. D., *see* Nereson, N. A. and others
- Wake, C. P., *see* Kang Shichang and others
- Waller, R. I., *see* Bennett, M. R. and others
- Waller, R. I., *see* Knight, P. G. and others
- Warner, R. C., *see* Fricker, H. A. and others
- Weber, J. E., (paper: Non-temperate glacier flow over wavy sloping ground) 453–458
- Whalley, W. B., *see* Rea, B. R. and others
- White Glacier (Canada):
Relating glacier mass balance to meteorological data by using a seasonal sensitivity characteristic, 1–6
- Wilch, E. and T. J. Hughes, (paper: Calculating basal thermal zones beneath the Antarctic ice sheet) 297–310
- Wilén, L. A., (paper: A new technique for ice-fabric analysis) 129–139
- Wilkins Ice Shelf (Antarctica):
The link between climate warming and break-up of ice shelves in the Antarctic Peninsula, 516–530
- Williams, R. S., Jr, *see* Hall, D. K. and others
- Willis, I., *see* Hubbard, A. and others
- Willis, I. C., *see* Brock, B. W. and others
- Wingham, D. J., (paper: Short fluctuations in the density and thickness of a dry firn column) 399–411
- Woelders, K., *see* Christensen, E. L. and others
- Wordie Ice Shelf (Antarctica):
The link between climate warming and break-up of ice shelves in the Antarctic Peninsula, 516–530
- Workman, W., *see* Kotzer, T. G. and others
- Wortmanns Glacier (Ak):
A model of ablation-dominated medial moraines and the generation of debris-mantled glacier snouts, 459–469
- Yala Glacier (Nepal):
Himalayan ice-core dating with snow algae, 335–340
- Yankielun, N. E., *see* Arcone, S. A.
- Yao Tandong, *see* Kang Shichang and others
- Yoshimura, Y. and others, (paper: Himalayan ice-core dating with snow algae) 335–340
- Young, N. W., *see* Fricker, H. A. and others
- Zachariae Isstrøm (Greenland):
Mass balance of the northeast sector of the Greenland ice sheet: a remote-sensing perspective, 265
- Zagorodnov, V. and others, (paper: Portable system for intermediate-depth ice-core drilling) 167–172
- Zhang, Y., *see* Braithwaite, R. J.
- Zheng, J., *see* Kotzer, T. G. and others
- Zirizzotti, A., *see* Frezzotti, M. and others