The Journal of Laryngology and Otology

EDITED BY G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

Contents

Some Facets of Hypopharyngeal Surgery · John F. Simpson
TREATMENT OF CANCER OF THE PHARYNX AND CERVICAL OESOPHAGUS H. D. FAIRMAN and H. T. John
Parosmia · · · · · · Kaj Zilstorff
MATERNAL DIABETES AND THE EARS OF NEWBORNS: HISTOPATHOLOGY NILS H. BUCH and M. Balslev Jørgensen
OBLITERATIVE OTOSCLEROSIS: AN ANALYSIS OF THE CLINICAL AND AUDIOMETRIC FINDINGS . R. E. GRISTWOOD
METHODS OF OBJECTIVE AUDIOMETRY · · · H. A. BEAGLEY and J. J. KNIGHT
A Note on the Use of Self-Recording Audiometry with Children M. E. Delany, L. S. Whittle and E. C. Knox
MEMBRANE PROBLEMS OF THE INNER EAR AND THEIR SIGNIFICANCE S. RAUCH
THE APPLICATION OF MODERN VESTIBULOMETRY TO THE PROBLEM OF VERTIGO T. J. WILMOT
CLINICAL RECORDS—
Tracheo-Oesophageal-Skin Fistula • • Maxwell Ellis
A Case of Malignant Melanoma of the Larynx · · · · · · · Christian Nsamba
GLIOMA OF THE NASOPHARYNX • • J. F. BIRRELL, J. TEVENDALE and A. D. BAIN

GENERAL NOTES

London

Headley Brothers

109 Kingsway WC2

Annual Subscription £5/5/0 net, U.S.A. \$15

Monthly, 12/6 net post free

The Journal of Laryngology and Otology

(Founded in 1887 by Morell Mackenzie and Norris Wolfenden)

EDITED BY

G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

1. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the understanding that they are contributed to this Journal solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgment must be made.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and

sent at the same time as original photographs, etc.

The Harvard system of recording references should be used, e.g. Green, C., and Brown, D. (1951) J. Laryng., 65, 33. Abbreviations of Journals should follow the style recommended in World Medical Periodicals, published by World Health Organization, 1952.

It is most important that authors should verify personally the accuracy of

every reference before submitting a paper for publication.

- 3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.
- 4. Illustration blocks will normally be held by the Printers for three years, after which they will be destroyed. Any author is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to Headley Brothers, 109 Kingsway, London, WC2. Coloured illustrations will be charged to authors, unless a special grant is authorized by the Editor.
- 5. Orders for reprints must be sent when returning galley proofs, and for this purpose special forms are supplied.
- Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the Journal of Laryngology. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.
- 7. Editorial communications may be addressed to THE EDITOR, Journal of Laryngology, c/o Headley Brothers, 109 Kingsway, London, WC2.
- 8. The annual subscription is five guineas sterling (U.S.A. \$15) post free, and payable in advance.
- 9. Single copies will be on sale at 12s. 6d. each; copies of parts up to Vol. LXIII which are available may be purchased at 7s. 6d. each.
- 10. All subscriptions, advertising and business communications should be sent to the publishers, Headley Brothers, 109 Kingsway, London, WC2.


United States of America

Orders for this Journal may be sent through local booksellers, or to STECHERT-HAFNER, INC., 31-33 East 10th Street, New York, or direct to the publishers, Headley Brothers, 109 Kingsway, London, WC2, England.

Dournal of Laryngology and Otology, 1966

CONTENTS

	PAGE
Some Facets of Hypopharyngeal Surgery. • John F. Simpson (London)	1077
TREATMENT OF CANCER OF THE PHARYNX AND CERVICAL OESOPHAGUS. H. D. Fairman (Bristol) and H. T. John (Bath)	
Parosmia. Kaj Zilstorff (Copenhagen) · · · · · · · ·	1102
Maternal Diabetes and the Ears of Newborns: Histopathology. Nils H. Buch and M. Balslev Jørgensen (Copenhagen)	1105
Obliterative Otosclerosis: An Analysis of the Clinical and Audio- metric Findings. R. E. Gristwood (Adelaide, South Australia)	1115
METHODS OF OBJECTIVE AUDIOMETRY. H. A. Beagley and J. J. Knight (London)	
A Note on the Use of Self-Recording Audiometry with Children. M. E. Delany and L. S. Whittle (London) and E. C. Knox (Glasgow)	
MEMBRANE PROBLEMS OF THE INNER EAR AND THEIR SIGNIFICANCE. S. Rauch (Düsseldorf) · · · · · · · · · · · · · · · · · · ·	1144
The Application of Modern Vestibulometry to the Problem of Vertigo. T. J. Wilmot (Omagh) · · · · · · · · · · · · · · · · · · ·	1156
Clinical Records—	
Tracheo-Oesophageal-Skin Fistula. Maxwell Ellis (London) ·	1173
A Case of Malignant Melanoma of the Larynx. Christian Nsamba (Portsmouth) · · · · · · · · · · · · · · · · · · ·	1178
GLIOMA OF THE NASOPHARYNX. J. F. Birrell, J. Tevendale and A. D. Bain (Edinburgh)	1182
General Notes · · · · · · · · · · · · · · · · · · ·	1187



beam-splitter for observation and photography during surgery

This new beam-splitter triples the versatility of the Zeiss Operating Microscope.

As illustrated it allows simultaneous binocular viewing by the surgeon, observation by an assistant and 35mm photography. All 3 images are identical, upright and unreversed. A second observation tube can replace the camera to permit 2 assistants or students to observe the operation. Alternatively, 2 cameras can be attached to provide stereo-photography or differently magnified photographic records. The observation tubes are adjustable to any angle for comfortable viewing without the observer impeding the surgeon. The Zeiss Beam Splitter is also suitable for use with the Zeiss Slit lamp, Photo Slit lamp and certain other optical instruments. Write for full data and demonstration to the sole UK importers Degenhardt & Co Ltd Carl Zeiss House 20/22 Mortimer St London W1 Museum 8050 (15 lines) Scottish Enquiries to Degenhardt & Co (Scotland) Ltd 22 Burnbank St Glasgow NW Douglas 0520 Z S239

egenhardt British Agents for Carl Zeiss, West Germany

'Otosporin' for ears and ears and ears...

'Otosporin' drops rapidly destroy virtually all organisms commonly found in infections of the ear, reduce irritation, inflammation and pain.

'Otosporin' brand Drops

contain polymyxin B sulphate, neomycin sulphate and hydrocortisone. Issued in bottles of 5 ml.







MIDDLE EAR PROSTHESES




Sultana's Stainless Steel Spring Middle Ear Prosthesis, 4 mm. long

Vide Journal of Laryngology Vol. LXXIX, No. 9



Smyth's "Clothes-Pin" Incus replacement Prosthesis





Shepard's Teflon Grommet Drain Tubes

Further details of these and other Middle Ear Prostheses on request

DOWN BROS. AND MAYER & PHELPS LTD CHURCH PATH, MITCHAM, SURREY, ENGLAND

In Canada

410 Dundas Street West, Toronto, 2B, Canada

WHEN A HEARING AID IS RECOMMENDED ONLY AMPLIVOX OFFERS THIS HEARING AID SERVICE

The design and manufacture of a complete range with individual fitting facilities.

Best of other makes in addition to Amplivox aids—Amplivox hearing aids are ONLY available from Amplivox.

Highest degree of training of any hearing aid organisation.

Master Hearing Aid evaluations.

Expert follow-up and rehabilitation advice.

15 branch offices with while-you-wait service.

FOR YOUR PATIENTS' GREATEST SATISFACTION SPECIFY AMPLIVOX

AMPLIVOX

World's most experienced and complete hearing service

AMPLIVOX HEARING ADVISORY SERVICE 80 NEW BOND STREET LONDON, W.1



for the girl who has everything...(even a stuffy nose)

Otrivine provides gentle, long-lasting relief from the misery of nasal congestion. One application maintains a clear nasal airway for up to eight hours, without subsequent rebound hyperaemia. Suitable for adults and children.

Nasal spray (0.1%) $\frac{1}{2}$ f1. oz, drops (0.1%) $\frac{1}{2}$ f1. oz, and paediatric drops (0.05%) 10 m1. (in special safety dropper-bottle)

Otrivine volometazoline hydrochloride





.CIBA

ONLY AMPLIVOX

provides a complete audiometric service!



Transistor Audiometers

with air conduction or air and bone conduction with masking, for hospitals, schools, consulting rooms, industrial medical departments, screening programmes, etc.




Clinical Audiometers

with narrow band masking for the highest accuracy and extended resolution.

Audiometer Booths

for ideal testing conditions in hospitals, clinics, factories. Standard and special sizes available.





PLUS An efficient and speedy calibration service

please write for technical literature to meet your needs

AMPLIVOX

HEARING ADVISORY SERVICE LTD.

80 New Bond Street, London, W.1. Tel: HYDe Park 9888

THE LARYNGOSCOPE

A Monthly Journal devoted to the disease of EAR, NOSE AND THROAT

Official organ for the American Laryngological Rhinological and Otological Society

Price \$18.00 per year Canada \$19.00 per year Foreign \$19.00 per year

ESTABLISHED 1896

THEODORE E. WALSH, M.D. EDITOR

517 SOUTH EUCLID AVENUE SAINT LOUIS MO. 63110.





PROFESSIONAL MODEL

AUDITORY TRAINING UNIT

- Robust Construction—high grade components
- ★ Portable, Battery Operated
- ★ Modern transistor circuitry
- ★ Versatile with many facilities
- ★ Improved Performance
- ★ Treble and Bass Tone Controls
- ★ Loop Drive
- * Reasonably priced—£55 Ex. Works

AN R·N·I·D-TESTED AID Nº 6449

Full report available from

The Royal National Institute for the Deaf-105 Gower St.,
London W.C.1.


For further details please write to:

G. C. JOHNSON, LEYTON INSTRUMENT CO. LTD.,


54 ALBERT ROAD, CAVERSHAM,

READING, BERKS.

Telephone: READING 72293



BÉKÉSY ACCESSORY



For the very first time a fully automatic unit has been made available for a Clinic Audiometer, making the combined units into a true BÉKÉSY Audiometer.

This automatic accessory is being made for the PETERS SPD/5 CLINIC AUDIOMETER, which has a continuous frequency range and continuously variable attenuation. The result is therefore a true Békésy audiometer.

It can be attached to or detached from the audiometer in under one minute so that the two units together offer the facilities of an advanced clinic audiometer or Békésy audiometry at will.

It can be fitted to any existing SPD/5 Audiometer.

Please write for full details and demonstration to:

ALFRED PETERS & SONS LIMITED
51 GELL STREET, SHEFFIELD 3

ORBENIN and its success in Gram-Positive Infections

Otitis media Otitis externa

Boils Carbuncles Infected dermatoses

Pneumonia Lung abscess Acute bronchitis

Osteomyelitis Osteitis Tonsillitis Pharyngitis

Post-operative wound infections Burns Skin graft protection

Septicaemia Acute endocarditis

Staphylococcal enterocolitis

Staphylococcal urinary tract infections

Staphylococcal meningitis

ORBENIN (cloxacillin sodium B.P.) is a penicillin active against virtually all Gram-positive organisms, including staphylococci, β-haemolytic streptococci and pneumococci, irrespective of their resistance to other antibiotics.

It is encouraging that the widespread use of Orbenin in hospitals has not resulted in the emergence of resistant strains.


More than 300 references in the World Literature testify to the excellent clinical results obtained with Orbenin in a wide variety of Gram-positive infections.

Detailed literature is available on request.

Dosage:

Two capsules (500mg.) or one vial (250mg.) q.i.d.

Availability:






Orbenin (cloxacillin sodium B.P.)(regd.) is a product of British research at Beecham Research Laboratories

Brentford, England Telephone: ISLeworth 4111
Published online by Cambridge University Press





—that is the triumph of the **MACKIE** hearing aid

The new Mackie post-aural hearing aid is made in several different circuit versions so that the gain and frequency response can be selected to match the patient's needs; and it is available where required with a unique form of automatic volume control which has proved highly successful in cases of recruitment. The forward pick-up microphone gives maximum

intelligibility of speech without background interference.


It is exported to many countries including the United States of America—a tribute to its high quality and advanced design.

In Great Britain it is available for trial by your patients from *independent* hearing aid dealers in all parts of the country.

MANUFACTURED BY A. G. R. MACKIE LTD. A member of the Bonochord Group of Companies FOR FURTHER DETAILS WRITE TO 48 WELBECK STREET, LONDON WI

KAMPLEX AUDIOMETRIC EQUIPMENT

AUDIOMETERS



SCREENING PORTABLE TA16 DIAGNOSTIC PORTABLE TA15

CLINIC AUDIOMETER
DOUBLE CHANNEL

DA1 DA2


AUTOMATIC AUDIOMETER
AB1

CLINIC AUDIOMETER DA1

ACOUSTIC IMPEDANCE APPARATUS ZA2

PNEUMOMETER ZP1

SKIN RESISTANCE APPARATUS RA4



SOUND LEVEL METER GA6
VISIBLE SPEECH APPARATUS
AUDITORY TRAINERS

Write for full catalogue and detailed descriptions to:

ALFRED PETERS & SONS LTD 51 GELL STREET, SHEFFIELD 3



the antistaphylococcal antibiotic that penetrates tissues at these levels...

bears the FUCIDIN label



The clinical significance of these levels has been confirmed in many publications during the past five years. Over 100 authors have endorsed the value of Fucidin in the treatment of staphylococcal Haves Middlesex

infection. Fucidin (Sodium Fusidate B.P.) 250 mgm capsules.

Leo Laboratories Limited


all clear


in sinusitis, otitis media and coryza

'Triogesic' provides decongestion with analgesia.

	TABLETS	ELIXIR
Revised Formula	scored (per tablet) Phenylpropanolamine-HCl 12.5 mg. Paracetamol B.P. 500 mg.	cherry flavoured (per 5 ml.) Phenylpropanolamine-HCl 3 mg. Paracetamol B.P. 125 mg. Alcohol (95%) 0.5 ml.
Packs	tubes of 30	bottles of 8 and 20 fl. oz.
Basic N.H.S. Price	3/6d. for 30 tablets	5/10d. for 8 fl. oz.

Literature and sample on request. A. Wander Ltd., 42 Upper Grosvenor Street, London W.1





Ceporin (cephaloridine) What makes this antibiotic so valuable in the E.N.T. unit?

Spectrum?

Exceptionally broad. It covers most gram-positive organisms—including the penicillin-resistant staphylococci—and many gram-negative organisms.

Action?

Intensely bactericidal. It gives a rapid kill of bacteria at or near the minimum inhibitory concentration, especially in the case of gram-negative organisms.

Safety?

Wide clinical usage has confirmed the laboratory evidence that Ceporin has a remarkably low toxicity—and it is usually well tolerated by patients allergic to penicillin.

Clinical record?

An analysis of results from over 1500 cases has recently shown high success rates in a wide variety of infections. Ear, nose and throat infections were included in a group of 87 soft tissue infections which showed an 85% satisfactory response. Many reports mention the rapid clinical response obtained with

Ceporin after other antibiotics had failed. Conditions which respond well to Ceporin include: mastoiditis, otitis media, sinusitis, quinsy, pharyngeal abscess, and intracranial infections secondary to infected sinuses and mastoiditis. Because of its three main attributesbroad spectrum, bactericidal action and low toxicity—Ceporin has proved valuable for use in place of more toxic drugs or combinations of antibiotics. It also has special value when a quick clinical response is essential in infections caused by penicillin-resistant staphylococci and in the penicillinhypersensitive patient.

Ceporin is issued in vials containing 250 mg, 500 mg and 1 gram of cephaloridine, packed singly and in boxes of five.

Detailed literature is available from Glaxo Laboratories Limited, Greenford, Middlesex.

Ceporin is a Glaxo trade mark