

## REFERENCES

1. Gaston MA. *Enterobacter*: an emerging nosocomial pathogen. *J Hosp Infect* 1988;11:197-208.
2. Flynn DM, Weinstein RA, Nathan C, Gaston MA, Kabins SA. Patients' endogenous flora as the source of 'nosocomial' *Enterobacter* in cardiac surgery. *J Infect Dis* 1987;156:363-368.
3. Griffith BP, Kormos RL, Hardesty RL, Armitage JM, Dummer JS. The artificial heart: infection-related morbidity and its effect on transplantation. *Ann Thorac Surg* 1988;45:409-414.
4. de Champs C, Guelon D, Joyon D, Sirot D, Chanal M, Sirot J. Treatment of a meningitis due to *Enterobacter aerogenes* producing a derepressed cephalosporinase and a *Klebsiella pneumoniae* producing an extended-spectrum beta-lactamase. *Infection* 1991;19:181-183.
5. Mellencamp MA, Roccaforte JS, Preheim LC, Sanders CC, Anene CA, Bittner MJ. Isolation of *Enterobacter aerogenes* susceptible to beta-lactam antibiotics despite high level beta-lactamase production. *Eur J Clin Microbiol Infect Dis* 1990;9:827-830.
6. de Champs C, Henquell C, Guelon D, Sirot D, Gazuy N, Sirot J. Clinical and bacteriological study of nosocomial infections due to *Enterobacter aerogenes* resistant to imipenem. *J Clin Microbiol* 1993;31:123-127.
7. Jaroux E, Collet L, Bellon O, et al. Emergence of *Enterobacter aerogenes* strains with an extended-spectrum betalactamase in Provence. 3rd International Conference on the Prevention of Infection, Nice, April 4-5, 1994, abstract MCB6.
8. Markowitz SM, Smith SM, Williams DS. Retrospective analysis of plasmid patterns in a study of bum unit outbreaks of infection due to *Enterobacter cloacae*. *J Infect Dis* 1983;148:18-23.
9. Haertl R, Bandlow G. Epidemiological fingerprinting of *Enterobacter cloacae* by small-fragment restriction endonuclease analysis and pulsed-field gel electrophoresis of genomic restriction fragments. *J Clin Microbiol* 1993;31:128-133.
10. Grattard F, Pozzetto B, Berthelot PH, et al. Arbitrarily primed PCR, ribotyping, and plasmid pattern analysis applied to investigation of a nosocomial outbreak due to *Enterobacter cloacae* in a neonatal intensive care unit. *J Clin Microbiol* 1994;32:596-602.
11. Garaizar J, Kaufmann ME, Pitt TL. Comparison of ribotyping with conventional methods for the type identification of *Enterobacter cloacae*. *J Clin Microbiol* 1991;29:1303-1307.
12. Poilane I, Cruaud P, Lachassinne E, et al. *Enterobacter cloacae* cross-colonization in neonates demonstrated by ribotyping. *Eur J Clin Microbiol Infect Dis* 1993;12:820-826.
13. Gaston MA, Strickland MA, Ayling-Smith BA, Pitt TL. Epidemiological typing of *Enterobacter aerogenes*. *J Clin Microbiol* 1989;27:564-565.
14. Grattard F, Pozzetto B, Ros A, Gaudin OG. Differentiation of *Pseudomonas aeruginosa* strains by ribotyping: high discriminatory power by using a single restriction endonuclease. *J Med Microbiol* 1994;40:275-282.
15. Williams JGK, Kubelick AR, Livak KJ, Rafalski JA, Tingey SV. DNA polymorphisms amplified by arbitrary primers are useful genetic markers. *Nucleic Acids Res* 1990;18:6531-6535.
16. Van Belkum A, Bax R, Peerbooms P, Goessens WHF, Van Leeuwen N, Quint WGV. Comparison of phage typing and DNA fingerprinting by polymerase chain reaction for discrimination of methicillin-resistant *Staphylococcus aureus* strains. *J Clin Microbiol* 1993;31:798-803.
17. MacPherson JM, Eckstein PE, Scoles GJ, Gajadhar AA. Variability of random amplified polymorphic DNA assay among thermal cyclers, and effects of primer and DNA concentration. *Mol Cell Probes* 1993;7:293-299.
18. Meunier JR, Grimont PAD. Factors affecting reproducibility of random amplified polymorphic DNA fingerprinting. *Res Microbiol* 1993;144:373-379.

## Transmission of Pertussis from Adult to Infant

by Gina Pugliese, RN, MS  
Medical News Editor

During 1993, a total of 6,586 pertussis cases were reported in the United States, including 675 cases (10%) among persons aged >19 years. However, the total number of cases probably was substantially higher, because only an estimated 10% of all pertussis cases are reported. Underreporting is greater among adults, who often have only a mild cough. Classic symptoms of pertussis include paroxysms of cough, posttussive whoop, and apnea.

The Centers for Disease Control and Prevention (CDC) recently reported pertussis transmission from an adult resident of Massachusetts who was visiting her four-month-old susceptible nephew in Michigan. The child had received only one dose of diphtheria and tetanus toxoids and

pertussis vaccine (DTP) prior to infection. The adult aunt had a cough that worsened upon her return to Massachusetts, and she developed paroxysms of cough with inspiratory whoop and posttussive apnea. A diagnosis of pertussis was considered only after learning of her history of close contact with her nephew, who developed culture-confirmed pertussis three weeks after her visit. A 14-day course of erythromycin was prescribed for both patients and their household contacts.

The CDC recommends that healthcare providers consider pertussis in the differential diagnosis of acute cough of >7 days duration in adults, particularly if the cough is paroxysmal and associated with posttussive vomiting and/or whooping.

Transmission of pertussis can be reduced with prompt diagnosis and treatment of cases and early

administration of chemoprophylaxis to close contacts. Administration of erythromycin for 14 days is recommended for persons with pertussis and all their household contacts (regardless of age or vaccination status) and is an important method of protecting children aged <6 months, who are too young to have received the initial three-dose series of DTP recommended during infancy. Chemoprophylaxis of household contacts with erythromycin should be administered as soon as possible after first contact with a primary case; chemoprophylaxis administered >21 days after first contact is considered of limited value.

FROM: CDC. Transmission of pertussis from adult to infant-Michigan, 1993. *MMWR* 1995;44(4):75-76.