



Philadelphia



The Society for
Healthcare
Epidemiology of
America



APRIL 17-20, 2004

14TH ANNUAL SCIENTIFIC MEETING
WYNDHAM FRANKLIN PLAZA HOTEL
PHILADELPHIA, PENNSYLVANIA

www.shea-online.org



0.67-mL Sepp®

For procedures such as:

- Injections
- Peripheral IVs
- Pediatrics

1.5-mL Frepp®

For procedures such as:

- Blood cultures
- Blood donations
- Peripheral IVs

3.0-mL

For procedures such as:

- Catheter line insertions
- Dialysis
- Site maintenance

10.5-mL

For procedures such as:

- Cardiac cath
- Interventional radiology
- Labor and delivery

One hospital One antiseptic Four applicators Chloraprep®

Chlorhexidine Gluconate (CHG) 2% w/v & Isopropyl Alcohol (IPA) 70% v/v

- **Fast** — Apply for 30 seconds, dry for 30 seconds
- **Long-lasting** — Persistent up to 48 hours
- **Effective** — 2% chlorhexidine is preferred by the CDC*
- **Easy to use** — one-step, back and forth application
- **Durable** — works in the presence of blood and organic matter
- **Clear solution** — allows for unobstructed site visualization

* See "Guidelines for the Prevention of Intravascular Catheter-Related Infections," MMWR 2002

Available in cartons of single-use sterile applicators



Medi-Flex, Inc.

1-800-523-0502 • www.chloraprep.com

©2003 Medi-Flex Hospital Products, Inc.