Vol. LXXXI No. 3

March 1967

The Journal of Laryngology and Otology

EDITED BY G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

Contents

SIR VICTOR NEGUS NASO-ALVEOLAR CYSTS: AETIOLOGY, PRESENTATION AND TREATMENT: SIX CASES ARE REPORTED. TRANSZYGOMATIC APPROACH TO THE PETROUS APEX WITH A RECORD OF THREE CASES . . . THE SOCIAL DEVELOPMENT OF DEAF CHILDREN . THE PLACE OF MYRINGOTOMY IN THE MANAGEMENT OF SECRETORY OTITIS MEDIA IN CHILDREN . SEVERE REDUCTION OF HEARING AND VESTIBULAR INVOLVEMENT CAUSED BY VIONACTAN® . NARROW FIELD MASTOIDECTOMY IN TYMPANOPLASTY. ON THE TREATMENT OF PAROTID FISTULAE . . MYRINGOPLASTY BY HOMOGENOUS VEIN GRAFT .

CLINICAL RECORDS-

MULTIPLE NEUROFIBROMATOSIS INVOLVING THE VIIITH NERVE

CONGENITAL UNILATERAL CHOANAL ATRESIA Associated with a Rhinolith.

MEMBRANOUS TOTAL CLOSURE OF THE EXTERNAL AUDITORY CANAL

GENERAL NOTES

London Headley Brothers

109 Kingsway WC2

Annual Subscription £5/5/0 net, U.S.A. \$15

G. P. WALSH-WARING

Lewis Labib Samy and Iskander H. Girgis Edmund Critchley

J. C. ARCHARD

Hans Jelert J. Nowinski A. C. Arulpragasam J. F. O. Mitchell

> H. A. BEAGLEY and J. J. KNIGHT

HELIO MARTINS and JAIME T. BENITEZ

DAOUD S. ALI

A. PALVA and S. SUURKARI

Montbly, 12/6 net post free

https://doi.org/10.1017/S0022215100067049 Published online by Cambridge University Press

ADVERTISEMENTS

The Journal of Laryngology and Otology

(Founded in 1887 by Morell Mackenzie and Norris Wolfenden)

EDITED BY

G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

1. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the understanding that they are contributed to this *Journal* solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgment must be made.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and sent at the same time as original photographs, etc.

The Harvard system of recording references should be used, e.g. GREEN, C., and BROWN, D. (1951) J. Laryng., 65, 33. Abbreviations of Journals should follow the style recommended in *World Medical Periodicals*, published by World Health Organization, 1952.

It is most important that authors should verify *personally* the accuracy of every reference before submitting a paper for publication.

3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.

4. Illustration blocks will normally be held by the Printers for three years, after which they will be destroyed. Any author is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to HEADLEY BROTHERS, 109 Kingsway, London, WC2. Coloured illustrations will be charged to authors, unless a special grant is authorized by the Editor.

5. Orders for reprints must be sent when returning galley proofs, and for this purpose special forms are supplied.

6. Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the *Journal of Laryngology*. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.

7. Editorial communications may be addressed to THE EDITOR, Journal of Laryngology, c/o HEADLEY BROTHERS, 109 Kingsway, London, WCz.

8. The annual subscription is five guineas sterling (U.S.A. \$15) post free, and payable in advance.

9. Single copies will be on sale at 12s. 6d. each; copies of parts up to Vol. LXIII which are available may be purchased at 7s. 6d. each.

to. All subscriptions, advertising and business communications should be sent to the publishers, HEADLEY BROTHERS, 109 KINGSWAY, LONDON, WC2.

United States of America

Orders for this *Journal* may be sent through local booksellers, or to STECHERT-HAFNER, INC., 31-33 East 10th Street, New York, or direct to the publishers, HEADLEY BROTHERS, 109 KINGSWAY, LONDON, WC2, England.

© Journal of Laryngology and Otology, 1967

CONTENTS

PAGE

Sir Victor Negus	259
NASO-ALVEOLAR CYSTS: AETIOLOGY, PRESENTATION AND TREATMENT: SIX CASES ARE REPORTED. G. P. Walsh-Waring (London)	263
TRANSZYGOMATIC APPROACH TO THE PETROUS APEX WITH A RECORD OF THREE CASES. Lewis Labib Samy and Iskander H. Girgis (Cairo)	277
THE SOCIAL DEVELOPMENT OF DEAF CHILDREN. Edmund Critchley (Mogerhanger)	291
The Place of Myringotomy in the Management of Secretory Otitis Media in Children. J. C. Archard (London)	309
Severe Reduction of Hearing and Vestibular Involvement Caused by Vionactan [®] . Hans Jelert (Aalborg, Denmark)	317
NARROW FIELD MASTOIDECTOMY IN TYMPANOPLASTY. J. Nowinski (Krakow, Poland).	325
ON THE TREATMENT OF PAROTID FISTULAE. A. C. Arulpragasam (Kandy, Ceylon)	329
Myringoplasty by Homogenous Vein Graft. J. F. O. Mitchell (Dundee)	339
Clinical Records—	
THE AUDITORY EVOKED CORTICAL RESPONSE AS AN INDEX OF HEARING IN PRACTICAL AUDIOMETRY, H. A. Beagley and J. J. Knight (London).	347
Multiple Neurofibromatosis Involving the VIIIth Nerve. Helio Martins and Jaime T. Benitez (Detroit, Michigan)	353
Congenital Unilateral Choanal Atresia Associated with a Rhinolith. Daoud S. Ali (Baghdad)	359

 RHINOLITH.
 Daoud S. Ali (Baghdad)
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .
 .

GENERAL NO	DTES.	•	•	•	•	•	•	•	•	•	•	367
------------	-------	---	---	---	---	---	---	---	---	---	---	-----

KAMPLEX AUDIOMETRIC EQUIPMENT

AUDIOMETERS



CLINIC AUDIOMETER DA1


ACOUSTIC IMPEDANCE APPARATUS ZA2

PNEUMOMETER ZP1

SKIN RESISTANCE APPARATUS RA4 SCREENING PORTABLE TA16 DIAGNOSTIC PORTABLE TA15

CLINIC AUDIOMETER DA1 DOUBLE CHANNEL DA2

AUTOMATIC AUDIOMETER AB1



SOUND LEVEL METER GA6

VISIBLE SPEECH APPARATUS

AUDITORY TRAINERS

Write for full catalogue and detailed descriptions to:

ALFRED PETERS & SONS LTD 51 GELL STREET, SHEFFIELD 3



'Otosporin' rapidly destroys virtually all bacteria commonly found in ear infections, reduces irritation, inflammation and pain. 'Otosporin' brand Drops contain polymyxin B sulphate, neomycin sulphate and hydrocortisone in a bland suspension designed to facilitate penetration. Issued in bottles of 5 ml.





Burroughs Wellcome & Co. London (The Wellcome Foundation Ltd.)

ORBENIN and its success in **Gram-Positive** Infections

Otitis media Otitis externa

Boils Carbuncles Infected dermatoses

Pneumonia Lung abscess Acute bronchitis

Osteomyelitis Osteitis

Tonsillitis Pharyngitis

Post-operative wound infections Burns Skin graft protection

Septicaemia Acute endocarditis

Staphylococcal enterocolitis Staphylococcal urinary tract infections Staphylococcal meningitis

ORBENIN (cloxacillin sodium B.P.) is a penicillin active against virtually all Gram-positive organisms, including staphylococci, *β*-haemolytic streptococci and pneumococci, irrespective of their resistance to other antibiotics.

It is encouraging that the widespread use of Orbenin in hospitals has not resulted in the emergence of resistant strains.


More than 300 references in the World Literature testify to the excellent clinical results obtained with Orbenin in a wide variety of Gram-positive infections.

Detailed literature is available on request.

Dosage:

Two capsules (500mg.) or one vial (250mg.) q.i.d.

Availability:



(250mg.)

(125mg



Orbenin (cloxacillin sodium B.P.) (regd.) is a product of British research at **Beecham Research Laboratories** Brentford, England. Telephone: ISLeworth 4111 Published online by Cambridge University Press



WHEN A HEARING AID IS RECOMMENDED ONLY AMPLIVOX OFFERS THIS HEARING AID SERVICE

The design and manufacture of a complete range with individual fitting facilities. Best of other makes in addition to Amplivox aids—Amplivox hearing aids are ONLY available from Amplivox. Highest degree of training of any hearing aid organisation. Master Hearing Aid evaluations. Expert follow-up and rehabilitation advice. 15 branch offices with while-you-wait service. FOR YOUR PATIENTS' GREATEST SATISFACTION SPECIFY AMPLIVOX **CANCELOSSION** World's most experienced and complete hearing service

AMPLIVOX HEARING ADVISORY SERVICE 80 NEW BOND STREET LONDON, W.1

BÉKÉSY Accessory

For the very first time a fully automatic unit has been made available for a Clinic Audiometer, making the combined units into a true BÉKÉSY Audiometer.

This automatic accessory is being made for the PETERS SPD/5 CLINIC AUDIOMETER, which has a continuous frequency range and continuously variable attenuation. The result is therefore a true Békésy audiometer.

It can be attached to or detached from the audiometer in under one minute so that the two units together offer the facilities of an advanced clinic audiometer or Békésy audiometry at will.

It can be fitted to any existing SPD/5 Audiometer.

Please write for full details and demonstration to :

ALFRED PETERS & SONS LIMITED 51 GELL STREET, SHEFFIELD 3

ONLY **AMPLIVOX** provides a <u>complete</u> audiometric service!



Transistor Audiometers

with air conduction or air and bone conduction with masking, for hospitals, schools, consulting rooms, industrial medical departments, screening programmes, etc.



Clinical Audiometers with narrow band masking for the highest accuracy and extended resolution.

Audiometer Booths

for ideal testing conditions in hospitals, clinics, factories. Standard and special sizes available.



PLUS An efficient and speedy calibration service please write for technical literature to meet your needs 80 New Bond Street, London, W.1. Tel: HYDe Park 9888



THE LARYNGOSCOPE

A Monthly Journal devoted to the disease of EAR, NOSE AND THROAT

Official organ for the American Laryngological Rhinological and Otological Society

Price \$18.00 per year Foreign \$19.00 per year

Established 1896

Theodore E. Walsh, M.D. Editor


517 SOUTH EUCLID AVENUE SAINT LOUIS MO. 63110.

all clear in sinusitis, otitis media and coryza


'Triogesic' provides decongestion with analgesia.

	TABLETS	ELIXIR				
Revised Formula	scored (per tablet) Phenylpropanolamine-HCl 12.5 mg. Paracetamol B.P. 500 mg.	cherry flavoured (per 5 ml.) Phenylpropanolamine-HCl 3 mg Paracetamol B.P. 125 mg Alcohol (95%) 0.5 ml				
Packs	tubes of 30	bottles of 8 and 20 fl. oz.				
Basic N.H.S. Price	3/6d. for 30 tablets	5/10d. for 8 fl. oz.				

Literature and sample on request. A. Wander Ltd., 42 Upper Grosvenor Street, London W.1



х



-that is the triumph of the **MACKIE** hearing aid

The new Mackie post-aural hearing aid is made in several different circuit versions so that the gain and frequency response can be selected to match the patient's needs; and it is available where required with a unique form of automatic volume control which has proved highly successful in cases of recruitment. The forward pick-up microphone gives maximum intelligibility of speech without background interference.

It is exported to many countries including the United States of America–a tribute to its high quality and advanced design.

In Great Britain it is available for trial by your patients from *independent* hearing aid dealers in all parts of the country.


MANUFACTURED BY A. G. R. MACKIE LTD. A member of the Bonochord Group of Companies FOR FURTHER DETAILS WRITE TO 48 WELBECK STREET, LONDON W1






FOR INFLAMMATION AND INFECTION IN THE EAR

ANTI-**INFLAMMATORY** ANTI-BACTERIAL ANTIFUNGAL





dispenser

٥



Δ


LEOIT

Please mention The Journal of Laryngology and Otology when replying to advertisements

xi

Basic NHS price 9s. 8d

ADVERTISEMENTS



What makes this antibiotic so valuable in the ENT unit ?

Spectrum?

Exceptionally broad. It covers most gram-positive organisms—including the penicillin-resistant staphylococci—and many gram-negative organisms.

Action?

Intensely bactericidal. It gives a rapid kill of bacteria at or near the minimum inhibitory concentration, especially in the case of gram-negative organisms.

Safety?

Wide clinical usage has confirmed the laboratory evidence that Ceporin has a remarkably low toxicity. It is usually well tolerated by patients allergic to penicillin.

Clinical record?

The latest analysis of clinical results obtained with Ceporin shows a total of 131 ENT infections treated : 109 (83%) responded satisfactorily. There were 28 cases of tonsillitis, of which 24 responded well. The majority of these infections was caused by beta-haemolytic streptococci, but there were several mixed infections involving streptococci, staphylococci and *Corynebacterium diphtheriae*. Otitis also responded well. Of the 28 cases treated, 24 responded satisfactorily. Most of these infections were caused by staphylococci but infections involving coliforms, pneumococci, Streptococcus pyogenes, proteus and Haemophilus influenzae were also represented. Often rapid clinical response was obtained with Ceporin after other antibiotics had failed. Conditions which respond well to Ceporin include : mastoiditis, otitis media, sinusitis, quinsy, pharyngeal abscess, and intracranial infections secondary to infected sinuses and mastoiditis. Because of its three main attributes broad spectrum, bactericidal action and low toxicity-Ceporin has proved valuable for use in place of more toxic drugs or combinations of antibiotics. **Presentation** Vials containing 250 mg, 500 mg and 1 gram of cephaloridine, packed singly and in boxes of five.

Ceporin is a Glaxo trade mark

Detailed literature is available from Glaxo Laboratories Limited Greenford, Middlesex