

3. Tomasz A. Multiple antibiotic resistant pathogenic bacteria. A report on the Rockefeller University workshop. *N. Eng J Med* 1994;330:1247-1251.
4. Billstein SA. How the pharmaceutical industry brings an antibiotic drug to market in the United States. *Antimicrob Agents Chemother* 1994;38:2679-2682.
5. Coronado VG, Edwards JR, Culver DH, Gaynes RP National Nosocomial Infections Surveillance system. Ciprofloxacin resistance among nosocomial *Pseudomonas aeruginosa* and *Staphylococcus aureus* in the United States. *Infect Control Hosp Epidemiol* 1995;16:71-75.
6. Blumberg HM, Rimland D, Carroll DJ, Terry FP Wachsmuth IK. Rapid development of ciprofloxacin resistance in methicillin-susceptible and resistant *Staphylococcus aureus*. *J Infect Dis* 1991;163:1279-1285.
7. Bryce EA, Smith JA Focused microbiological surveillance and gram-negative β -lactamase-mediated resistance in an intensive care unit. *Infect Control Hosp Epidemiol* 1995;16:331-334.
8. Yu VL, Oakes CA, Axnick KJ, Merigan TC. Patient factors contributing to the emergence of gentamicin-resistant *Serratia marcescens*. *Am J Med* 1979;44:468-473.
9. Weinstein RA, Kabins SA. Strategies for prevention and control of multiple drug-resistant nosocomial infection. *Am J Med* 1981;70:449-452.
10. Centers for Disease Control and Prevention. Nosocomial enterococci resistant to vancomycin - United States, 1989-1993. *MMWR* 1993;42:597-599.
11. Burwen DR, Banerjee SN, Gaynes RP National Nosocomial Infections Surveillance system. Ceftazidime resistance among selected nosocomial gram-negative bacilli in the United States *J Infect Dis* 1994;170:1622-1625.
12. Gaynes R, Culver D, National Nosocomial Infections Surveillance (NNIS) system. Resistance to imipenem among selected gram-negative bacilli in the United States. *Infect Control Hosp Epidemiol* 1991;13:1014.
13. Weinstein RA, Nathan C, Gruensfelder R, Kabins SA. Endemic aminoglycoside resistance in gram-negative bacilli: epidemiology and mechanisms. *J Infect Dis* 1980;141:338-342.
14. Centers for Disease Control and Prevention. Nosocomial infection rates for interhospital comparison: limitations and possible solutions. *Infect Control Hosp Epidemiol* 1991;12:609-621.
15. McGowan JE Jr. Do intensive hospital antibiotic control programs prevent the spread of antibiotic resistance? *Infect Control Hosp Epidemiol* 1994;15:478-483.
16. Kritchewsky SB, Simmons BP Toward better antibiotic use in hospitals. *Infect Control Hosp Epidemiol* 1994;15:68&690.
17. McGowan JE. Antibiotic-resistant bacteria and healthcare systems: four steps for effective response. *Infect Control Hosp Epidemiol* 1995;16:67-70.
18. HICPAC. Recommendations for preventing the spread of vancomycin resistance. *Infect Control Hosp Epidemiol* 1995;16:105113.
19. Pallares R, Dick R, Wenzel RP Adams JR, Nettleman MD. Trends in antimicrobial utilization at a tertiary teaching hospital during a 15-year period (1978-1992). *Infect Control Hosp Epidemiol* 1993;14:37&382.
20. Ballou CH, Schentag JJ. Trends in antibiotic utilization and bacterial resistance. *Diagn Microbiol Infect Dis* 1992;15:37S-42S.
21. Evans RS, Larsen RA, Burke JP, et al. Computer surveillance of hospital-acquired infections and antibiotic use. *JAMA* 1986;256:1007-1011.

Community Exposure Predicts Healthcare Worker TB Skin-Test Conversion

by **Gina Pugliese, RN, MS**
Medical News Editor

Dr. Thomas Bailey and colleagues from Barnes Hospital in St. Louis, Missouri, a large urban teaching hospital, recently completed a study that assessed the risk for positive tuberculin skin tests among employees. The annual incidence of tuberculosis is approximately five cases per 100,000 persons in Missouri and approximately 11 cases per 100,000 persons in St. Louis. During the period January 1989 to July 1991, three patients with pulmonary tuberculosis were admitted to Barnes hospital.

Of the 6,070 active employees for whom TB screening data was available, 684 (11.3%) had a positive tubercu-

lin skin test (TST) during the study period. Risk factors associated with a positive skin test were age >35 years, minority group status (black, Asian, Hispanic), and percentage of low-income persons within the employee's residential postal zone. Of 3,106 employees who had at least two skin tests, 29 had TST conversion; 15 (52%) occurred among employees who had no direct patient contact. Skin-test conversion was independently associated only with the percentage of low-income persons in the employee's residential postal zone. Stratifying the employees according to the degree of contact with patients or according to departmental group was not useful in determining risk for a positive TST or for TST conversion.

For certain groups of employees, an exposure to tuberculosis in the community poses a greater risk than exposure in a hospital setting. The authors concluded that their findings support the Centers for Disease Control and Prevention recommendation to include healthcare workers from "risk groups with increased prevalence of tuberculosis ... even if they do not have potential occupational exposure ... " in a periodic TST program.

FROM: Bailey TC, Fraser VJ, Spitznagel EL, and Dunagan WC. Risk factors for a positive tuberculin skin test among employees of an urban, midwestern teaching hospital. *Ann Intern Med* 1995;122(8):580-585.