

Human-caused and natural mortality of giant tortoises in the Galapagos Islands during 1995–2004

Cruz Márquez, David A. Wiedenfeld, Sandra Landázuri and Juan Chávez

Abstract Although the killing of giant tortoises in the Galapagos Islands has been prohibited since 1933, poaching of tortoises still occurs. Personnel of the Galapagos National Park Service and the Charles Darwin Research Station regularly survey populations of tortoises throughout the archipelago and report all dead tortoises found. For the 10-year period 1995–2004 the field personnel reported evidence of 190 giant tortoises killed, primarily on the southern portion of Isabela Island. For the first 6 years the number of tortoises found killed was <15 per year, but since 2001 the number killed has increased dramatically, with 49 tortoises poached in 2004. During the same 10 years the number of tortoises found dead

from natural causes was 131. Many of these deaths can be attributed to events associated with the 1997–1998 El Niño or with outbreaks of disease on Santa Cruz Island in 1996 and 1999. The results indicate that poaching exceeds natural mortality, and is a significant factor affecting these long-lived and slow-reproducing animals. Environmental education efforts in the human population of southern Isabela appear to have had little effect. Because tortoise poaching takes place at a small number of sites, effective enforcement at those sites could reduce killing of tortoises.

Keywords Galapagos Islands, Galapagos tortoise, *Geochelone*, mortality, poaching.

Introduction

Giant land tortoises, although at one time abundant on all continents except Antarctica, now reside only on Aldabra atoll in the Seychelles Islands of the Indian Ocean (*Geochelone gigantea*) and the Galapagos Islands (*Geochelone* spp.; Gunther, 1875; MacFarland *et al.*, 1974). In the Galapagos, giant tortoises were abundant in the 16th–18th centuries (Porter, 1815; VanDenburgh, 1914) but were decimated by buccaneers, whalers, seal hunters, merchants and colonists, who exploited the animals for meat and oil (Townsend, 1925; Slevin, 1959). The tortoises also suffered from depredation on eggs, juveniles and adults inflicted by feral mammals and introduced fire ants *Solenopsis* sp. (Beck, 1903; MacFarland *et al.*, 1974).

Of the original 13 species of giant land tortoises on the Galapagos Islands (Ernst & Barbour, 1989) one, the Floreana Island species *Geochelone elephantopus* was extinguished by human pressure in the 1800s (MacFarland *et al.*, 1974). The species from Fernandina Island, *G. phantastica*, was last encountered in 1906, having presumably fallen victim to extensive volcanic activity of the Fernandina volcano. By 1974 MacFarland *et al.* (1974) listed 11 surviving species, one of which (*G. abingdoni* from Santa Cruz Island) was extinct in the wild, with only one surviving male, Solitary George, maintained in captivity since 1972 at the Charles Darwin Research Station and the Galapagos National Park. Seven or eight of the other remaining species were reduced to minimal numbers, and three or four maintained healthy populations, although these were also threatened by predators or introduced mammalian competitors (MacFarland *et al.*, 1974). Since the analysis by MacFarland *et al.* (1974) efforts by the Charles Darwin Research Station and Galapagos National Park Service to protect and restore these populations have resulted in increases for all species except *G. abingdoni* (Solitary George; unpublished reports of Charles Darwin Research Station).

Although killing of Galapagos tortoises was legally prohibited by the Ecuadorian government in 1933, killing of tortoises for local meat consumption continued, albeit at a lower level. Cayot & Lewis (1994) documented an alarming increase in poaching of tortoises

Cruz Márquez and David A. Wiedenfeld* (Corresponding author) Charles Darwin Research Station, Puerto Ayora, Galapagos, Ecuador. E-mail dwiedenfeld@dwiedenfeld.org

Sandra Landázuri Charles Darwin Research Station, Puerto Villamil, Galapagos, Ecuador.

Juan Chávez Galapagos National Park Service, Puerto Villamil, Galapagos, Ecuador.

*Current address: 619 McFarland Avenue, Dalton, Georgia 30720, USA.

Received 12 February 2006. Revision requested 3 August 2006.

Accepted 6 October 2006.

during 1980–1994. The poaching has continued to the present day, especially in the areas of Sierra Negra and Cerro Azul volcanoes on Isabela Island, although poaching has occurred on other islands as well. Here we document and examine change in levels of tortoise poaching in the period since the report by Cayot & Lewis (1994), covering events during 1995–2004, and make recommendations for the future conservation of these species.

Methods

During 1995–2004 populations of Galapagos tortoises were monitored by personnel of the Galapagos National Park Service and the Charles Darwin Research Station, on Isabela, Española, Santiago, Santa Cruz, San Cristóbal, and Pinzón islands (Fig. 1). These surveys were carried out several times per year for some sites (especially Roca Unión and Cerro Cazuela on Isabela Island and on Santa

Cruz Island), and at least twice per year at other sites on the coast of Isabela Island. More remote sites, such as the interior of Pinzón, Española and Santiago islands were usually visited only once per year. During these surveys tortoises found dead were recorded along with information on the cause of death (natural or human-induced) and additional information such as date, location (recorded with a global positioning system if possible), estimated date of death, sex (if possible to determine), age class, and a photograph of the remains. In some cases the curved carapace length of the dead tortoise was measured and recorded. In addition to more formal surveys, personnel of the Park Service and Research Station kept incidental records of dead tortoises whenever they encountered them.

Tortoises dying from natural causes can easily be distinguished from those killed by humans for meat. To extract the meat from a tortoise shell, the plastron is removed, usually by chopping along the sides of the




Fig. 1 The Galapagos Archipelago, showing locations of sites referred to in Tables 1 and 2 where dead tortoises, either killed by humans or a result of natural causes, have been found. Islands (the largest is Isabela Island) and volcanoes (V.) are identified along with the species name of the tortoise (all are *Geochelone*) occurring at that site.

Table 1 Number and percentages of Galapagos tortoises found killed by humans, by site and year, during 1995–2004. All are on Isabela Island except Punta Pitt, which is on San Cristóbal (Fig. 1).

Year	Caleta Iguana	Cinco Cerros	Caleta San Pedro	Roca Unión	Cerro Paloma	Cerro Cazuela	Bahía Urbina	Piedra Blanca	Puerto Bravo	Punta Pitt	Total (%)*
1995				2		1					3 (2)
1996			1	5		1					7 (4)
1997	2			5		3					10 (5)
1998	4			5		5					14 (7)
1999	3			4	2	2					11 (6)
2000	2			3		4	4				13 (7)
2001	7		4	11		6					28 (15)
2002	9		2	5		1					17 (9)
2003	5		4	17		12					38 (20)
2004	3	3	3	19		8	3	4	5	1	49 (26)
Total (%)*	35 (18)	3 (2)	14 (7)	76 (40)	2 (1)	43 (23)	7 (4)	4 (2)	5 (3)	1 (1)	

*Percentages do not add up to 100% because of rounding errors

plastron of an overturned tortoise using a machete or hatchet. Natural causes of death do not cause the removal of the plastron, even in cases of violent death (usually by falling). Usually even a highly decayed carcass from a natural mortality will be upright, with the plastron intact beneath and the limb bones relatively in place.

Results

From 1995 to 2004 personnel recorded the remains of 189 Galapagos tortoises killed by humans on Isabela Island and one on San Cristóbal (Table 1). Tortoises that had been killed by poachers were found at nine sites on Isabela Island (Fig. 1). Of these, three sites account for the large majority of the poached tortoises, with 154 (81%) of the 190 tortoises killed at Roca Unión, Cerro Cazuela and Caleta Iguana (Table 1).

The number of tortoises found killed by humans did not surpass 14 until 2001 (Fig. 2). Prior to 2001 the average number of tortoises found poached each year was 9.7 tortoises; for the 4 years 2001–2004 the average was 33.0 tortoises per year (Table 1), an increase of more than three-fold. The average number of tortoises found poached for the entire 10-year period was 19.0 tortoises per year.

During the same period natural tortoise mortality was recorded on the islands Santa Cruz, Española, Santiago, and Pinzón, and on Isabela Island on volcanoes Wolf, Darwin and Alcedo (Fig. 1), with 131 tortoises found dead from natural causes (Table 2). The greatest number of natural tortoise mortalities (56 individuals) was recorded from Santa Cruz Island (primarily adult males), and the second largest number (49 individuals) from Alcedo Volcano on Isabela Island (adult males and

females, and juveniles). The 12 dead tortoises found on Española Island were all juveniles, as would be expected, because all tortoises on that island are the result of a captive breeding and repatriation project begun in the 1970s. The average number of tortoises found dead from natural causes was 13.1 per year.

High mortality on Alcedo was recorded in 1997 and 1998 during the very strong El Niño of 1997–1998 but high mortality on Santa Cruz occurred in 1996 and 1999, the years before and after the El Niño event.

Discussion

The count of 190 tortoises killed by humans for food during the 10 years 1995–2004 should be considered a minimum estimate. The remoteness, difficult terrain, and dense vegetation of areas where Galapagos tortoises occur and therefore where poaching takes place, coupled




Fig. 2 Numbers of Galapagos tortoises found dead each year during 1995–2004 from natural mortality and from poaching.

Table 2 Natural mortality of Galapagos tortoises, by site and year, during 1995–2004. Santa Cruz, Española, Santiago and Pinzón are islands, and Alcedo, Wolf and Darwin are all volcanoes on Isabela Island (Fig. 1). NV means the site was not visited.

Year	Santa Cruz	Española	Santiago	Pinzón	Alcedo Volcano	Wolf Volcano	Darwin Volcano	Total (%)*
1995	0	NV	NV	NV	1	NV	NV	1 (1)
1996	21	NV	NV	NV	2	NV	NV	23 (18)
1997	0	NV	NV	1	16	NV	NV	17 (13)
1998	0	NV	NV	NV	20	NV	NV	20 (15)
1999	22	NV	NV	NV	1	2	3	28 (21)
2000	6	3	NV	NV	0	NV	NV	9 (7)
2001	0	NV	NV	NV	0	NV	NV	0
2002	2	6	4	NV	0	NV	NV	12 (9)
2003	0	3	NV	NV	6	NV	NV	9 (7)
2004	5	NV	NV	3	3	NV	1	12 (9)
Total (%)*	56 (43)	12 (11)	4 (3)	4 (3)	49 (37)	2 (2)	4 (3)	131

*Percentages do not add up to 100% because of rounding errors

with efforts by poachers to hide the remains of butchered tortoises, mean it is likely that many tortoises killed by human poachers are never found and recorded. Overall, tortoises dying from natural causes are probably more likely to be found than poached tortoises, because no effort is made to hide the remains.

The average number of tortoises found dead each year from natural mortality (13.1) is less than the average number poached each year (19.0). This is especially pronounced when only the period 2001–2004 is considered. Therefore, it is clear that killing of adult and juvenile tortoises by humans has a more significant impact on tortoise populations than natural mortality. Although many more hatchling tortoises are killed by natural (Galapagos hawks *Buteo galapagoensis* and great blue herons *Ardea herodias*) and introduced (cats *Felis catus*, dogs *Canis familiaris*, pigs *Sus scrofa* and rats *Rattus rattus*) predators, the tortoises are long-lived species with low reproductive rates and loss of adults has a much greater effect on population recovery. The wild populations of the most heavily poached species (*Geochelone guntheri* and *G. vicina*) are being augmented by juveniles produced in the Galapagos National Park Service captive breeding centre (Fritts *et al.*, 2000), which offsets and exceeds the losses to poaching. However, because adults are selectively killed by the poachers, natural reproduction in the wild is significantly reduced, precluding any natural reproduction and population recovery.

The only human settlements on Isabela Island are Puerto Villamil, on the southern coast, and small villages and farms on the southern slope of Sierra Negra Volcano. It is therefore not surprising that the tortoises on Sierra Negra Volcano suffer the greatest levels of human depredation, with 64% of all poached tortoises found at its Roca Unión, Cerro Cazuela and Cerro Paloma

sites. The first of these sites is frequented by local fishermen, while Cerros Cazuela and Paloma are inland, and most tortoise poaching is probably carried out by local people while hunting feral goats *Capra hircus*, cattle *Bos taurus* and pigs.

The tortoises of Cerro Azul are primarily depredated at coastal sites (Caleta Iguana, Cinco Cerros and Caleta San Pedro) by local fishermen. These sites, especially Roca Unión, caletas San Pedro and Iguana and Cinco Cerros, are easily accessed by boat from Puerto Villamil. These are also sites where the tortoises range near to the coast, and where they are therefore more available and accessible to crews of fishermen coming ashore. On 10 September 2004 wardens of the Park Service on patrol encountered a Galapagos fishing boat at Caleta Iguana with tortoise meat in the cooking pot. At the time of writing the results of legal action against the fishermen involved in this incident have not been finalized. The more remote sites, also coastal, of Bahía Urbina, Puerto Bravo and Piedra Blanca have experienced lower levels of human depredation on their tortoise populations. These depredations are also probably being carried out by fishermen.

Cayot & Lewis (1994) reported high numbers of poached tortoises at Piedra Blanca, a site that had a low level of poaching in our study, but only one tortoise killed at Caleta Iguana, the site with the second-highest level of poaching in our study. However, overall, Cayot & Lewis (1994) reported findings similar to those of the present study, with poaching of tortoises in approximately the same areas, including high levels of poaching at Cerro Cazuela and Roca Unión. Little has therefore changed since 1994 with regard to the main sites where tortoise killing has been taking place.

Natural mortality was noticeably high in the Santa Cruz population (*Geochelone nigrita*), with 43 mortalities

between 1996–1999, and in the Alcedo Volcano, Isabela Island population (*G. vandenburghi*), with 36 tortoises dying in 1997–1998 and a total of 49 during the decade. On Santa Cruz the mortality of the 43 tortoises was apparently a result of a minor epidemic caused by bacterial septicemia from *Aeromonas hydrophila* and *Pseudomonas* sp. (G. Trueba, pers. comm., 1999; Ontaneda *et al.*, 2001), associated with nematodal parasitosis by *Atractis marquezii* and *Labiduris* sp. (Bursey & Flanagan, 2002). These parasites and environmental factors combined to cause the deaths of 21 tortoises in 1996, with a second outbreak in 1999 causing the death of 23 tortoises. The majority of these mortalities were of adult males. On Alcedo Volcano the large majority of mortalities were associated with the effects of the very strong El Niño of 1997–1998. During this event 36 tortoises were killed, mostly as a result of torrential rains causing tortoises to be washed into and down ravines on the slopes of the volcano. During the same event a few of the tortoises died of unknown causes, but probably related to the heavy rains.

The inhabitants of southern Isabela have inherited a history of predation on tortoises. Prior to the establishment by Antonio Gil of a town and a tortoise processing facility in 1897 at Puerto Villamil, thousands of *Geochelone guntheri* and *G. vicina* lived on the coast and southern slopes of nearby Sierra Negra and Cerro Azul volcanoes (VanDenburgh, 1897; Beck, 1903). Gil's colonists and workers slaughtered most of these animals for meat and oil, such that afterwards so many plastrons were found on the flanks of the volcano that the workers of the colony used them for roofing materials (P. Cartagena, pers. comm., 1986). Local beliefs, such as that tortoise meat is an important nutritional need for post-parturition mothers for 40 days, have reinforced the harvesting of the tortoises.

A further cause of the increased poaching in recent years is resentment towards the Park Service and conservationists in general, which has on occasions produced negative reactions in the local population. Many households in Puerto Villamil in particular remain dependent on fishing, and have felt that regulation of fishing harvests (especially of sea cucumbers, a lucrative item) has reduced their ability to earn a living. The negative backlash to the regulation has been directed towards the regulating agency (Galapagos National Park Service) and towards conservationists, who have generally supported the fishing regulations, and ultimately towards the Park and conservation efforts. Providing economic alternatives to fishing, such as opportunities in tourism business, may reduce the effect of regulations on fishermen and reduce this backlash.

Conservation and environmental education efforts by the Charles Darwin Research Station and Galapagos

National Park Service have apparently had little effect, and poaching of tortoises may be increasing. Improved environmental education is necessary from the earliest ages to enhance the understanding of the value of the tortoises by the inhabitants and reduce the ongoing poaching. In addition, it is necessary to improve enforcement through monitoring of the most active sites where tortoises are killed. The number of these sites is relatively small, with most of the poaching occurring at Cerro Cazuela, Roca Unión and Caleta Iguana, and therefore concentrating enforcement at these sites could have great effect. A redoubled effort to prevent future poaching of Galapagos tortoises is necessary so that similar levels of harvesting are not reported over the next 10 years.

Acknowledgements

We would like to thank the multitude of wardens of the Galapagos National Park Service, and volunteers, scholarship students and personnel of the Charles Darwin Research Station who have helped acquire information on the giant tortoise populations and who have reported incidents of tortoise mortality. Our gratitude also to the Frankfurt Zoological Society, Friends of Galapagos – Switzerland, Galapagos Conservation Foundation, the GEF Project ECU/00/G31 *Especies Invasoras de las Galápagos*, the Government of Ecuador and others that have financed studies of the giant tortoises of Galapagos, without whose help and support it would not have been possible to complete this study. We also wish to express our appreciation to the Galapagos National Park Service for allowing us to work on the islands, and to the Charles Darwin Foundation and Charles Darwin Research Station for the logistical support and administration of funds. An anonymous reviewer helped to improve the manuscript.

References

- Beck, R. (1903) In the home of the giant tortoise. *Annual Report of the New York Zoological Society*, 7, 1–17.
- Bursey, C.R. & Flanagan, J.P. (2002) *Atractis marquezii* N. Sp. (Nematoda: Atractidae) and a revision of *Atractis* Dujardin, 1845, sensu Baker, 1987. *Journal of Parasitology*, 88, 320–324.
- Cayot, L.J. & Lewis, E. (1994) Recent increase in killing of giant tortoises on Isabela Island. *Noticias de Galápagos*, 54, 2–7.
- Ernst, C. & Barbour, R. (1989) *Turtles of the World*. Smithsonian Institution Press, Washington, DC, USA.
- Fritts, T.H., Snell, H.L., Cayot, L., MacFarland, C., Earsom, S., Márquez, C., Llerena, W. & Llerena, F. (2000) Progress and priorities in research for the conservation of reptiles. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique*, 70, 39–45.
- Gunther, A. (1875) Description of the living and extinct races of gigantic land tortoises. Part I–II, introduction, and the

- tortoises of the Galapagos Islands. *Philosophical Transactions of the Royal Society*, **165**, 251–284.
- MacFarland, C.G., Villa, J. & Toro, B. (1974) The Galapagos giant tortoises (*Geochelone elephantopus*). I. Status of the surviving populations. *Biological Conservation*, **6**, 118–133.
- Ontaneda, S., Trueba, G., Sevilla, A.M. & Sevilla, E. (2001) *Informe del análisis microbiológico de muestras de tortugas*. Unpublished Report to Charles Darwin Research Station, Puerto Ayora, Galapagos, Ecuador.
- Porter, D. (1815) *Journal of a Cruise Made to the Pacific Ocean by Captain David Porter, in the United States Frigate Essex, in the Years 1812, 1813, 1814*. Bradford & Inskeep, Philadelphia, USA.
- Slevin, J.R. (1959) Islands of the tortoises. *Pacific Discovery*, **12**, 4–10.
- Townsend, C.H. (1925) The Galapagos tortoises in their relation to the whaling industry: a study of old logbooks. *Zoologica*, **4**, 55–135.
- VanDenburgh, J. (1897) The reptiles of the Pacific coast and Great Basin. *Occasional Papers of the California Academy of Science*, **5**.
- VanDenburgh, J. (1914) The gigantic land tortoises of the Galapagos archipelago. *Proceedings of the California Academy of Science*, **2**, Part 1, 203–374.

Biographical sketches

Cruz Márquez is a herpetologist at the Charles Darwin Research Station. His work focuses on Galapagos tortoises and land and marine iguanas.

David A. Wiedenfeld is the former head of the Department of Vertebrate Ecology and Monitoring at the Charles Darwin Research Station. Although his primary research interests are in bird population studies, he has also carried out investigations of tortoises and iguanas.

Sandra Landázuri is Herpetology Associate at the Charles Darwin Research Station and is based at Puerto Villamil on southern Isabela. Her work involves monitoring giant tortoise and introduced frog populations.

Juan Chávez is the head of the Technical Office of the Galapagos National Park Service in Puerto Villamil, Isabela.