

Methods: A case study of 36 year old female presented with clinical manifestations of autoimmune encephalitis syndrome.

Results: Diagnosis confirmed by presence of NMDA receptor antibodies in serum and cerebrospinal fluid.

Conclusions: Early recognition of clinical features of Anti-NMDA receptor encephalitis and early initiation of treatment has shown to improve outcomes, speed recovery and reduce the risk of relapses.

Disclosure: No significant relationships.

Keywords: encephalitis; anti-nmda receptor encephalitis

EPV0397

The right temporoparietal junction and cooperation dilemma

S. Tei^{1,2,3,4,*} and J. Fujino^{1,4,5}

¹Department Of Psychiatry, Kyoto University, Kyoto, Japan; ²Institute Of Applied Brain Sciences, Waseda University, Saitama, Japan; ³School Of Human And Social Sciences, Tokyo International University, Saitama, Japan; ⁴Medical Institute Of Developmental Disabilities Research, Showa University, Tokyo, Japan and ⁵Department Of Psychiatry And Behavioral Sciences, Tokyo Medical and Dental University, Tokyo, Japan

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1952

Introduction: Cooperation is a key component of our lives. When we identify people in need, we are frequently motivated to cooperate by overcoming selfishness. However, we may also become selfish to pursue greater gains by putting ourselves at risk and exploiting others. Such cooperation dilemmas are ubiquitous in real life. Although functional magnetic resonance imaging studies have repeatedly reported the involvement of right temporoparietal junction (rTPJ) in cooperation dilemmas, a causal link between the two has been rarely explored.

Objectives: To investigate a causal role of rTPJ in resolving cooperation dilemmas in ecologically valid settings.

Methods: Twenty-two healthy volunteers were examined. We combined repetitive transcranial magnetic stimulation (rTMS) with a snowdrift cooperation dilemma game task (cross-the-traffic intersection version) wherein either cooperation or defection should be chosen. Participants and opponents jointly faced a problem at the intersection where their cooperation could diffuse the situation (stopping/avoiding a car-crash). This conflicted with a choice in the participant's self-interest which was more rewarding, but risky (not stopping/defection). We also included explicit-cue condition that showed elderly/pregnant passengers in the opponent's car. Furthermore, we measured participants' empathic-traits (e.g., perspective-taking) to study personality-cooperation associations.

Results: The cooperation-ratio did not statistically differ between the sham stimulation and inhibitory continuous theta burst stimulation (cTBS) in both the no-cue and with-cue conditions. However, after cTBS, only in the no-cue condition, the strength of the relationship between cooperation-ratios and empathic-traits decreased significantly ($p < 0.05$).

Conclusions: These results contribute to our understandings of rTPJ's role in spontaneous social cognition, which may be considerably complex and require further examination.

Disclosure: No significant relationships.

Keywords: temporoparietal junction; cooperation; transcranial magnetic stimulation; decision making

EPV0398

Sense of coherence, spontaneous brain activity, and burnout severity

S. Tei^{1,2,3,*} and J. Fujino^{1,4}

¹Department Of Psychiatry, Kyoto University, Kyoto, Japan; ²Institute Of Applied Brain Sciences, Waseda University, Saitama, Japan; ³School Of Human And Social Sciences, Tokyo International University, Saitama, Japan and ⁴Department Of Psychiatry And Behavioral Sciences, Tokyo Medical and Dental University, Tokyo, Japan

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1953

Introduction: Burnout has become a critical issue in health care systems during the COVID-19 pandemic. Several studies report on the importance of peoples' sense of coherence (SOC) or control over work for dealing with burnout. SOC implicates a stress-coping capacity involving comprehensibility, manageability, and meaningfulness. However, little is known on how SOC cognitively modulates burnout experiences.

Objectives: To investigate neurocognitive mechanisms of SOC and burnout in medical professionals.

Methods: Forty-one registered nurses were enrolled. We used functional magnetic resonance imaging and measured resting-state brain activity. We identified brain regions associated with SOC and burnout levels by correlating these trait scores to regional fractional amplitude of low frequency fluctuations (fALFF). Subsequently, we investigated whether participants' levels of SOC impacted their fALFF-burnout association by mediation analysis.

Results: SOC and depersonalization dimension of burnout were negatively correlated ($p < 0.01$). The fALFF in the mid-dorsolateral prefrontal cortex (DLPFC) correlated positively with SOC scores, and negatively with depersonalization dimension of burnout ($p < 0.05$). Furthermore, SOC mediated the negative relationship between DLPFC activity and burnout severity ($p < 0.05$).

Conclusions: Our data suggested that SOC alleviates burnout experience and supports prefrontal activity to prompt cognitive control; they may facilitate flexible shifting of perspective and optimistic reappraisal of work-stress. In effect, workplace-stressors may be acknowledged as being more meaningful than distressing. Without sufficient SOC, frequent exposures to stressors can lead to maladaptive coping to exhibit emotional numbing or depersonalization.

Disclosure: No significant relationships.

Keywords: sense of coherence; burnout; medical professionals; fMRI

EPV0399

An fMRI study of decision-making under conflict in individuals with autism spectrum condition

S. Tei^{1,2,3,4,*} and J. Fujino^{1,4,5}

¹Department Of Psychiatry, Kyoto University, Kyoto, Japan; ²Institute Of Applied Brain Sciences, Waseda University, Saitama, Japan; ³School Of Human And Social Sciences, Tokyo International University,

Saitama, Japan; ⁴Medical Institute Of Developmental Disabilities Research, Showa University, Tokyo, Japan and ⁵Department Of Psychiatry And Behavioral Sciences, Tokyo Medical and Dental University, Tokyo, Japan

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1954

Introduction: Individuals with autism spectrum condition (ASC) frequently report difficulties in social communications, combined with restricted inflexible behaviors. However, it is unclear whether this rigidity is pervasive across cognitive flexibility (CF) and affective flexibility (AF) in situations which resolve different social conflicts.

Objectives: To study CF and AF levels and associated brain activity in individuals with ASC.

Methods: Individuals with ASC and with typical development (TD) performed a moral dilemma task during functional magnetic resonance imaging. For CF, participants made decisions on (1) whether to enforce result-oriented actions to prioritize social/public benefits; and (2) judged whether these actions are right or wrong. For AF, participants made decisions on (1) whether to permit social norm/rule violations in sympathy-evoking situations; and (2) permit these identical violations in no sympathy-evoking situations. We calculated participants' CF/AF levels by computing the switching-rate of decisions in CF/AF sessions (switching was defined as: CF, judging the actions as wrong but choosing to enforce the action in the same vignette; AF, judging the violations as not permissible in a no sympathy-evoking circumstance, but permissible in a sympathy-evoking circumstance).

Results: For CF, ASC participants showed a marked decrease in CF switching-rates compared to TD participants ($p < 0.05$), and in corresponding brain activity for executive functioning. For AF, although the AF switching rate difference was non-significant, we observed unique brain activities in each group (e.g., TD activation of the greater dorsomedial-prefrontal cortex and ASC activation of the cingulate cortex).

Conclusions: Our results suggest ASC inflexibility may be further characterized by both CF and AF.

Disclosure: No significant relationships.

Keywords: flexibility; decision making; autism; fMRI

EPV0400

Functional connectivity alterations between default mode network and occipital cortex in patients with obsessive-compulsive disorder (OCD)

T. Geffen^{1*}, J. Smallwood², C. Finke^{3,4}, Z. Sjoerds^{5,6} and F. Schlagenhauf^{1,7,8}

¹Psychiatry, Charité, Berlin, Germany; ²Psychology, Queen's University, Kingston, Canada; ³Neurology, Charité, Berlin, Germany; ⁴Berlin School Of Mind And Brain, Humboldt-Universität, Berlin, Germany; ⁵Institute Of Psychology - Cognitive Psychology Unit, Leiden University, Leiden, Netherlands; ⁶Leiden Institute For Brain & Cognition, Leiden University, Leiden, Netherlands; ⁷Institute For Human Cognitive And Brain Sciences, Max-Planck, Leipzig, Germany and ⁸Bernstein Center For Computational Neuroscience, Humboldt-Universität, Berlin, Germany

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1955

Introduction: A meta-analysis by Gürsel et al. (2018) found altered functional connectivity in OCD patients within and between default mode (DMN), salience (SN), and frontoparietal networks (FPN), as well as evidence for aberrant fronto-striatal circuitry.

Objectives: Testing the replicability of meta-analysis rsfMRI findings in OCD patients.

Methods: We measured functional connectivity during resting-state fMRI in a sample of OCD patients ($n=24$) and controls matched for age and sex ($n=33$). The CONN toolbox implemented in SPM was used to perform seed-to-voxel analysis using 30 seed regions based on the previous meta-analytic findings.

Results: OCD patients showed reduced functional connectivity between SN and DMN compared to controls, replicating previous findings. We did not observe significant group differences of functional connectivity within the DMN, SN, or FPN. The strongest finding consisted of altered connectivity between DMN and SN to the visual network. OCD patients showed reduced functional connectivity between the left lateral parietal seed (LPI) and the inferior lateral occipital pole left (iLOCL) compared to controls. Furthermore, the LPI was found to be hyperconnected with the right superior lateral occipital cortex (sLOCr) and the right precuneus. This finding was positively correlated to OCD symptom severity, especially compulsions.

Conclusions: Our findings replicated partly the meta-analysis findings, specifically reduced connectivity between SN and DMN. Using seeds based on the meta-analysis, we identified aberrations between the SN and, in particular, the DMN to the visual network. This raises the question about the visual system's involvement in OCD symptoms and the abnormal connectivity of a unimodal region to the multimodal DMN.

Disclosure: No significant relationships.

Keywords: neuropsychiatry; functional-connectivity; DMN_Visual-Network; ocd

EPV0401

Effects of acute stress on probabilistic reversal learning in healthy participants

L. Wieland^{1*}, C. Ebrahimi¹, T. Katthagen¹, M. Panitz², A. Heinz¹, Z. Sjoerds³ and F. Schlagenhauf¹

¹Psychiatry And Psychotherapy, Charité Universitätsmedizin, Berlin, Germany; ²Neurologie, Max-Planck-Institut für Kognitions- und Neurowissenschaften, Leipzig, Germany and ³Institute Of Psychology - Cognitive Psychology Unit, Leiden University, Leiden, Netherlands

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1956

Introduction: Altered reward-based learning and stress play an important role in psychiatric illnesses, such as psychosis or addiction. Stress sometimes increases learning from rewards, other times it does not show an effect (Starcke & Brand, 2016). A task addressing reward-based learning is the reversal learning task, which uses probabilistic rewards as feedback and incorporates sudden changes in reward contingencies. The effects of acute stress on reversal learning have rarely been addressed.

Objectives: Here, we investigated the effect of acute social stress in a within-subject design in healthy participants.

Methods: A sample of $n = 28$ male non-clinical participants performed the task in a control condition versus the Trier Social Stress