



The p -Rank of Ramified Covers of Curves

IRENE I. BOUW

Department of Mathematics, 209 South 33rd Street, Philadelphia, PA 19104, U.S.A.;
e-mail: bouw@math.upenn.edu

(Received: 10 November 1999; accepted: 12 May 2000)

Abstract. In this paper we study the p -rank of Abelian prime-to- p covers of the generic r -pointed curve of genus g . There is an obvious bound on the p -rank of the cover. We show that it suffices to compute the p -rank of cyclic prime-to- p covers of the generic r -pointed curve of genus zero. In that situation, we show that, for large p , the p -rank of the cover is equal to the bound.

Mathematics Subject Classification (2000). 14H30.

Key words. covers, p -rank.

1. Introduction

This paper is concerned with computing the p -rank of (ramified) covers of curves in characteristic p . Mainly we are interested in Abelian covers of prime-to- p order of the generic r -pointed curve of genus g . This question can be reformulated in terms of quotients of tame fundamental groups.

Let X be a nonsingular connected projective curve over an algebraically closed field k of characteristic $p > 0$ and let $g = g(X)$. Put $U = X - S$. Not much is known about the tame fundamental group $\pi_1^t(U)$. Its structure is known only for $2g + r \leq 2$. By a result of Grothendieck [6, XII.2.12], $\pi_1^t(U)$ is a quotient of $\hat{\Gamma}_{g,r}$, the fundamental group of a curve over \mathbb{C} with same g and r . The prime-to- p parts of both groups are equal, but $\pi_1^t(U) \simeq \hat{\Gamma}_{g,r}$ only in ‘trivial’ cases. For $g \geq 1$ this is seen by considering the p -cyclic quotients of $\pi_1^t(U)$; they correspond to the étale p -cyclic covers of X . The maximal elementary Abelian p -quotient of $\pi_1^t(U)$ is $(\mathbb{Z}/p)^{\sigma(X)}$, where $0 \leq \sigma(X) \leq g(X)$ is the p -rank of X . The maximal elementary Abelian p -quotient of $\hat{\Gamma}_{g,r}$ is $(\mathbb{Z}/p)^{2g}$. This implies that there are less étale p -cyclic covers in characteristic p than in characteristic zero. In fact, the p -part of $\pi_1^t(U)$ is a free pro- p group on $\sigma(X)$ generators. This result is related to the Deuring–Shafarevich formula. This is a formula for the p -rank $\sigma(Y)$ in terms of $\sigma(X)$ and the ramification indices for a (possibly ramified) cover $Y \rightarrow X$ whose Galois group is a p -group, [2]. We will see that the situation for prime-to- p covers is more complicated.

The p -part and the prime-to- p part of $\pi_1^t(U)$ are known; the next case to consider is quotients G which are an extension of a prime-to- p group H by a p -group P . We may

suppose that P is elementary Abelian, since $\pi^p(X)$ is a free pro- p group. Suppose we are given a quotient $\pi_1^t(U) \twoheadrightarrow G$. To this quotient corresponds a G -cover $Z \rightarrow X$; it factors through $Y := Z/P$. The cover $Y \rightarrow X$ is prime-to- p ; these covers we know exactly. The cover $Z \rightarrow Y$ is étale. We can reformulate the question of G -quotients of $\pi_1^t(U)$ as follows. For convenience, we consider only quotients which lie over a fixed H -quotient, i.e. we fix the H -cover $Y \rightarrow X$. Then G -quotients of $\pi_1^t(U)$ correspond to $\mathbb{F}_p[H]$ -submodules P of $H^1(Y, \mathcal{O}_Y)^F$, where F is the Frobenius morphism. This means that in order to solve our problem, we have to compute the structure of $H^1(Y, \mathcal{O}_Y)^F$ as $\mathbb{F}_p[H]$ -module. The dimensions of the isotypical spaces are called the generalized Hasse–Witt invariants. For a more precise definition, see Section 2. The generalized Hasse–Witt invariants can be viewed as generalizations of the p -rank by taking into account the H -Galois action. It is easy to see that if we consider a family of covers of curves with fixed (g, r) , then the p -rank will depend on the bottom curve. Here we are interested in computing the p -rank in case the bottom curve is the generic r -pointed curve of genus g , i.e. corresponds to the generic point of the moduli space $M_{g,r} \otimes \mathbb{F}_p$.

The question of the p -rank of a prime-to- p cover has been considered previously in the étale case, [11, 14, 17]. In these papers it is shown that many étale covers of the generic curve are ordinary, most importantly this holds for Abelian covers. But in [17] it is shown that this is not true for all groups: there exist étale nonordinary covers of the generic curve X_g for every $g \geq 2$. In this paper we consider what happens for ramified covers. It is easy to see that it is not to be expected that all (Abelian) covers of the generic r -pointed curve of genus g are ordinary. It turns out that there is an obvious bound $B(\mathbf{a}, g)$ on the p -rank coming from the structure of $H^1(Y, \mathcal{O}_Y)$ as $k[H]$ -module, here \mathbf{a} denotes the monodromy. The $k[H]$ -module structure of $H^1(Y, \mathcal{O}_Y)$ is known by a results of Chevalley–Weil, see Section 3.

The question whether all étale covers of the generic curve of genus g are ordinary translates for ramified covers into the question whether all (prime-to- p) covers of the generic r -pointed curve of genus g have p -rank equal to $B(\mathbf{a}, g)$. The situation for ramified covers is analogous to the situation for étale covers, with this modification. We will show that there exists a non-Abelian cover of the generic curve whose p -rank is less than the bound (Example 3.5). It seems reasonable to expect that the p -rank of Abelian covers of the generic r -pointed curve of genus g is equal to the bound. We will show this, under some mild hypothesis on the characteristic.

THEOREM. *Let (X, S) be the generic r -pointed curve of genus g . Suppose that p is sufficiently large. Let H be an Abelian group of order prime-to- p , and $Y \rightarrow X$ an H -Galois cover, unbranched outside S . Then the p -rank of Y is equal to the bound.*

The proof will proceed in several steps. We show that we can restrict to the case that H is cyclic. In Section 6 we prove the theorem for cyclic covers of \mathbb{P}^1 . We prove this by computing the coefficients of the Hasse–Witt matrix of Y . In Section 8 we

show that the statement for genus greater than zero can be reduced to the case of genus zero. This is shown by degenerating covers in a suitable way.

The outline of the paper is as follows. In Section 2 we define the generalized Hasse–Witt invariants and study basic properties. We relate the generalized Hasse–Witt invariants to quotients of tame fundamental groups. In Section 3 we describe the $k[H]$ -module structure of $H^1(Y, \mathcal{O}_Y)$ and use this to define a bound on $\sigma(Y)$. In Section 3 we also compute the generalized Hasse–Witt invariants for some non-Abelian covers. We give an example, due to Raynaud, of a cover of \mathbb{P}^1 branched at three points whose p -rank is unequal to the bound. In Section 4 we concentrate on the case of cyclic covers. In Section 5 we compute the coefficients of the Hasse–Witt matrix of a cyclic cover of \mathbb{P}^1 . In Section 6 we use this to compute the p -rank of a cyclic cover of \mathbb{P}^1 in case p is large and the branch points are general. In Section 7 we explain how we can get information on the p -rank of a cover by degenerating it to a cover of semistable curves. We use this to reprove the main theorem in case $p \equiv \pm 1 \pmod{\ell}$ or $r \leq 4$. In these cases we can drop the assumption p large. In Section 8 we use the same method to show that the computation of the generalized Hasse–Witt invariants of covers of the generic curve of g can be reduced to the computation of the generalized Hasse–Witt invariants of covers of \mathbb{P}^1 .

2. Generalized Hasse–Witt Invariants

In this section we will define generalized Hasse–Witt invariants and study their basic properties. Most importantly, we will give the relation between quotients of $\pi_1^t(U)$ which are an extension of a prime-to- p group H by a p -group P , and the generalized Hasse–Witt invariants of H -covers of X , unbranched outside S .

The generalized Hasse–Witt invariants were first introduced by Katsurada in [9] in the case of an étale ℓ -cyclic cover with $\ell | (p - 1)$. In that paper they were used to show that $\pi_1(X)$ is not determined by the genus, the characteristic and the p -rank of X . Namely, it was shown that $\pi_1(X)$ also depends on the generalized Hasse–Witt invariants. The reason being that the number of (étale) covers of X whose Galois group is an extension of \mathbb{Z}/ℓ by \mathbb{Z}/p can be expressed in terms of the generalized Hasse–Witt invariants. Later this was generalized by Nakajima [11], Rück [18] and Pacheco [13]. Of these authors, Pacheco gave the most general definition, namely in the case of possibly ramified Galois covers whose Galois group is of order prime to the characteristic. Also in this case the generalized Hasse–Witt invariants of a G -Galois cover $Y \rightarrow X$ can be used to give a formula for the number of tame Galois covers of X factoring through Y , whose Galois group is an extension of G by an elementary Abelian p -group. In the étale case this was proved by Pacheco.

Let $\pi: Y \rightarrow X$ be a Galois cover of nonsingular projective irreducible curves defined over an algebraically closed field k of characteristic $p > 0$. We allow the cover to be ramified. Let H be the Galois group of π , suppose that its order is prime to the characteristic. The group H acts naturally on $H^1(Y, \mathcal{O}_Y)$. As was explained

in the introduction, we are interested in the dimension over \mathbb{F}_p of the isotypical spaces of $H^1(Y, \mathcal{O}_Y)^F$, where F is the Frobenius morphism. For each irreducible k -character χ of G , we denote by $L(\chi)$ the χ -isotypical part of $H^1(Y, \mathcal{O}_Y)$, i.e. the largest subspace which, as a $k[G]$ -module, is a sum of irreducible representation with character χ . The Frobenius morphism F sends $L(\chi)$ to $L(\chi^p)$, here χ^p is the Frobenius twist of χ . For each irreducible character χ , we denote by $f(\chi)$ the minimal positive integer such that $F^{f(\chi)}$ sends $L(\chi)$ to itself. Recall that the p -rank $\sigma(X)$ of X is equal to the k -dimension of the largest subspace of $H^1(X, \mathcal{O}_X)$ on which F is a bijection. We mimic this to define the generalized Hasse–Witt invariants.

DEFINITION 2.1. Let $\pi: Y \rightarrow X$ be a G -Galois cover of curves over k and suppose that $p \nmid |G|$. For each irreducible character χ of G , we define the *generalized Hasse–Witt invariant* $\gamma_\pi(\chi)(= \gamma(\chi))$ of the cover $\pi: Y \rightarrow X$ as the dimension of the largest subspace of $L(\chi^{-1})$ on which $F^{f(\chi^{-1})}$ is a bijection.

We consider the inverse character in the above definition to make our definition consistent with the literature. Let q be a sufficiently large power of p . We identify k -characters with \mathbb{F}_q -characters. The space $J(Y)[p] \otimes_{\mathbb{F}_p} \mathbb{F}_q$ is dual to $H^1(Y, \mathcal{O}_Y)^F \otimes_{\mathbb{F}_p} \mathbb{F}_q$ as $\mathbb{F}_q[G]$ -module, [20, p. 38]. In [13] the generalized Hasse–Witt invariants are defined as the dimensions of the χ -isotypical part of $J(Y)[p] \otimes_{\mathbb{F}_p} \mathbb{F}_q$. One easily checks that this definition coincides with our definition. The following lemma gives some elementary properties of the generalized Hasse–Witt invariants (cf. [13]).

LEMMA 2.2. Let $\pi: Y \rightarrow X$ be a G -Galois cover of curves and suppose that $p \nmid |G|$. Let χ be an irreducible character of G and let n_χ be its dimension. Then

- (i) $\gamma(\chi) = \gamma(\chi^p)$,
- (ii) $\sigma(Y) = \sum_\chi \gamma(\chi)$, where the sum is taken over the irreducible k -characters of H ,
- (iii) if χ is the trivial character, then $\gamma(\chi) = \sigma(X)$,
- (iv) $\gamma(\chi)$ is a multiple of n_χ
- (v) let G_χ be the kernel of the representation corresponding to χ and write $\bar{\pi} = \pi/G_\chi$.

Then χ may be considered as a character of G/G_χ and $\gamma_\pi(\chi) = \gamma_{\bar{\pi}}(\chi)$.

Proof. We can write $L(\chi)$ as $L(\chi)^s \oplus L(\chi)^n$, where $F^{f(\chi)}$ is a bijection on $L(\chi)^s$ and is nilpotent on $L(\chi)^n$, [20, no. 9]. The Frobenius morphism is a bijection from $L(\chi)^s$ to $L(\chi^p)^s$. This proves the first statement. The second statement follows immediately from the first, since $H^1(Y, \mathcal{O}_Y)^s = \bigoplus_{\chi \in X} L(\chi)^s$. Let χ be the trivial character, then

$$L(\chi) = H^1(Y, \mathcal{O}_Y)^G \simeq H^1(X, \mathcal{O}_X).$$

This proves the third statement. The fourth statement follows from the fact the $H^1(Y, \mathcal{O}_Y)^s$ is a G -module. The fifth statement is obvious. □

We will now relate the generalized Hasse–Witt invariants to the quotients of $\pi_1^t(U)$. The p -rank of a curve X is related to the number of étale p -cyclic covers of X . If σ denotes the p -rank of X , then the number of étale p -cyclic covers of X is equal to the number of p -cyclic quotients of $(\mathbb{Z}/p)^\sigma$. In fact, something more general holds. The p -part of the fundamental group $\pi^p(X)$ is a free pro- p group on σ generators. Hence, a p -group P occurs as Galois group of an étale cover over X iff P can be generated by σ elements. A similar statement holds for the generalized Hasse–Witt invariants.

LEMMA 2.3. *Let $\pi: Y \rightarrow X$ be an H -Galois cover, with H prime-to- p . There is a 1-1 correspondence between Galois covers $Z \rightarrow X$ which dominate π with Galois group an extension of H by an elementary Abelian p -group and H -submodules of $\text{Hom}(\pi_1(Y), \mathbb{Z}/p)$.*

Proof. Suppose given a Galois cover $Z \rightarrow X$ which dominates π , whose Galois group G is an extension of H by an elementary Abelian p -group $P \simeq (\mathbb{Z}/p)^n$. Let $V = \text{Hom}(\text{Gal}(Z, Y), \mathbb{Z}/p)$. We may regard V as a subspace of $\text{Hom}(\pi_1(Y), \mathbb{Z}/p)$ via the exact sequence

$$1 \rightarrow \pi_1(Z) \rightarrow \pi_1(Y) \rightarrow \text{Gal}(Z, Y) = P \rightarrow 1.$$

Note that V as an $\mathbb{F}_p[H]$ -module is dual to P .

Conversely, suppose given $V \subset \text{Hom}(\pi_1(Y), \mathbb{Z}/p)$ with $\dim_{\mathbb{F}_p} V = n$. Then V corresponds to an étale $(\mathbb{Z}/p)^n$ -Galois cover $Z \rightarrow Y$. The cover $Z \rightarrow X$ is Galois iff V is H -equivariant. The space V with the action of H can be identified with $\text{Hom}(\text{Gal}(Z, Y), \mathbb{Z}/p)$. □

Let $X(H)$ be the set of irreducible k -characters of H . Write $\chi \sim \chi'$ if $\chi' = \chi^{p^j}$ for some j . Let $\bar{X}(H) = X(H)/\sim$. Recall that the set $\bar{X}(H)$ corresponds to the set of \mathbb{F}_p -irreducible characters of H .

For an $\mathbb{F}_p[H]$ -module P , write $\sum_{[\chi] \in \bar{X}(H)} m_P(\chi)(\chi + \chi^p + \dots + \chi^{p^{f-1}})$ for its character and let $n_P = \dim_{\mathbb{F}_p} P$. Let $P^* = \text{Hom}(P, \mathbb{Z}/p)$ be the dual $\mathbb{F}_p[H]$ -module. The proof of the above lemma shows that $P^* \subset H^1(Y, \mathcal{O}_Y)^F$ corresponds to a tame $P \rtimes H$ cover dominating π . The character of $H^1(Y, \mathcal{O}_Y)^F$ is

$$\sum_{[\chi] \in \bar{X}(H)} (\gamma(\chi^{-1})/n(\chi))(\chi + \chi^p + \dots + \chi^{p^{f-1}}).$$

Therefore $P^* \subset H^1(Y, \mathcal{O}_Y)^F$ iff $m_P(\chi) \leq \gamma(\chi)/n(\chi)$ for all χ .

Let $\phi \in \text{Aut}(H)$. For an irreducible k -character χ , denote by χ^ϕ the character obtained by twisting by ϕ . Let P be as above and put $G = P \rtimes H$. Then there exists a tame G -cover dominating π iff there exists a $\phi \in \text{Aut}(H)$ such that $m_P(\chi) \leq \gamma(\chi^\phi)/n(\chi)$. We need to twist by ϕ because for two different $\mathbb{F}_p[H]$ -modules P and P' , the groups $P \rtimes H$ and $P' \rtimes H$ might be isomorphic. This proves the following proposition.

PROPOSITION 2.4. *Let $\pi: Y \rightarrow X$ be a tame H -Galois cover of nonsingular curves. Let P be an $\mathbb{F}_p[H]$ -module and put $G = P \rtimes H$. Then there exists a tame G -Galois cover of X dominating π if and only if there exists an automorphism ϕ of H such that*

$$m(\chi) \leq \frac{\gamma(\chi^\phi)}{n(\chi)} \text{ for each } [\chi] \in \bar{X}(H),$$

where $n(\chi)$ is the dimension of the representation with character χ .

Results similar to the proposition above can be found in [11] and [13]. The result of [11] is a special case of the result in [13] which is a special case of the present case. Our proof follows the proof of [11].

We can phrase this proposition differently. Let G_{\max} be the group which is an extension of H by the elementary Abelian p -group $P = (\mathbb{Z}/p)^{\sigma(Y)}$, where H acts on P via $\sum_{\chi \in X(H)} (\gamma(\chi)/n(\chi))\chi$. Then G_{\max} is the largest extension of H by an elementary Abelian p -group for which there is a Galois cover dominating π . A group G as in the proposition exists if and only if G is a quotient of G_{\max} .

PROPOSITION 2.5. *Let G be a group which is an extension of a prime-to- p group H by a p -group P . Let $\Phi(P) = P^p[P, P]$ be the Frattini subgroup of P and $\bar{P} = P/\Phi(P)$. Write $\bar{G} = \bar{P} \rtimes H$. Suppose given a tame \bar{G} -cover $f: Z \rightarrow X$. Then there exists a tame G -cover $g: W \rightarrow X$ dominating f .*

Proof. The proposition is proved in [14] for étale covers, but the proof carries over immediately to the case of tamely ramified covers. The results follows from the fact that, for $Y = Z/\bar{P}$, the p -part of the fundamental group $\pi_1^p(Y)$ is a free pro- p group. □

3. A Bound on the p -Rank

From Lemma 2.2 it follows that

$$\gamma(\chi^{-1}) \leq \min_j \dim L(\chi^{p^j}).$$

For the p -rank of Y , this implies

$$\sigma(Y) - \sigma(X) \leq \sum_{\chi \neq 1} \min_j \dim L(\chi^{p^j}). \tag{1}$$

Therefore knowledge of the dimensions of the $L(\chi^j)$ gives nontrivial information on the generalized Hasse–Witt invariants and the p -rank of the cover. These dimensions are known by a classical result of Chevalley–Weil [1] (Proposition 3.1 below). The results was originally stated for $k = \mathbb{C}$, but can be extended to our case, since $p \nmid |G|$ (see [8, 12]).

For each $y \in Y$ write G_y for the decomposition group of y , let e_y be the order of G_y . Choose a local parameter u_y at y . Define a character

$$\theta_y: G_y \rightarrow k^*, \theta_y(g) = \frac{g \cdot u_y}{u_y} \pmod{(u_y)}. \tag{2}$$

Note that if $y_1, y_2 \in Y$ both map to $x \in X$ then the characters θ_{y_1} and θ_{y_2} are conjugate.

PROPOSITION 3.1 (Chevalley–Weil). *There exists a unique $k[G]$ -module R such that*

$$|G| \cdot R \simeq \bigoplus_{y \in Y} \left(\bigoplus_{d=0}^{e_y-1} d \cdot \text{Ind}_{G_y}^G \theta_y^d \right) \tag{3}$$

as $k[G]$ -modules. The $k[G]$ -module structure of $H^1(Y, \mathcal{O}_Y)$ is given by

$$H^1(Y, \mathcal{O}_Y) \simeq k \oplus k[G]^{g(X)-1} \oplus R.$$

Proof. [1]. □

DEFINITION 3.2. Let x_1, \dots, x_r be the branch points of the G -Galois cover $\pi: Y \rightarrow X$. For each branch point $x_i \in X$ choose $y_i \in \pi^{-1}(x_i)$. Let θ_{y_i} be the character defined in (2). Then the set of characters $(G; \theta_{y_1}, \dots, \theta_{y_r})$ we will call the *type* of $\pi: Y \rightarrow X$.

Note that the Galois module structure of $H^1(Y, \mathcal{O}_Y)$ can be described in terms of the type. Let $\theta = (G; \theta_{y_1}, \dots, \theta_{y_r})$ be a type corresponding to some group G of order prime-to- p . Let $R = R_\theta$ be the module defined in (3). For an irreducible character χ of G , let V_χ be the G -module over k with character χ and write n_χ for the dimension of V_χ . By $f(\chi)$ we denote the smallest positive integer f such that the p^f th twist of χ is equal to χ . Write $R = \bigoplus_\chi m_\chi V_\chi$. Then for $\chi \neq 1$ the multiplicity of V_χ in $k \oplus k[G]^{g-1} \oplus R$ is $m_\chi + (g-1)n_\chi$. For each nonnegative integer g define

$$B(\theta, g) := \sum_{1 \neq \chi \in \bar{X}} f(\chi) \min_j [m_{\chi^{p^j}} + (g-1)n_\chi] \cdot n_\chi.$$

Here $\bar{X} = \bar{X}(H)$ as before corresponds to the set of characters of the irreducible $\mathbb{F}_p[H]$ -characters.

Let $Y \rightarrow X$ be a G -Galois cover with type θ and $g(X) = g$. Then, using the notation explained above, the multiplicity of a nontrivial irreducible character χ of G in $H^1(Y, \mathcal{O}_Y)$ is $m_\chi + (g-1)n_\chi$ by the result of Proposition 3.1. The result of (1) becomes with the new notation

$$\sigma(Y) - \sigma(X) \leq B(\theta, g). \tag{4}$$

QUESTION 3.3. Suppose (X, S) is the generic r -pointed curve of genus g . Let H be a group of order prime-to- p . Fix the monodromy θ . Under what conditions is it true that all H -covers with monodromy θ have p -rank equal to $B(\theta, g)$?

EXAMPLE 3.4. (i) Let E be an elliptic curve with an automorphism ϕ of order three, defined over an algebraically closed field k of characteristic not three. Then $g(E/\langle\phi\rangle) = 0$. This implies that ϕ acts nontrivially on the 1-dimensional space $H^1(E, \mathcal{O}_E)$. Let χ be the character of this representation. Then $\dim L(\chi) = 1$ and $\dim L(\chi^2) = 0$. We find

$$\sum_{\chi \in \bar{X}} f(\chi) \min_j \dim L(\chi^{-p^j}) = \begin{cases} 1 & \text{if } p \equiv 1 \pmod{3}, \\ 0 & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

Of course, it is well known that $\sigma(E) = 1$ if $p \equiv 1 \pmod{3}$ and $\sigma(E) = 0$ if $p \equiv 2 \pmod{3}$. So in this case, $\sigma(E)$ is equal to the bound.

(ii) Let E_λ be the 2-cyclic cover of \mathbb{P}^1 branched at $0, 1, \infty, \lambda$ and suppose that $p \neq 2$. The $\sigma(E_\lambda)$ is one for almost all λ and zero for finitely many λ . This illustrates that the p -rank varies in a family of covers. This explains the condition (X, S) generic in Question 3.3.

In this paper we will mainly consider Question 3.3 for cyclic covers. Here the results are quite general. The case of Abelian covers can be reduced to the case of cyclic covers. In the rest of this section we will consider the question for non-Abelian covers. We will see that the answer to the question is no, in general: there exists a non-Abelian cover $Y \rightarrow \mathbb{P}^1$ branched at three points with $\sigma(Y) < B(\theta, 0)$.

This situation is similar to the situation for étale covers. For $Y \rightarrow X$ étale, the statement of Proposition 3.1 becomes

$$H^1(Y, \mathcal{O}_Y) \simeq k \oplus k[G]^{g(X)-1},$$

as $k[G]$ -modules. Therefore the expression for the bound $B(\theta, g)$ in (4) becomes very easy:

$$B(\theta, g) = g(Y) - g(X).$$

Question 3.3 becomes in this case: are all étale Galois covers of prime-to- p order of the generic curve ordinary? Nakajima [11] has proved that this is true for all Abelian covers. Raynaud [17] has shown that there exist non-Abelian étale covers of the generic curve which are nonordinary. In fact, (some of) the nonordinary covers given in that paper have nilpotent Galois group; the nonordinary cover discussed below has a solvable Galois group.

EXAMPLE 3.5. This example was suggested to me by M. Raynaud. Consider Galois covers $\pi: Y \rightarrow \mathbb{P}_{\mathbb{Q}}^1$ branched at three points $0, 1, \infty$, with Galois group S_4 and

ramification of order 4, 4, 3. We are going to show that for infinitely many primes p the curve Y has good supersingular reduction mod p . Note that for S_n covers of \mathbb{P}^1 in characteristic $p > n$ the bound is equal to the genus, since all irreducible characters of S_n are defined over the prime field.

Let C_1 (resp. C_2) be the conjugacy class in S_4 of a 4-cycle (resp. a 3-cycle). The triple (C_1, C_1, C_2) is rational and rigid and therefore $Y \rightarrow \mathbb{P}^1$ is defined over \mathbb{Q} , [21]. Note that $g(Y) = 3$. Define $H_1 = \langle (12)(34) \rangle$, $H_2 = \langle (13)(24) \rangle$, $H_3 = \langle (14)(23) \rangle$ as subgroups of S_4 and let $E_i = Y/H_i$. The E_i are isomorphic elliptic curves defined over \mathbb{Q} . (They have j -invariant $2^4 \cdot 13^3 \cdot 3^{-2}$.) This implies that $J(Y) \sim E_1 \times E_2 \times E_3 \simeq E_1^3$. To compute the p -rank of the reduction Y_p of Y to characteristic p it suffices to compute the p -rank of $E_{1,p}$. By the result of Elkies [3] the elliptic curve E_1 has infinitely many primes of supersingular reduction. Since $J(Y) \sim E_1^3$, the same holds for Y .

This shows that there exists a p and a type θ such that for all covers $Y \rightarrow \mathbb{P}^1$ branched at $0, 1, \infty$ of type θ we have $\sigma(Y) < B(\theta, 0)$.

4. Cyclic Covers

In the previous section it is shown that there exists a non-Abelian cover of $\mathbb{P}^1 - \{0, 1, \infty\}$ for which the p -rank is strictly less than the bound. Therefore in the rest of the paper we will concentrate on the case of Abelian covers. Here our results are quite general. In this section we specialize the results of the previous two sections to Abelian covers. Note that by part (v) of Lemma 2.2 we may reduce the computation of the generalized Hasse–Witt invariants of an Abelian cover to the generalized Hasse–Witt invariants of suitable cyclic subcovers. In case of a cyclic cover the result of Chevalley–Weil becomes easier to formulate.

Let $\pi: Y \rightarrow X$ be a H -Galois cover with $p \nmid |H|$ and H cyclic of order ℓ . Note that ℓ is not supposed to be prime. We fix a generator ϕ of H and a primitive ℓ th root of unity $\zeta \in k$. The definition of type can be reformulated as follows in this case. This reformulation is less canonical, but will facilitate the formulas in what follows. Let x_1, \dots, x_r be the branch points of π . Write $n_i = |\pi^{-1}(x_i)|$. For each i , choose some $y_i \in \pi^{-1}(x_i)$ and a local parameter u_i at y_i . Let χ be the character of H which sends ϕ to ζ . We will write L_i and γ_i instead of $L(\chi^i)$ and $\gamma(\chi^i)$.

DEFINITION 4.1. We say that π is of type $(\ell; a_1, \dots, a_r)$ if for all $i \in \{1, \dots, r\}$ we have that $0 \leq a_i < \ell$ and $n_i | a_i$ and $(\phi^{n_i})^* u_i = \zeta^{n_i b_i} u_i \pmod{u_i^2}$, where $b_i \cdot a_i / n_i \equiv 1 \pmod{\ell / n_i}$.

We will suppose the branch points to be ordered. In the statement ‘ $Y \rightarrow X$ is a cover of type $\mathbf{a} = (\ell; a_1, \dots, a_r)$ branched at x_1, \dots, x_r ’ we will assume the a_i corresponds to the x_i as in the definition above.

We can interpret the type of an ℓ -cyclic cover $\pi: Y \rightarrow X$ in terms of the map $f: \pi_1^i(X - \{x_1, \dots, x_r\}) \twoheadrightarrow \mathbb{Z}/\ell$ corresponding to the cover. Choose generating

elements $\alpha_1, \dots, \alpha_g, \beta_1, \dots, \beta_g, \delta_1, \dots, \delta_r$ of $\pi_1'(X - \{x_i\})$ such that $\prod[\alpha_i, \beta_i] \prod \delta_i = 1$ and such that for each continuous $\pi_1'(X - \{x_i\}) \twoheadrightarrow G$, with G finite, the images of the δ_i generate a decomposition group above x_i ([6, XIII.2.12]). The cover π is of type $(\ell; f(\delta_1), \dots, f(\delta_r))$. If $(\ell; a_1, \dots, a_r)$ is a type, then $\sum a_i \equiv 0 \pmod{\ell}$. Conversely, if $\sum a_i \equiv 0 \pmod{\ell}$ then there exists a (not necessarily connected) cover of type $(\ell; a_1, \dots, a_r)$.

The type of a cyclic cover as defined above, contains the same information as the type defined in the previous section. The two definitions can be related as follows. Let $\pi: Y \rightarrow X$ be as above. Let y_i be some point mapping to a branch point x_i . Then ϕ^{n_i} generates the decomposition group of y_i . The character θ_{y_i} , which was defined in (2), sends ϕ^{n_i} to $\zeta^{n_i b_i}$. Conversely, given the character θ_{y_i} we find a_i as the integer $0 \leq a_i < \ell$ which is a multiple of n_i and satisfies $a_i b_i / n_i \equiv 1 \pmod{\ell / n_i}$.

The definition of type is independent of the choice of y_i and u_i . Replacing ϕ or ζ changes $(\ell; a_1, \dots, a_r)$ into $(\ell; na_1, \dots, na_r)$ for some $n \in (\mathbb{Z}/\ell)^*$. Hence, we should consider the type as an element of

$$\{(a_1, \dots, a_r) \mid 0 \leq a_i < \ell, \sum a_i \equiv 0 \pmod{\ell}\} / (\mathbb{Z}/\ell)^*.$$

We will always choose a representative of such a class.

EXAMPLE 4.2. Suppose $g(X) = 0$ and choose $x_1, \dots, x_r \in X$. If $\sum a_i \equiv 0 \pmod{\ell}$ then, up to isomorphism, there is a unique cover $Y \rightarrow \mathbb{P}^1$ of type $(\ell; a_1, \dots, a_r)$ with Y nonsingular, branched at $x_1, \dots, x_r \in \mathbb{P}_k^1$. If none of the x_i is ∞ , this curve is the nonsingular curve associated to the equation

$$y^\ell = (x - x_1)^{a_1} (x - x_2)^{a_2} \cdots (x - x_r)^{a_r}.$$

The curve Y is connected iff $\gcd(\ell, a_1, \dots, a_r) = 1$.

LEMMA 4.3. *Let $\pi: Y \rightarrow X$ be of type $(\ell; a_1, \dots, a_r)$. Then*

$$\dim_k L_i = \begin{cases} g(X), & \text{if } i = 0, \\ \left(\sum_{j=1}^r \left\langle \frac{ia_j}{\ell} \right\rangle \right) - 1 + g(X), & \text{otherwise.} \end{cases}$$

Here $\langle \cdot \rangle$ denotes the fractional part.

Proof. The lemma follows from Proposition 3.1; in a slightly different terminology, it is a special case of [8, Proposition 1]. But we can also prove it directly.

We can write $\pi_* \mathcal{O}_Y = \bigoplus_{i=0}^{\ell-1} \mathcal{L}_i$, where locally \mathcal{L}_i is the eigenspace of ϕ with eigenvalue ζ^i . This is Kummer theory. Choose $y_j \in \pi^{-1}(x_j)$ for each j . Let u_j be a local parameter at y_j such that $\phi^{n_j}(u_j) = \zeta^{n_j b_j} u_j$, with notation as in Definition 4.1. Then, by the definition of type, $\pi_*((u_j \phi(u_j) \cdots \phi^{n_j-1}(u_j))^{(ia_j/\ell)n_j})$ generates \mathcal{L}_i locally

around x_j . It follows that

$$\mathcal{L}_i^{\otimes \ell} \simeq \mathcal{O}_X \left(- \sum_{j=1}^r \left\langle \frac{ia_j}{\ell} \right\rangle \ell x_j \right).$$

In fact $\mathcal{L}_i = \mathcal{L}^i$. Hence,

$$\deg \mathcal{L}_i = - \frac{1}{\ell} \sum_{j=1}^r \left\langle \frac{ia_j}{\ell} \right\rangle \ell.$$

Note that $H^1(X, \mathcal{L}_i) \simeq L_i$. We find that

$$\dim_k L_i = \dim_k H^1(X, \mathcal{L}_i) = \left(\sum_{j=1}^r \left\langle \frac{ia_j}{\ell} \right\rangle \right) + g(X) - 1,$$

by the Riemann–Roch Theorem. □

NOTATION 4.4. For a type $\mathbf{a} = (\ell; a_1, \dots, a_r)$ we will denote

$$\|i\| = \|i\|_{\mathbf{a}} = \left(\sum_{j=1}^r \left\langle \frac{ia_j}{\ell} \right\rangle \right) - 1 \quad \text{for } i \in (\mathbb{Z}/\ell) - \{0\}.$$

In the case $\|i\|_{\mathbf{a}} \geq 0$, it is equal to the dimension of the i th eigenspace L_i of $H^1(Y, \mathcal{O}_Y)$, where $Y \rightarrow \mathbb{P}^1$ is a cover of type \mathbf{a} . If all a_j 's are zero (mod $\ell/(i, \ell)$) then $\|i\|_{\mathbf{a}} = -1$ and then $\|i\|$ does not have an interpretation as a dimension of an isotypical space of a cover of \mathbb{P}^1 , but we will use the notation in this case nonetheless.

LEMMA 4.5. Fix a type $(\ell; a_1, \dots, a_r)$. Let s be the number of a_j unequal to $0 \pmod{\ell/(i, \ell)}$. Then

$$\begin{aligned} \|i\| + \|\ell - i\| &= s - 2, \quad \text{if } \ell/(i, \ell) \neq 2, \\ \|i\| &= \frac{s}{2} - 1 \quad \text{if } \ell/(i, \ell) = 2. \end{aligned}$$

In particular, $\|i\| \leq s - 2$.

Proof. Immediate. □

LEMMA 4.6. Let \mathbf{a} be a type and X any curve. Denote by $\Delta_r \subset X^r$ the generalized diagonal. Define $U_{\mathbf{a}} \subset X^r - \Delta$ as the set of $(x_1, \dots, x_r) \in X^r - \Delta$ such that for all ℓ -cyclic covers $Y \rightarrow X$, unbranched outside the x_i , of type \mathbf{a} , we have $\sigma(Y) - \sigma(X) = B(\mathbf{a}, g(X))$. Then $U_{\mathbf{a}}$ is open.

Proof. For each $(x_1, \dots, x_r) \in X^r - \Delta$ we define $Y_{x_1, \dots, x_r} \rightarrow X$ as the smallest cover which has all covers of X of type \mathbf{a} branched at the x_i as subcovers. Choose generators $\alpha_1, \dots, \alpha_g, \beta_1, \dots, \beta_g, \delta_1, \dots, \delta_r$ of $\pi_1^t(X - \{x_1, \dots, x_r\})$ such that for each

finite quotient the images of the δ_i generate a decomposition group above x_i . The cover $Y_{x_1, \dots, x_r} \rightarrow X$ corresponds to the maximal quotient

$$\pi'_1(X - \{x_1, \dots, x_r\}) \twoheadrightarrow (\mathbb{Z}/\ell)^n$$

which sends δ_i to a_i, \dots, a_i . For suitable ℓ -cyclic subcovers $Y_j \rightarrow X$ of $Y_{x_1, \dots, x_r} \rightarrow X$ we have

$$J(Y_{x_1, \dots, x_r}) \sim J(X) \times \prod_{j=1}^n (J(Y_j)/J(X)).$$

Here, for a curve Z we denote by $J(Z)$ its Jacobian. This implies

$$\sigma(Y_{x_1, \dots, x_r}) \leq \sigma(X) + nB(\mathbf{a}, g(X)) =: s.$$

We obtain a family $f: \mathcal{A} \rightarrow X^r - \Delta_r$, with $\mathcal{A}_{x_1, \dots, x_r} = J(Y_{x_1, \dots, x_r})$. Let $W_{\mathbf{a}} = X^r - (\Delta_r \cup U_{\mathbf{a}}) = \{(x_1, \dots, x_r) \mid \sigma(Y_{x_1, \dots, x_r}) \leq s - 1\}$. From [15, Cor. 1.5] it follows that $W_{\mathbf{a}}$ is closed. This proves the lemma. \square

5. Coefficients of the Hasse–Witt Matrix

Let $\pi: Y \rightarrow \mathbb{P}^1$ be an ℓ -cyclic cover, defined over an algebraically closed field k of characteristic p with $(\ell, p) = 1$. In this section we will compute the coefficients of the Hasse–Witt matrix of Y and relate these coefficients to the generalized Hasse–Witt invariants. We fix the following notations. The branch points of π we will denote by x_1, \dots, x_r , and $\mathbf{a} = (\ell; a_1, \dots, a_r)$ will be the type of the cover π (defined in Section 4). We choose a coordinate on \mathbb{P}^1 such that none of the x_i is ∞ . We will suppose that none of the a_i is congruent to zero mod ℓ . We denote by L_i the i th eigenspace of $H^1(Y, \mathcal{O}_Y)$ with respect to a fixed primitive ℓ th root of unity ζ and a fixed generator ϕ of $\text{Gal}(Y, \mathbb{P}^1)$ as in Section 4, i.e. $L_i = \{\xi \mid \phi \cdot \xi = \zeta^i \xi\}$. The dimension of L_i we denote by $\|i\|$; it is equal to $(\sum_{j=1}^r \langle ia_j/\ell \rangle) - 1$. The integer f denotes the order of p in $(\mathbb{Z}/\ell)^*$. Note $F^f: L_i \rightarrow L_i$. Put $m = (p^f - 1)/\ell$. The curve Y is the nonsingular projective curve defined by the equation

$$y^\ell = (x - x_1)^{a_1} (x - x_2)^{a_2} \dots (x - x_r)^{a_r}.$$

We are going to calculate the coefficients of the matrix of F^f on L_i . We will compute them using Čech cohomology; we first describe a basis of $H^1(Y, \mathcal{O}_Y)$ (cf. [22, Section 5], note that the $\|i\|$ from that paper is +1 more than our $\|i\|$). A special case of this result is proved in [4], using a different method. Write $U_1 = \mathbb{P}^1 - \{\infty\}$ and $U_2 = \mathbb{P}^1 - \{0\}$. As open sets we take $V_s = \pi^{-1}(U_s) \subset Y$ for $s = 1, 2$. For $i = 1, \dots, \ell - 1$ let

$$v_i = y^i (x - x_1)^{-\lfloor \frac{a_1 i}{\ell} \rfloor} \dots (x - x_r)^{-\lfloor \frac{a_r i}{\ell} \rfloor}.$$

Note

$$\begin{aligned} \Gamma(V_1, \mathcal{O}_{V_1}) &= \bigoplus_{i=1}^{\ell-1} k[x] v_i, \\ \Gamma(V_2, \mathcal{O}_{V_2}) &= \bigoplus_{i=1}^{\ell-1} k[x^{-1}] x^{-\|i\|-1} v_i, \\ \Gamma(V_1 \cap V_2, \mathcal{O}) &= \bigoplus_{i=1}^{\ell-1} k[x, x^{-1}] v_i. \end{aligned}$$

Define

$$\xi_{i,j} = x^{-j} v_i \quad \text{for } j = 1, \dots, \|i\|$$

as an element of $H^1(Y, \mathcal{O}_Y) = \Gamma(V_1 \cap V_2) / (\Gamma(V_1) + \Gamma(V_2))$. Then the $\{\xi_{i,j} | 0 < i < \ell, 0 < j \leq \|i\|\}$ form a basis of $H^1(Y, \mathcal{O}_Y)$. More precisely, $L_i = \langle \xi_{i,j} \rangle$. Let B_i be the matrix of $F^f : L_i \rightarrow L_i$ with respect to the basis $\xi_{i,j}$, for $1 \leq j \leq \|i\|$. We will compute the coefficients of B_i . To calculate the (j, j') th coefficient of B_i we have to find the coefficient of $\xi_{i,j'}$ in $(\xi_{i,j})^{p^f}$. We have

$$(\xi_{i,j})^{p^f} = x^{-jp^f} y^{ip^f} (x - x_1)^{-p^f \lfloor \frac{ia_1}{\ell} \rfloor} \dots (x - x_r)^{-p^f \lfloor \frac{ia_r}{\ell} \rfloor}.$$

Note

$$-p^f \lfloor ia/\ell \rfloor = -[p^f ia/\ell] + [p^f (ia/\ell)] = -[ia/\ell] - iam + \ell m \langle ia/\ell \rangle$$

and

$$y^\ell (x - x_1)^{-a_1} \dots (x - x_r)^{-a_r} = 1.$$

Hence

$$(\xi_{i,j})^{p^f} = (x)^{-jp^f} (x - x_1)^{\ell m \langle \frac{ia_1}{\ell} \rangle} \dots (x - x_r)^{\ell m \langle \frac{ia_r}{\ell} \rangle} v_i. \tag{5}$$

The coefficient of $\xi_{i,j'}$ in $(\xi_{i,j})^{p^f}$ is

$$(-1)^N \sum_{n_1 + \dots + n_r = N} \binom{m \ell \langle \frac{ia_1}{\ell} \rangle}{n_1} \dots \binom{m \ell \langle \frac{ia_r}{\ell} \rangle}{n_r} x_1^{n_1} \dots x_r^{n_r}, \tag{6}$$

where $N = (\|i\| + 1 - j)\ell m + j' - j$. This proves the first part of the following lemma. The other part is proved analogously.

LEMMA 5.1. Let $(\xi_{i,j})_{i,j}$ be the basis of $H^1(Y, \mathcal{O}_Y)$ which was described above.

(i) The $(i, j), (i, j')$ th coefficient of the matrix of F^f on $H^1(Y, \mathcal{O}_Y)$ is

$$(-1)^N \sum_{n_1 + \dots + n_r = N} \binom{m \ell \langle \frac{ia_1}{\ell} \rangle}{n_1} \dots \binom{m \ell \langle \frac{ia_r}{\ell} \rangle}{n_r} x_1^{n_1} \dots x_r^{n_r}, \tag{7}$$

where $N = (\|i\| + 1 - j)\ell m + j' - j$.

(ii) If $i' \equiv pi \pmod{\ell}$, then the $(i, j), (i', j')$ th coefficient of the Hasse–Witt matrix of Y is equal to

$$(-I)^N \sum_{n_1 + \dots + n_r = N} \binom{[p^{(ia_1)/\ell}]}{n_1} \dots \binom{[p^{(ia_r)/\ell}]}{n_r} x_1^{n_1} \dots x_r^{n_r}, \tag{8}$$

where $N = p(\|i\| + 1 - j) - (\|i'\| + 1 - j')$. If $i' \not\equiv pi \pmod{\ell}$, then the $(i, j), (i', j')$ th coefficient of the Hasse–Witt matrix is zero.

6. Main Theorem

THEOREM 6.1. *Suppose $p \geq \ell(r - 3)$. Then for each $\mathbf{a} = (\ell; a_1, \dots, a_r)$ there exist x_1, \dots, x_r such that for the ℓ -cyclic cover $Y \rightarrow \mathbb{P}^1$ of type \mathbf{a} branched at x_1, \dots, x_r we have $\sigma(Y) = B(\mathbf{a}, 0)$.*

To prove the theorem, we have to prove that there exist $x_1, \dots, x_r \in \mathbb{P}_k^1$ such that for every $1 \leq a \leq \ell$ we have $\gamma_{-a} = \min_i \|p^i a\|$. It is no restriction to suppose that $a = 1$ and $n := \|1\| \leq \|p^i\|$ for all i . Let B be the matrix of $F^f : L_1 \rightarrow L_1$ with respect to the basis $\zeta_{1,j}$. Let A be the matrix of $F : \bigoplus_{i=0}^{f-1} L_{p^i} \rightarrow \bigoplus_{i=0}^{f-1} L_{p^i}$ and A_i the matrix of $F : L_{p^i} \rightarrow L_{p^{i+1}}$. To prove the theorem, we will prove that $\det(B)$ is not identically zero as polynomial in the x_i . We will prove this by showing that a certain monomial occurs in $\det(B)$ with a nonzero coefficient. The strategy of the proof is the following. We define an ordering on monomials in x_1, \dots, x_r . We give an expression for $\det(B)$ as a sum over an index set \mathcal{J} of terms which are products of determinants of $n \times n$ submatrices of the A_i (Lemma 6.2). We find for each $J \in \mathcal{J}$ the largest monomial T_J with respect to the ordering (Lemma 6.5). We conclude the proof by showing that for $J \neq J'$ we have $T_J \neq T_{J'}$ (Lemma 6.6).

Recall that each L_{p^i} has a basis $\zeta_{p^i,j}$ with $0 < j \leq \|p^i\|$ (see the previous section). Number the rows and columns of A as (i, j) with $0 \leq i \leq f - 1$ and $1 \leq j \leq \|p^i\|$. Here (i, j) corresponds to the basis vector $\zeta_{p^i,j}$ of L_{p^i} . Put $I = \{(i, j) \mid 0 \leq i \leq f - 1, 1 \leq j \leq \|p^i\|\}$. Let \mathcal{J} be the set of $J \subset I$ such that J contains for each i exactly n indices $(i, j_i(x))$ and $j_i(1) < j_i(2) < \dots < j_i(n)$. For $i = 0$, we have $j_0(1) = 1, \dots, j_0(n) = n$. To a $J \in \mathcal{J}$ we associate for each i a matrix $M_{J,i}$. This $M_{J,i}$ is the $n \times n$ submatrix of A_i consisting of the columns $(i, j_i(1)), \dots, (i, j_i(n))$ and the rows $(i + 1, j_{i+1}(1)), \dots, (i + 1, j_{i+1}(n))$. For $i = f - 1$, the matrix $M_{J,i}$ has columns $(f - 1, j_{f-1}(1)), \dots, (f - 1, j_{f-1}(n))$ and rows $(0, j_0(1) = 1), \dots, (0, j_0(n) = n)$.

LEMMA 6.2. *We have*

$$|B| = \sum_{J \in \mathcal{J}} \left(\prod_{i=0}^{f-1} |M_{J,i}|^{p^{f-1-i}} \right).$$

Proof. The lemma follows from the fact that

$$B = A_{f-1}A_{f-2}^{(p)} \cdots A_0^{(p^{f-1})},$$

here $A^{(p)}$ is the matrix obtained by raising all the coefficients of A to the p th power. \square

NOTATION 6.3. Define an ordering on monomials $x_1^{n_1} \cdots x_r^{n_r}$ by $x_1^{n_1} \cdots x_r^{n_r} > x_1^{k_1} \cdots x_r^{k_r}$ if there exists an i such that $k_j = n_j$ for all $j < i$ and $n_i > k_i$.

In the following lemma we find the largest monomial in $|M_{J,i}|$ (cf. Lemma 6.2). Let $M = M_{J,i}$. Recall that the columns of M are $(i, j_i(\alpha))$ and the rows are $(i + 1, j_{i+1}(\beta))$, where $1 \leq \alpha, \beta \leq n$ and $1 \leq j_i(1) < \cdots < j_i(n) \leq \|p^i\|$ and $1 \leq j_{i+1}(1) < \cdots < j_{i+1}(n) \leq \|p^{i+1}\|$. If J and i are understood we can number the rows and columns by β and α .

The coefficients of $M_{J,i}$ are

$$m_{\beta,\alpha}^{J,i} = (-1)^{N_{\beta,\alpha}^{J,i}} \sum_{n_1 + \cdots + n_r = N_{\beta,\alpha}^{J,i}} \binom{\lfloor p \langle \frac{p^i a_1}{\ell} \rangle \rfloor}{n_1} \cdots \binom{\lfloor p \langle \frac{p^i a_r}{\ell} \rangle \rfloor}{n_r} x_1^{n_1} \cdots x_r^{n_r},$$

with $N_{\beta,\alpha}^{J,i} = p(\|p^i\| + 1 - j_i(\alpha)) - (\|p^{i+1}\| + 1 - j_{i+1}(\beta))$, by the second part of Lemma 5.1. The following notation is introduced to describe the largest monomial $T_{J,i}$ in $|M_{J,i}|$.

NOTATION 6.4. Define $c_{\beta,\alpha}^{J,i}$ as the smallest number c in $\{1, \dots, r\}$ such that

$$\left\lfloor p \left\langle \frac{p^i a_1}{\ell} \right\rangle \right\rfloor + \cdots + \left\lfloor p \left\langle \frac{p^i a_c}{\ell} \right\rangle \right\rfloor > N_{\beta,\alpha}^{J,i}.$$

And let

$$C_{\beta,\alpha}^{J,i} = N_{\beta,\alpha}^{J,i} - \left(\left\lfloor p \left\langle \frac{p^i a_1}{\ell} \right\rangle \right\rfloor + \cdots + \left\lfloor p \left\langle \frac{p^i a_{c-1}}{\ell} \right\rangle \right\rfloor \right),$$

where $c = c_{\beta,\alpha}^{J,i}$.

Note that

$$\left\lfloor p \left\langle \frac{p^i a_1}{\ell} \right\rangle \right\rfloor + \cdots + \left\lfloor p \left\langle \frac{p^i a_r}{\ell} \right\rangle \right\rfloor - N_{\beta,\alpha}^{J,i} = pj_i(\alpha) - j_{i+1}(\beta) \geq p - \|p^{i+1}\| \geq 0,$$

since $j_{i+1}(\beta) \leq \|p^{i+1}\| \leq r - 2$ (Lemma 4.5) and $p \geq r - 2$ (by assumption). This implies that $c_{\beta,\alpha}^{J,i}$ is well defined. Moreover, $0 \leq C_{\beta,\alpha}^{J,i} < \lfloor p \langle p^i a_c / \ell \rangle \rfloor$.

LEMMA 6.5. *The largest monomial $T_{J,i}$ in $|M_{J,i}|$ is the largest monomial in the product of the diagonal elements of $M_{J,i}$.*

Proof. In this proof we fix J and i and drop them from the notation. The largest monomial in $m_{\beta,\alpha}$, which we denote by $T_{\beta,\alpha}$, is

$$T_{\beta,\alpha} := (-1)^{N_{\beta,\alpha}} \binom{[p\langle \frac{p^i a_c}{\ell} \rangle]}{C} x_1^{[p\langle \frac{p^i a_1}{\ell} \rangle]} \cdots x_{c-1}^{[p\langle \frac{p^i a_{c-1}}{\ell} \rangle]} x_c^C. \tag{9}$$

Here $c = c_{\beta,\alpha}$ and $C = C_{\beta,\alpha}$. Note that $0 \leq C_{\beta,\alpha} < [p\langle p^i a_c / \ell \rangle] < p$, therefore the coefficient of $T_{\beta,\alpha}$ is unequal to zero. One checks that

$$c_{\beta,\alpha} > c_{\beta,\gamma}, \quad \text{if } \alpha < \gamma, \tag{10}$$

$$c_{\beta,\alpha} \leq c_{\gamma,\alpha}, \quad \text{if } \beta < \gamma. \tag{11}$$

We will prove the lemma by induction. For $n = 1$ there is nothing to prove. We will prove the lemma for $n = 2$.

Choose $\alpha < \alpha'$ and $\beta < \beta'$. We want to compare $T_{\beta,\alpha} T_{\beta',\alpha'}$ and $T_{\beta',\alpha} T_{\beta,\alpha'}$. To determine which of the two is smaller, we have to look at the smallest $1 \leq s \leq r$ where the power of x_s in one of the monomials is less than $[p\langle p^i a_s / \ell \rangle]$. From (10) and (11) it follows that this happens for $s = c_{\beta,\alpha'}$. One checks that in case $c_{\beta,\alpha'} = c_{\beta',\alpha'}$, then $C_{\beta,\alpha'} - C_{\beta',\alpha'} = N_{\beta,\alpha'} - N_{\beta',\alpha'} < 0$. This implies that

$$T_{\beta,\alpha} T_{\beta',\alpha'} > T_{\beta',\alpha} T_{\beta,\alpha'}.$$

This proves the lemma for $n = 2$, by taking $\alpha = \beta = 1$ and $\alpha' = \beta' = 2$.

For $n > 2$ we have

$$|M| = \sum_{\alpha=1}^n (-1)^{\alpha+1} m_{1\alpha} |M_{1\alpha}|, \tag{12}$$

where $M_{1\alpha}$ is the minor of M obtained by omitting the α th column and the first row. Therefore, by the induction hypothesis

$$T_\alpha := T_{1\alpha} T_{21} \cdots T_{\alpha,\alpha-1} T_{\alpha+1,\alpha+1} \cdots T_{nn} \text{ for } \alpha = 1, \dots, n$$

is the largest monomial in $m_{1\alpha} |M_{1\alpha}|$ (cf. (12)). One checks that

$$T_\alpha - T_{\alpha+1} = T_{21} \cdots T_{\alpha,\alpha-1} T_{\alpha+2,\alpha+2} \cdots T_{nn} (T_{1\alpha} T_{\alpha+1,\alpha+1} - T_{1\alpha+1} T_{\alpha+1,\alpha}).$$

Note that $T_{1\alpha} T_{\alpha+1,\alpha+1} > T_{1,\alpha+1} T_{\alpha+1,\alpha}$ by the $n = 2$ case applied to the minor of M consisting of the rows 1 and $\alpha + 1$ and the columns α and $\alpha + 1$. As remarked above, the coefficients of the T_{xy} do not vanish. This implies that $T_\alpha > T_{\alpha+1}$, and hence that the largest monomial $T_{J,i}$ in $|M_{J,i}|$ is T_1 . □

The lemma gives an expression for

$$T_J = \prod_{i=0}^{f-1} T_{J,i}^{p^{f-1-i}}. \tag{13}$$

Note, T_J is not identically zero as polynomial in the x_i . To finish the proof of the

Main Theorem, we have to find a nonvanishing monomial in the expression for $\det(B)$ given in Lemma 6.2. This is done if for $J_1 \neq J_2$ we show that the monomials T_{J_1} and T_{J_2} are nonidentical. In that case the largest of the T_J does not cancel in $\det(B)$. This is proved in the next lemma. This is the only place where we use the assumption $p \geq \ell(r - 3)$. Up to now we only used $p \geq r - 2$.

LEMMA 6.6. *Suppose $p \geq \ell(r - 3)$. Let $J_1, J_2 \in \mathcal{J}$. If T_{J_1} and T_{J_2} are equal up to a (nonzero) constant, then $J_1 = J_2$.*

Proof. We want to compare T_J for different J . In the previous lemma we showed that T_J is a product of powers of $T_{j_i(\alpha), j_{i+1}(\alpha)}$, (see (13)). Note that the monomials $T_{j_i(\alpha), j_{i+1}(\alpha)}$ actually depend on i , but we suppressed this in the notation; in this proof we will therefore refer to these monomials as $T_{\cdot, \cdot}^i$. An expression for $T_{j_i(\alpha), j_{i+1}(\alpha)}^i$ is given in (9). Note that now β will be always equal to α in (9); therefore we will suppress β from the notation of $c_{\beta, \alpha}^{J, i}, C_{\beta, \alpha}^{J, i}, N_{\beta, \alpha}^{J, i}$.

CLAIM. *Suppose $p \geq \ell(r - 3)$. Then $c_{\alpha+1}^{J, i} < c_{\alpha}^{J, i}$.*

Proof. Note that

$$\begin{aligned} N_{\alpha}^{J, i} &= p(\|p^i\| + 1 - j_i(\alpha)) - (\|p^{i+1}\| + 1 - j_{i+1}(\alpha)) \\ &= \left[p \left\langle \frac{p^i a_1}{\ell} \right\rangle \right] + \dots + \left[p \left\langle \frac{p^i a_r}{\ell} \right\rangle \right] + j_{i+1}(\alpha) - pj_i(\alpha). \end{aligned}$$

Therefore the integers $c_{\alpha}^{J, i}$, defined in Notation 6.4, may also be defined for given J, i, α as the unique integer c such that

$$\left[p \left\langle \frac{p^i a_{c+1}}{\ell} \right\rangle \right] + \dots + \left[p \left\langle \frac{p^i a_r}{\ell} \right\rangle \right] < pj_i(\alpha) - j_{i+1}(\alpha) \leq \left[p \left\langle \frac{p^i a_c}{\ell} \right\rangle \right] + \dots + \left[p \left\langle \frac{p^i a_r}{\ell} \right\rangle \right].$$

Suppose $p \geq \ell(r - 3)$, then $\|p^{i+1}\| \leq r - 2 \leq p/\ell + 1$. Since

$$\begin{aligned} pj_i(\alpha + 1) - j_{i+1}(\alpha + 1) &\geq p(j_i(\alpha) + 1) - \|p^{i+1}\| \\ &> pj_i(\alpha) > pj_i(\alpha) - j_{i+1}(\alpha) \end{aligned}$$

we have $c_{\alpha+1}^{J, i} \leq c_{\alpha}^{J, i}$. Suppose $c_{\alpha+1}^{J, i} = c_{\alpha}^{J, i} = c$, i.e.

$$\left[p \left\langle \frac{p^i a_{c+1}}{\ell} \right\rangle \right] + \dots + \left[p \left\langle \frac{p^i a_r}{\ell} \right\rangle \right] < pj_i(\beta) - j_{i+1}(\beta) \leq \left[p \left\langle \frac{p^i a_c}{\ell} \right\rangle \right] + \dots + \left[p \left\langle \frac{p^i a_r}{\ell} \right\rangle \right]$$

for $\beta = \alpha, \alpha + 1$. Then

$$pj_i(\alpha + 1) - j_{i+1}(\alpha + 1) - pj_i(\alpha) + j_{i+1}(\alpha) < \left[p \left\langle \frac{p^i a_c}{\ell} \right\rangle \right]. \tag{14}$$

We assumed that

$$1 \leq j_x(\alpha) \leq j_x(\alpha + 1) - 1 \quad \text{for any } x.$$

Hence

$$p(j_i(\alpha + 1) - j_i(\alpha)) - j_{i+1}(\alpha + 1) + j_{i+1}(\alpha) > p - j_{i+1}(\alpha + 1) + 1. \tag{15}$$

Equations (14) and (15) imply that $[p\langle p^i a_c / \ell \rangle] > p - j_{i+1}(\alpha + 1) + 1$. Hence

$$\begin{aligned} j_{i+1}(\alpha + 1) &> p + 1 - \left[p \left\langle \frac{p^i a_c}{\ell} \right\rangle \right] = p + 1 - p \left\langle \frac{p^i a_c}{\ell} \right\rangle + \left\langle \frac{p^{i+1} a_c}{\ell} \right\rangle \\ &\geq p + 1 - p \left(\frac{\ell - 1}{\ell} \right) + \frac{1}{\ell} \\ &= p + 1 - p + \frac{p + 1}{\ell} = 1 + \frac{p + 1}{\ell} > 1 + \frac{p}{\ell} \geq \|p^{i+1}\|. \end{aligned}$$

This is impossible since $j_{i+1}(\alpha + 1) \leq \|p^{i+1}\|$. Hence $c_{\alpha+1}^{J,i} < c_{\alpha}^{J,i}$. This proves the claim.

Let J_1 and J_2 be as in the statement of Lemma 6.6. We will write $x_1^{n_1} \dots x_r^{n_r} \sim_s x_1^{k_1} \dots x_r^{k_r}$ if $k_t = n_t$ for all $t \leq s$, where we ignore constants. Write $J_1 = \{j_i(\alpha)\}$ and $J_2 = \{k_i(\alpha)\}$.

CLAIM. *The following two statements are equivalent:*

- (1) $T_{J_1} \sim_s T_{J_2}$ for s in $\{1, \dots, r\}$,
- (2) for each α and $t \leq s$ the sets $\{i \mid c_{\alpha}^{J_1,i} = t\}$ and $\{i \mid c_{\alpha}^{J_2,i} = t\}$ are equal. Furthermore, for each i, α with $c_{\alpha}^{J_1,i} = s$ we have that $p j_i(\alpha) - j_{i+1}(\alpha) = p k_i(\alpha) - k_{i+1}(\alpha)$.

Proof. The second statement implies the first statement. Suppose that $T_{J_1} \sim_s T_{J_2}$ and suppose that the second statement holds for all $t < s$. For $s = 1$ this is an empty assumption. If $c_{\alpha}^{J_1,i} < s$ then $c_{\alpha}^{J_2,i} < s$, by assumption. Since $c_{\alpha+1}^{J,i} < c_{\alpha}^{J,i}$, there is an α_0 such that

$$\begin{aligned} c_{\beta}^{J_1,i} &< s \quad \text{for all } \beta > \alpha_0, \\ c_{\alpha_0}^{J_1,i} &\geq s. \end{aligned}$$

If $c_{\beta}^{J,i} < s$ for all $\beta \in \{1, \dots, n\}$, put $\alpha_0 = 0$. Write

$$T_{J_x} = (\text{constant}) x_1^{n_1(J_x)} \dots x_r^{n_r(J_x)} \quad \text{for } x = 1, 2.$$

The assumption that $J_1 \sim_s J_2$ implies that $n_t(J_1) = n_t(J_2)$ for $t \leq s$. Consider $\sum_{i=0}^{f-1} p^{f-1-i} \eta_{i,s}$ with

$$\eta_{i,s} = \begin{cases} 0, & \text{if } \alpha_0 = 0, \\ 0, & \text{if } c_{\alpha_0}^{J_1,i} > s \text{ and } c_{\alpha_0}^{J_2,i} > s, \\ C_{\alpha_0}^{J_1,i} - \left[p \left\langle \frac{p^i a_s}{\ell} \right\rangle \right] & \text{if } c_{\alpha_0}^{J_1,i} = s \text{ and } c_{\alpha_0}^{J_2,i} > s, \\ \left[p \left\langle \frac{p^i a_s}{\ell} \right\rangle \right] - C_{\alpha_0}^{J_2,i} & \text{if } c_{\alpha_0}^{J_1,i} > s \text{ and } c_{\alpha_0}^{J_2,i} = s, \\ C_{\alpha_0}^{J_1,i} - C_{\alpha_0}^{J_2,i} & \text{if } c_{\alpha_0}^{J_1,i} = c_{\alpha_0}^{J_2,i} = s. \end{cases} \tag{16}$$

The sum $\sum_{i=0}^{f-1} p^{f-1-i} \eta_{i,s}$ is equal to $n_s(J_1) - n_s(J_2)$. Note that

$$-p < -\left[p \left\langle \frac{p^i a_s}{\ell} \right\rangle \right] \leq \eta_{i,s} \leq \left[p \left\langle \frac{p^i a_s}{\ell} \right\rangle \right] < p.$$

Let i_0 be the smallest i such that $\eta_{i,s} \neq 0$, then

$$\eta_{i_0,s} < 0 \implies n_s(J_1) > n_s(J_2)$$

and

$$\eta_{i_0,s} > 0 \implies n_s(J_1) < n_s(J_2).$$

(This is seen by noting that if $\eta_{i_0,s} \geq 1$, then

$$\begin{aligned} n_s(J_1) - n_s(J_2) &\geq p^{f-1-i_0} - \sum_{i=i_0+1}^{f-1} p^{f-1-i} (p-1) \\ &= p^{f-1-i_0} - p^{f-1-i_0} + p^{f-2-i_0} \dots + p - p + 1 = 1. \end{aligned}$$

For the other inequality, interchange J_1 and J_2 .) Our assumptions imply that $n_s(J_1) = n_s(J_2)$. In particular, $\eta_{i,s} = 0$ for all i . Looking at the explicit formula (16) for $\eta_{i,s}$ yields that one of the following holds

$$\begin{aligned} \alpha_0 &= 0, \\ c_{\alpha_0}^{J_1,i} > s \quad \text{and} \quad c_{\alpha_0}^{J_2,i} > s, \\ c_{\alpha_0}^{J_1,i} = c_{\alpha_0}^{J_2,i} = s \quad \text{and} \quad C_{\alpha_0}^{J_1,i} = C_{\alpha_0}^{J_2,i}. \end{aligned}$$

This last condition implies

$$pj_i(\alpha) - j_{i+1}(\alpha) = pk_i(\alpha) - k_{i+1}(\alpha).$$

This proves the claim.

For each J and each i, α there exists an $s \leq r$ with $c_{\alpha}^{J,i} = s$, as was remarked below Notation 6.4. Hence, the claim implies that if $T_{J_1} \sim_r T_{J_2}$ then for each i, α we have

$$pj_i(\alpha) - j_{i+1}(\alpha) = pk_i(\alpha) - k_{i+1}(\alpha).$$

Furthermore, we have $1 \leq j_0(1) < \dots < j_0(n) \leq \|p^0\| = n$, hence $j_0(\alpha) = \alpha$. The same holds for the $k_0(\alpha)$. This shows that $j_i(\alpha) = k_i(\alpha)$ for all i, α . We conclude $J_1 = J_2$. This finishes the proof of the lemma. \square

Proof of the theorem. Consider the set of all $J \in \mathcal{J}$ for which T_J is maximal. Lemma 6.6 implies that this set consists of one element, i.e. there is a unique $J' \in \mathcal{J}$ such that $T_{J'}$ is maximal among the T_J 's. Therefore, in the expression for $\det(B)$ given in Lemma 6.2, the term $T_{J'}$ does not cancel. Furthermore,

$\deg(T'_j) \neq 0$. This implies that $\det(B)$ is a polynomial in x_1, \dots, x_r of positive degree. \square

It is not clear to me if the condition p large in the Main Theorem is really necessary. I do not know of any counter example for small p . In the case that $\|i\| = 1$ the proof can be simplified considerably, and we can improve on the condition on p .

PROPOSITION 6.7. *Fix a type $\mathbf{a} = (\ell; a_1, \dots, a_r)$. Suppose $p \geq r - 2$. Suppose furthermore that $\|i\| = 1$ and $\|ip^j\| \geq 1$ for all j and some i . Then there exist x_1, \dots, x_r such that $\gamma_{-i} = 1$.*

Proof. We will use the same notation as before. It is no restriction to suppose that $i = 1$. We have $\mathcal{J} = \{(i, j_i) | 0 \leq i < f, 1 \leq j_i \leq \|p^i\|\}$. To prove the lemma, we have to show that the matrix of $F^f : L_1 \rightarrow L_1$ is invertible, for some choice of the branch points x_1, \dots, x_r . By assumption L_1 is one dimensional. The matrix is given by

$$\sum_{J \in \mathcal{J}} \prod_{i=0}^{f-1} b_{J,i}^{p^{f-1-i}},$$

with

$$b_{J,i} = (-1)^N \sum_{n_1 + \dots + n_r = N} \binom{\lfloor p \langle \frac{p^i a_1}{\ell} \rangle \rfloor}{n_1} \dots \binom{\lfloor p \langle \frac{p^i a_r}{\ell} \rangle \rfloor}{n_r} x_1^{n_1} \dots x_r^{n_r}.$$

Here $N = p(\|p^i\| + 1 - j_i) - (\|p^{i+1}\| + 1 - j_{i+1})$. As before, the $b_{J,i}$ are not identically zero as polynomial in the x_i . The proof that for $J_1 \neq J_2$ the largest of $b_{J_1,i}$ does not cancel against the largest monomial of $b_{J_2,i}$ is obvious in this case. We do not need the assumption $p \geq \ell(r - 3)$, which was needed in the general case. \square

COROLLARY 6.8. *Let $Y \rightarrow \mathbb{P}^1$ be a cover of type \mathbf{a} branched at three points. Then $\sigma(Y) = B(\mathbf{a}, 0)$.*

Proof. This follows immediately from the above proposition, since Lemma 4.5 implies that the dimension of the eigenspaces are at most one in this case. \square

This result can also be deduced from a result of Yui [24] by using the formula for the zeta function of Y , where $Y \rightarrow \mathbb{P}^1$ is an ℓ -cyclic cover unbranched outside $0, 1, \infty$, [5, Section 1]. Actually one can calculate not only the p -rank of Y , but also the isogeny type of the p -divisible group of the Jacobian of Y .

EXAMPLE 6.9. Let k be an algebraically closed field of characteristic p . Let E be an elliptic curve defined over k and P a k -point of E .

CLAIM. *Then*

$$\pi_1'(\mathbb{P}_k^1 - \{0, 1, \infty\}) \not\cong \pi_1'(E - \{P\}).$$

For simplicity we will suppose $p \neq 2, 3$, but the argument can easily be extended to any characteristic. Much more general results of this type have been obtained by A. Tamagawa (forthcoming).

Choose $\ell|p - 1$ with $\ell > 2$ but ℓ not necessarily prime. Choose an ℓ -cyclic cover $Y \rightarrow \mathbb{P}^1$ branched at $0, 1, \infty$ with $g(Y) > 0$. Corollary 6.8 implies that Y is ordinary, since $p \equiv 1 \pmod{\ell}$. Proposition 2.4 implies that $\pi_1^t(\mathbb{P}_k^1 - \{0, 1, \infty\})$ has a quotient of order $p\ell$; this quotient is non-Abelian. On the other hand, every quotient of $\pi_1^t(E - \{P\})$ of order $p\ell$ is Abelian. This proves the claim.

7. Degeneration

A way to get information on the p -rank of a cover is to degenerate the cover, and deduce information on the original cover from the, hopefully easier, degenerate cover. In the case of Galois covers of degree prime-to- p , this method works very well, as one knows exactly which covers one gets by degenerating it, namely *admissible covers*. In this section and the next, we use this idea to strengthen the results from the previous section, under certain extra hypotheses. Unfortunately, this method does not work in general. Under specialization, the p -ranks drops. It is possible to find examples where all degenerations of a given cover $Y_K \rightarrow X_K$ have p -rank strictly less than the p -rank that one want to show the original cover has, namely $B(\mathbf{a}, g)$.

The following notations will be fixed throughout this section and the next. Suppose $k = \bar{k}$ and the characteristic of k is $p > 0$. Let $A := k[[t]]$ have maximal ideal m_A and field of fractions K . Put $S := \text{Spec}(A)$. For a scheme \mathcal{X} over S we denote by $\mathcal{X}_{\bar{k}} = \mathcal{X} \otimes_A \bar{K}$ the generic fiber and $\mathcal{X}_s = \mathcal{X} \otimes_A k$ the special fiber. A curve \mathcal{X}/S is a flat scheme such that the geometric fibers are projective, connected, reduced and of dimension one. A semistable curve \mathcal{X}/S is a curve over S such that the geometric fibers of \mathcal{X} have at most ordinary double points as singularities. A morphism $\pi: \mathcal{Y} \rightarrow \mathcal{X}$ of curves over S is a cover if it is finite and generically étale. Let G be a subgroup of $\text{Aut}(\mathcal{Y}/\mathcal{X})$. The cover π is called G -Galois if $\mathcal{Y}/G \xrightarrow{\sim} \mathcal{X}$ and the order of G is equal to the degree of π .

DEFINITION 7.1. Let $\pi: Y \rightarrow X$ be a tame G -Galois cover of semistable curves over k . Suppose the decomposition group of τ is cyclic. Let ϕ be a generator of the decomposition group of τ . Let u_i for $i = 1, 2$ be a uniformizing parameter at τ of the branches of Y meeting at τ . Then π is called *admissible* if for each singular point τ of Y the following condition is satisfied: $\phi^*u_1 = \zeta u_1 \pmod{(u_1^2)}$ and $\phi^*u_2 = \zeta^{-1}u_2 \pmod{(u_2^2)}$, for some root of unity $\zeta \in k$.

Let \mathcal{X} be a semistable curve over S , whose generic fiber $\mathcal{X}_{\bar{k}}$ is smooth. Suppose we are given a G -Galois cover $\pi_K: Y_K \rightarrow \mathcal{X}_K$ of smooth curves and suppose $p \nmid |G|$. Denote the branch locus of π_K by x_1, \dots, x_r . Let \mathcal{Y} be the normalization of \mathcal{X} in the function field of Y_K . After replacing A by a finite extension of A , we may assume

\mathcal{Y} is reduced, by Abhyankar’s Lemma. Suppose that x_1, \dots, x_r specialize to distinct points in the smooth locus of \mathcal{X}_s .

PROPOSITION 7.2. *Let $\mathcal{Y} \rightarrow \mathcal{X}$ be as above. The cover $\mathcal{Y}_s \rightarrow \mathcal{X}_s$ is admissible and unbranched outside $\{x_1(s), \dots, x_r(s)\} \cup \text{Sing}(\mathcal{X}_s)$.*

Proof [19, Théorème 3.2]. Note that what is called admissible here is called *kummérien* in that paper. □

In fact, a stronger result is true. In the situation as above, one can show that admissible covers of \mathcal{X}_s can be lifted to covers of \mathcal{X} , unbranched outside x_1, \dots, x_r , see [7] or [23]. There is an isomorphism between the prime-to- p part of the fundamental group of $\mathcal{X}_{\bar{k}}$ and the prime-to- p part of a suitably defined *admissible fundamental group* of \mathcal{X}_s , but we do not need this here.

If \mathcal{X}/S is a curve with semistable fibers then $\sigma(\mathcal{X}_{\bar{k}}) \geq \sigma(\mathcal{X}_s)$, [16, p. 1281]. The analogous statement for generalized Hasse–Witt invariants is also true, as follows from the next proposition.

PROPOSITION 7.3. *Let $\mathcal{X} \rightarrow S$ be a semistable curve with $\mathcal{X}_{\bar{k}}$ nonsingular. Let $\pi: \mathcal{Y} \rightarrow \mathcal{X}$ be a G -Galois cover of order prime-to- p . For each irreducible character χ of G we have $\gamma_{\pi_{\bar{k}}}(\chi) \geq \gamma_{\pi_k}(\chi)$.*

Proof. Let Y be a semistable curve over k on which acts a group G of prime-to- p order. Then G acts on $H^1(Y, \mathcal{O}_Y)^F$. Using the identification

$$H^1(Y, \mathcal{O}_Y)^F \simeq H^1_{\text{ét}}(Y, \mathbb{Z}/p) \simeq (\mathbb{Z}/p)^{\sigma(Y)}$$

[10, III. 4.12], we can define an action of G on $(\mathbb{Z}/p)^{\sigma(Y)}$.

Now let χ be an irreducible character of G . Then the generalized Hasse–Witt invariant $\gamma(\chi)$ is equal to the product of the dimension of χ and the multiplicity of $\chi^{-1} \oplus (\chi^p)^{-1} \oplus \dots \oplus (\chi^{p^f-1})^{-1}$ in $(\mathbb{Z}/p)^{\sigma(Y)} \otimes_{\mathbb{F}_p} \overline{\mathbb{F}}_p$.

Let $\pi: \mathcal{Y} \rightarrow \mathcal{X}$ be as in the statement of the proposition. There exists an injective, canonical map

$$H^1(\mathcal{Y}_s, \mathbb{Z}/p) \hookrightarrow H^1(\mathcal{Y}_{\bar{k}}, \mathbb{Z}/p),$$

[16, p. 1281]. Since the map is G -equivariant, $\gamma_{\pi_k}(\chi) \leq \gamma_{\pi_{\bar{k}}}(\chi)$. □

Suppose that $p \equiv 1 \pmod{\ell}$. Let $Y \rightarrow \mathbb{P}^1$ be an ℓ -cyclic cover over K . Decompose $H^1(Y, \mathcal{O}_Y) = \bigoplus_{i=1}^{\ell-1} L_i$ into isotypical spaces. Then $F: L_i \rightarrow L_i$. Hence in this case $B(\mathbf{a}, 0) = g(Y)$. Question 3.3 boils down to: for a fixed type \mathbf{a} , do there exist x_1, \dots, x_r such that the cover of \mathbb{P}^1 branched at the x_i of type \mathbf{a} is ordinary? This is the case as is shown in the following proposition.

PROPOSITION 7.4. *Let $\ell|(p-1)$. There exists $x_1, \dots, x_r \in \mathbb{P}^1$ such that all ℓ -cyclic covers $\pi: Y \rightarrow \mathbb{P}^1_k$, unbranched outside x_1, \dots, x_r , are ordinary.*

Proof. We use induction on r . For $r = 2$ there is nothing to prove since each cover of \mathbb{P}^1 branched at 2 points has genus zero. For $r = 3$ the result follows from Corollary 6.8. So suppose $r \geq 4$.

Suppose the statement holds for all $s < r$. Let $A = k[[t]]$, $S = \text{Spec} A$ and let K be the quotient field of A . Construct a semistable curve $\mathcal{X} \rightarrow S$, with semistable special fiber consisting of two curves X_1 and X_2 of genus 0 meeting in one point τ , and generic fiber a nonsingular curve of genus 0. Choose sections $x_1, \dots, x_r: S \rightarrow \mathcal{X}$, such that x_1, x_2 meet $X_1 - \{\tau\}$ and x_3, \dots, x_r meet $X_2 - \{\tau\}$. By induction, we can choose the x_i such that for all $m|\ell$, the m -cyclic covers of X_2 , unbranched outside x_3, \dots, x_r, τ , are ordinary. Also the m -cyclic covers of X_1 , unbranched outside x_1, x_2, τ , are ordinary, by the results for $r = 3$ branch points.

Let $\pi_K: Y_{\bar{K}} \rightarrow \mathcal{X}_{\bar{K}}$ be an ℓ -cyclic cover, unbranched outside x_1, \dots, x_r . This cover will be defined over some finite extension of K , so, after replacing A by a finite extension, we may assume that the cover is defined over K . We now proceed as in the beginning of this section. Let \mathcal{Y} be the normalization of \mathcal{X} in the function field of $Y_{\bar{K}}$. Assume \mathcal{Y} is reduced (replace A by a finite extension, if necessary). Proposition 7.2 implies that $\mathcal{Y}_s \rightarrow \mathcal{X}_s$ is an admissible ℓ -cyclic cover which is unbranched outside $x_1(s), \dots, x_r(s), \tau$.

Let Z be an irreducible component of \mathcal{Y}_s . Let m be the order of the decomposition group of Z . Then $Z/\langle \phi^{\ell/m} \rangle$ is isomorphic to X_i for i equal to 1 or 2 hence $Z \rightarrow X_i$ is an m -cyclic cover of X_i and Z is ordinary by assumption. Since all components of \mathcal{Y}_s are ordinary, the curve \mathcal{Y}_s itself is ordinary. It follows that

$$g(\mathcal{Y}_{\bar{K}}) \geq \sigma(\mathcal{Y}_{\bar{K}}) \geq \sigma(\mathcal{Y}_s) = p_a(\mathcal{Y}_s).$$

Because $\mathcal{Y} \rightarrow S$ is flat, $p_a(\mathcal{Y}_s) = g(\mathcal{Y}_{\bar{K}})$ and $\mathcal{Y}_{\bar{K}}$ is ordinary. □

LEMMA 7.5. *Suppose that \mathbf{a} is a type such that $\| - p^i \| = r - 2$ for all i . Then there exist $x_1, \dots, x_r \in \mathbb{P}^1$ such that for all covers of type \mathbf{a} , unbranched outside the x_i , we have $\gamma_1 = r - 2$.*

Proof. This lemma follows easily by induction, using an argument similar to the argument used in the previous proposition. □

The situation of the lemma above is very special. Here is a case where it occurs.

EXAMPLE 7.6. Let $\ell = (p^r - 1)/(p - 1)$ and let $a := (\ell; 1, p, p^2, \dots, p^{r-1})$. Let $\pi: Y \rightarrow \mathbb{P}^1$ be an ℓ -cyclic cover of type a branched at x_1, \dots, x_r . Then

$$\| p^j \| = \| 1 \| = \frac{1}{\ell} (1 + \dots + p^{r-1}) - 1 = 0 \quad \text{for all } j.$$

Hence, $\| - p^j \| = \| - 1 \| = r - 2$ for all j by Lemma 4.5. From the lemma above it follows that $\gamma_1 = r - 2$, in case x_1, \dots, x_r are sufficiently general.

PROPOSITION 7.7. *Let $Y_x \rightarrow \mathbb{P}^1$ be the cover of type \mathbf{a} branched at $0, 1, \infty, x$. Then there exists an $x \in \mathbb{P}^1$ such that $\sigma(Y_x) = B(\mathbf{a}, 0)$.*

Proof. To prove the proposition, it suffices to show that for each $j = 1, \dots, \ell - 1$ we have $\gamma_{-j} = \min_i \|p^i j\|$. As before we may assume $j = 1$ and $\|p^i\| \geq \|1\|$. From Lemma 4.5 we know that $\|p^i\| \leq r - 2 = 2$ for all i . Therefore, distinguish three cases:

- (i) $\|1\| = 0$,
- (ii) $\|1\| = 1$,
- (iii) $\|p^i\| = 2$ for all i .

We have to show for a suitable choice of x we have $\gamma_{-1} = 0$ in the first case, $\gamma_{-1} = 1$ in the second case and $\gamma_{-1} = 2$ in the third case.

In Case (i), the action of Frobenius on $\oplus_i L_{p^i}$ is nilpotent, so $\gamma_{-1} = 0$. Case (ii) follows from Proposition 6.7. Case (iii) follows from Lemma 7.5. □

PROPOSITION 7.8. *Suppose $p \equiv -1 \pmod{\ell}$. Let $\mathbf{a} = (\ell; a_1, \dots, a_r)$ be a type. There exist x_1, \dots, x_r such that for the ℓ -cyclic cover $Y \rightarrow \mathbb{P}^1$ of type \mathbf{a} , branched at the x_i , we have $\sigma(Y) = B(\mathbf{a}, 0)$.*

Proof. Let $\pi: Y \rightarrow \mathbb{P}^1$ be a cover of type $\mathbf{a} = (\ell; a_1, \dots, a_r)$ with $0 < a_i < \ell \neq 2$. Suppose that $p \equiv -1 \pmod{\ell}$. The coset of $\langle p \rangle$ in $(\mathbb{Z}/\ell)^*$ containing i consists of two elements: i and $-i$. We want to show

$$\gamma_{-i} = \min(\|i\|, \|-i\|).$$

We may suppose $i = 1$.

From Lemma 4.5 we know that $\|1\| + \|-1\| = r - 2$. Put $n = \|1\|$. We may suppose $0 \leq n \leq [(r - 2)/2]$. If $n = 0$ the matrix of Frobenius on $L_1 \oplus L_{-1}$ is nilpotent and $\gamma_1 = 0$. Suppose that $n > 0$.

We will prove the proposition by induction. For $r = 3, 4$ the statement follows from Corollary 6.8 and Proposition 7.7. Suppose the statement holds for $r' < r$.

Case 1. Suppose $n < (r - 2)/2$. For r odd this always holds. Choose $i \neq j$ with $a_i + a_j < \ell$. We can do this since if $a_{2s-1} + a_{2s} \geq \ell$ for each s , then $a_1 + \dots + a_r \geq [r/2]\ell$. But $a_1 + \dots + a_r = n\ell < (r - 2)\ell/2 < [r/2]\ell$, so we get a contradiction. Put $\mathbf{a}_1 = (a_i, a_j, -a_i - a_j)$ and $\mathbf{a}_2 = (a_i + a_j, a_1, \dots, \hat{a}_i, \dots, \hat{a}_j, \dots)$, i.e. in \mathbf{a}_2 we omit a_i and a_j . For a cover $f: X \rightarrow \mathbb{P}^1$ of type \mathbf{a}_1 we find

$$\|1\|_{\mathbf{a}_1} = \frac{a_i + a_j + (\ell - a_i - a_j)}{\ell} - 1 = 0 \quad \text{and} \quad \|-1\|_{\mathbf{a}_1} = 1.$$

For a cover $g: Z \rightarrow \mathbb{P}^1$ of type \mathbf{a}_2 we find

$$\|1\|_{\mathbf{a}_2} = \frac{a_1 + \dots + a_r}{\ell} - 1 = n \quad \text{and} \quad \|-1\|_{\mathbf{a}_2} = r - 3 - n.$$

Note $k \leq r - n - 3$. By induction $\gamma_1(f) = 0$ and $\gamma_1(g) = n$. As in the proof of the previous proposition, Proposition 7.3 implies $\gamma_1(\pi) \geq \gamma_1(g) = n$. Hence $\gamma_1(\pi) = n$.

Case 2. Suppose $r = 2s$ with $s > 2$ and $n = s - 1$. Then $a_1 + \dots + a_r = s\ell$.

CLAIM. *There exist distinct h, i, j , such that $\ell < a_h + a_i + a_j < 2\ell$.*

To prove the claim, put $x_i = a_i/\ell - 1/2$. Note

$$-1/2 < x_i < 1/2 \quad \text{and} \quad x_1 + \dots + x_r = 0$$

We will see that for any m there exist distinct $i_1, \dots, i_m \in \{1, \dots, r\}$ such that $-1/2 < \sum_{j=1}^m x_{i_j} < 1/2$. The claim follows from this by taking $m = 3$.

Use induction. For $m = 1$ it is trivial. Let $m > 1$. Choose $I = \{i_1, \dots, i_{m-1}\}$ such that $-1/2 < S_{m-1} := \sum_{i \in I} x_i < 1/2$. If $S_{m-1} \geq 0$ then $(\sum x_i) - S_{m-1} < 0$, so there is an $j \notin I$ with $x_j < 0$. We can add j to I to fulfill the condition for m . Analogously for $S_{m-1} < 0$. This proves the claim.

Choose h, i, j as in the claim. Put $\mathbf{a}_1 = (a_h, a_i, a_j, -a_h - a_i - a_j)$, and $\mathbf{a}_2 = (a_h + a_i + a_j, \text{rest of the } a_i)$. Then $\|1\|_{\mathbf{a}_1} = \|-1\|_{\mathbf{a}_1} = 1$ and $\|1\|_{\mathbf{a}_2} = \|-1\|_{\mathbf{a}_2} = s - 1$. As in the previous case, the proposition follows by induction. \square

Remark 7.9. In this section, we showed several results of the form: under this condition there exists a position of the branch points such that all covers have maximal p -rank. Recall that by Lemma 4.6 the set U of all branch points with this property is open. So in fact, we could reformulate, e.g. Proposition 7.4 as follows. Suppose $p \equiv 1 \pmod{\ell}$. Then there exists a dense open subset $U \subset (\mathbb{P}^1)^r - \Delta$ such that all ℓ -cyclic covers branched at $(x_1, \dots, x_r) \in U$ are ordinary. Similar for the other results.

8. Cyclic Covers of Generic Curves

Let X be an ordinary elliptic curve or the generic curve of genus ≥ 2 . In this section we will show that we can reduce Question 3.3 on the p -rank of an ℓ -cyclic cover of X to the case of ℓ -cyclic covers of \mathbb{P}^1 .

PROPOSITION 8.1. *Let X be the generic curve of genus $g \geq 2$ or an ordinary elliptic curve. Let \mathbf{a} be a type such that there exists a (not necessarily connected) cover of \mathbb{P}^1 of type \mathbf{a} with p -rank $B(\mathbf{a}, 0)$. Then there exist $x_1, \dots, x_r \in X$ such that each cover of X of type \mathbf{a} , unbranched outside the x_i , has p -rank $B(\mathbf{a}, g) + g(X)$.*

We need to consider nonconnected covers of \mathbb{P}^1 in the statement of the proposition because of the following. Suppose $\gcd(\ell, a_1, \dots, a_r) \neq 1$ and suppose $g(X) \geq 1$. Then

there exist connected covers of X of type $\mathbf{a} := (\ell; a_1, \dots, a_r)$, but the covers of \mathbb{P}^1 of type \mathbf{a} are nonconnected.

Proof. Let $g \geq 1$ be an integer. Let X be the generic curve of genus g if $g \geq 2$ or an ordinary elliptic curve if $g = 1$ and let k be some algebraically closed field of characteristic p over which X is defined. Put $S = \text{Spec } k[[t]]$ and $K = Q(k[[t]])$. Construct a semistable curve $\mathcal{X} \rightarrow S$ as follows. The generic fiber $\mathcal{X}_{\bar{k}}$ is a trivial deformation of X and the special fiber X_s consist of two components X_1 and X_2 meeting in one point τ , where $X_1 \simeq X$ and X_2 is a nonsingular curve of genus zero. A curve like this can be constructed by blowing up the trivial deformation of X over S in one point of the special fiber. Choose distinct sections x_1, \dots, x_r of $S \rightarrow \mathcal{X}$ such that the x_i meet the special fiber \mathcal{X}_s in $X_2 - \{\tau\}$. Let $n_2 = \text{gcd}(\ell_1, a_1, \dots, a_r)$ and write $\mathbf{a}' = (\ell_2; a'_1, \dots, a'_r)$, where $a'_i = a_i/n_2$ and $\ell_2 = \ell/n_2$. We assume that the ℓ_2 -cyclic cover $Y'_2 \rightarrow X_2$ of type \mathbf{a}' with branch locus $x_1(s), \dots, x_r(s)$ has p -rank $B(\mathbf{a}', 0)$. (This is equivalent to the assumption in the statement of the proposition.)

Recall that starting from the cover $Y'_2 \rightarrow X_2$ we can form a (nonconnected) \mathbb{Z}/ℓ -cover by taking the induced cover, i.e. take $n_2 = \ell/\ell_2$ copies of Y'_2 and let \mathbb{Z}/ℓ permute the copies suitably. Call the resulting cover $Y_2 \rightarrow X_2$; it is of type \mathbf{a} . Consider all admissible covers $Y \rightarrow \mathcal{X}_s$ whose restriction to X_2 is isomorphic to $Y_2 \rightarrow X_2$. (These covers are what one could call ‘admissible covers of \mathcal{X}_s of type \mathbf{a} with branch locus $x_1(s), \dots, x_r(s)$ ’.) We would like to show that all these covers have p -rank equal to $B(\mathbf{a}, g) + g$. The proposition follows from this, as in the proof of Proposition 7.4.

We will now compute the p -rank of the ‘admissible covers of \mathcal{X}_s of type \mathbf{a} with branch locus $x_1(s), \dots, x_r(s)$ ’. Let $Y \rightarrow \mathcal{X}_s$ be such a cover. Write $Y_1 = Y|_{X_1}$ and let n_1 be the number of connected components of Y_1 and put $\ell_1 = \ell/n_1$. Note $\text{gcd}(n_1, n_2) = 1$ since Y is supposed to be connected. Let Y'_1 be a connected component of Y_1 . If $g(X_1) = 1$ then $Y'_1 \rightarrow X_1$ is an étale cover of an ordinary elliptic curve and Y'_1 is ordinary. If $g(X_1) \geq 2$ it follows from [11, Section 5.III] that Y'_1 is ordinary.

By assumption, the curve Y'_2 has p -rank

$$\sigma(Y'_2) = \sum_{i \in I'} f'_i \min_j \left(\sum_{n=1}^r \left\langle \frac{ip^j a'_n}{\ell_2} \right\rangle - 1 \right),$$

where I' is a set of representatives of the cosets of $\langle p \rangle$ in $\mathbb{Z}/\ell_2 - \{0\}$ and f'_i is the number of elements of the coset containing i . Of course, since all components of Y_2 are isomorphic, the same statement holds for the other components of Y_2 . Note that $H^1(Y_2, \mathcal{O}_{Y_2}) = \text{Ind}_{\mathbb{Z}/\ell_2}^{\mathbb{Z}/\ell} H^1(Y'_2, \mathcal{O}_{Y'_2})$ as a $k[\mathbb{Z}/\ell]$ -module. This yields

$$\sigma(Y_2) = n_2(\sigma(Y'_2)) = \sum_{i \in I} f_i \min_j \left(\sum_{n=1}^r \langle ip^j a_n \ell_2 \rangle - 1 \right) + n_2 - 1,$$

where I is a set of representatives of the cosets of $\langle p \rangle$ in $\mathbb{Z}/\ell - \{0\}$ and f_i is the number of elements of the coset containing i .

The curve Y contains $1 + \ell - n_1 - n_2$ loops. Therefore, we find for the p -rank of Y that

$$\begin{aligned} \sigma(Y) &= n_1 \cdot g(Y'_1) + 1 + \ell - n_1 - n_2 + \sigma(Y_2) \\ &= \ell g + \sum_{i \in I} f_i \min_j \left(\sum_{n=1}^r \left\langle \frac{ip^j a_n}{\ell} \right\rangle - 1 \right) = B(\mathbf{a}, g) + g. \end{aligned}$$

Recall that

$$B(\mathbf{a}, g) = \sum_{i \in I} f_i \min_j \left(\sum_{n=1}^r \left\langle \frac{ip^j a_n}{\ell} \right\rangle - 1 \right) + (\ell - 1)g.$$

Note that for i divisible by ℓ_2 , we have

$$\sum_{n=1}^r \left\langle \frac{ip^j a_n}{\ell} \right\rangle - 1 = -1,$$

since all a_n 's are divisible by n_2 . This finishes the proof of the proposition. □

In fact, the proof applies to every curve X such that all étale cyclic covers $Y \rightarrow X$ whose order m divides ℓ , have p -rank $\sigma(X) + (m - 1)(g - 1)$. In [11] it is proved that this holds for curves in a dense open subset of the moduli space \mathcal{M}_g .

COROLLARY 8.2. *Let X be the generic curve of genus $g \geq 2$ or an ordinary elliptic curve. Let ℓ be an integer prime-to- p , and suppose one of the following holds:*

- (i) $p \geq \ell(r - 3)$,
- (ii) $r \leq 4$,
- (iii) $p \equiv \pm 1 \pmod{\ell}$.

Then there exists $x_1, \dots, x_r \in X$ such that all ℓ -cyclic covers $Y \rightarrow X$, unbranched outside the x_i of type a satisfy

$$\sigma(Y) = B(a, g) + g.$$

Acknowledgements

This paper is a rewritten version of part of my PhD thesis. I would like to thank Kate Stevenson and Marco Garuti for their comments on an earlier version of this paper.

References

1. Chevalley, C. and Weil, A.: Über das Verhalten der Integrale erster Gattung bei Automorphismen des Funktionenkörpers, *Abh. Math. Sem. Univ. Hamburg* **10** (1934), 358–361.
2. Crew, R. M.: Etale p -covers in characteristic p , *Compositio Math.* **52** (1984), 31–45.

3. Elkies, N. D.: The existence of infinitely many supersingular primes for every elliptic curve over \mathbb{Q} , *Invent. Math.* **89** (1987), 561–567.
4. González, J.: Hasse–Witt matrices for the Fermat curves of prime degree, *Tôhoku Math. J.* **49** (1997), 149–163.
5. Gross, B. H. and Rohrlich, D. H.: Some results on the Mordell–Weil group of the Jacobian of the Fermat curve, *Invent. Math.* **44** (1978), 201–224.
6. Grothendieck, A. et al.: *Revêtements étales et géométrie algébrique* (SGA 1), Lecture Notes in Math. 224, Springer-Verlag, Heidelberg, 1971.
7. Harbater, D. and Stevenson, K. F.: Patching and thickening problems, *J. Algebra* **212** (1999), 272–304.
8. Kani, E.: The Galois-module structure of the space of holomorphic differentials of a curve, *J. reine Angew. Math.* **367** (1986), 187–206.
9. Katsurada, H.: On generalized Hasse–Witt invariants and unramified Galois extensions of an algebraic function field, *J. Math. Soc. Japan* **31** (1979), 101–125.
10. Milne, J. S.: *Étale Cohomology*, Princeton Univ. Press, Princeton, 1980.
11. Nakajima, S.: On generalized Hasse–Witt invariants of an algebraic curve, In: Y. Ihara (ed.), *Galois groups and their representations (Nagoya 1981)*, Adv. Stud. Pure Math. 2, North-Holland, Amsterdam, 1983, pp. 69–88.
12. Nakajima, S.: Galois module structure of cohomology groups for tamely ramified coverings of algebraic varieties, *J. Number Theory* **22** (1986), 115–123.
13. Pacheco, A.: Unramified coverings of algebraic curves, *J. Number Theory* **53** (1995), 211–228.
14. Pacheco, A. and Stevenson, K. F.: Galois modules and the algebraic fundamental group of projective curves in positive characteristic, to appear in *Pacific J. Math.*
15. Oort, F.: Subvarieties of moduli spaces, *Invent. Math.* **24** (1974), 95–119.
16. Raynaud, M.: Mauvaise réduction des courbes et p -rang, *C.R. Acad. Sci. Paris Série I*, **319** (1994), 1279–1282.
17. Raynaud, M.: Revêtements des courbes en caractéristique p et ordinarité, preliminary version.
18. Rück, H.-G.: Class groups and L -series of function fields, *J. Number Theory* **22** (1986), 177–189.
19. Sadi, M.: Revêtements modérés et groupe fondamental de graphe de groupes, *Compositio Math.* **107** (1997), 321–340.
20. Serre, J-P.: Sur la topologie des variétés algébriques en caractéristiques p , *Sympos. Int. Top. Algebra*, Mexico, 1958, pp. 24–53.
21. Serre, J-P.: *Topics in Galois Theory*, Jones and Bartlett, Boston, 1992.
22. Stienstra, J., van der Put, M. and van der Marel, B.: On p -adic monodromy, *Math. Z.* **208** (1991), 309–325.
23. Wewers, S.: Deformation of tame admissible covers of curves, In: H. Völklein (ed.), *Aspects of Galois Theory (Gainesville 1996)*, Cambridge Univ. Press, 1999.
24. Yui, N.: On the Jacobian variety of the Fermat curve, *J. Algebra* **65** (1980), 1–35.