

28. Boelaert JR. Staphylococcus *aureus* infection in hemodialysis patients. Mupirocin as a topical strategy against nasal carriage: a review. *J Chemother*1994;6:19S-24S.
29. Connolly S, Noble WC, Phillips I. Mupirocin resistance in coagulase-negative staphylococci. *J Med Microbiol*1993;39:450-453.
30. Naguib MH, Naguib MT, Flournoy DJ. Mupirocin resistance in methicillin-resistant *Staphylococcus aureus* from a veterans hospital. *Chemotherapy* 1993;39:400-404.
31. Neu HC. The use of mupirocin in controlling methicillin-resistant *Staphylococcus aureus*. *Infect Control Hosp Epidemiol* 1990;11:11.
32. Rahman M, Connolly S, Noble WC, Cookson B, Phillips I. Diversity of staphylococci exhibiting high-level resistance to mupirocin. *J Med Microbiol* 1990;33:97-100.
33. Capobianco JO, Doran CC, Goldman RC. Mechanisms of mupirocin transport into sensitive and resistant bacteria. *Antimicrob Agents Chemother*1989;33:156-163.
34. Cookson B. Failure of mupirocin-resistant staphylococci to inactivate mupirocin. *Eur J Clin Microbiol Infect Dis*1989;8:1038-1040.
35. Farmer TH, Gilbert J, Elson SW. Biochemical basis of mupirocin resistance in strains of *Staphylococcus aureus*. *J Antimicrob Chemother* 1992;30:587-596.
36. Dyke KGH, Curnock SF, Golding M, Noble WC. Cloning of the gene conferring resistance to mupirocin in *Staphylococcus aureus*. *FEMS Microbiol Lett*1991;77:195-198.
37. Gilbert J, Perry CR, Slocombe B. High-level mupirocin resistance in *Staphylococcus aureus* evidence for two distinct isoleucyl-tRNA synthetases. *Antimicrob Agents Chemother*1993;37:32-38.
38. Hodgson JE, Curnock SP, Dyke KGH, Morris R, Sylvester DR, Gross MS. Molecular characterization of the gene encoding high-level mupirocin resistance in *Staphylococcus aureus* J2870. *Antimicrob Agents Chemother* 1994;38:1205-1208.
39. Noble WC, Rahman M, Cookson B, Phillips I. Transferable mupirocin-resistance. *J Antimicrob Chemother*1988;22:771-776.
40. Janssen DA, Zarins LT, Schaberg DR, Bradley SE, Terpenning MS, Kauffman CA. Detection and characterization of mupirocin resistance in *Staphylococcus aureus*. *Antimicrob Agents Chemother* 1993;37:2003-2006.
41. Rahman M, Noble WC, Dyke KGH. Probes for the study of mupirocin resistance in staphylococci. *J Med Microbiol*1993;39:446-449.
42. Morton T, Patterson J, Schaberg D, Archer G. Appearance of a mupirocin resistance gene on a staphylococcal conjugative plasmid: a model for IS-mediated recombination. Taken from Interscience Conference on Antimicrobial Agents and Chemotherapy; 1993. Abstract.
43. Ramsey MA, Zarins LT, Bradley SE, Kauffman CA, Morton TM. Identification of the mupirocin resistance gene on the staphylococcal chromosome: evidence to support IS-mediated recombination. Taken from American Society for Microbiology; 1994. Abstract.
44. Slocombe B, Perry C. The antimicrobial activity of mupirocin-an update on resistance. *J Hosp Infect*1991;19B:19-25.
45. Cookson BD. Mupirocin resistance in staphylococci. *J Antimicrob Chemother*1990;25:497-503.

## Diphtheria Epidemic in the Newly Independent States of the Former Soviet Union

by Gina Pugliese, RN, MS  
Medical News Editor

Although diphtheria was controlled for approximately 30 years after the institution, in the late 1950s, of childhood vaccination with diphtheria toxoid, epidemic diphtheria has reemerged in the newly independent states (NIS) of the former Soviet Union. Overall, reported cases of diphtheria in the NIS increased from 839 in 1989 to 47,802 in 1994. In 1994, a total of 1,746 persons died; case-fatality rates ranged from 2.8% to 23.0%

The World Health Organization considers the rapidly expanding diphtheria epidemic in the NIS a public health emergency. In the Russian Federation, the epidemic has intensified each year. Although the reasons for the diphtheria epidemic in the NIS are not fully understood, one important factor is the presence of a large number of susceptible children and adults.

Spread of the organism may have been facilitated by crowding and population migration resulting from the dissolution of the Soviet Union. In addition, adequate control measures (particularly aggressive mass vaccination in affected areas) were not implemented during the early phase of the epidemic.

Increases in the number of susceptible children in the NIS probably resulted from a combination of low vaccination coverage in many areas and inappropriate primary vaccination of substantial numbers of infants with Td, a formulation for adults containing decreased amounts of diphtheria toxoid. The existence of large numbers of susceptible adults is a new phenomenon. In the prevaccine era, most persons acquired immunity to diphtheria naturally before adulthood through exposure to *Corynebacterium diphtheriae*. Following the introduction of childhood vacci-

nation diphtheria toxoid, circulation of toxigenic *C diphtheriae* decreased substantially. In addition, vaccine-induced immunity wanes over time unless periodic boosters are administered. Lack of effectiveness of diphtheria toxoid is not considered to be an important contributing factor for this epidemic

Control of the epidemic in the NIS requires immediate efforts to raise levels of immunity through extensive mass vaccination of adolescents and adults. Shortages of vaccine, antitoxin, and antibiotics exist in the NIS (except the Russian Federation); these needs must be addressed immediately through coordinated efforts of international public health and donor agencies.

FROM: CDC. Diphtheria epidemic-new independent states of the former Soviet Union, 1990-1994. *MMWR* 1995;44(10):177-80.