

# Report on the recent advances performed in the determination of radiative parameters for spectral lines of astrophysical interest in heavy elements

Pascal Quinet

Physique Atomique et Astrophysique, Université de Mons, B-7000, Mons, Belgium  
email: [Pascal.Quinet@umons.ac.be](mailto:Pascal.Quinet@umons.ac.be)

**Abstract.** We present an overview of the advances made during the past 15 years by the Atomic Physics and Astrophysics Group of Mons University regarding the analysis of the spectra, the transition probabilities, and the radiative lifetimes in heavy elements. More precisely, this review is focused on neutral and lowly ionized atoms belonging to the lanthanide group, the fifth row, and the sixth row of the periodic table ( $Z = 37\text{--}86$ ), for which a very large amount of new data has been obtained.

**Keywords.** atomic data, atomic processes

---

The study of complex atomic structures has known a considerable development over the past years, the new improvements arising from both theory and experiment. During the period 1998–2015, part of our research activities has been focused on the combination of computational and experimental techniques for providing new accurate atomic data for neutral and lowly ionized atoms of astrophysical interest. In the present contribution, we report on the recent advances we have made concerning the determination of new radiative parameters (transition probabilities, oscillator strengths, radiative lifetimes) in heavy elements belonging to the fifth row and the sixth row including the lanthanide group of the periodic table. In astrophysics, these elements present a great interest for the investigation of the atmospheres of chemically peculiar stars, deriving chemical abundances, and computing the opacity of the stellar matter, and consequently for building stellar models. In stellar nucleosynthesis, a detailed analysis of the  $r$ - and  $s$ -processes is not possible without a great deal of reliable atomic data for not only the strongest lines of heavy elements, but also a huge number of weak absorption features, which are now observable in the high-resolution astrophysical spectra.

All the data obtained so far in our work correspond to transition decay rates for more than 200 000 spectral lines in about 100 heavy atoms and ions. These were generally deduced from a combination of experimental radiative lifetimes measured with the time-resolved laser-induced fluorescence (LIF) (over 700 lifetimes were measured) and theoretical branching fractions calculated with a relativistic Hartree-Fock (HFR) approach taking core-polarization effects into account. This extensive work and the numerous subsequent comparisons have provided us with a broad experience concerning the evaluation of radiative parameters. On that basis, we have been able to deduce some criteria regarding the accuracy estimate of these atomic data on both the theoretical and experimental side. Our research team has published about 100 papers in relation with this work. A comprehensive list of those references has been presented at the IAU General Assembly.