

TANGENT BUNDLES, MONOIDAL THEORIES AND WEIL ALGEBRAS

POON LEUNG

(Received 17 December 2017; first published online 3 May 2018)

2010 Mathematics subject classification: primary 18D99; secondary 53A99.

Keywords and phrases: tangent structure, Weil algebra.

In any discussion of differential geometry (for instance, see [4]), a key role is played by the *tangent space* TM to a given space (say, smooth manifold) M ; this is equipped with a projection map, $p_M: TM \rightarrow M$, and various other structures.

These structures have also been studied from a category theoretic point of view, in which T becomes a functor in its own right, and the projection p and the other structure become natural transformations. These notions have been given the name *tangent structure*. The precise structure of the tangent bundles then varies depending on the choices made in relation to the ‘other structure’.

Rosický [6], for instance, detailed a set of choices which resulted in the tangent bundles having the structure of abelian groups, motivated by examples from computer science. Cockett and Cruttwell [3] later gave a set of choices resulting in the tangent bundles having the structure of commutative monoids.

In this thesis, we present a different perspective. We first introduce the Weil algebras, which have a well-established role in the discussion of differential geometry (for instance, see [5] or [7]). However, we follow up by restricting to a subcategory we shall call \mathbf{Weil}_1 , and describe a way to characterise the Weil algebras of this (sub)category using (co)graphs (see [2] for more details). We develop this idea even further by showing how the maps between such Weil algebras can also be described using the language of graphs.

This part of the thesis concludes with a demonstration that to give a tangent structure to some category \mathcal{M} is equivalent to giving a strong monoidal functor

$$F: \mathbf{Weil}_1 \rightarrow \text{End}(\mathcal{M})$$

Thesis submitted to Macquarie University in February 2017; degree approved on 14 July 2017; supervisors Stephen Lack and Richard Garner.

© 2018 Australian Mathematical Publishing Association Inc.

preserving certain limits; or more broadly, a strong monoidal functor

$$F: \mathbf{Weil}_1 \rightarrow \mathcal{G}$$

for some monoidal category \mathcal{G} . This also suggests that we may think of \mathbf{Weil}_1 as a sort of monoidal theory (see [1]) for tangent structures. Further, the choice of structure for the tangent bundles (monoid/group/vector space) is determined by the choice of the set of coefficients for the Weil algebras.

In the penultimate chapter of the thesis, we discuss a method for extending the definition of tangent structure to include such things as higher-order derivatives and more complicated relationships between these derivatives. Namely, we propose a candidate category \mathbf{Weil}_∞ intended to capture all finite-order derivatives. We then detail a methodical approach to characterise each Weil algebra in this category as an appropriate limit.

We conclude this chapter by suggesting that an ‘extended’ tangent structure could be given as a strong monoidal functor

$$F: \mathbf{Weil}_\infty \rightarrow \mathcal{G}$$

preserving the limits described.

In the final chapter, we give a brief discussion of what is needed to have an appropriate notion of ‘addition’ of tangent vectors, and then give some examples of possible subcategories other than \mathbf{Weil}_1 and \mathbf{Weil}_∞ which may also prove to be useful in the broader discussion of tangent structure. We conclude by suggesting some ideas for extending the ideas described beyond the category \mathbf{Weil}_∞ .

References

- [1] M. Barr and C. Wells, ‘Toposes, triples, and theories’, in: *Grundlehren der Mathematischen Wissenschaften* (Springer, New York, 1985).
- [2] A. J. Bondy and U. S. R. Murty, *Graph Theory with Applications* (Wiley, New York, 1991).
- [3] R. Cockett and G. Cruttwell, ‘Differential structure, tangent structure and synthetic differential geometry’, *Appl. Categ. Structures* **22**(2) (2014), 331–417.
- [4] N. Hitchin, *Differentiable Manifolds*, Oxford UK Course C3.2b Lecture Notes (Oxford University, 2012), available at <https://people.maths.ox.ac.uk/hitchin/hitchinnotes/manifolds2012.pdf>.
- [5] I. Kolar, P. W. Michor and J. Slovák, *Natural Operations in Differential Geometry* (Springer, Berlin Heidelberg, 2010).
- [6] J. Rosický, ‘Abstract tangent functors’, *Diagrammes* **12**(3) (1984), JR1–JR11.
- [7] A. Weil, ‘Théorie des points proches sur les variétés différentielles’, in: *Colloque de Topologie et Géométrie Différentielle, Strasbourg, 1953* (Centre National de la Recherche Scientifique, Paris, 1953), 111–117.

POON LEUNG, School of Mathematics, Macquarie University,
North Ryde 2113, New South Wales, Australia
e-mail: poon.leung@mq.edu.au