

HEAD INJURY: ORGANIZATIONAL ASPECTS OF CARE

**Corrado Manni, M.D., F. Della Corte, M.D. and
A. Proietti, M.D., Rome, Italy**

Head injury still accounts for 1% of the deaths in Italy and a larger percentage of severe disability. Thus, the technological, organizative, economical and therapeutical efforts appear truly justifiable. The best outcome results can be obtained by instituting the following: 1) Continuous and appropriate information to the general population about rescue and first aid; 2) The most immediate intervention at the site with an appropriate means of rescue; 3) Experienced general and medical assistants; 4) Ability to carry out instrumental and pharmacological therapies aimed at maintaining vital functions during transportation; 5) Uniformity and simplicity in exchanging information; 6) Selection of the most appropriate hospital.

ENDOTOXIN BLUNTS THE CHRONOTROPIC RESPONSE TO ISOPROTERENOL IN THE DOG MODEL

**Thomas J. Iberti, M.D., David R. Gentili, M.D.,
Saul R. Berger, M.D., Thomas A. Paluch, M.D.,
Ernest Benjamin, M.D., New York,
New York, U.S.A.**

Previous studies have demonstrated myocardial depression associated with endotoxemia. In addition, we recently described a significantly slower heart rate (HR) seen in canine endotoxic shock than that observed with hemorrhagic shock of similar severity. The purpose of this study was to examine the chronotropic effects of isoproterenol (ISO) in endotoxic dogs to determine if the negative chronotropy seen in endotoxemia could be eliminated by beta agonist therapy. Eighteen dogs were anesthetized, ventilated and hemodynamically monitored. After stabilization, they were randomized to 3 groups. Group I received isoproterenol 0.1 mg/kg/min; Group II 4 mg/kg E. coli endotoxin (ENDO); and Group III received both ISO and ENDO.

	HEART RATE (\pm SD)	BASELINE	Rx
I (ISO)		150 \pm 27	201 \pm 18 ($p < 0.05$ vs. Control II, III)
II (ENDO)		158 \pm 13	120 \pm 15 ($p < 0.05$ vs. Control I, III)
III (ISO + ENDO)		161 \pm 26	140 \pm 21 ($p < 0.05$ vs. Control I, II)

The results of this study demonstrate that the chronotropic effects of ISO are depressed in the presence of endotoxin. Thus, in the endotoxic animal model, beta adrenergic stimuli does not overcome the negative chronotropic response to endotoxin. The pathophysiology of this phenomenon remains to be defined, but may involve nodal ischemia, diffuse myocardial depression, or direct action of endotoxin.

EMERGENCY MANAGEMENT OF PENETRATING MISSILE HEAD INJURIES

A.A. Ameen, M.D., Basrah, Iraq

The experience of the neurosurgical unit at Basrah General Teaching Hospital (Iraq) during the 6 years period of the Iraqi Iranian War in the treatment of penetrating craniocerebral injuries is reported, with particular reference to the triage and management of large combat casualties received during busy periods that follow a major battle or bombardment of the city.

Attention is drawn also to the diagnostic difficulties that occur in assessing patients with head and associated injuries. The role of CAT scanning in the immediate management is discussed, as compared to other studies in the pre-CAT scanner era.

TRAINING, TEACHING, ORGANIZATION AND MANAGEMENT OF TRAUMA IN AN URBAN SETTING

**Ayuyao Adolfom, M.D., New York,
New York, U.S.A.**

Much has been written about the training of a surgeon, the proper qualifications, and what it means to be a surgeon. However, there has been no emphasis given to the important topic of trauma. A training surgical resident rotates through different surgical divisions and subspecialties, but seldom will you find a rotation in an organized trauma service.

Providing care to the injured was one of the earliest recorded roles of a surgeon. However, this aspect of medicine has not been emphasized in most training programs while other areas of surgery have been favored. It is only in the past few decades that attention has been focused on the care of the injured. The

responsibilities of a trauma surgeon are not limited to the mechanical repair of the various injuries, but include physiological support of the patient, organization of the team and hospital resources, coordination of pre-hospital care, research and education.

Harlem Hospital Center was established in 1887, serving a population that is depressed. People in the area are poor and deprived and suffer from severe illnesses and an extraordinary high number of interpersonal injuries. Starting as a 20 bed hospital 100 years ago, it has grown into a relatively modern 800 bed institution. Its experience in trauma management has been unique. It has contributed a great deal to the management of various trauma which has brought this center into international focus. This also led to the development of an organized trauma service as a separate surgical division and was later designated as one of the four certified centers for the treatment of trauma in New York City. Our overall experience over the past five years in trauma management, as well as special types of injuries encountered and their management, will be presented. Planning, teaching and organization of a trauma service will also be discussed so as to share the information and experience with others whose population and type of injuries are similar to ours.

PATTERNS OF DYING FROM SEVERE BLOOD LOSS IN AWAKE RATS

David Crippen, M.D., Peter Safar, M.D. and Catherine Snyder, Pittsburgh, Pennsylvania, U.S.A.

The RRC has established volume controlled hemorrhagic models in awake rats (McGlew et al, Circ Shock 16:35, 1985) and lightly anesthetized monkeys (Bar-Joseph et al, JWAEDM 1, SI:169, 1985). In this study, we determined in awake spontaneously breathing rats, responses and dying patterns in the same model over 3 h post-hemorrhage (H), while monitoring art. and CV pressures, breathing movements, EEG, and art. blood gases. Four groups of 10 rats each were studied. After cannulation under light anesthesia, and awakening, the rats were bled over 20 min: Gr I, 2 ml/100 g H; Gr II, 2.5 ml/100 g H; Gr III, 3 ml/100 g H; Gr IV, 3.5 ml/100 g H. Death was defined as syst. BP \geq 25 mmHg. *Results:* Survival rates were 90% after H 2 ml/100 g; 80% after H 2.5

ml/100 g; 40% after H 3 ml/100 g with mean survival time 73 (7-162) min; and 0% after H 3.5 ml/100 g with mean survival time 32 (1-122) min. Art. press. decreased at end of H to a minimum of 30 mmHg, rose transiently in all groups (attempted self-resuscitation), and then either recovered to near normotension or declined to death. Art. PO₂ increased. Art. PCO₂, pH, Hct decreased. Hyperventilation changed to gasping until pulselessness. EEG depression correlated with hypotension. *Conclusions:* The H 3.5 ml/100 g model is suitable (and will be used next) for the study of responses to field resuscitation potentials suitable for use by laymen, with the aim to delay secondary cardiac arrest from severe hemorrhage while waiting for transport (e.g., external stimuli, rectal fluids, hypothermia, O₂ inhalation). (Assisted by Lisa Porter. Supported by A.S. Laerdal Foundation.)

EFFECTS OF SODIUM BICARBONATE IN CANINE HEMORRHAGIC SHOCK

Thomas J. Iberti, M.D., New York, New York, U.S.A.

We studied the use of sodium bicarbonate (NaHCO₃) administration in a canine model of hemorrhagic shock to determine its effect on lactic acidosis, serum bicarbonate, pH, and cardiac output. Thirteen dogs were anesthetized, paralyzed, mechanically ventilated and hemodynamically monitored. Hypotension was induced and maintained at a MAP of 40-45 mmHg by controlled hemorrhage and reinfusion. After 2.5 hr of shock, the dogs were randomized in 2 groups: A=6 control dogs received NaCl infusion; B=7 dogs received NaHCO₃ 1 mEq/kg followed by a continuous infusion of 2.5 mEq/kg/hr for 2.5 hr.

VARIABLES (MEAN \pm SD)		BASILINE	HEMORRHAGE ONLY	HEMORRHAGE + TREATMENT	<i>Results:</i> *p<0.05
Cl	A	4.3 \pm 0.9	1.1 \pm 0.1	1.2 \pm 0.3	
Cl	B	4.2 \pm 1.3	1.0 \pm 0.2	1.3 \pm 0.3	
LACT	A	0.9 \pm 0.4	8.8 \pm 0.9	5.1 \pm 1.2	
LACT	B	0.7 \pm 0.4	7.1 \pm 3.1	10.1 \pm 3.2	
HCO ₃ ⁻	A	20.1 \pm 1.8	13.4 \pm 1.6	13.6 \pm 2.0	
HCO ₃ ⁻	B	20.1 \pm 1.8	11.1 \pm 3.0	13.3 \pm 2.3	
pH	A	7.34 \pm .03	7.09 \pm .07	7.17 \pm .04	
pH	B	7.37 \pm .02	7.03 \pm .10	7.16 \pm .10	

As in other models of lactic acidosis (hypoxic and phenformin induced), this model demonstrates an increase in lactic acid associated with NaHCO₃ administration. Although the pathophysiologic