The Journal of Laryngology and Otology

EDITED BY
WALTER HOWARTH

ASSISTANT EDITOR G. H. BATEMAN

Contents

VASODILATORS IN THE TREATMENT OF LABYRINTHIN.	
Vascular Insufficiencies	. J. C. SEYMOUR
INFANT TRACHEOSTOMY	. LINDSAY P. GRAY
A New Concept of the Cause of Costen's Temporo	
MANDIBULAR ARTHROSIS	. FRANK COFFIN
RESPIRATORY OBSTRUCTION IN HEMOPHILIA .	. Betsy Brown,
	F. Nour-Eldin
	and John F. WILKINSON
CLINICAL RECORDS—	
TUBERCULOSIS OF THE PALATE—AN UNUSUA	L
Presentation	. R. F. McNab Jones
Hæmangioma of the Mandible Presenting a	s
Epistaxis	. R. FRANCE ROHAN
A Case of Metastatic Amelanotic Melanom.	A STATE OF THE STA
OF THE LARYNX	. OMAR H. SHAHEEN
A Case of Foreign Body in the Naso-Pharyn	G. S. IRVINE

REVIEW OF BOOK

GENERAL NOTES

London

Headley Brothers Ltd

109 Kingsway WC2

Annual Subscription £4/4/0 net, U.S.A. \$13

Monthly, 10/- net post free

The Journal of Laryngology and Otology

(FOUNDED in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY

WALTER HOWARTH

ASSISTANT EDITOR G. H. BATEMAN

1. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the understanding that they are contributed to this Journal solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgement must be made.

2. Manuscripts should be typewritten, on one side only of the paper, and

well spaced. Captions to illustrations should be typed on a separate sheet and

sent at the same time as original photographs, etc.

The Harvard system of recording references should be used, e.g. Green, C., and Brown, D. (1951) J. Laryng., 65, 33. Abbreviations of Journals should follow the style recommended in World Medical Periodicals, published by World Health Organization, 1952.

It is most important that authors should verify personally the accuracy of

every reference before submitting a paper for publication.
3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.

4. Illustration blocks are provided free up to the limit of f to per article; beyond this authors are expected to pay half the cost. Coloured illustrations will

be charged in full to authors.

Blocks will normally be held by the Printers for three years after which they will be destroyed. Any author who has borne a part of the cost of his blocks is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to HEADLEY BROTHERS LTD., 109 Kingsway, London, WC2.

5. Orders for reprints must be sent when returning galley proofs, and for this

purpose special forms are supplied.

6. Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the Journal of Laryngology. If they are willing, at the same time, to submit their own abstract (in English,

French, Italian, or German) it will be welcomed.

7. Editorial communications may be addressed to The Editor, Journal of Laryngology, c/o Headley Brothers Ltd., 109 Kingsway, London, WC2.

8. The annual subscription is four guineas sterling (U.S.A. \$13) post free, and is payable in advance.

9. Single copies will be on sale at 10s. od. each; copies of parts up to Vol.

LXIII which are available may be purchased at 7s. 6d. each.

10. All subscriptions, advertising and business communications should be sent to the publishers, Headley Brothers Ltd., 109 Kingsway, London,

United States of America

Orders for this Journal may be sent through local bookseller, or to STECHERT-HAFNER, INC., 31-33 East 10th Street, New York, or direct to the publishers, Headley Brothers Ltd., 109 Kingsway, London, WC2, England.

O Journal of Laryngology and Otology, 1960

The Journal of Laryngology and Otology

CONTENTS

	PAGE
Vasodilators in the Treatment of Labyrinthine Vascular Insufficiencies. J. C. Seymour (London)	133
INFANT TRACHEOSTOMY. Lindsay P. Gray (Perth, Western Australia)	145
A New Concept of the Cause of Costen's Temporo-Mandibular Arthrosis. Frank Coffin (London)	155
RESPIRATORY OBSTRUCTION IN HÆMOPHILIA. Betsy Brown, F. Nour-Eldin and John F. Wilkinson (Manchester)	
Clinical Records—	
Tuberculosis of the Palate—An Unusual Presentation. R. F. McNab Jones (Manchester)	174
Hæmangioma of the Mandible Presenting as Epistaxis. R. France Rohan (Manchester)	. 178
A Case of Metastatic Amelanotic Melanoma of the Larynx. Omar H. Shaheen (London)	182
A Case of Foreign Body in the Naso-Pharynx. G. S. Irvine, R.N. (Surg. Com.) (Haslar)	188
REVIEW OF BOOK—	
La Chirurgie de la Surdite	192,
General Notes	. 194

For Advertisement space in this Journal apply to: HEADLEY BROTHERS Ltd, 109 Kingsway London WC 2

PERDILATAL

in the treatment of sudden perceptive deafness and Ménière's syndrome.


- * Vascular spasm may well be responsible for the majority of these cases.1
- * Vascular changes in the labyrinth are an important factor in the physiopathology of Ménière's disease.2
- * In the treatment of Ménière's disease emphasis must be put upon early restoration of blood flow to the inner ear.3
- * Perdilatal in doses of at least 6 mg. 4 times a day is the most effective vasodilator for the treatment of these conditions.4
- 1 Fowler, E. P. Ann. Otorhinolaryng., 1950, 59, 980.
- 2 Atkinson, M. Proc. roy. Soc. Med., 1946, 39, 807
- 3 Wilmot, T. J. Brit. med. J., 1957, 2, 1047.
- 4 Wilmot, T. J. Paper read on February 6th, 1959 at a Meeting of the Section of Otology, Roy. Soc. Med.

Perdilatal is available in tablets of 6 mg. as PERDILATAL FORTE in 50's, 250's and 1,000's.



Smith & Nephew Pharmaceuticals Limited

WELWYN GARDEN CITY · HERTFORDSHIRE



- * They can try the different makes of hearing aids
- ★ They get impartial advice on all makes of hearing aids

to S. C. INGRAM

- ★ Their personal needs are given individual and continued attention
- ★ A full report on each fitting recommended is sent to the patient's ear specialist
- ★ They can compare at one consultation the different makes one with another

S. C. INGRAM Independent Hearing Aid Consultant

On the approved List of the National Institute for the Deaf

2 SHEPHERD STREET, SHEPHERD MARKET, LONDON, W.I Hyde Park 9042

In schools for the DEAF— multitone Telesonic INDUCTION LOOP SYSTEM brings new advantages

- 1 Can be installed in adjoining classrooms without risk of 'spill over'.
- 2 Combines good hearing with complete freedom of movement. Unaffected by distance between pupil and teacher.
- 3 Clear speech at high amplification.
- 4 Any number of pupils may use the same installation.
- 5 The child can use the instrument as a normal hearing aid outside school.




The Kindergarten Class at the London Residential School for Jewish Deaf Children where all classrooms have been installed with the Multitone Telesonic Induction Loop System

Write for further details to:

multitone

ELECTRIC COMPANY LTD., 12-20 UNDERWOOD STREET, LONDON, N.I.



It is an excellent idea to advise your patients who are hard of hearing to visit a Philips Audiometric Centre. There is a network of them throughout Britain, where patients can discuss their individual needs with a skilled audiometrician entirely free of charge.

Yours faithfully,

PHILIPS ELECTRICAL LTD

Century House · Shaftesbury Avenue London . W.C.2



'OTAMIDYL'

trade mark

brand

ANTIBACTERIAL | ANTIFUNGAL

EAR DROPS

INDICATED in the treatment of otitis externa and chronic suppurative otitis media.

EFFECTIVE, in the presence of blood, pus and tissue debris against many organisms including pyogenic cocci, some Gram-negative bacilli and a number of pathogenic fungi.

CONVENIENT, for topical application in the form of drops or as a ribbon gauze dressing, in 15 c.c. dropper bottles and 100 c.c. bottles.

Detailed information is available on request

An M&B brand Medical Product



MANUFACTURED BY

MAY & BAKER LTD


Distributors:

PHARMACEUTICAL SPECIALITIES (MAY & BAKER) LTD.


DAGENHAM

15 c.c.
'OTAMIDYL'
EAR DROPS
containing
immograpamydra
hethionate 0151
Nidnodgharrylarane
Dlystochloride 0551
salt in Pricaso N
15 BAKEP LTD DACEN

MA6608/100



all nasal sprays have a DECONGESTANT



most nasal sprays have an ANTIBIOTIC



some nasal sprays have an ANTIHISTAMINIC

but only



Biomydrin

nacal ennav

has all these plus a MUCOLYTIC as well

Thonzonium Bromide, the new mucolytic agent in Biomydrin, penetrates thick mucus and so ensures that all the active principles reach the affected areas.

that is why Biomydrin is the only COMPLETE nasal spray


In a plastic self-sterilizing pack (\(\frac{1}{2}\) fluid ounce). *Supplied on prescription only, so there is no danger of indiscriminate self-medication.

FORMULA: Neomycin Sulphate 0.1%, Gramicidin 0.005%, Thonzylamine Hydrochloride 1.0%, Phenylephrine Hydrochloride 0.25%, Thonzonium Bromide 0.05%. Preserved with Thiomersal 0.002%.



WILLIAM R. WARNER & CO. LTD. EASTLEIGH \cdot HAMPSHIRE

B10 443



USES

DIAGNOSIS

The excellent optical performance of the Hallpike Ear Microscope gives the user greatly increased confidence in the diagnosis of abnormal conditions; and many disorders may be recognized at an earlier stage than has hitherto been possible with "headlamp and speculum" techniques. Thus slight pulsation of swellings and vascular tumours—of high diagnostic significance and almost impossible to detect by conventional methods—can readily be detected under this microscope.

Use of the Siegle attachment permits observation of small bubbles present in fluid collections within the tympanum; it is also invaluable for demonstrating minute perforations. Where defects of the tympanic membrane are present, the middle-ear cavity can be scrutinized minutely; the detailed structure of the stapes, stapedius tendon, round window and Eustachian orifice—not normally visible by ordinary methods of examination—are seen with quite remarkable clarity when the Ear Microscope is used.

TREATMENT

Manipulative procedures may be undertaken with greater confidence and precision if the Ear Microscope and its special instruments are used.

The curettes enable adherent wax and debris to be removed with the greatest precision from the walls of the deep meatus or from the tympanic membrane itself. Cholesteatomatous masses or neoplasms may be examined critically and biopsy specimens obtained by means of the curettes. These may also be used as delicate probes for palpation of the incus and head of the stapes in certain cases of otosclerosis with retraction and atrophy of the tympanic membrane; in this way unmistakable evidence may be obtained of bony fixation of the footplate. Or again, it is sometimes possible to identify by palpation defects of the bony tympanic ring or meatal wall which may underlie granulations.

defects of the bony tympanic ring or meatal wall which may underlie granulations.


Myringotomy or removal of foreign bodies can be undertaken with surprisingly high precision because the optics of the Ear Microscope and the specially angled myringotome blade eliminate all shadows so that the surgeon can see clearly where and what he is cutting. The outfit is invaluable also for post-operative examinations and treatment of mastoid cavities. Granulations can be cauterized and exposed areas of dura mater carefully palpated.

*J. Laryng. (1953) 67, 108.

39 WIGMORE STREET LONDON W.1



5241 WHITBY AVENUE PHILADELPHIA 43, Pa, U.S.A.



Please mention The Journal of Laryngology and Otology when replying to advertisements

the question of topical choice...

in ear infections, 'Otosporin' for:

rapid relief of inflammation and pain wide bactericidal action

no cross-resistance or cross-sensitisation to other antibiotics

'Otosporin' contains 5 mgm. neomycin sulphate, 10,000 units polymyxin B sulphate, and 5 mgm. hydrocortisone per c.c. of aqueous vehicle. **Bottles of 5 c.c. with dropper.**

in skin infections, 'Cortisporin' for:

rapid relief of inflammation and pain, with wide antibacterial activity

negligible risk of skin sensitisation or bacterial resistance

no cross-resistance or cross-sensitisation to other

- 'Cortisporin' Ointment contains 400 units zinc bacitracin, 5 mgm. neomycin sulphate, 5000 units polymyxin B sulphate, and 10 mgm. hydrocortisone per gm. of petrolatum base.

 Tubes of 10 gm. with nozzle.
- 'Cortisporin' Lotion contains 5 mgm. neomycin sulphate, 10,000 units polymyxin B sulphate, and 10 mgm. hydrocortisone per c.c. of aqueous base. *Plastic squeeze-packs of 10 c.c.*



