

GENERAL INDEX.

The numerals in heavy type indicate that the species to which they refer is figured.

- Acacia arabica*, var. *kraussiana*, 116.
 „ *spirocarpoides*, 116.
Acari in N. Nigeria, 322, 328, 346.
acarta, *Olfersia*, 172 note.
accraensis, *Culex*, 247.
 „ *Culicelsa*, 247, 256, 261.
actinella, *Tachardia*, in Natal, 100; compared with *T. longisetosa*, 103.
acuminata, *Kirkia*, 360.
Adansonia digitata, puparium of *Glossina morsitans*, at base of, 359.
Aedides, 242, 243.
Aedinae, 241.
aegypti, *Culex*, 265.
aegypticus, *Gonicocotes*, sp. n., 148.
aegyptium, *Hyalomma*, 197, 212, 230.
aethiopicus, *Phacochoerus*, louse of, 145.
africana, *Anthonina*, 100, 101.
 „ *Atthebella*, 129, 130, 135.
 „ *Fiorinia*, sp. n., 90.
 „ *Lecaniodiaspis*, sp. n., 100-102.
 „ *Mansonia*, 254.
 „ *Stegomyia*, 275.
africanum, *Menopon*, sp. n., 149, 150.
africanus, *Ceroplastes*, common in Egypt, 97.
 „ *Linognathus*, sp. n., 146.
 „ *Mucidus*, 246.
 „ *Tabanus*, 112, 279, 339, 346.
 „ *Taeniorhynchus*, 251.
Afzelia, 117.
aganippea, *Leptopsylla*, 271.
ager, *Culex*, 259; var. *sinensis*, 259.
 „ *Taeniorhynchus*, 256, 259.
- Agromyza salicifolii*, sp. n., 79; bred from mines in willow leaves, Egypt, 80.
aikeni, *Anopheles*, a true *Stethomyia*, 141.
Akidoproctus maximus, 148.
 „ *stenopygus*, from spurwing goose, Sudan, 148; distinct from *A. maximus*, 149.
albicollis, *Corvultur*, lice on, 147, 149, 151.
albida, *Tachardia*, in Natal, 100.
albifasciatus, *Ochlerotatus*, 247.
albipalpus, *Tabanus*, 325, 327, 346.
albitarsis, *Megaculex*, 245.
alboannulatus, *Culicelsa*, 247.
albovirgatus, *Culex*, 259.
Algeria, *Culicinae* in, 248, 249, 251, 265.
alternans, *Mucidus*, 246.
ambigua, *Grabhamia*, 259.
Amblyomma nuttalli, in N. Nigeria, 320.
 „ *variegatum*, in Gambia, 197; in N. Nigeria, 319, 322, 325, 330.
anarmostus, *Culex*, 259, 260.
Anglo-Egyptian Sudan, two new spp. of *Tabanus* from, 173-177; *Cimex rotundatus*, in, 179-180; Anoplura and Mallophaga from, 145-152.
Angola, *Culicinae* in, 252, 259, 262.
annettii, *Chrysoconops*, 253.
 „ *Taeniorhynchus*, 253.
annulata, *Reedomyia*, 320.
 „ *Theobaldia*, 251.
annulifera, *Mansonia*, 253.
annulipes, *Mansonia*, 253.
annulioris, *Culex*, 259.
annulirostris, *Culicelsa*, 259.
Anopheles, new W. African spp. with notes on nomenclature, 141-143.

- Anopheles aitkeni*, appears to be a true *Stethomyia*, 141.
- „ *argyrotarsis*, type of *Nyssorhynchus*, 141.
- „ (*Myzomyia*) *flavicosta*, sp. n., 142-143.
- „ *ludlowi*, considered a var. of *A. rossi*, 143.
- „ *rossi*, type of *Myzomyia*, 141.
- „ *umbrosa*, distinct from *A. funesta*, 142.
- „ *watsoni*, sp. n., 143; occurrence at Minna, 330.
- Anophelina, 241, note; spp. in N. Nigeria, 321, 328, 345.
- Anophelinae, notes on nomenclature, 141.
- Anophelines, infected, where to find, 1; method of capture, 1; identification of species, 1-2; dissection of, 2-5; endo- and ecto-parasites, 6; infection of, 6; problems concerning infection of, 7-8; properties of salivary secretion, 8; suggested lines of research, 8; "black spores" (Ross) in, 5; breeding in native domestic water-vessels, 133; *Haplochilus grahami* eats larvae of, 137; occurrences in Gambia, 201, 205, 217; larvae and pupae abundant in pools N. of Minna, 330.
- Anoplura, from African hosts, 145-147; instructions for collecting, 238-239.
- Antelope, sable, in relation to tsetse, 43, 44, 45.
- antennatum*, *Menopon*, sp. n., 150-151.
- anthropophaga*, *Cordylobia*, 206.
- Antonina africana*, published inadvertently, 100, 101.
- „ (see *Lecaniodiaspis*).
- Anidia glandulosa*, 103-104; in Egypt on *Acacia arabica*, 104.
- Aponomma exornatum*, in N. Nigeria, 320.
- Aporoculex*, included in *Culex*, 256.
- „ *punctipes*, synonym of *Culex quasigelidus*, 258.
- aquilus*, *Culex*, 264.
- arabica*, *Acacia*, 116.
- „ *Mansonia*, 249.
- arabicus*, *Culex*, 249.
- Arachis hypogaea*, 193.
- Argasidae, instructions for collecting, 232; occurrence in N. Nigeria, 346.
- Argas persicus*, 232.
- argyrotarsis*, *Anopheles*, 141.
- arundinum*, *Cervicapra*, lice on, 147.
- Ashanti, *Glossina* taken in, 18, 34; *Culex* and allied genera in, 245, 246, 248, 259, 263, 265.
- asini*, *Haematopinus*, 145.
- Aspidiotus cydoniae*, synonym of *A. lataniae*, 86.
- „ *hederae*, in German E. Africa, 85; on *Magnolia* and *Datura alba*, in Cairo, 86.
- „ *lataniae*, *A. cydoniae* synonym of, 87; on *Antigonon octopus*, in Uganda, 86.
- „ *trilobitiformis*, occurrence in German E. Africa on Oleander, 85.
- Aspidoproctus maximus*, a pest in Rhodesia, 100; occurrence in German E. Africa, 100.
- Aswan (Egypt), Coccidae from, 102, 104.
- Atthegella africana*, in native water-vessels at Lagos, 129, 130, 135.
- Auchmeromyia luteola*, occurrence in Gambia, 205, 206, 212.
- Aulacaspis* (see *Diaspis*).
- aureosquamiger*, *Anopheles*, 143.
- aurites*, *Chrysoconops*, 252.
- „ *Taeniorhynchus*, 252, 253.
- austeni*, *Anopheles*, 143.
- azoriensis*, *Culex*, 263.
- Baboons, 206, 223.
- baculus*, *Lipeurus*, 149.
- Bagana (N. Nigeria), Sleeping Sickness and Horse Trypanosomiasis at, 339.
- Banksinella chrysothorax*, synonym of *B. luteolateralis*, 245.
- „ *luteolateralis*, distribution and synonyms, 245; in N. Nigeria, 317, 337.
- „ *metallicus*, referred to *Taeniorhynchus*, 352.
- „ *punctocostalis*, in Ashanti, 245.
- „ *taeniarostris*, may be a var. of *luteolateralis*, 245.
- barbirostris*, *Myzorhynchus*, 268.
- Baro (N. Nigeria), importance of, 304, 327; mosquito breeding area near, 328; distribution of *Glossina* near, 329, 330; recommendations with regard to, 355; biting flies and ticks at, 328; trypanosomiasis in, 328.

- Bassa Province (N. Nigeria), Sleeping Sickness in, 339; Horse Sickness in, 340-345, 353; recommendations with regard to, 355.
- Bathurst (Gambia), residential quarters, *Stegomyia fasciata* and tsetse infest, 195.
- beckii*, *Mytilaspis*, synonym of *M. citricola*, 88.
- Bed-bugs, in native houses, 212; instructions for collecting, 234.
- Beira (Mozambique), *Glossina morsitans* and *G. brevipalpis* near, 41, and note.
- Bembex capensis*, preying on *Holcoceria nobilis*, 182.
- " *tricolor*, preying on *Tabanus secedens*, 182.
- biclavis*, *Chionaspis*, mining habits of, 88.
- bicolor*, *Chrysops*, sp. n., 162, 163.
- " *Culex*, 266.
- " *Culicada*, 256, 266.
- bifoliata*, *Culex*, 261.
- biguttatus*, *Tabanus*, 112.
- " var. *croceus*, *Tabanus*, 321, 324, 325, 328, 330, 346.
- Bintang Creek (Gambia), in regard to tsetse, 208-211.
- Bironella*, considered distinct from *Anopheles*, 141.
- bisignata*, *Myzomyia*, 268.
- "Black spores" (Ross) in Anophelines, 5.
- blanchardi*, *Parlatoria*, male puparia of, 88, 89; possible variety of, 92.
- Blood-sucking Arthropods, from Gambia, 214-218; from N. Nigeria, 345, 346.
- " Diptera, in Dowa District, Nyasaland, 111, 112; in Ilesha District, S. Nigeria, 273-276.
- Bornu Province (N. Nigeria), haunts of *Glossina tachinoides* in, 119-126.
- bostocki*, *Culex*, 264.
- Boycia mimomyiaformis*, belongs to *Megaculex*, 244, note; should probably be included in *Ludlowia*, 266.
- Brachystegia pectinata*, 117.
- Breeding grounds of *Phlebotomus*, 48-52; destruction of, 55, 56.
- brevicomis*, *Linognathus*, 147.
- brevipalpis*, *Glossina*, group characters, 9; male armature 10-12; male and female, description, 19-21; occurrences in Nyasaland, 19; erroneously referred to as *fusca*, 41, note; colour conventions for, 298, and note.
- British E. Africa, *Glossina* taken in, 31.
- brucei*, *Trypanosoma*, 353.
- Buffalo, in relation to tsetse, 43, 44.
- bullatifrons*, *Haematopota*, 324-327.
- cacti*, *Diaspis*, on *Opuntia* in Egypt, 87.
- Cairo, Coccidae from 85, 86, 97, 102.
- calcitrans*, *Stomoxys*, empty pupae of, in caves in Malta, 48; distribution, 197, 207, 208, 319-321, 323, 324, 327, 333, 346.
- caliginea*, *Glossina*, 10-12, 14, 23-25.
- caliginosus*, *Ochlerotatus*, 250.
- Calosoma sycophantæ*, preying on Gipsy Moth, 369.
- Calvertina*, considered distinct from *Anopheles*, 141.
- calyptroides*, *Diaspis*, in Egypt, 87.
- camelarius*, *Tabanus*, sp. n., 173-175.
- camelina*, *Hippobosca*, 170, 171, 346.
- cana*, *Chrysops*, sp. n., 166.
- canis*, *Ctenocephalus*, 321, 324.
- Cape Colony, Culicidae from, 251, 253.
- capensis*, *Bembex*, 182.
- Cape St. Mary (Gambia), residential quarters of Bathurst to be moved to, 195, 196; site for Sleeping Sickness camp at, 197, 224; area could be easily freed from tsetse, 198.
- capitis*, *Pediculus*, 145.
- Casa Leoni (Malta), *Phlebotomus* in rabbit hutches at, 49, 70; *Phlebotomus perniciosus* abundant at, 72.
- cassiae*, *Chionaspis*, sp. n., 89, 90.
- Cellia*, included in *Nyssorhynchus*, 141; type sp. incorrectly cited, 141.
- centurionis*, *Chrysops*, sp. n., 164-166.
- Ceroplastes africanus*, common in Egypt, 97.
- " *floridensis*, 95, 96.
- " *galeatus*, sp. n., 95.
- " *quadrilineatus*, 95; produces crimson dye, 95.

- Ceroplastes ruscii*, on *Crataegus*, in Cairo, 97.
 " *ugandae*, sp. n., 94; produces crimson dye, 95.
 " *vinsonioites*, sp. n., 96.
Ceroplastodes gowdeyi, sp. n., 98, 99; associated with *Lecanium nigrum*, 100; presence of large paired glands in, 99, 100.
Cervicapra arundinum, Anoplura parasitic on, 147.
 Ceylon, Coccidae on trees imported from, 86, 100.
Chagasia, considered distinct from *Anopheles*, 141.
 Chalcidid, parasitic on *Lecanium tenuivalvatum*, 93; on *Tachardia longisetosa*, 103.
chamberlaini, *Mimomyia*, 244.
Chimaeropsylla potis, n. g. et n. sp. from Nyasa, 269-272.
Chionaspis biclavis, mining habits of, 88.
 " *cassiae*, sp. n., 89, 90; on *Cassia floribunda* in Uganda, 90.
 " *dentilobis*, in Uganda, 90.
 " *longispina*, sp. n., 88; on *Justicia alba* in Egypt, 88; mining habits, 88.
 " *substriata*, on Palm in Uganda, 90.
 Chironomidae, 57; in Gambia, 218.
Chironomus, larvae of, in domestic water-vessels at Lagos, 129, 130, 134.
chloroventer, *Culex*, 264.
choeropotamus, *Potamochoerus*, 145.
chorea, *Dicranomyia*, 263.
Chrysoconops, purely an American genus, 251, 252; African species described as, belong to *Taeniorhynchus*, 252, 253.
 " *nigra*, 252.
 " *pseudoconopas*, 253.
 " *taeniorhynchus*, *Culex titilans*, wrongly determined as, 252.
Chrysops bicolor, 162-163.
 " *cana*, sp. n., 166-168.
 " *centurionis*, sp. n., 164-166.
 " *dimidiata*, 165, 166.
 " *distinctipennis*, in N. Nigeria, 333.
 " *longicornis*, in Gambia, 207.
Chrysops magnifica, sp. n., 161-163.
 " " var. *inornata*, nov. 163.
 " *nearvei*, 163.
 " *silacea*, 166; in N. Nigeria, 341-344.
 " *wellmannii*, 167.
chrysothorax, *Banksinella*, 245.
Cimex rotundatus, in Anglo-Egyptian Sudan, 179; distribution of, 180; common in Gambia, 212.
 Cimicidae, in Gambia, 218; new, from British E. Africa, 363, 364.
cinerea, *Culicomyia nebulosa*, var., 255.
cinereus, *Culex*, 254.
cinnamomi, *Diaspis*, 86; var. *mangiferae*, nov., 86; var. on imported mango trees in Egypt, 86.
cirnei, *Rhynchocyon*, 272.
citricola, *Mytilaspis*, 88.
claritibialis, *Tabanus*, 282.
clavipalpus, *Radioculex*, 244, and note.
 Climate, in Gambia, 191-193; in N. Nigeria, 308-314.
climax, *Trichochetes*, 147.
 Clinocoridae, n. g. et n. sp. of, 363, 364.
Clinocoris pilosellus, 364, note.
 " *pipistrelli*, 364, note.
 Coccidae, 85-104.
 Collectors, hints for, 227-239; collecting and preserving fleas, 231, 232; collecting ticks, 232-234; collecting and preserving Cimicidae, 234; collecting Diptera, 234-237; tsetse, points on which observations are wanted, 238; collecting Anoplura and Mallophaga, 238, 239; notes on bionomics wanted, 230; points to be mentioned on labels, 229, 230.
Colpocephalum flavescens, from vulture, S. Nigeria, 149.
 " *semicinctum*, from *Colvultur albicollis*, S. Nigeria, 149.
 " *sjoestedti*, from spotted rat, Uganda, 149.
Combretum microphyllum, 117.
commovens, *Culex*, 246.
conchiformis, *Inglisia*, in Uganda, 98.
concolor, *Culex*, 261, 262.
condylodesmus, *Culex*, 259, 260, 268.

- Congo Free State, *Glossina* taken in, 18 ;
Culicinae in, 245, 256, 259, 264, 265.
- Conosia irrorata*, African darker than
 Oriental, 262.
- consimilis*, *Culex*, 259.
- " " *tigripes*, var., 259.
- Copaifera mopani*, puparium of *Glossina*
morsitans in hollow at base of trunk of,
 358.
- copemani*, *Tabanus*, sp. n., 281, 282.
- Coquillettidia*, not separable from *Taenior-*
hynchus, 252.
- Cordylobia anthropophaga*, 206.
- Corethrinae, in Gambia, 218 ; sub-family
 or family, 241.
- Corvultur albicollis*, Mallophaga parasitic
 on, 147, 149, 151.
- costalis*, *Pyretophorus* (see under *My-*
zomyia).
- " *Myzomyia* (see *Myzomyia*)
- Cottonera (Malta), *Phlebotomus* found at,
 48 ; breeding places at, 56.
- cristatus*, *Chrysoconops*, 253.
- " *Taeniorhynchus*, 253.
- croceus*, *Tabanus biguttatus*, var. (see
Tabanus).
- crocodilinus*, *Tabanus*, sp. n., 284.
- Crustacea, in native water-vessels at
 Lagos, 129, 130, 134, 135 ; breeding
 in native yards, 136.
- Ctenocephalus canis*, abundant on dogs in
 N. Nigeria, 321, 324.
- " *felis*, in Gambia, 200.
- Culex*, undetermined spp., 7 ; larvæ in
 in water-vessels at Lagos, 129-
 131, 132, 133 ; and allied
 genera, 241-268 ; legs of various
 species compared, 200 ; index
 of specific names of, 266-268 ;
 occurrences in N. Nigeria, 319,
 320, 321, 328, 337, 345.
- " *accraensis*, 247, 261.
- " *aegypti*, 265.
- " *ager*, Oriental, 259.
- " " var., *sinensis*, 259.
- " *albopirgatus*, synonym of *C.*
duttoni, 259.
- " *anarmostus*, synonym of *C. duttoni*
 259 ; remarkable abnormality
 of a, 260.
- " *annulioris*, has been confused with
C. duttoni, 259 ; distribution,
 259.
- Culex aquilus*, synonym of *C. invidiosus*,
 264.
- " *arabicus*, probably a homonym of
Mansonia arabica, 249.
- " *azoriensis*, included as a var. of
C. pipiens, 263.
- " *bicolor*, 266.
- " *bifoliata*, synonym of *C. thalassius*,
 261.
- " *bostocki*, synonym of *C. salis-*
huriensis, 264.
- " *chloroventer*, synonym of *C. in-*
vidiosus, 264.
- " *cinereus*, synonym of *Culiciomyia*
nebulosa, 254.
- " *commovens*, a var. of *Mucidus*
mucidus, 246.
- " *concolor*, compared with *C.*
tigripes, 261, 262.
- " *condylodesmus*, probably only *C.*
duttoni, 259, 260, 268.
- " *consimilis*, synonyms and distri-
 bution of, 259.
- " *decens*, in Gambia, 205 ; synonyms
 and distribution of, 263 ; in N.
 Nigeria, 319, 328, 345.
- " *didieri*, 266.
- " *dissimilis*, synonym of *C. duttoni*,
 259, 260.
- " *dorsalis*, 247, 248, 249.
- " *drymoecius*, a var. of *Taenior-*
hynchus fuscopennatus, 253.
- " *duttoni*, larvæ in native water
 vessels at Lagos, 129 ; number
 of vessels in which were larvæ
 of, 130 ; seasonal variation of
 larvæ of, 131 ; distribution of
 larvæ, 132 ; sensitive to amount
 of chlorine in water, 132 ;
 possible carrier of *Filaria*
diurna, 133 ; synonyms of,
 259 ; distribution of, 260 ;
 mid leg of, 260 ; common at
 Oshogbo in October, 275 ; in
 N. Nigeria, 345.
- " *euclastus*, synonym of *C. in-*
vidiosus, 264.
- " *fatigans*, included as a var. of *C.*
pipiens, 262 ; in N. Nigeria,
 262, note, 345.
- " *freetownensis*, 254 ; specimens
 wrongly determined as, 255.
- " *grahami*, 263, 264 ; in N. Nigeria,
 319, 345.

- Culex* *guiarti*, 263; green colour not a specific character of, 263; in N. Nigeria, 319, 345.
- „ *hirsutipalpis*, synonym of *C. duttoni*, 259.
- „ *insignis*, 265.
- „ *invenustus*, 252.
- „ *invidiosus*, 241, 264; in N. Nigeria, 319, 321, 337, 345.
- „ *jamaicensis*, is type of *Grabhamia*, 247, 256.
- „ *laurenti*, 266.
- „ *leucacanthus*, synonym of *Ochlerotatus pulcritarsis*, 249.
- „ *lividocostalis*, synonym of *C. decens*, 263.
- „ *longefurcatus*, 265.
- „ *maculicrura*, 261.
- „ *maculiventris*, 265.
- „ *masculus*, synonym of *C. decens*, 263.
- „ *melanorhinus*, 265.
- „ *microannulatus*, 261.
- „ *minutus*, synonym of *C. decens*, 263.
- „ *mundulus*, 266, 268.
- „ *neaxei*, synonym of *C. guiarti* 263.
- „ *nigeriensis*, 250.
- „ *nigrochaetae*, 252, 254, 255, 256.
- „ *nigrocostalis*, larvae in water vessels at Lagos, 129-131; larvae favourite food of *C. tigripes*, 133; synonym of *C. decens*, 263.
- „ *ornatothoracis*, 263; thorax of, **264**.
- „ *pallidocephala*, included as a var. of *C. pipiens*, 262.
- „ *pallidostriatus*, synonym of *Ochlerotatus ochraceus*, 250.
- „ *pallidothoracis*, synonym of *C. pruina*, 263.
- „ *pallipes*, 265.
- „ *par*, allied to or identical with *C. consimilis*, 266.
- „ *parascelos*, synonym of *Ochlerotatus ochraceus*, 250.
- „ *pettigrewi*, synonym of *C. theileri*, 250, 262.
- „ *viliferus*, 265.
- Culex* *pipiens*, *Ochlerotatus* compared with, 247; varieties and distribution, 248, 262, 263.
- „ *pruina*, 263; thorax of, **264**.
- „ *pruinus*, synonym of *C. pruina*, 263.
- „ *pseudoannulioris*, synonym of *C. consimilis*, 259.
- „ *pseudocinereus*, synonym of *Culicomyia nebulosa*, 254.
- „ *pulcritarsis*, 249.
- „ *pullatus*, synonym of *C. grahami*, 264.
- „ *pusillus*, 265.
- „ *pygmaeus*, 266.
- „ *quasigelidus*, synonyms of, 258; palpi of, **258**; distribution of, 259; hind leg of, **260**; thorax of, **264**.
- „ *quasiguarti*, synonym of *C. zombaensis*, 264; in N. Nigeria, 321, 328, 345.
- „ *rima*, distribution, 265; synonym of *Melanoconion rimus*, 265.
- „ *rubinotus*, 265.
- „ *rufinus*, 265.
- „ *salisburyensis*, synonym of *C. bostocki*, 264.
- „ *salsus*, synonym of *C. somaliensis*, 261.
- „ *salus*, synonym of *C. somaliensis*, 261.
- „ *sergenti*, in Algeria, 265.
- „ *simpsoni*, 263.
- „ *somaliensis*, possibly a form of *C. microannulatus*, 261.
- „ *stoehri*, included as a var. of *C. pipiens*, 263.
- „ *taeniorhynchoides*, synonym of *C. quasigelidus*, 258.
- „ *thalassius*, synonyms and distribution, 261.
- „ *theileri*, mid leg of, **260**; *C. pettigrewi*, synonym of, 262; distribution of, 262.
- „ *tigripes*, closely related to *C. concolor*, 261.
- „ *tigripes*, var. *consimilis*, is a distinct species, 259.
- „ *tigripes*, var. *fusca*, larvae in water vessels at Lagos, 129-131;

- wholly carnivorous, 133; includes darkest forms, 262; occurrence in N. Nigeria, 321, 345.
- Culex titillans*, wrongly determined as *Culicelsa taeniorhynchus*, 252.
- „ *transvaalensis*, synonym of *Ochlerotatus hirsutus*, 249.
- „ *univittatus*, hind leg of, 260; synonyms and distribution, 250, 262, 345.
- „ *vigilax*, 247, 250.
- „ *viridis*, synonym of *C. guiarti*, 263.
- „ *zeltneri*, 266.
- „ *zombaensis*, 264.
- Culicada*, synonym of *Ochlerotatus*, 246.
- „ *bicolor*, included in *Culex*, 256, 266.
- „ *fuscopalpalis*, synonym of *Ochlerotatus cumminsii*, 248.
- „ *lateralis*, 250.
- „ *mediopunctata*, var. of *Ochlerotatus cumminsii*, 248.
- „ *morsitans*, included in *Theobaldia*, 251.
- „ *nemorosa*, referred to *Ochlerotatus*, 248.
- „ *quasimodesta*, seems to belong *Culex*, 256, 266.
- Culicales, separate from Megalorhini, 241; table of genera, 242; table of females, 243; contiguous eyes in nearly all, 256.
- Culicelsa*, synonym of *Ochlerotatus*, 246, 247.
- „ *accraensis* (*neotaeniorhynchus*), is a *Culex*, 247, 256, 261.
- „ *alboannulatus*, is a true *Ochlerotatus*, 247.
- „ *annulirostris*, included in *Culex*, 256.
- „ *neotaeniorhynchus*, same as *Culex thalassius*, 261.
- „ *taeniorhynchus*, type of genus, 247; included in *Ochlerotatus*, 247, 249; *Culex titillans*, wrongly determined as, 252.
- Culicidae, in water-vessels at Lagos, 129-134; in Gambia, 217; classification of, 241; African forms often much darker than the Oriental, 262; six new spp. from Kamerun and Togo, 268; spp. from S. Nigeria, 274, 275; in N. Nigeria, 231, 238, 345.
- Culicides, 242, 243.
- culicifacies*, *Myzomyia*, 6.
- Culicina, 241, note.
- Culicinae, 241, 242.
- Culicines, where to find, 1; black spores in oocysts of *C. fed* on *Proteosoma*, 5; investigation of commoner house frequenting species should not be neglected, 6; can they transmit malaria, 7; five spp. of *C. larvæ* eaten by *Haplochilus grahami*, 137; *C. larvæ* and pupæ abundant in pools north of Minna, 330.
- Culiciomyia*, only one variable African sp. of, 255; synonyms of, 254-256.
- „ *dalzielii*, synonym of *Ochlerotatus quasiunivittatus*, 250, 255.
- „ (?) *insignis*, sp. n. 37, 38; referred to *Culex*, 265.
- „ *nebulosa*, synonyms and varieties of, 254-256; common in W. Africa, 256.
- „ *nebulosa*, var. *cinerea*, 255.
- „ „ „ *freetownensis*, 255.
- „ „ „ *fusca*, 255.
- „ „ „ *pseudocinerea*, 255; sp. described as *Culex invenustus* is probably, 256.
- „ *uniformis*, synonym of *C. nebulosa*, 255.
- Culicoides*, in N. Nigeria, 319, 324.
- Culiseta absobrina*, included in *Theobaldia*, 251.
- cumminsii*, *Ochlerotatus*, 248.
- cuniculi*, *Spilopsyllus*, 269.
- Cyclops similimus*, present in water-vessels at Lagos, 129, 130, 134; of importance in relation to transmission of Guinea Worm, 134; distribution of, 134.
- cydoniae*, *Aspidiotus*, synonym of *A. lataniae*, 86.
- Cyniphes molestus*, synonym of *Phlebotomus papatasii*, 73.

Cyprinodontidae, feeding on mosquito larvae, 137-139, 182.

Cypris, in water-vessels at Lagos, 129, 130, 135; harbour *Taenia*, Trematode larvae and *Acanthocephalus*, 135.

Dactylopius longispinus, on vine in Egypt, 100.

dalzieli, *Culiciumyia*, 250, 255.

Daphnia, in water-vessels at Lagos, 129, 130, 135; in relation to *Taenia*, Trematode larvae and *Acanthocephalus*, 135.

decens, *Culex*, 205, 263, 319, 328, 345.

decora, *Haematopota*, 197, 275, 321, 327, 331, 334.

decorella, *Tachardia*, new to Africa, 85, 100; compared with *T. longisetosa*, 103.

defassa, *Cobus*, *Hippobosca* on, 170, 171.

Delagoa Bay, Culicinae in, 246, 250, 261.

Dendromyinae, 241.

dentatus, *Ochlerotatus*, 248.

dentilobis, *Chionaspis*, 90.

Dermatophilus penetrans, in N. Nigeria, 325.

Diaspinæ, 86, 87, 90.

Diaspis cacti, on *Opuntia* in Egypt, 87.

„ *calyptroides*, 87.

„ (*Aulacaspis*) *cinnamomi*, var. *mangiferae*, nov., 86; in Egypt on imported mango trees, 86.

„ *cinnamomi*, 86.

„ *regularis*, sp. n., 86-87.

Dicranomyia chorea, 263.

didieri, *Culex*, 266.

digitata, *Adansonia*, 359.

dimidiata, *Chrysops*, 165, 166.

Diptera, photography of, 153-160; instructions for collecting, 234-237; occurring in N. Nigeria, 321, 328, 345-346; pupæ among exposed roots of trees in S. Rhodesia, 357, 359.

dissimilis, *Culex*, 259, 260.

distinctipennis, *Chrysops*, 333.

ditaeniatus, *Tabanus*, 111, 200, 207, 321, 324, 346.

diurna, *Filaria*, in S. Nigeria, 132; possibly carried by *Culex duttoni*, 133.

Dogs, piroplasmiasis of, 323-354; trypanosomiasis of, 344.

dorsalis, *Culex*, 247-249.

dorsalis, *Grabhamia*, 248.

dorsalis, *Ochlerotatus*, 248, 249.

Dowa district (Nyasa), habits of blood-sucking flies in, 111-112.

draperi, *Sphaerococcus*, synonym of *S. marlatti*, 104.

Drosophila, in water-vessels at Lagos, 129, 130, 134.

drymoecius, *Culex*, 253.

dukei, *Olfersia*, sp. n., 171, 172.

durandi, *Enyalopsis*, 180.

durbanensis, *Grabhamia*, 250.

durbanensis, *Ochlerotatus*, 250.

duttoni, *Culex* (see *Culex*).

Ebute Metta (S. Nigeria), Mosquito larvae in water-vessels at, 129-136; other larvae and Crustacea, 134, 135; analysis of water containing larvae of *Culex duttoni* at, 132.

Eggs, of Mosquitos, 7; of *Phlebotomus*, 48, 50, 65, 67; of *Phlebotomus papatasi*, 75; of *Pseudoeripopsylla*, 106; of Mallophaga, 239.

Egg parasites of Gipsy Moth and Brown-tail Moth, 369.

Egypt, Coccidæ from, 85-88, 91, 92, 97, 100, 102, 103-104.

eknomicus, *Eoctenes*, 152.

elongatum, *Lecanium*, on *Albizzia* in Uganda, 92; antenna and spines of, 92.

Entebbe (Uganda), new mosquito from, 38, Coccidæ from, 86-88, 90, 92-95, 97, 98, 103.

Enyalopsis durandi, injurious to man, 180.

„ *petersi*, emits a yellow fluid when handled, 180.

Eoctenes eknomicus, 152.

Ephemeridæ, occurrence in native water-vessels in S. Nigeria, 129, 130, 134.

Epiaburgi, 241.

Eretmapodites, contiguous eyes not found in, 256.

Essau (Gambia), swarms with *Glossina palpalis*, 213; removal of town recommended, 213, 224.

- Etorleptomyia* probably a synonym of *Mansonia*, 253.
 „ *mediolineata*, 253, 254.
euclastus, *Culex*, 264.
Eugenia guineensis, 117.
Eulecanium (see *Lecanium*).
Eumelanomyia inconspicua, wrongly determined as *Culex freetownensis*, 255.
exornatum, *Aponomma*, 320.
- falcatus*, *Rhipicephalus*, 330.
fasciata, *Stegomyia* (see under *Stegomyia*).
fasciatus, *Tabanus*, 202, 321, 324, 346.
fatigans, *Culex*, 262, and note, 345.
felis, *Ctenocephalus*, 200.
Ficus, puparium of *Glossina morsitans* between roots of, 355.
Filaria diurna, occurrence in S. Nigeria, 132; possibly carried by *Culex duttoni*, 133.
Fiorinia africana, sp. n., 90.
 Fish and mosquito larvae, 137-139, 182.
 Flagellata, found in Anophelines, 6.
flavescens, *Colpocephalus*, 149.
flavicosta, *Anopheles*, sp. n., 142.
 Flea, new, occurring on *Rhynchocyon cirnei*, 272.
 Floriana (Malta), *Phlebotomus* breeding at, 51; *Phlebotomus minutus* taken at, 70; *P. perniciosus* abundant at, 71.
floridensis, *Ceroplastes*, 95, 96.
 Fly disease (see Trypanosomiasis of Cattle).
Forcipomyia regulus, 57.
forficula, *Limnathus*, sp. n. 147.
freetownensis, *Culex*, 254, 255.
 „ *Culicomyia nebulosa*, var., 255.
fulvus, *Chrysoconops*, 252.
Fundulus taeniopygus, preys on mosquito larvae, 182.
funesta, *Anopheles*, 142.
 „ *Myzomyia*, 6, 320, 321, 337.
fusca, *Culex tigripes*, var., 129-131, 133, 262, 321, 345.
 „ *Culicomyia nebulosa*, var., 255.
 „ *Glossina* (see under *Glossina*).
fuscipes, *Glossina* (see under *Glossina*).
fuscopalpalis, *Culicada*, 248.
fuscopennatus, *Chrysoconops*, 253.
fuscopennatus, *Taeniorhynchus*, 253.
fuscum, *Protomelanocnion*, 244, 245, 264.
fuscus, *Nirmus*, 147.
 „ *Pectinopalpus*, 129-132, 244, 245, 255.
galeatus, *Ceroplastes*, sp. n. 95.
gambensis, *Lipeurus*, 149.
 „ *Plectropterus*, lice on, 148, 149, 151.
 Gambia (British W. Africa), research in, 187-239; insect-borne diseases in, 188; geography and vegetation of, 188-191; climate and rainfall, 191-193; tribes inhabiting the Colony and Protectorate, 193-195; blood-sucking Arthropods from, 214-218; native names of blood-sucking insects and other Arthropods, 219; bionomics of *Glossina* in, 220-223; remedial measures and recommendations, 223-226; *Culex* and allied genera in, 258, 259, 261, 264.
 „ River, description of, 189-190; islands in, swarming with tsetse, 190; possibility of disease being carried from or into Senegal along, 195; *Glossina palpalis* haunts on banks of, 201-204; clearing spaces at wharves on, 225.
gambiense, *Trypanosoma* (see under *Trypanosoma*).
 Game, in relation to tsetse, 43-46, 113-118, 223.
 Garara, River (N. Nigeria), *Glossina* and mosquitos but no Tabanidae found near, 335-337; Sleeping Sickness not found in villages on, 352.
gelida, var. *sinensis*, *Leucomyia*, 259.
 German E. Africa, Coccidae from 85, 98, 100; new *Chrysopa* from, 162; *Glossina brevipalpis* transmitting cattle-disease in, 181, 182; *Culex* and allied genera in, 251, 262.
gigas, *Goniocotes*, 148.
Girardinus poeciloides, preys on mosquito larvae, 182.
glandulosa, *Aomidia*, sp. n., 103, 104.
Glossina, scheme of classification, 9, 10; general characters of male armature, 10-12; methods of preparing male genitalia for microscopical examination, 13, 14; synopsis of spp., 14,

- 15; description of spp., **15-36**; game in relation to, **43-46, 113**; disappearance from Transvaal of, **115**; can exist without game, **40, 117**; colour of clothing attracting or repelling, **210, 211**; instructions for collecting, points on which observations are wanted, **238**; colour conventions for, **297-299**; distribution in N. Nigeria, **347-351**; information desired concerning, **356**.
- Glossina brevipalpis*, group characters of, **9, 14**; male armature of, **10-12**; male and female, description, **19-21**; in Mozambique, **40-42**; erroneously referred to as *fusca*, **41, note**; probable vector of cattle trypanosomiasis, **181**.
- " *caliginea*, group characters of, **10, 14**; male armature of, **23-25**.
- " *fusca*, group characters of, **9**, and *note*; male armature of, **15-17**; purely western, **41, note**; not found in Gambia, **212**; colour conventions for group of, **297-299**; distribution in N. Nigeria, **347**, and *note*, **350**.
- " *fucipes*, group characters of, **10, 15**; description of male, **26, 27**.
- " *grossa*, **9**, and *note*, **17**.
- " *longipalpis*, group characters of, **10, 15**; male armature, **29, 30**; compared with *G. pallidipes*, **31**; occurrence in N. Nigeria, **328, 329, 336, 337, 341, 344, 346, 347, 349**.
- " *longipennis*, group characters of, **9, 14**; male armature of, **21, 22**.
- " *maculata*, female only known of, **9**; group characters of, **14**; distinct from *palpalis*, **22**.
- " *medicorum*, male armature of, **107, 109, 110**.
- " *morsitans*, group characters of, **10, 15**; male armature of, **32, 33**; dark and light varieties of, **35**; in Mozambique, **40-42**; habits in Nyasa, **43-46, 112**; extension of area of, **112**; in relation to human trypanosomiasis, **112**; in relation to cattle trypanosomiasis, **181, note**; occurrences in Gambia, **199-201, 206, 207, 212, 213, 217**; bionomics of, **222, 223**; predilection for baboons, **206, 207, 223**; preliminary stages of, **291-295**; preserving and feeding captive, **292**; breeding habits, **292, 293**; larva of, **293, 294**; puparium of, **294, 295**; duration of pupal stage of, **295**; colour conventions for group of, **297, 299**; breeding haunts of, **357-361**.
- Glossina nigrofusca*, group characters of, **9**, and *note*, **14**; description of male and female of, **17-19**.
- " *palligera*, group characters, **10, 15**; male armature of, **27-29**.
- " *pallidipes*, group characters, **10, 15**; male armature, **30-32**; compared with *longipalpis*, **31**; in Mozambique, **40-42**; colour conventions for, **297**.
- " *palpalis*, group characters of, **10, 14**; male armature of, **22, 23**; male from Rhodesia possibly distinct from, **107**; distribution in Gambia, **197-199, 200, 203-213, 216**; hunted by a spider, **210**; bionomics of, **220-222**; common in Ilesha district, S. Nigeria, **276**; colour conventions for group of, **297-299**; distribution in N. Nigeria, **317-323, 326, 328, 331-345, 347, 349-353, 355**.
- " *palpalis* var. *wellmani*, **23**.
- " *submorsitans*, distinct from *morsitans*, **9**; group characters, **10, 15**; male armature, **34, 35**; compared with *morsitans*, **33, 36**; two varieties of, **33**, occurrences in Ashanti, **34**; distribution in N. Nigeria, **320-322, 324, 326, 328-330, 338, 339, 346, 349, 350**.
- " *tabaniformis*, relationship, **9** and *note*; male armature, **108**.
- " *tachinoides*, alliances and distinctions, **9**; group characters, **10, 15**; male armature, **25**; haunts in N. Nigeria, **119-126**; var. of *G. palpalis* compared with, **202, 203, 204, note**; colour conventions for, **297-299**; distribution in N. Nigeria, **317-322, 326, 328, 329, 332-339, 343, 344, 347, 349-351**.
- Gold Coast, *Glossina tabaniformis* in, **109**; outbreak of Yellow Fever in, **127**;

- Crustacea found in water-vessels in, 134, 135; *Culex* and allied genera in, 246, 263, 264.
- Goniocotes aegypticus*, sp. n., from blue pigeon, Sudan, 148.
- " *gigas*, from guinea - fowl, Sudan, 148.
- " *major*, compared with *aegypticus*, 148.
- " *procerus*, compared with *aegypticus*, 148.
- gowdeyi*, *Ceroplastodes*, gen. et sp. n., associated with *Lecanium nigrum*, 92, 100; described, 98-100; presence of large paired glands in, 99, 100.
- Grabhamia*, synonym of *Culex*, 247.
- " *ambigua*, 259.
- " *dorsalis*, 248.
- " *durbanensis*, 250.
- " *jamaicensis*, 256.
- " *maculosa*, an extreme form of *Ochlerotatus nemorosus*, 248.
- " *ocellata*, 250.
- " *pulcripalpis*, probably synonym of *Culex dorsalis*, 248, 249.
- " *pulcritarsis*, 249.
- " *subtilis*, pale var. of *Culex dorsalis*, 248.
- " *taeniarostris*, 259.
- " *wilcocksii*, pale var. of *Culex dorsalis*, 248.
- gracilis*, *Haematopota*, 324.
- " *Palaeopsylla*, 269.
- grahami*, *Culex*, 263, 264, 319, 345.
- " *Haplochilus*, preys on mosquito larvae, 137-139; description of, 137, note, 139.
- " *Mucidus*, 246.
- Grassia*, type *Anopheles rossi*, 141.
- gratus*, *Tabanus*, 175, 321, 330, 334, 346.
- grossa*, *Glossina*, 9 and note, 17.
- guiarti*, *Culex*, 263, 319, 345.
- guineensis*, *Eugenia*, 117.
- Haliaëtus vocifer*, *Olfersia* on, 172.
- Haplochilus grahami*, preys on mosquito larvae, 137; description of, 137, note, 139.
- " *pumilus*, preys on mosquito larvae, 182.
- Haplochilus senegalensis*, 137, note.
- Haemaphysalis leachi*, occurring on dogs in Gambia, 212; on dogs in N. Nigeria, 324; biting man, 326.
- Haematopinus asini*, from horse, Sudan, 145.
- " *peristictus*, sp. n., from wild pigs, Nyasa, 145-146.
- " *suus*, compared with *H. peristictus*, 145.
- Haematopota bullatifrons*, in N. Nigeria, 324-327.
- " *decora*, in Gambia, 197; common in Oshogbo, S. Nigeria, 275; does not bite man if animals present, 275; in N. Nigeria, 321, 327, 331, 333, 334.
- " *gracilis*, in N. Nigeria, 324.
- " *laccessens*, in N. Nigeria, 321, 324-326.
- " *nactans*, method of biting, 111.
- " *pallidipennis*, in N. Nigeria, 321, 320-334.
- " *pertinens*, habitat and habits, 111.
- " *puniens*, in N. Nigeria, 326.
- " *tenuicrus*, in N. Nigeria, 321.
- " *vittata*, habitat and habits, 111.
- hederae*, *Aspidiotus*, in German E. Africa, 85; at Cairo on *Magnolia* and *Datura alba*, 86.
- Helea (Forcipomyia) regulus*, larval skin remains attached to pupa of, 57.
- Hemiptera, 238.
- Heptaphlebomyia*, included in *Culex*, 256.
- " *montforti*, synonym of *Culex univittatus*, 262.
- " *simplex*, ♀ synonym of *Culex univittatus*, 262; ♂ of *C. decens*, 263.
- Heptaphlebomyinae, 242.
- Haemasson minutus*, 73.
- hildegardeae*, *Taphozous*, 364.
- Hippobosca camelina*, compared with *H. hirsuta*, 170; in N. Nigeria, 346.
- " *hirsuta*, sp. n., on *Kobus defassa*, Uganda, 169.

- Hippobosca hirsuta*, var. *neavei*, nov., on *Kobus defassa*, N.E. Rhodesia, 171.
- " *maculata*, compared with *H. hirsuta*, 169, 170; in N. Nigeria, 319, 321, 322, 324, 325, 330, 346.
- Hippoboscidae, not recorded from Gambia, 215; in Oshogbo, S. Nigeria, 275; in N. Nigeria, 322, 346.
- Hippocentrum versicolor*, in Oshogbo, S. Nigeria, 275; in N. Nigeria, 319, 320, 324, 325, 327, 330, 346.
- hirsuta*, *Hippobosca*, 169.
- " var. *neavei*, *Hippobosca*, 171.
- hirsutipalpis*, *Culex*, 259, 260.
- hirsutus*, *Ochlerotatus*, 249.
- hispida*, *Hispidimya*, 244, 245.
- " *Ludlowia*, 245.
- Hispidimya hispida*, little different from *Ludlowia chamberlaini*, 244, 245.
- Holocereria nobilis*, preyed upon by *Bembex*, 182.
- Horse Sickness, enormous increase in Ilesha and Oshogbo districts of, 276; in N. Nigeria, 319, 320, 323, 324, 326, 330, 332, 334, 336-339, 341-345, 353-354; segregation camps for, 320, 323, 355; investigation required into forms of, 356.
- Hyalomma aegyptium*, in Gambia, 197, 212; in N. Nigeria, 330.
- Hydrachnidae, larvae found in Anophelines, 6.
- Hymenoptera, parasites of Gipsy Moth and Brown-tail Moth, 369, 370.
- hypogaea*, *Arachis*, 193.
- Ibo (Port. Nyasa), *Glossina morsitans* near, 39, 40.
- Icerya longisetosa*, at Lake Nyasa, 100; on *Acacia* in German E. Africa, 100.
- Ilesha district (S. Nigeria), blood-sucking flies of, 273-276; Mallophaga from, 147.
- immaculata*, *Anopheles*, 141.
- inconspicua*, *Eumelanomyia*, 255.
- Inglisia conchiformis*, in Uganda, 98.
- Insecticides, kerosene for, 104, 355.
- insignis*, *Culex*, 265.
- " *Culicomyia*, 37.
- intertropica*, *Olfersia*, 172, and note.
- invenustus*, *Culex*, 256.
- invidiosus*, *Culex*, 241, 264, 321, 337, 345.
- Inyamandzi river (Mozambique), *Glossina morsitans* near, 41.
- Inyambane district, disappearance of tsetse from, 42.
- irrorata*, *Conosia*, 262.
- irroratus*, *Tabanus*, 286.
- Ixodidae, in N. Nigeria, 322, 328, 346.
- Ixodoidea, in Gambia, 218.
- Izon (N. Nigeria), tsetses swarm at important ford near, 334.
- jacksoni*, *Pulvinaria*, in Uganda and W. Africa, 93; food plants, 93.
- jamaicensis*, *Culex*, 247, 256.
- " *Grabhamia*, 256.
- Kala-azar, *Cimex rotundatus* in relation to, 179; case in Sudan of, 180.
- Katagum R. (N. Nigeria), *Glossina submorsitans* from, 34; Sleeping Sickness reported from, 352.
- Kateri (N. Nigeria), Sleeping Sickness prevalent in, 330-333; blood-sucking flies at, 331-333; removal recommended, 332, 355.
- Kerawan (Gambia), tsetse troublesome at, 198; swarms of *Glossina palpalis* a mile from, 198; elephantiasis prevalent in, 199; clearing to be done at, 225.
- Kerosene, emulsion for destroying Coccidae, 104; use in pools for mosquito larvae, 355.
- Khartoum, Coccidae from, 97.
- Kingia*, 268.
- kingii*, *Mimetecculex*, 250.
- kingsleyi*, *Tabanus*, 320.
- Kirkia acuminata*, puparia of *Glossina morsitans* at base of, 360.
- Kobus defassa*, Hippoboscidae on, 170, 171.
- Kurimis, description of, 307; distribution of, 307, 323, 330-333, 336, 338, 340-344; *Glossina palpalis* generally found in, 337; villages should be moved from, 344; rest camps should be moved further from, 355.
- laccensens*, *Haematopota*, 321, 324-326.
- Lagos, Anophelines occurring at, 6; examination of native water receptacles at, 127-136, analysis of water at, 132; mosquito larvae at, 129-136.

- Lake Chad, of great interest in connection with *Glossina*, 305; system of, 306; area free from *G. palpalis*, 353.
- Lake Nyasa, Coccidae from, 100.
- Lake Salisbury (Uganda), *Pulvinaria jacksoni* found on Cotton near, 93.
- Lasioconops*, included in *Culex*, 256.
- " *poicilipes*, synonym of *Culex quasigelidus*, 258.
- lataniae*, *Aspidiotus*, 86.
- lateralis*, *Culicada*, 250.
- " *Ochlerotatus*, 250.
- latipes*, *Tabanus*, 318, 321, 328, 346.
- laurenti*, *Culex*, 266.
- laverani*, *Tabanus*, 205, 346.
- Laverania*, type, *Anopheles argyrotarsis*, 141.
- leachi*, *Haemaphysalis*, 212, 324, 326.
- Lecaniodiaspis africana*, sp. n., on *Acacia arabica* in Egypt, 100-102.
- " *mimosae*, 102.
- " *prospodidis*, 102.
- Lecanium elongatum*, (= *L. longulum*), on *Albizia* in Uganda, 92.
- " *longulum*, 92.
- " (*Saissetia*) *nigrum*, on *Ficus* in Uganda, 92; associated with *Ceroplastodes gowdeyi*, 100.
- " *persicae*, 93.
- " *tenuivalvatum*, on Citronella grass in Uganda, 92; parasitised by a Chalcidid, 93.
- " (*Eulecanium*) *tremae*, in Uganda, 93.
- Lepidoptera, pupae in hollow of a Baobab, 359; parasites of Gipsy Moth and Brown-tail Moth, 365-371; controlling agencies for, 367, 368, 371, and note.
- Leptopsylla aganippes*, 271.
- " *musculi*, 269, 270.
- leucacanthus*, *Culex*, 249.
- leucarthritis* *Ochlerotatus*, 251.
- Leucomyia*, included in *Culex*, 247, 256.
- " *gelida*, var. *sinensis*, 259.
- " *quasigelida*, 258.
- " *scapularis*, referred to *Ochlerotatus*, 247.
- " *sinensis*, var. of *Culex uger*, 259.
- leucostomus*, *Tabanus*, 176, 177.
- Liberia, *Glossina medicorum* from, 110.
- Light, deterrent for *Phlebotomus*, 54; artificial light attracts, 54.
- Limatinae, 241.
- lineatopennis*, *Taeniorhynchus*, 245.
- Linognathus africanus*, sp. n., from sheep, S. Nigeria, 146.
- " *brevicomis*, 147.
- " *forficula*, sp. n., from reed-buck, Nyasa, 147.
- " *tibialis*, 147.
- " *vituli*, from calf, Nyasa, 146.
- Lipeurus baculus*, from blue pigeon, Sudan, 149.
- " *gambensis*, from spurwing goose, Sudan, 149.
- lividocostalis*, *Culex*, 263.
- Lokoja (N. Nigeria), importance of, 304, 318-320; *Myzomyia costalis* prevalent in, 319; ticks troublesome in, 319; mosquito larvae near, 319; trypanosomiasis in, 320, 353; recommendations with regard to, 355.
- longefurcatus*, *Culex*, 265.
- longicornis*, *Chrysops*, 207.
- longifolia*, *Rhus*, 117.
- longipalpis*, *Glossina* (see under *Glossina*).
- " *Stegomyia*, 268.
- longipennis*, *Glossina* (see under *Glossina*).
- longisetosa*, *Icerya*, 100.
- " *Tachardia*, sp. n., 102, 103.
- longispina*, *Chionaspis* sp. n., 88.
- longispinus*, *Dactylopius*, 100.
- longisquamosus*, *Ochlerotatus*, 249.
- longulum*, *Lecanium* (see *L. elongatum*).
- lowisii*, *Reedomyia*, 250.
- Loxaspis mirandus*, g. et sp. n., from British E. Africa, 363, 364.
- Ludlowia*, essential characters and table of spp. of, 244.
- " *hispidia*, 245.
- " *mimomyiaformis*, in N. Nigeria, 337.
- " *pincerna*, 245.
- " *plumosa*, 245.
- Lujenda River (Nyasa), tsetse-flies plentiful near, 39, 40, 44, 45; tsetse absent west of, 118.
- luridum*, *Trinoton*, 151.
- luteola*, *Auchmeromyia*, 205, 206, 212.

- luteolateralis*, *Banksinella*, 245, 317, 337.
Lutzia, included in *Culex*, 256.
Lyperosia minuta, in Gambia, 197, 207.
- MacCarthy Id. (Gambia), 189, 204, 205 ;
 not suitable as a site for Sleeping Sick-
 ness Camp, 224 ; clearing to be done
 at, 225.
- mactans*, *Haematopota*, habitat and method
 of biting, 111.
- maculata*, *Glossina* (see under *Glossina*).
 " *Hippobosca* (see under *Hippo-*
bosca).
- maculatissimus*, *Tabanus*, 286.
maculicrura, *Culex*, 261.
maculipalpis, *Anopheles*, 143.
maculipes, *Taeniorhynchus tenax*, var., 258.
maculiventris, *Culex*, 265.
maculosa, *Grabhamia*, 248.
 Madagascar, *Culex* in, 259, 262.
madagascariensis, *Pseudoheptophlebomyia*,
 258.
 Maganja da Costa (Mozambique), *Glossina*
 spp. in district of, 41.
magnifica, *Chrysops*, **161-163**.
 " var. *inornata*, *Chrysops*, 163.
magretti, *Pangonia*, 176.
Maillotia pilifera, 265.
major, *Goniocotes*, 148.
 " *Mansonia*, 254.
- Malaria, common in Gambia, 188 ; com-
 mon but diminishing in N. Nigeria, 351.
- Mallophaga, from African hosts, 145-147 ;
 instructions for collecting, 239.
- Malta, *Phlebotomus perniciosus* almost as
 abundant as *P. papatasi* in, 47 ; larvae
 in catacombs near centre of, 48 ; other
 investigators in, 49 ; means for de-
 struction of *Phlebotomus* in, 53-56 ; *P.*
perniciosus, widely distributed in, 71.
- mangiferae*, *Diaspis cinnamomi*, var., **86**.
- Mansonia*, synonym of *Taeniorhynchus*,
 252.
 " *africana*, a var. of *Mansonioides*
uniformis, 254.
 " *arabica*, 249.
 " *annulifera*, 253.
 " *annulipes*, 253.
 " *major*, a var. of *Mansonioides*
uniformis, 254.
- Mansonia nigerrima*, a var. of *Man-*
sonioides uniformis, 254
 " *nigra*, 254.
 " *uniformis* (see *Mansonioides*).
- Mansonioides*, characters of genus and
 table of spp., 253.
 " *mediolineata*, 254.
 " *nigra*, 254.
 " *uniformis*, varieties and
 synonyms of, 254 ; struc-
 ture of palpi, **254** ;
 distribution, 254, 275 ;
 affected with immature
 parasitic mite, 275 ; in
 N. Nigeria, 319, 321,
 324, 328, 330, 336, 337,
 345.
- marlatti*, *Phoenicoccus*, 104.
 " *Sphaerococcus*, 104.
- masculus*, *Culex*, 263.
- Mashonaland, Culicidae in, 245, 256.
- maximus*, *Akidoproctus*, 148.
 " *Aspidoproctus*, 100.
- medicorum*, *Glossina*, 107, **109**.
- mediolineata*, *Etorleptomyia*, 253, 254.
 " *Mansonioides*, 254.
- mediopunctata*, *Culicada*, 248.
- Megaculex*, subgenus of *Ludlowia*, 244,
 note.
 " *albitarsis*, synonym of *Lud-*
lowia plumosa, 245.
- Megalorhini, 241.
- Megalorhinina, 241, note.
- Megarhininae, 241.
- Melanoconion*, included in *Culex*, 256.
 " *rimus*, 265.
- melanorhinus*, *Culex*, 265.
- Menopon africanum*, sp. n., from spurwing
 goose, Sudan, **149**.
 " *antennatum*, sp. n., from guinea
 fowl, Sudan, **150**.
 " *mesoleucum*, from owl and star-
 ling, Sudan, 149, 151.
 " *nigrum*, sp. n., from *Corvultur*
albicollis, **151**.
 " *pallidum*, from chicken, S.
 Nigeria, 149.
 " *picae*, 151.
 " *spinotum*, from starling, Sudan,
 149.
- mesoleucum*, *Menopon*, 149, 151.

- metallicus*, *Banksinella*, 252.
 „ *Taeniorhynchus*, 252.
Metanototricha, 241; contiguous eyes in most of the, 256.
Metanototrichina, 241, *note*.
microannulatus, *Culex*, 261.
Micro-organisms found in Anophelines, 6.
Mimeteculex, synonym of *Ochlerotatus*, 247, 248, 250.
 „ *kingii*, synonym of *Ochlerotatus ochraceus*, 250.
Mimomyia (?) *chamberlani*, *Ludlowia*, founded on, 244.
mimomyiaformis, *Boyca*, 244, *note*, 266.
 „ *Ludlowia*, 337.
mimosae, *Lecaniodiaspis*, 102.
Minna (N. Nigeria), blood-sucking flies and ticks obtained at, 330.
minor, *Palaeopsylla*, 269.
minuta, *Lyperosia*, 197, 207.
minutus, *Culex*, 263.
 „ *Haemasson*, 73.
 „ *Phlebotomus* (see under *Phlebotomus*).
mirandus, *Loxaspis*, sp. n., 363, 364.
mitrata, *Numida*, 148, 150.
mobola, *Parinarium*, 358.
molestus, *Cyniphes*, 73.
mopani, *Copaiifera*, 358.
Mopea (Zambesi), cattle trypanosomiasis in 1908-9 at, 41.
mordax, *Tabanus*, sp. n., 175.
morsitans, *Culicada*, 251.
 „ *Glossina* (see under *Glossina*).
Mosquitos, infected with malaria, 1-8; where to find, 1; method of capture, 1; identification of species, 1, 2; dissection of, 2-5; gland examination, 3-4; staining of sporozoites, 4; dissection of mid-gut, 4-5; staining of zygotes, 5; endo- and ecto-parasites of, 6; problems concerning infection of, 7, 8; properties of the salivary secretion of, 8; suggested lines of research, 8; larvae in water-vessels at Lagos, 127-136; seasonal variation of larvae, 131; *Filaria* carried by, 133; larvae and pupae killed by raised temperature of water, 135.
Mosquito larvae, search in domestic water-vessels for, 127-136; seasonal variation in water-vessels of, 131; raised temperature of water kills, 135; fish and, 137-139; *Girardinus paeciloides* preys on, 182; green gas-producing plant that kills, 276.
Mosquito pupae, raised temperature of water kills, 135.
Mozambique, distribution of tsetse-flies in, 39-42.
Mucidus africanus, synonym of *M. mucidus*, 246.
 „ *alternans*, 246.
 „ *grahami*, var. of *M. mucidus*, 246.
 „ *mucidus*, 246, 319, 345.
 „ *scatophagoides*, 246.
 „ *sudanensis*, synonym of *M. scatophagoides*, 246.
mundulus, *Culex*, 266, 268.
Musca, 56.
Muscidae, blood-sucking in Gambia, 216, 217; in S. Nigeria, 274, 275; in N. Nigeria, 321, 328, 346, 347-351.
musculi, *Leptopsylla*, 269.
Mytilaspis beckii, 88.
 „ *citricola*, in Uganda, 88.
Myzomyia, synonym and spp. of, 141, 142; type sp. incorrectly cited, 141.
 „ *disignata*, probably synonym of *M. rhodesiensis*, 268.
 „ *costalis*, infected with sporozoites, 6; distribution in N. Nigeria, 319-321, 325, 328, 330.
 „ *culicifacies*, infected with sporozoites, 6.
 „ *funesta*, infected with sporozoites, 6; occurrences in N. Nigeria, 320, 321, 337.
 „ *nili*, same as *M. umbrosa*, 268.
 „ *rhodesiensis*, 268
 „ *rossi*, infected with sporozoites, 6.
 „ *umbrosa*, in N. Nigeria, 319, 321, 328, 345.
 „ *unicolor*, same as *M. umbrosa*.
Myzorhynchus, sporozoites found in, 7; type sp. incorrectly cited, 141.
 „ *barbirostris*, 268.

- Myzorhynchus obscurus*, may be a var. of *M. barbirostris*, 268.
 „ *strachani*, synonym of *M. obscurus*, 268.
- Natal, Culicinae in, 246, 250, 251, 259, 264.
- Native names, of blood-sucking insects and other Arthropods, 219, 223, 340; of insects and diseases carried by them, 347.
- Natives, instruction of, 226.
- naevei*, *Chrysops*, 163.
 „ *Culex*, 263.
 „ *Hippobosca hirsuta*, var., 171.
- nebulosa*, *Culicomyia*, 254–256.
nebulosus, *Trichorhynchus*, 320, 328.
necopinus, *Tabanus*, sp. n., 279, 339.
- Nematodes, found in Anophelines, 6.
- nemorosa*, *Culicada*, 248.
nemorosus, *Ochlerotatus*, 248.
Neocellia, included in *Nyssorhynchus*, 141.
Neomelanconion palpale, 226.
 „ *rima*, 254, 255, 256.
- Neostethopheles*, synonym of *Stethomyia*, 141.
- neotaeniorhynchus*, *Culicelsa*, 261.
- Nets for *Phlebotomus*, 55.
- Niger River, research along, 315–320.
- niger*, *Taeniorhynchus*, 247.
- Nigeria, N., haunts of *Glossina tachinoides* in, 119–126; new *Anopheles* from, 142, 143; variety of *Stegomyia fasciata* from, 179; *Cimex rotundatus* in, 180; *Culex* and allied genera in, 244, note, 246, 250, 253, 259, 262, 263, 264; research in, 301–356; geography of, 303–308; climate and rainfall, 308–314; population and tribes, 314–315; records of blood-sucking insects and ticks in, 34, 35, 345, 346; legislation in, 356; subjects requiring further investigation in, 356.
- Nigeria, S., examination of native water-receptacles in, 127–136; fish that prey on mosquito larvae in, 137–139; Anoplura from, 146; Mallophaga from, 147, 149, 151; *Cimex rotundatus* in, 180; *Culex* and allied genera in, 245, 246, 250, 252, 253, 259, 261, 263–265; blood-sucking flies in, 109, 273–276.
- nigeriensis*, *Culex*, 250.
 „ *Ochlerotatus*, 250.
- nigerrima*, *Mansonia*, 254.
nigerrimus, *Phlebotomus*, 47, 62, 68.
nigra, *Chrysoconops*, 252.
 „ *Mansonia*, 254.
 „ *Mansonioides*, 254.
 „ *Stomoxys* (see under *Stomoxys*).
- nigrochaetae*, *Culex*, 252, 254–256.
nigrocostalis, *Culex*, 129–131, 133–263.
nigrofusca, *Glossina* (see under *Glossina*).
nigrothorax, *Taeniorhynchus*, 252.
nigrum, *Lecanium*, 92, 100.
 „ *Menopon*, sp. n., 151.
- nili*, *Myzomyia*, 268.
- Nirmus fuscus*, from owl, Sudan, 147.
 „ *varius*, from raven, S. Nigeria, and starling, Sudan, 147.
 „ *vulgatus*, from owl and starling, Sudan, 148.
- Niumi Province (Gambia), in regard to tsetse, 211–14, 222; vegetation in, 190.
- nobilis*, *Holcoceria*, 182.
- nocturnus*, *Ochlerotatus*, 250.
- Nosema*?, found in Anophelines, 6.
- Notabile (Malta), *Phlebotomus* found in catacombs at, 48, 51.
- Numida mitrata*, Mallophaga parasitic on, 148, 150.
- nuttalli*, *Amblyomma*, 320.
- Nyasa, *Glossina* taken in, 19; reports of tsetse from, 39, 40; Anoplura from, 145–147; new Psyllidae from, 105–106; no official record of sleeping sickness in N., 181; cattle trypanosomiasis in, 181, and note; sleeping sickness in Dowa district, 112; habits of blood-sucking flies in, 111–112; new *Chrysops* from, 163; *Cimex rotundatus* in, 180; a Locustid injurious to man in, 180; *Culex* and allied genera in, 246, 248, 250, 259, 260, 264, 265; new flea from, 269–272.
- nyasae*, *Pseudoeropsylla*, sp. n., 105.
- Nycteribiidae, pseudo-joints found in some, 364, note.
- Nyssomyzomyia*, synonym of *Myzomyia*, 141.

- Nyssorhynchus*, genera included in, 141; type sp. incorrectly cited, 141.
- " *pharoensis*, in Gambia, 201; in N. Nigeria, 319, 328, 345.
- obscurus*, *Myzorhynchus*, 268.
- ocellata*, *Grabhamia*, 250.
- Ochlerotatus*, synonyms of, 246, 247; table of spp. of, 247.
- " *albifasciatus*, type of genus, 247.
- " *caliginosus*, 250; unguis of, 246.
- " *cumminsi*, synonyms of, 248.
- " *dentatus*, 248.
- " *dorsalis*, unguis formula and synonyms of, 248.
- " *durbanensis*, 250.
- " *hirsutus*, proboscis and palpi, 249.
- " *lateralis*, in Algeria, 250.
- " *leucarthrius*, additional characters, 251.
- " *longisquamosus*, 249.
- " *nemorosus*, varieties of, 248.
- " *nigeriensis*, 250.
- " *nocturnus*, 250.
- " *ochraceus*, 250; unguis of, 246.
- " *pulcritarsis*, unguis formula and synonyms, 249.
- " *quasiunivittatus*, 250.
- " *scapularis*, 247.
- " *vigilax*, 250.
- ochraceus*, *Ochlerotatus*, 246, 250.
- " *Taeniorhynchus*, 253.
- Oculeomyia*, included in *Culex*, 256.
- " *sarawaki*, a var. of *Culex* *ager*, 259.
- Olfersia acarta*, synonym of *O. intertropica*, 172, note.
- " *dukei*, sp. n., on a fish-eagle in Uganda, 171.
- " (*Ornithomyia*) *intertropica*, 172 and note.
- ornatothoracis*, *Culex*, 263, 264.
- Ornithodoros*, instructions for collecting, 232.
- Ornithomyia* (see *Olfersia*).
- Oshogbo district (S. Nigeria), blood-sucking flies of, 273-276.
- Palaeopsylla gracilis*, 259.
- " *minor*, 269.
- pallicera*, *Glossina* (see under *Glossina*).
- pallidipennis*, *Haematopota* 321, 330, 334.
- pallidipes*, *Glossina* (see under *Glossina*).
- pallidocephala*, *Culex*, 262.
- pallidostriatus*, *Culex*, 250.
- pallidothoracis*, *Culex*, 263.
- pallidum*, *Menopon*, 149.
- pallipes*, *Culex*, 265.
- palpale*, *Neomelanoconion*, 266.
- palpalis*, *Glossina* (see under *Glossina*).
- " var. *wellmani*, *Glossina*, 23.
- Pandanus*, supposed by natives to poison drinking water, 330.
- Pangonia magretti*, said to be fatal to camels, 176.
- Papataci-flies (see under *Phlebotomus*).
- papatasi*, *Phlebotomus* (see under *Phlebotomus*).
- Papio sphinx*, predilection of *Glossina morsitans* for, 206-207, 223; potential reservoir of trypanosomes, 223.
- par*, *Culex*, 266.
- " *Tubanus*, 207, 321, 324, 326, 346.
- parascelos*, *Culex*, 250.
- Parasites, control of pests by introduction of, 365-367.
- Parinarium mobola*, puparium of *Glossina morsitans* at base of trunk of, 358.
- Parlatoria blanchardi*, 88-89, 92.
- " (*Websteriella*) *zzyphi*, in Egypt, 91.
- paykulli*, *Plexippus*, 210.
- Pecomysia*, 250.
- Pectinopalpus fuscus*, larvae in native water-vessels at Lagos, 129, 130; seasonal variation of larvae of, 131; usually in foul water, 131; associated with *Stegomyia fasciata*, 132; synonym of *Culicomyia nebulosa*, 255.
- Pediculus capitis*, in Sudan, 145.
- penetrans*, *Dermatophilus*, 325.
- peristictus*, *Haematopinus*, sp. n., 145.

- pernicius*, *Phlebotomus* (see under (*Phlebotomus*)).
- persicae*, *Lecanium*, 93.
- persicus*, *Argas*, 232.
- pertinens*, *Haematopota*, 111.
- " *Tabanus*, sp. n., **286-290**, 321, 328, 330, 332-334, 346.
- petersi*, *Enyaliopsis*, 180.
- pettigrewi*, *Culex*, 250, 262.
- Phacochoerus aethiopicus*, Anoplura parasitic on, 145.
- pharoensis*, *Nyssorhynchus*, 201, 319, 328, 345.
- Phlebotominae, 57.
- Phlebotomus*, of Maltese Is., 47-78; search for breeding places, 48-50; habits and occurrence of adults of, 50-52; seasonal prevalence of, 52-53; prophylactic measures against, 53-56; morphology of **57-67**; oviposition in captivity, 67; synopsis of, 68; list of references to, 77-78; description of spp., **68-77**; adults found associated with rabbits, 49.
- " *minutus*, apparently rare in Malta, 47; described, **69-70**; much more active than other Maltese spp., 70; wing venation of, **62**.
- " *nigerrimus*, sp. n., **68**; wing venation, **62**.
- " *papatasi*, abundant in Maltese Is., 47; breeding places, 48, 49; head and appendages of, **57-60**; external genitalia, **61, 74**; wing venation, **62**; male and female described, **73, 74**; much more restless in captivity than *P. perniciosus*, 75; ovum of, **75**; larva of, **75, 76**; pupa of, **76, 77**; eggs only hatch in moist atmosphere, 75.
- " *pernicius*, sp. n., abundant in Malta, 47; possible carrier of Papataci fever, 47; breeding places, 49; wing venation, 62; male and female described, 70-72; pupa of, 72-73.
- Phoenicoccus*, synonym of *Sphaerococcus*, 104.
- Photography of Diptera, 153-160; apparatus for, 153, 154; preparation of insects for, 155, 156; illumination, 156, 157; magnification, 157, 158; exposure, 158, 158; development, 159; retouching, 159; printing, 160.
- picae*, *Menopon*, 151.
- pilifera*, *Maillotia*, 265.
- piliferus*, *Culex*, 265.
- pilosellus*, *Clinocoris*, 364, note.
- pincerna*, *Ludlowia*, 245, 266.
- pipistrelli*, *Clinocoris*, 364, note.
- pipiens*, *Culex*, 247, 248, 262, 263.
- Piroplasmiasis, in dogs in N. Nigeria, 323, 354.
- pitchfordi*, *Anopheles*, 143.
- Plectropterus gambensis*, Mallophaga on, 148, 149, 151.
- Plexippus paykulli*, preying on *Glossina palpalis*, 210.
- plumosa*, *Ludlowia*, 245.
- poeciloides*, *Girardinus*, 182.
- poecilipes*, *Lasioconops*, 258.
- pollinator*, *Stegomyia*, 268.
- Polyctenidae, 152; pseudo-joints found in, 364, note.
- Portuguese E. Africa, distribution of *Glossina* in, 39-42; relation between game and *Glossina* in, 116-118.
- Potamochoerus choeropotamus*, Anoplura on, 145.
- potis*, *Chimaeropsylla*, g. et sp. n., **269**.
- pretoriensis*, *Anopheles*, 143.
- procerus*, *Goniocotes*, 148.
- Prophylactic measures against *Phlebotomus*, 53-56; repellents, 53, 54; fumigation, 54; light, 54; artificial air movement, 54; traps, 54; nets, 55; destruction of breeding grounds, 55, 56
- prospodidis*, *Lecaniodiaspis*, 102.
- Proteosoma*, birds infected with, 7.
- Protomelanoconion fuscum*, 244.
- pruinus*, *Culex*, 263.
- pseudoannulioris*, *Culex*, 259.
- pseudocinerea*, *Culicomyia nebulosa* var., 255, 256.
- pseudocinereus*, *Culex*, 254.
- pseudoconopas*, *Chrysoconops*, 253.
- Pseudoeriopsylla*, g.n., 105.

Pseudoeriopsylla nyasae, sp.n., 105.
Pseudoheptophlebomyia, included in *Culex*, 256.
 " *madagascariensis*,
 synonym of *Culex quasigelidus*, 258.
psidii, *Pulvinaria*, 94.
psusemisi, *Tabanus*, 176, note.
Psychoda, larvae in foul water at Lagos, 129, 130, 134.
 Psychodidae, 57; in Gambia, 218.
 Psychodinae, 57.
 Psyllidae, n. g. of, 105.
pulcritarsis, *Culex*, 249.
 " *Grabhamia*, 249.
 " *Ochlerotatus*, 249.
pulcripalpis, *Grabhamia*, 248, 249.
 Pulicidae, in Gambia, 218; in N. Nigeria, 322, 346.
pullatus, *Culex*, 264.
pullulus, *Tabanus*, sp. n., 282.
Pulvinaria jacksoni, in Uganda and W. Africa, 93; food plants, 93.
 " *psidii*, injurious to coffee in Uganda, 94.
pumilus, *Haplochilus*, 182.
punctipes, *Aporoculex*, 258.
punctocostalis, *Banksinella*, 245.
puniens, *Haematopota*, 326.
pusillus, *Culex*, 265.
pygmaeus, *Culex*, 266.
Pyrethophorus, type sp. incorrectly cited, 141.
 " *costalis*, breeding in native domestic water-vessels at Lagos, 133; occurrences in water-vessels at Lagos, 130, 131; found at lowest elevation above sea level, 134; breeding in native yards, 136; occurrence in Gambia, 205.
quadrilineatus, *Ceroplastes*, 95-98.
quasigelida, *Leucomyia*, 258.
quasigelidus, *Culex*, 258-260, 264.
quasiguiarti, *Culex*, 264.
quasimodesta, *Culicada*, 256, 266.
quasinivittatus, *Ochlerotatus*, 250.
Radioculex clavipalpus, synonym of *Ludlowia chamberlaini*, 244, and note.

Railways, recommendations with regard to, 354.
 Rainfall, in Gambia, 191-193; in N. Nigeria, 308-314.
Reedomyia annulata, in N. Nigeria, 320.
 " *lowisii*, 250.
regularis, *Diaspis*, sp. n., 86, 87.
regulus, *Helea* (*Forcipomyia*), 57.
 Rest camps, in Gambia, 204, 205, 225; in N. Nigeria should be moved further from kurimis, 344, 345.
Rhipicephalus falcatus, in N. Nigeria, 330
 " *sanguineus*, on dogs in Gambia, 212; in N. Nigeria, 319, 322, 324; biting man, 326.
 " *simus*, abundant on dogs in N. Nigeria, 321, 322, 324; on horses in N. Nigeria, 325; biting man, 326; found at Baro, 328.
 Rhodesia, N.E., new var. of *Hippobosca* from. 171; *Cimex rotundatus* in, 180.
 " N.W., tsetse able to exist without game in, 40.
 " S., tsetse found in, 41, 357-361; Culicinae in, 249, 250, 259, 262, 264.
Rhus longifolia, 117.
Rhynchocyon cirnei, new flea occurring on, 272.
rima, *Culex*, 265.
 " *Neomelanoconion*, 254, 255, 265.
rimus, *Melanoconion*, 265.
 Binderpest and *Glossina*, 114, 115.
rossi, *Anopheles*, 141, 143.
rossii, *Myzomyia*, 6.
rotundatus, *Cimex*, 179, 180, 212.
 Rovuma river, fly-belts along, 40, and note; diary of H. T. Barrett's journey down the, 43-46; tsetse abundant along affluents of, 43, 44.
rubinotus, *Culex*, 265.
rufinus, *Culex*, 265.
rusci, *Ceroplastes*, 97
Saissetia (see *Lecanium*).
salicifolii, *Agromyza*, sp. n., 79.
salisburyensis, *Culex*, 264.
salsus, *Culex*, 261.
salus, *Culex*, 261.

- Sand-flies** (see *Phlebotomus*).
- sanguineus*, *Rhipicephalus*, 212, 319, 322, 324, 326.
- sarawaki*, *Oculeomyia*, 259.
- Sarcopsyllidae**, in N. Nigeria, 325, 346.
- Scale Insects**, African, observations on, 85-104.
- scapularis*, *Leucomyia*, 247, 256.
- " *Ochlerotatus*, 247.
- scataphagoides*, *Mucidus*, 246, 319, 345.
- semicinctum*, *Colpocephalum*, 149.
- secedens*, *Tabanus*, 182, 202, 320, 346.
- Sená** (Mozambique), cattle trypanosomiasis in vicinity of, 41.
- Senegal**, French, possibility of diseases being carried from or into, 195.
- senegalensis*, *Haplochilus*, 137, note.
- sergenti*, *Culex*, 265.
- Sierra Leone, outbreak of Yellow Fever in, 127; Culicinae in, 246, 259, 263, 265.
- silacea*, *Chrysops*, 166, 341-344
- simillimus*, *Cyclops*, 129, 130, 134.
- simpsoni*, *Culex*, 263.
- Simulium*, in N. Nigeria, 317, 319, 324, 326.
- simus*, *Rhipicephalus*, 321, 322, 324-326, 328.
- sinensis*, *Culex ager*, var., 259.
- " *Leucomyia*, 259.
- " " *gelida*, var., 259.
- Siphonaptera, new, from Nyasa, 269-272; in N. Nigeria, 322, 325, 346.
- sjoestedti*, *Colpocephalum*, 149.
- Sleeping Sickness**, occurrence in Dowa district, Nyasa, 112; in Gambia, 187, note, 188, 205, 210, 214, 223; camps for, 224, 353, 355; method of minimizing spread of, 225; not in Ilesha and Oshogbo districts, S. Nigeria, 276; colour convention for, 297, 298; occurrence in N. Nigeria, 331-333, 337-339, 344, 352, 353; native names for, 347; no case discovered in villages on Garara River, 352; sporadic in Baro, 352, 353; natives of W. Africa though immune may act as reservoirs of, 353.
- socialis*, *Tabanus*, 208, 209.
- somaliensis*, *Culex*, 261.
- Somaliland, British, occurrence of *Stegomyia fasciata* in, 179; new *Tabanus* in, 176; *Culex* in, 261.
- spathipalpis*, *Theobaldia*, 251.
- Sphaerococcus draperi*, synonym of *S. marlatti*, 104.
- " *marlatti*, common in Lower Egypt, 104; remedy for, 104.
- sphinx*, *Papio*, 206, 207, 223.
- Spilopsyllus cuniculi*, 269.
- spinosum*, *Menopon*, 149.
- spirocarpoides*, *Acacia*, 116.
- Sporozoa, found in Anophelines, 6.
- Sporozoites in mosquitos, methods for detecting, 1-8.
- Steamers and Launches, tsetse invading, 190, 198, 202, 204, 205, 207-210; effective mosquito proofing for, 226; mosquitos breed in "bilge water" on, 354.
- Stegomyia africana*, bites only at dusk, 275.
- " *fasciata*, identification of larva of, 127; distribution of larva at Lagos, 131, 135; number of vessels in which were found larvae of, 130; seasonal variation of larvae of, 131; occurs in Sudan, 179; may transmit virus of horse-sickness, 179; vector of dengue fever, 179; abundant in coast towns of British Somaliland, 179; variations of, 179; in Gambia, 188, 195; pale var. common in Algeria, 248; in N. Nigeria, 318, 319, 328, 351, 354, 355.
- " *longipalpis*, 268.
- " *pollinator*, synonym of *S. longipalpis*, 268.
- stenopygus*, *Akidoproctus*, 148.
- Stethomyia*, considered distinct, 141; *Neostethopheles*, synonym of, 141.
- Stomoxys*, cattle trypanosomiasis spread by, 41; as a protozoal carrier, 223; inoculation experiments should be carried out with, 323.
- " *calcitrans*, empty pupae of in caves of Notabile, Malta, 48; in Gambia, 197, 207, 208; in N. Nigeria, 319-321, 323, 324, 327, 333, 346.
- " *nigra*, in Gambia, 201, 202, 207, 208; common in Oshogbo, S. Nigeria, 275; in N. Nigeria, 317, 319-321, 323, 324, 327, 346.

- strachani*, *Myzorhynchus*, 268.
- subangustus*, *Tabanus*, 275, 320, 325-328, 346.
- submorsitans*, *Glossina* (see under *Glossina*).
- substriata*, *Chionaspis*, 90.
- subtilis*, *Grabhamia*, 248.
- Sudan, *Culicinae* in, 245, 246, 250, 252-254, 259, 262, 263, 265.
- sudanensis*, *Mucidus*, 246.
- suis*, *Haematopinus*, 145.
- sycophanta*, *Calosoma*, 369.
- Tabanidae, cattle trypanosomiasis spread by, 41; three new spp. of, 161-168; two new spp. from Sudan, 173-177; *Bembez* preying on, 182; list of species in Gambia, 215-216; in S. Nigeria, 274, 275; new African species, 279-290; abundant at Zungeru, N. Nigeria, 323; spp. occurring in N. Nigeria, 321, 328, 339, 346; none seen along River Garars, 337; native names of, 347.**
- tabaniformis*, *Glossina*, 9, and note, 108.
- Tabanus africanus*, habits and habitats, 112, 346.
- „ *albipalpus*, in Gambia, 215; in N. Nigeria, 325, 327, 346.
- „ *biguttatus*, habits, 112.
- „ var. *croceus*, in N. Nigeria, 321, 324, 325, 328, 330, 346.
- „ *camelarius*, sp. n., 173; attacking camels, 175.
- „ *claritibialis*, 282.
- „ *copemani*, sp. n., 281.
- „ *crocodilinus*, sp. n., 284.
- „ *ditaeniatus*, habitats of, 111; in Gambia, 200, 207; in N. Nigeria, 321, 324, 346.
- „ *fasciatus*, in Gambia, 202; in N. Nigeria, 320, 346.
- „ *gratus*, in N. Nigeria, 321, 330, 334, 346.
- „ *irroratus*, 286.
- „ *latipes*, in N. Nigeria, 318, 321, 328, 346.
- „ *laverani*, in Gambia, 202; in N. Nigeria, 346.
- „ *leucostomus*, 176, note, 177.
- „ *maculatissimus*, 286.
- Tabanus mordax*, sp. n., 175; said to be fatal to camels, 176.
- „ *necopinus*, sp. n., 279, 339.
- „ *par*, in Gambia, 207; in Nigeria, 321, 324, 326, 346.
- „ *pertinens*, sp. n., 286; distribution, 289-290; troublesome to human beings, 290; in N. Nigeria, 321, 328, 330, 332-334, 346.
- „ *psusemnia*, synonym of *leucostomus*, 176, note.
- „ *pullulus*, sp. n., 282; distribution, 283-284.
- „ *secedens*, preyed upon by *Bembez*, 182; possible vector of *Trypanosoma pecorum*, 182; in Gambia, 202; in N. Nigeria, 320, 346.
- „ *socialis*, in Gambia, 208, 209.
- „ *subangustus*, habits in S. Nigeria, 275; in N. Nigeria, 320, 325-328, 346.
- „ *sufis*, 290.
- „ *taeniola*, in N. Nigeria, 318, 320, 321, 324, 327, 328, 330, 339, 346.
- „ *taeniola*, var. *variatus*, habits and habitats, 111, 112; in Gambia, 200-202, 204-207, 216.
- „ *thoracinus*, in S. Nigeria, 275; in N. Nigeria, 321, 346.
- „ *unitaeniatus*, habits, 111; males common, 112.
- Tachardia actinella*, in Natal, 100.
- „ *albida*, in Natal, 100.
- „ *decorella*, new to Africa, 85, 100; on *Anona muricata*, 100.
- „ *longisetosa*, sp. n., 102; infested with Chalcidids, 103; on *Ficus*, 103.
- Tachinidae, parasites on larvae of Gipsy Moth, 369, 370.
- tachinoides*, *Glossina* (see under *Glossina*).
- taeniarostris*, *Banksinella*, 245.
- taeniarostris*, *Grabhamia*, 259.
- taeniola*, *Tabanus* (see under *Tabanus*).
- taeniopygus*, *Fundulus*, 182.
- taeniorhynchoides*, *Culex*, 258.
- Taeniorhynchus*, characters of, 251-252; tables of spp. of, 252; *Mansonia* a synonym of, 252.

- Taeniorhynchus africanus*, not a true *Taeniorhynchus*, 266.
- " *ager*, included in *Culex*, 256; not African, 259.
- " *annettii*, 253.
- " *aurites*, 253; wrongly recorded from Malay States, 253.
- " *cristatus*, 253.
- " *fasciolatus*, 251.
- " *fuscopennatus*, 253; *Culex drymoecius* a var. of, 253.
- " *lineatopennis*, synonym of *Banksinella luteolateralis*, 245.
- " *metallicus*, 252.
- " *niger*, included in *Ochlerotatus*, 247.
- " *nigrithorax*, 252.
- " *ochraceus*, 253.
- " *taeniorhynchus*, 251.
- " *tenax*, included in *Culex*, 256.
- " *tenax*, var. *maculipes*, synonym of *Culex quasigelidus*, 258.
- " *titillans*, type of genus, 251, 252.
- " *violaceus*, synonym of *T. metallicus*, 252.
- taeniorhynchus*, *Chrysoconops*, 252.
- " *Culicelsa*, 247, 249.
- " *Taeniorhynchus*, 251.
- Taphozous hildegardeae*, 364.
- tenax*, *Taeniorhynchus*, 256, 259.
- Tendaba (Gambia), blood-sucking flies abundant at, 207-208; clearing needed at, 225.
- tenuicrus*, *Haematopota*, 321.
- tenuivalvatum*, *Lecanium*, sp. n., 92-93.
- Tete district (Mozambique), fly areas in, 41.
- thalassius*, *Culex*, 260, 261.
- theileri*, *Culex*, 260, 262.
- Theobaldia annulata*, 251.
- " *spathipalpis*, 251.
- thoracimus*, *Tabanus*, 275, 321, 346.
- Thysanoptera, 238.
- tibialis*, *Linognathus*, 147.
- igripes*, *Culex*, 251, 261, 321.
- igripes*, var. *consimilis*, *Culex*, 259.
- " var. *fusca*, *Culex* (see under *Culex*).
- titillans*, *Culex*, 252
- " *Taeniorhynchus*, 251, 252.
- Transvaal, relation between game and tsetse in, 114-116; *Culex* and allied genera in, 248, 249, 251, 254, 259-264.
- Traps for *Phlebotomus*, 54.
- tremae*, *Lecanium*, 93.
- Trematodes, found in Anophelines, 6.
- Trichochetes climax*, from goats, S. Nigeria, 147.
- Trichoprosoponinae, 241.
- Trichorhynchus (Culicomyia) nebulosus*, in N. Nigeria, 320, 328.
- trilobitiformis*, *Aspidiotus*, 85.
- Trinoton luridum*, from spurwing goose, Sudan, 151.
- Trypanosoma brucei*, found in blood of horses in N. Nigeria, 353; more virulent than *T. vivax*, 353.
- " *gambiense*, natives of Gambia infected with, 188; 5.4 per cent of population of Essau infected with, 213.
- " *vivax*, found in blood of horses in N. Nigeria, 353.
- Trypanosomes, investigation of, in wild animals, 170; in bulls, 181; blood of baboons potential reservoir of, 223; not certain that horses and cattle are infected with the same, 213; found in horses, 353.
- Trypanosomiasis of Cattle, spread by *Stomoxys* or Tabanidae, 41; *Glossina morsitans* in relation to, 181, note; *G. brevipalpis* in relation to 181, 182; cases in district between Karonga and Songwe rivers, Nyasa, 181, note; in Gambia, 188, 200, 212, 214, 223, 226; investigation required into forms of, 356; prevalent in N. Nigeria, 353.
- Trypanosomiasis of Horses, in N. Nigeria, 319, 320, 323, 330, 332, 334, 336, 338-342, 344; not always carried from horse to horse, 345, 353; and of dogs, 344; camps for, 226, 320, 323, 332.
- Tsetse-flies (see *Glossina*).
- Uganda, *Glossina* taken in, 26, 27; new mosquito from Entebbe, 38; Coccidae from, 85, 86, 87, 88, 90, 92-100, 103

- Culex* and allied genera in, 245, 248, 250, 252, 253, 256, 259, 263-265; *Chrysops centurionis*, n. sp. from, 164; *Hippobosca hirsuta*, n. sp. from, 169; *Olfersia dukei*, n. sp. from, 171; fish preying on mosquito larvae in, 182; *Bembex* preying on Tabanidae in, 182.
- ugandae*, *Ceroplastes*, 94, 95.
- umbrosa*, *Anopheles*, 142.
- " *Myzomyia*, 268, 319, 321, 328, 345.
- unicolor*, *Myzomyia*, 268.
- uniformis*, *Culicomyia*, 255.
- " *Mansonia*, 253.
- " *Mansonioides*, 253.
- unitaeniatus*, *Tabanus*, 111, 112.
- univittatus*, *Culex*, 250, 260, 262, 345.
- Uranotaeniinae*, 218, 241.
- Valletta, searched for breeding places of *Phlebotomus*, 48, 49; *Phlebotomus*, in cellars and prison cells of, 51.
- variegatum*, *Amblyomma*, 197, 319, 325, 330.
- varius*, *Nirmus*, 147.
- versicolor*, *Hippocentrum*, 275, 319, 320, 324, 325, 327, 330, 346.
- vigilax*, *Culex*, 263.
- " *Ochlerotatus*, 250.
- Vinsonia*, 97.
- visonioides*, *Ceroplastes*, sp. n., 96.
- violaceus*, *Taeniorhynchus*, 252.
- viridis*, *Culex*, 263.
- vittata*, *Haematopota*, 111.
- vituli*, *Linognathus*, 146.
- vivax*, *Trypanosoma*, 353.
- vocifer*, *Haliaëtus*, 172.
- vulgatus*, *Nirmus*, 148.
- Waterbuck, in relation to *Glossina morsitans*, 43-46; *Hippobosca* infesting, 169-172.
- Water supply, danger of wells for, 197, 205, 206, 331, 333; *Glossina palpalis* swarming in N. Nigeria near, 343; clearing to be done round wells and springs, 344.
- watsoni*, *Anopheles*, 143, 330.
- Websteriella* (see *Parlatoria*).
- wellmani*, *Glossina palpalis*, var., 23.
- wellmanni*, *Chrysops*, 167, 168.
- Wells, danger of, 197, 205, 206, 331, 333.
- wilcocksii*, *Grabhamia*, 248.
- Yellow fever, outbreak in Sierra Leone and Gold Coast, 127; in Bathurst, Gambia, 188, 195, 224; not reported from N. Nigeria, 351.
- Zambesi river, fly areas of, 41.
- Zambesia district, considerable fly belts in, 40.
- zeltneri*, *Culex*, 266.
- zizyphi*, *Parlatoria*, 91.
- zombaensis*, *Culex*, 264.
- Zungeru (N. Nigeria), importance of, 304; biting insects and ticks in, 321, 322; tsetse nearly always present in, 322; clearing required near, 322; trypanosomiasis common amongst horses in, 323; piroplasmiasis in dogs in, 323, 354; game near, 323; trypanosomes found in pony at, 324; recommendations with regard to, 355.
- Zygotes in mosquitos, methods for detecting, 1-8.

INDEX TO NAMES OF PERSONS.

- Aders, Dr. W. M., 373.
 Alcock, Lieut.-Col. A., 141, 241.
 Anderson, T. J., 373.
 Annandale, N., 59, 61.
 Anson, Dr. W. G., 255.
 Archibald, Capt., 179.
 Archibald, Dr. J. W., 184, 321.
 Arribalzaga, 252.
 Ashworth, Dr. J. H., 289.
 Austen, E. E., 9, *note*, 10, *note*, 14, *note*,
 31, 40, 41, 107, 122, 161-177, 181, 249,
 279-290, 298, 339, 357.
- Babbington, Capt., 77.
 Bagshawe, Dr. A. G., 298.
 Balfour, Dr. Andrew, 179.
 Barclay, Dr. A. H., 277, 283.
 Barnshaw, A. L., 183.
 Barrett, H. T., 40, *note*, 43-46, 184.
 Beaman, Capt., 77.
 Bell, Sir Hesketh, 356.
 Beresford, M. H. de la P., 356.
 Best, Dr., 215.
 Bevan, L. E. W., 41, *note*.
 Birt, Lieut.-Col. C., 49.
 Bissell, Capt. F. E., 350, 353.
 Blair, Capt. A. H., 273, 274, 277.
 Blair, Dr. M. Cameron, 341, 356.
 Blakeney, Major, 338.
 Böderer, Dr. H. A., 277, 373.
 Boulenger, G. A., 137, *note*, 139.
 Boyce, Sir Rubert, 351.
 Brown, Henry, 81.
 Bruce, Col. Sir David, 27, 31, *note*, 170.
 Budgen, T. A., 81.
 Burdett, Dr., 217.
- Campbell-Irons, A., 81.
 Carand, C. A., 81.
 Carlyle, T. F., 81.
 Carpenter, Dr. G. D. H., 278.
- Carter, H. F., 2, *note*, 37-38.
 Champion, A. M., 373.
 Chartres, Dr. E. A., 319, 354, 373.
 Cherrett, Dr. B. W., 373.
 Chisholm, James A., 183.
 Christophers, Dr., 5, 6, 7.
 Clark, Dr. W. S., 82.
 Cockerell, T. D. A., 100, 102, 104.
 Cody, W. S., 277.
 Collett, Dr., 184.
 Collin, James E., 79, 80.
 Connal, Dr. A., 184, 277.
 Copeman, E. A., 282, 284.
 Copland, Dr. J. G., 184.
 Corsellis, M. H., 81.
 Cox, Rev. H. A. M., 81.
 Crawford, Major G. S., 53, 77.
 Currie, Dr. J., 373.
- Dalziel, Dr. J. M., 262, 289, 347, 350.
 Davey, Dr. J. B., 19, 180, 181, 145, 146,
 283, 285.
 David, R., 82.
 Dayrell, E., 184.
 Denton, Sir George, 209, 226.
 Dibblee, Major F. L., 70, 77.
 Dibblee, Mrs., 77.
 Dickie, James, 81.
 Drake-Brockman, Dr. R. E., 82, 176, 179,
 277.
 Dudgeon, G. C., 217.
 Duke, Dr. H. L., 172, 373.
 Duncan, N. C., 81.
 Dutton, Dr. J. E., 18, 187, 188, 195, 215,
 216, 217, 218.
 Dyar, H. G., 141, 246, 247, 251, 252, 253.
- Eaton, Rev. A. E., 248.
 Edwards, F. W., 141-143, 215, 241-268.
 Edwards, William, 81.

- Falk, G., 81.
 Farquhar, J. H. J., 82, 109, 277.
 Fattah ud Din, Dr., 184.
 Fell, T. E., 373.
 Fisher, Dr. Walter, 81.
 Fiske, W. F., 365.
 Forsyth, H., 183.
 Foster, A. N., 215-217, 277.
 Foulkes, Capt. H. D., 81
 Foy, Dr. H. A., 280, 339.
 Franklin, Dr. J. C., 217.
 Fraser, Capt. A. D., 161, 165, 184, 245.
 Fraser, Donald, 81.
 Fülleborn, Dr., 100.

 Gallagher, Dr. G. H., 184.
 Gaye, J. A. de, 82, 277.
 Gouldsberg, Henry C., 183.
 Gowdey, C. C., 82, 85-104, 182, 184, 277,
 373.
 Graham, Dr. W. M., 82, 127-136, 137-
 139, 153-160, 255, 273.
 Grassi, Prof. B., 49, 56-61, 66.
 Gray, Dr. G. M., 82.
 Gray, Dr. B. W., 180, 184.
 Greaves, J. A. Ley, 82, 184, 373.
 Green, E. E., 85, 93, 97.
 Grünberg, Dr. K., 268.
 Gush, Dr. Howard W., 81.

 Haddon, E. B., 183.
 Hailstone, Dr., 184.
 Hamilton, Major J. S., 39, 40, 113-118.
 Hannington, Dr., 109.
 Harrington, Hubert T., 183.
 Hearsey, Dr. H., 180.
 Henderson, Dr. F. L., 373.
 Hiscock, Dr. R. C., 184, 320.
 Hood, Dr. T., 197, 215-217, 224, 226.
 Hopkinson, Dr. E., 215-217.
 Howard, C. W., 39-42, 117.
 Howard, Dr. L. O., 365.
 Hughes, F., 102, 104.
 Humfrey, Capt. L. E. H., 82, 273, 274,
 276.
 Hutchins, E., 184, 373.
 Hynde, B. A., 81.

 Ingram, Dr. A., 35, 82, 328, 352.
 Inness, Dr. W. D., 326, 373.

 Jack, Rupert W., 357-361.
 Jackson, F. J., 277, 284, 364.
 James, Capt., 141.
 Jemmett, C. W., 184.

 Kellogg, Prof. V. L., 145-152.
 Kilroy, Surg. Lancelot, 77.
 King, Harold H., 82, 97, 145, 147-152,
 173, 175, 176, & note, 179, 285, 289.
 Kinghorn, Dr. Allan, 18, 34, 107, 109,
 291-295.
 Kleine, Dr. 292.
 Knab, Dr. F., 141, 251-253.

 Lamborn, W. A., 184, 277.
 Lane, Capt G., 183.
 Leal, Dr. Amaral, 40.
 Lefroy, H. Maxwell, 371.
 Lloyd-Jones, Capt., 77.
 Lloyd, Ll., 295.
 Lounsbury, C. P., 371.
 Luhn, Dr. O. S. F., 185.

 McCloy, Surg. A., 110.
 McConnell, Dr. Robt. E., 373.
 Macdonald, Dr. P. H., 277, 373.
 Macfarlane, Dr. A. M., 77.
 Macfie, Dr. J. W. S., 35, 82, 277, 289,
 328, 335, 352, 356, 374.
 McKay, Dr. F. W., 34, 185.
 Mackay, Rev. J., 276.
 Mackie, Capt. F. P., 27, 170.
 McKinney, Dr. 325.
 McLeay, Dr. C. W., 350.
 Manning, G. F., 81.
 Manuk, Capt. M. W., 319.
 Marett, Capt. P. J., 47-49, 51, 55, 68, 77.
 Marshall, Guy A. K., 107, 142, 145, 241,
 248, 298.
 Marshall, Dr. C. H., 185.
 Marshall, Hugh C., 183.
 Maskell, W. M., 102, 104.
 Maugham, R. C. F., 117
 Mayer, Dr. T. F. G., 82, 142, 185, 255,
 273-276, 277.

- Moffat, Rev. M., 183.
 Moiser, Dr. B., 83, 119-126.
 Morrison, Dr. W., 35, 36, 83, 323, 324,
 328, 329, 330, 352.
 Mouat, Dr. A., 374.
 Murphy, Dr. J. C., 280, 374.
 Murray, L., 81.
 Murray, Dr. Wm. Ansley, 81.
- Nair, T. D., 374.
 Neave, S. A., 83, 85-104, 106, 162, 166,
note, 180-182, 185, 284, 289, 374.
 Newstead, Prof. R., 9-36, 47-78, 85-104,
 105-110, 203, 259, 348.
- O'Brien, Col. C. W., 226.
 Old, Dr. J. E. S., 81, 83, 185, 278, 282,
 374.
 Owen, Dr. H. B., 180, 185.
- Paine, J. H., 145-152.
 Pask, Dr. E. H. Allon, 81, 83, 147.
 Patterson, W. H., 278, & *note*.
 Percival Bey, El-Kaimakam, 179.
 Percival, J., 81.
 Perkins, Dr. R. C. L., 370.
 Piaget, A., 148.
 Pirie, Dr. G. J., 185, 344.
 Powell, W. A., 183.
 Prentice, George, 81.
 Prichard, Dr. W. Owen, 374.
 Pugh, Dr. J., 374.
 Putron, P. de, 317, 326.
- Bendle, Dr. A. C., 185, 374.
 Boss, H. Malcolm, 374.
 Rothschild, Hon. N. Charles, 269-272,
 363-364.
 Roubaud, E., 292, *note*.
 Roupell, Hon. E. C., 72, 77.
 Bundle, Gen. Sir Leslie, 77.
- Sanderson, Dr. Meredith, 83, 111, 112,
 181, 278.
 Sant' Anna, Dr., 40, 41.
 Schalkwijk, Rev. J. van, 183.
 Schaudinn, Dr., 8.
 Scholefield, S. W. J., 374.
- Scicluna, Hon. C. Caruana, 77.
 Sergeant, E., 7.
 Shircore, Dr. J. O., 26, 27, & *note*.
 Silberrad, H., 374.
 Simpson, C. B., 278.
 Simpson, Jas. J., 83, 122, 146, 147, 149,
 151, 179, 180, 185, 187-239, 278, 289,
 290, 297-299, 301-356, 374.
 Sjöstedt, Prof. Y., 251, 253.
 Smythe, Dr. A. W., 185.
 Someren, Dr. R. A. L. van, 149, 170, 171,
 186, 278, 374.
 Speiser, Dr. P., 57, 251, 253.
 Stannus, Dr. Hugh S., 81, 163, 180, 181,
note, 185, 272, 278.
 Stephens, Dr. J. W. W., 1-8, 356.
 Steward, Capt., 77.
 Strathairn, Dr. G. C., 278.
 Stuhlmann, Dr. F., 181.
 Surcouf, B., 286.
 Swann, Dr. A. J. T., 81.
 Swynnerton, C. F. M., 374.
- Taschenberg, 148.
 Theobald, F. V., 141, 143, 185, 246-251,
 253, 255, 259, 260-263, 268.
 Thompstone, Dr. S. W., 302, 330, 356.
 Thornicroft, H., 183.
 Todd, Dr. J. L., 18, 187, 188, 213, 224.
 Traill, Capt. H. L. Norton, 81.
 Turnbull, A. M. D., 180, 278.
- Ventrillon, Dr., 258.
 Vosseler, Prof., 100.
- Watson, Dr. C. E. S., 83, 186, 289.
 Wesché, W., 10, *note*.
 Wheelwright, 276.
 Wiggins, Dr. C. A., 27, 38, 180, 186.
 Willcocks, F. C., 80, 83, 85-104.
 Wolbach, Dr., 188, 213.
 Woosnam, R. P., 374.
- Young, R. A., 183.
- Zammitt, Prof. T., 77.
 Zimmermann, Prof. A., 93.

INDEX TO PLANTS ATTACKED BY INSECTS.

- | | |
|---|--|
| <p><i>Acacia</i>, 97, 100.
 <i>Acacia arabica</i>, 102, 104.
 <i>Albizzia</i>, 92.
 <i>Alternanthera versicolor</i>, 94.
 Amakebe, 94.
 <i>Anona muricata</i>, 97, 98, 100.
 <i>Antigonon octopus</i>, 86.
 Apricot, 92.</p> <p><i>Baikea eminii</i>, 97.
 Bark-cloth tree, 98, 100, 103.</p> <p>Cacao, 93.
 <i>Cassia floribunda</i>, 90.
 Citronella grass, 93.
 <i>Citrus</i>, 88, 91.
 Coffee, 94, 97.
 Cotton, 93.
 <i>Crataegus</i>, 97.</p> <p>Date palm, 104.
 <i>Datura alba</i>, 86.</p> | <p><i>Ficus</i>, 92, 93, 97, 100, 102, 103.</p> <p><i>Justicia alba</i>, 88.</p> <p><i>Magnolia</i>, 86.
 Mango trees, 86, 100.
 <i>Markhamia platycalyx</i>, 97.
 Mish-Mish, 92.</p> <p>Nsambyia, 93.</p> <p>Oleander, 85.
 <i>Opuntia</i>, 87.
 Orange, 91.</p> <p>Palm, 90.
 <i>Phoenix</i>, 104.
 Poplar, 91.
 Prickly Pear, 87.
 <i>Psidium guyava</i>, 98</p> <p>Sunt, 104.
 Vine, 100</p> |
|---|--|

ERRATA.

- P. 68, lines 9 and 12, transpose the characters of the claspers of the male given for *Phlebotomus perniciosus* and *P. papatasi*.
- P. 73 (in synonymy of *Phlebotomus papatasi*), for "*Hermasson minutus*" read "*Haemasson minutus*."
- P. 145, for "*Pediculis capitis*" read "*Pediculus capitis*."
- P. 148, for "*Coniocotes gigas*" read "*Goniocotes gigas*."
- P. 149, for "*Liperus baculus*" and "*Liperus gambensis*" read "*Lipeurus baculus* and "*Lipeurus gambensis*" respectively.
- P. 214, line 25, for "protozoal" read "protozoal."
- P. 215, table, for "*Chrysops longicornis*, F.," read "*Chrysops longicornis*, Macq."
- P. 248, under *O. nemorosus*, for "Rev. E. A. Eaton" read "Rev. A. E. Eaton."
- P. 345, table of Culicina, for "*Mucidus mucidus*, Theo.," "*Stegomyia fasciata*, Theo.," and "*Stegomyia sugens*, Theo.," read "*Mucidus mucidus*, Karsch," "*Stegomyia fasciata*, F.," and "*Stegomyia sugens*, Wied.," respectively.
- P. 346, table of Tabanidae, for "*Haematopota cordigera*, Aust." and "*Tabanus pluto*, Wied.," read "*Haematopota cordigera*, Big." and "*Tabanus pluto*, Walk.," respectively.
- P. 346, table of Siphonaptera, for "*Ctenocephalus canis*, Dugès," read "*Ctenocephalus canis*, Curtis."
- P. 363, title, for "Uganda" read "British East Africa."

LONDON :

PRINTED FOR HIS MAJESTY'S STATIONERY OFFICE,
By DARLING & SON, LTD., 34-40, BACON STREET, E.

1912.

2