

From the Editor's desk

By Kamaldeep Bhui

Brexit, social division and discrimination: impacts on mortality and mental illness?

The British public's vote to leave the European Union has sent ripples and shock waves of worry among all people fearing social division, financial insecurity, and uncertainty about residential rights of nationals of non-UK countries. Scientific, security and social policy cooperation had fostered stability and innovation to improve the lives of the disempowered, including better collective interventions to tackle health inequalities, and there has been a period of careful scrutiny of the care systems and processes, and progress towards protecting the human rights of people living with mental illness. Greater trust and cooperation were at the heart of the European project, alongside greater trade and political connection.

Preceding the referendum, a culture of extremism and intolerance became visible in political debate including discussions about the future of the NHS and publicly funded services, specifically on the subject of recruiting professionals from around the world. Following the referendum, stigma, prejudice and discrimination seem to be on the rise, with some egregious examples of racism and assault against migrants and religious and ethnic minorities, as well as British citizens of non-UK heritage. These overt expressions of hostility appear to thrive off latent ideologies of national identity, superiority, destiny, self-protection and survival. A higher mortality in minorities is associated with experiences of discrimination in older adults¹ and area levels of racism,² although the same has not been shown in a younger cohort of Black women.3 Low levels of segregation in the USA correlate with lower Black/White differences in mortality of infants and young adults.4 Mortality, suicide, homicide/violence, and cardiovascular diseases are substantially elevated among sexual minorities in high-prejudice communities.⁵ In the UK, discrimination is associated with common mental disorders.⁶ Employment, social action and anti-stigma interventions including social contact rather than segregation help improve outcomes.⁷⁻⁹ Better knowledge is needed on how social adversity (prejudice, discrimination, stigma, low income, socioeconomic position) influences cumulative risks of poor mental health 10-13 and biological mechanisms of disease, for example, by accelerated ageing, 14 epigenetics, 15,16 and explanations for higher rates of heart disease, 17 respiratory disease, 18 poorer outcomes of HIV treatment, 19 and experiences of pain.²⁰ Young people and children are also witnessing more violence and open prejudice, as well as being victims of it, and such experiences can become biologically embedded.²¹ Encouragingly, early life experiences of adversity might be buffered by learning and education, reflecting neuronal plasticity.¹⁶

This month's *BJPsych* contributes several new studies of mortality and potential mechanisms by which adversity influences biology. As further evidence of biological embedding of adversity, Demakakos *et al* (pp. 135–141) found that poor parenting style was associated with a higher mortality at follow-up, and with cancer but not heart disease. Bipolar disorder is associated with a higher risk of dementia and increased mortality (Almeida *et al*, pp. 121–126), and increased risk of death related to suicide, accidents, pneumonia, influenza, and liver and digestive system disease. Anhedonia is associated with higher mortality among patients with type 2 diabetes (Nefs *et al*, pp. 142–149), and relevant to the notion of social connectedness being critical in

all societies, loneliness is associated with higher mortality in older adults, the effect being mediated by physical activity (Holwerda *et al*, pp. 127–134).

The biosocial fabric in which mental illness emerges is further illustrated by several studies; for example, on the impact of childhood trauma on later mental illness (Petros et al, pp. 169-170). The mental health of Syrian refugees in Jordan is influenced by environmental (financial, housing, employment) and psychosocial outcomes such as loss of role and social support, and inactivity (Wells et al, pp. 99-106). Area deprivation may also explain higher rates of compulsory treatment in mental health services in urban areas (Keown et al, pp. 157-161, and see relevant editorials by Burns & Rugkåsa, pp. 97-98, and Lepping et al, pp. 95-96). Other groundbreaking studies improve our knowledge about classification and diagnosis (Böhnke & Croudace, pp. 162-168, and Reininghaus et al, pp. 107-113), and reflect on emerging new genetic markers of mental illness (Tesli et al, pp. 114-120, and Curtis, pp. 93-94). Despite more people receiving psychological and pharmacological treatments, levels of common mental disorders remain constant, and a substantial proportion of people are still not receiving treatments that we know reduce premature mortality in those with mental illness.²²

Social and political divisions are powerful sources of discrimination, as are aggressive biological drives and instincts and motivational mechanisms related to limbic system and amygdala. But the reflective human mind can moderate the forces of instinct and perceptions of difference, reconciling apparent inconsistencies in our socially connected world, and potentially altering the biological imprint of social adversity. Much more research is needed on the way social adversity can be encoded in biology, and the role of social, psychological, psychiatric and neuroscience-based interventions in promoting human behaviour that improves population mental health while traducing extremist sentiments that are hazardous for health.

- 1 Barnes LL, de Leon CF, Lewis TT, Bienias JL, Wilson RS, Evans DA. Perceived discrimination and mortality in a population-based study of older adults. Am J Public Health 2008: 98: 1241–7.
- 2 Chae DH, Clouston S, Hatzenbuehler ML, Kramer MR, Cooper HL, Wilson SM, et al. Association between an internet-based measure of area racism and Black mortality. *PloS One* 2015; 10: e0122963.
- 3 Albert MA, Cozier Y, Ridker PM, Palmer JR, Glynn RJ, Rose L, et al. Perceptions of race/ethnic discrimination in relation to mortality among Black women: results from the Black Women's Health Study. Arch Intern Med 2010; 170: 896–904.
- 4 Polednak AP. Segregation, discrimination and mortality in US blacks. *Ethn Dis* 1996; **6**: 99–108.
- 5 Hatzenbuehler ML, Bellatorre A, Lee Y, Finch BK, Muennig P, Fiscella K. Structural stigma and all-cause mortality in sexual minority populations. Soc Sci Med 2014; 103: 33–41.
- 6 Hatch SL, Gazard B, Williams DR, Frissa S, Goodwin L, Hotopf M. Discrimination and common mental disorder among migrant and ethnic groups: findings from a South East London Community sample. Soc Psychiatry Psychiatric Epidemiol 2016; 51: 689–701.
- 7 Hamilton S, Corker E, Weeks C, Williams P, Henderson C, Pinfold V, et al. Factors associated with experienced discrimination among people using mental health services in England. *J Ment Health* 8 Feb 2016 (doi: 10.3109/09638237.2016.1139068).
- 8 Thornicroft G, Mehta N, Clement S, Evans-Lacko S, Doherty M, Rose D, et al. Evidence for effective interventions to reduce mental-health-related stigma and discrimination. *Lancet* 2016: 387: 1123–32.
- 9 Perez-Garin D, Molero F, Bos AE. The effect of personal and group discrimination on the subjective well-being of people with mental illness: the role of internalized stigma and collective action intention. *Psychol Health Med* 17 Mar 2016 (doi: 10.1080/13548506.2016.1164322).
- 10 Wallace S, Nazroo J, Becares L. Cumulative effect of racial discrimination on the mental health of ethnic minorities in the United Kingdom. Am J Public Health 2016; 106: 1294–300.
- 11 Nazroo JY. The structuring of ethnic inequalities in health: economic position, racial discrimination, and racism. *Am J Public Health* 2003; 93: 277–84.

- 12 Simons AM, Groffen DA, Bosma H. Income-related health inequalities: does perceived discrimination matter? Int J Public Health 2013; 58: 513–20.
- 13 Veling W, Selten JP, Susser E, Laan W, Mackenbach JP, Hoek HW. Discrimination and the incidence of psychotic disorders among ethnic minorities in The Netherlands. *Int J Epidemiol* 2007; **36**: 761–8.
- 14 Chae DH, Epel ES, Nuru-Jeter AM, Lincoln KD, Taylor RJ, Lin J, et al. Discrimination, mental health, and leukocyte telomere length among African American men. Psychoneuroendocrinology 2016; 63: 10–6.
- 15 Labonte B, Turecki G. The epigenetics of suicide: explaining the biological effects of early life environmental adversity. Arch Suicide Res 2010; 14: 291– 310
- 16 McGowan PO. Epigenetic clues to the biological embedding of early life adversity. *Biol Psychiatry* 2012; 72: 4–5.
- 17 Harris R, Tobias M, Jeffreys M, Waldegrave K, Karlsen S, Nazroo J. Effects of self-reported racial discrimination and deprivation on Maori health and inequalities in New Zealand: cross-sectional study. *Lancet* 2006; 367: 2005–9.

- 18 Cykert S. A new paradigm: approaching racism as a method of disease prevention. Chest 2014; 145: 442–3.
- 19 Kerrigan D, Vazzano A, Bertoni N, Malta M, Bastos FI. Stigma, discrimination and HIV outcomes among people living with HIV in Rio de Janeiro, Brazil: the intersection of multiple social inequalities. *Global Public Health* 10 Aug 2015 (doi: 10.1080/17441692.2015.1064459).
- 20 Mathur VA, Kiley KB, Haywood C Jr, Bediako SM, Lanzkron S, Carroll CP, et al. Multiple levels of suffering: discrimination in health-care settings is associated with enhanced laboratory pain sensitivity in sickle cell disease. Clin J Pain 1 Apr 2016 (doi: 10.1097/AJP.000000000000361).
- 21 Nelson CA. Biological embedding of early life adversity. JAMA Pediatrics 2013; 167: 1098–100.
- 22 Baxter AJ, Harris MG, Khatib Y, Brugha TS, Bien H, Bhui K. Reducing excess mortality due to chronic disease in people with severe mental illness: meta-review of health interventions. Br J Psychiatry 2016; 208: 222.0