with their parenting style, had a higher number of unprotected sexual contacts with unfamiliar persons, practiced group sex, had a history of sexual abuse, illicit drug use experience, were smokers, had piercing and severe body modifications. Lifetime history of suicide plan, attempts and NSSI were significantly more common in people with SSE (all: <0.05).

Conclusions: A number of suicide risk factors were found to be more prevalent in people with SSE. Homosexual experience in people with mental disorders is associated with an increased risk of NSSI, suicide plan development and suicide attempts.

Disclosure: No significant relationships.

Keywords: Suicide; NSSI; LGBT; non-psychotic mental disorders

O301

Rational suicide: The paradigm of survival

S. Torres

Psychiatry And Mental Health Department, Centro Hospitalar Barreiro-Montijo, Barreiro, Portugal
doi: 10.1192/j.eurpsy.2021.467

Introduction: Suicide is an intriguing act of the human being. The reasons behind the violation of an instinct for survival is far from being understood. Besides, the emergence of assisted dying is raising even more questions about the concept of rational suicide, defined as a well-thought-out decision to die by whom is mentally competent. **Objectives:** Understand the concept of rational suicide, in parallel with suicide, by exploring the views on this debate over the years and elucidating the relationship with mental disorders, mental capacity and patient's rights.

Methods: Literature review performed on PubMed and Google Scholar databases, using the keywords "rational suicide", "assisted death", "suicide", "phenomenology", "mental capacity" and "responsibility for life".

Results: The theological condemnations of suicide – as sin or crime – were put aside with psychiatric development in the last century. Durkheim was the first important precursor of the contemporary view - suicide is a form of mental illness (psychosis or depression) not compatible with rational deliberation. With the increasingly open debate on assisted dying, this vision is being tested by cases of terminally ill patients subjected to experiences that many wouldn't choose to tolerate. Moral right to self-determination and needless suffering are examples of arguments in favor of rational suicide.

Conclusions: The need for an open discussion about rational suicide is raising, specifically in relation to psychiatric disorders, mainly to resolve the conflict between the duty of care of psychiatrists and the autonomy of patients.

Disclosure: No significant relationships.

Keywords: rational suicide; assisted death; suicide; phenomenology

O302

Effect of group psychotherapy on the annual incidence of self-harm and suicide attempts in borderline personality disorder: A pilot study

I. Figuerio^{1*}, J. Labad², A. González-Rodríguez³, M. Santamaria¹, A.I. Cebrià¹, D. Palao Vidal¹ and J.A. Monreal⁴

¹Mental Health., Parc Tauli University Hospital. I3PT. UAB, CIBERSAM, Sabadell, Spain; ²Mental Health, Hospital of Mataró.

Consorti Sanitari del Maresme. CIBERSAM., Mataró, Spain; ³Mental Health, Parc Tauli University Hospital. Autonomous University of Barcelona (UAB). I3PT, Sabadell, Spain and ⁴Mental Health., Hospital Universitari Mútua de Terrassa. CIBERSAM, Terrassa, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.468

Introduction: Borderline personality disorder (BPD) has been characterized by mood instability, impulsive behavior and eventual dissociative and psychotic symptoms. Around 70% of patients present repeated self-injury behavior which is associated with high risk of completed suicide.

Objectives: To investigate the effect of group psychotherapy on the annual incidence of self-harm behavior and suicide attempts in BPD. **Methods:** We carried out a retrospective longitudinal study by selecting BPD patients who received group psychotherapy during 2016. Systems Training for Emotional Predictability and Problem Solving (STEPPS) or Mentalization-Based Treatment (MBT) psychotherapies were applied. Patients without any self-harm/suicidal attempt before the intervention, those with comorbid diagnosis and those who did not engage at least half of total sessions were excluded for final analyses. Number of self-harm events, suicide attempts and other clinical events were recorded and compared one-year before and one-year post-intervention. SPSS software version 21.0 (IBM) was used for statistical analyses. Nonparametric tests and Survival tests were performed.

Results: Eight women out of 35 fulfilled our inclusion criteria. After group psychotherapy, a significant reduction in the number of self-harm events and suicidal attempts was found (mean 1.9 $\pm$ 1.4 vs 0.5 $\pm$ 1.1; $p=0.042$). Survival tests revealed significant differences in the occurrence of suicidal attempts. We did not find significant differences in the other clinical events.

Conclusions: Our results show a clear effectiveness of group psychotherapy in reducing self-harm events and/or suicidal attempts in BPD patients. If these findings are confirmed in future studies including larger samples, group psychotherapy could be indicated for diminishing suicide risks in BPD.

Disclosure: No significant relationships.

Keywords: Borderline personality disorder; psychotherapy; Suicide; Self-harm behavior

O303

Behavioural addictions as risk factors for incidence and reoccurrence of suicide ideation and attempts in a prospective cohort study among young swiss men

M. Wicki^{1*}, M. Andronicos¹, J. Studer¹, S. Marmet¹ and G. Gmel^{1,2,3,4}

¹Addiction Medicine, Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland; ²Institute For Mental Health Policy Research, Centre for Addiction and Mental Health, Toronto, Canada; ³Frenchay Campus, University of the West of England, Bristol, United Kingdom and ⁴Research Institute, Addiction Switzerland, Lausanne, Switzerland

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.469

Introduction: Substance use disorder, depression and sexual minority are well documented risk factors for suicidal behaviour, far less is known about behavioural addictions.

Objectives: First, to explore associations between behavioural addictions (gaming, gambling, cybersex, internet, smartphone, work) at

age 25 and the incidence and reoccurrence of suicide ideation (SID), suicide attempts (SAT), and suicide attempts among those with suicide ideation (SATID) at age 28. Second, to test whether these associations were impacted by adjusting for cannabis and alcohol use disorder, nicotine dependence, sexual orientation and depression.

Methods: Based on two waves of a prospective cohort study of 5,428 young Swiss men, nested models with and without adjustment for risk factors were used to regress SID, SAT and SATID on preceding behavioural addictions.

Results: Without adjustment, each of the behavioural addictions at age 25 significantly predicted the incidence of SID and SAT at age 28. Gambling and cybersex addiction furthermore predicted SATID. When adjusting for other risk factors, associations with behavioural addictions were reduced, whereas depression and cannabis use disorder were the most important and consistent predictors for the incidence and recurrence of SID, SAT and SATID.

Conclusions: Among young Swiss men, behavioural addictions are important predictors of SID and SAT, however a large part of their association is shared with depression and cannabis use disorder. Treatment for addictive behaviors, especially cannabis use can open the door to larger mental health screening and targeted intervention. Crisis intervention among men presenting addictive behaviours with or without substance may therefore be key to preventing suicidal behaviour.

Disclosure: No significant relationships.

Keywords: suicide ideation and attempts; cohort study; addictive behaviours; Depression

O304

The reciprocal relation between stigma and suicidality in a sample of patients with affective disorders

L. Giannelli*, G. Cristilli, L. Marone, G. De Felice, A. Carello, M. Di Vincenzo, V. Sollo, A. Di Cerbo, V. Giallonardo, G. Sampogna, V. Del Vecchio, M. Luciano and A. Fiorillo

Department Of Psychiatry, University of Campania “Luigi Vanvitelli”, Naples, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.470

Introduction: Suicide is one of the major public health concerns worldwide, currently listed as the 15th most common cause of death. Mental illness stigma may contribute to suicidality and is associated with social isolation and low self-esteem among people with affective disorders.

Objectives: The aim of the present study is to assess, in a sample of people with affective disorders, whether high levels of internalized stigma are associated to suicidal thoughts and behaviours.

Methods: 60 outpatients diagnosed with depression or bipolar disorder according to DSM-5 have been recruited. Suicidal behaviours and ideation were assessed through the Columbia Suicide Severity Rating Scale (C-SSRS); internalized stigma through the Internalized Stigma of Mental Illness (ISMI) scale. Socio-demographic characteristics have been collected through an ad hoc schedule.

Results: 62.9% of the sample was female, with a mean age of 45.7 (± 14) years. About half of the sample had a diagnosis of major depression (54.8%). Patients with suicidal ideation reported higher score at ISMI “alienation” subscale ($p < 0.05$), compared to those without suicidal ideation. Patients with a previous history suicide attempts reported higher score at “alienation” and “social withdrawal”

ISMI subscales ($p < 0.05$). Moreover, “alienation” ISMI subscale significantly correlated with suicidal ideation and behaviours ($p < 0.01$).

Conclusions: These results are in line with the available literature, highlighting that stigma and suicidality are strongly correlated. This underline the importance of interventions at addressing internalizing stigma, in particular to those with previous suicidal attempts and with an active suicidal ideation.

Disclosure: No significant relationships.

Keywords: Stigma; Suicide; suicide prevention

O305

Outcomes of a regional suicide prevention systems intervention study: Suicide prevention by monitoring and collaborative care (SUPREMOCOL) in noord-brabant in the netherlands

C. Van Der Feltz-Cornelis*

Health Sciences, University of York, Heslington, York, United Kingdom

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.471

Introduction: Since 2007, suicide rates increased in the Netherlands and the province of Noord-Brabant ranked second nationally with a 64% increase. 60% of people who died by suicide did not receive treatment at the time of their death. Gap analysis showed 1) lack of expertise to explore suicide risk in health care or community settings where persons at risk presented; 2) lack of swift access to specialist care addressing suicidality; 3) lack of oversight of the care process and 4) lack of follow up.

Objectives: We developed a regional suicide prevention systems intervention with chain partners at community, general health and mental health care level to address these gaps in Noord-Brabant, aiming at a 20% decrease in the number of suicides.

Methods: The project started October 2016 and lasted 4 years. The intervention has four pillars: 1) Online decision aid for health care professionals to assess suicidal risk and to communicate with chain partners; 2) swift access to care; 3) facilitation of care through the care chain by a dedicated nurse; and 4) 12 months follow up monitoring if the patient still receives appropriate care. We examined the effect of SUPREMOCOL on suicides in a pre-post design.

Results: During the implementation year of the intervention, suicides in Noord-Brabant dropped 17% whereas nationally they dropped 5%, and this effect was sustained after one year.

Conclusions: This suicide prevention systems intervention is effective in reducing suicide rates. Long-term follow-up and implementation is warranted.

Disclosure: No significant relationships.

Keywords: SUPREMOCOL; Digital monitoring; Suicide prevention; Systems intervention

O306

Association of traumatic events in childhood, impulsivity and decision-making with previous suicide attempt and/or current suicidal ideation in adult patients with major depressive disorder

A. Roman^{1*}, J. Rodríguez-Revuelta^{2,3,4,5,6}, F. Dal Santo^{2,3,4,5,6}, L. García-Alvarez^{2,3,4,5,6}, L. De La Fuente-Tomás^{2,3,4,5,6}