



RESEARCH ARTICLE

Cohomological χ -dependence of ring structure for the moduli of one-dimensional sheaves on \mathbb{P}^2

Woonam Lim¹, Miguel Moreira² and Weite Pi³

¹Utrecht University – Department of Mathematics, Budapestlaan 6, 3584 CD Utrecht, The Netherlands; E-mail: w.lim@uu.nl.
²Massachusetts Institute of Technology – Department of Mathematics, 77 Massachusetts Avenue, Cambridge MA, 02139, USA; E-mail: miguel@mit.edu.
³Yale University – Department of Mathematics, 219 Prospect St, New Haven, CT 06511, USA; E-mail: weite.pi@yale.edu (corresponding author).

Received: 21 June 2023; Revised: 16 February 2024; Accepted: 27 February 2024

2020 Mathematics subject classification: Primary – 14D20; Secondary – 14C15

Abstract

We prove that the cohomology rings of the moduli space $M_{d,\chi}$ of one-dimensional sheaves on the projective plane are not isomorphic for general different choices of the Euler characteristics. This stands in contrast to the χ -independence of the Betti numbers of these moduli spaces. As a corollary, we deduce that $M_{d,\chi}$ are topologically different unless they are related by obvious symmetries, strengthening a previous result of Woolf distinguishing them as algebraic varieties.

Contents

1	Introduction	1
1.1	Moduli of sheaves	1
1.2	BPS invariants and χ -independence	3
1.3	Structure of the proof of Theorem 1.2	4
1.4	Relations to other works	4
2	Tautological classes and relations in $A^d(M_{d,\chi})$	5
2.1	Tautological classes	5
2.2	Tautological relations	6
2.3	Three relations in $A^d(M_{d,\chi})$	8
3	Truncated relations and proof of the main result	10
3.1	Type I	13
3.2	Type II	14
3.3	Type II – extended matrices	14

1. Introduction

We work over the complex numbers \mathbb{C} .

1.1. Moduli of sheaves

Fix two integers d and χ with $d \geq 1$. We consider the moduli space $M_{d,\chi}$ of semistable one-dimensional sheaves \mathcal{F} on \mathbb{P}^2 with

$$[\text{supp}(\mathcal{F})] = d \cdot H \in H_2(\mathbb{P}^2, \mathbb{Z}), \quad \chi(\mathcal{F}) = \chi.$$

© The Author(s), 2024. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

Here, H is the class of a line, $\text{supp}(\mathcal{F})$ denotes the Fitting support and the stability condition is with respect to the slope

$$\mu(\mathcal{E}) = \frac{\chi(\mathcal{E})}{c_1(\mathcal{E}) \cdot H} \in \mathbb{Q}.$$

Le Potier [LeP] first studied this moduli space and showed that $M_{d,\chi}$ is an irreducible projective variety of dimension $d^2 + 1$, nonsingular at all stable points. In particular, when d and χ are coprime, semistability coincides with stability and $M_{d,\chi}$ is nonsingular. Geometry and topology of the moduli space $M_{d,\chi}$ have been intensively studied from various perspectives; see [PS, Introduction] for a brief overview. In this paper, we are interested in the cohomology rings $H^*(M_{d,\chi}, \mathbb{C})$ for a fixed d and *different* Euler characteristics χ coprime to d . Under the coprime assumption, the cycle class map

$$\text{cl} : \text{CH}^*(M_{d,\chi}) \longrightarrow H^{2*}(M_{d,\chi}, \mathbb{Z})$$

is an isomorphism by [Mar, Theorem 2]. Hence, in this case we use $A^*(-)$ to denote the even cohomology $H^{2*}(-, \mathbb{C})$, or equivalently the Chow ring $\text{CH}^*(-, \mathbb{C})$ with complex coefficients.

The moduli spaces $M_{d,\chi}$ admit two types of symmetries:

(a) The first type of symmetry is given by the isomorphism

$$\psi_1 : M_{d,\chi} \xrightarrow{\sim} M_{d,\chi+d}, \quad \mathcal{F} \mapsto \mathcal{F} \otimes \mathcal{O}_{\mathbb{P}^2}(1).$$

(b) The second type of symmetry [Mai, Theorem 13] is given by the duality isomorphism

$$\psi_2 : M_{d,\chi} \xrightarrow{\sim} M_{d,-\chi}, \quad \mathcal{F} \mapsto \text{Ext}^1(\mathcal{F}, \omega_{\mathbb{P}^2}).$$

Thus, given two Euler characteristics χ, χ' satisfying $\chi \equiv \pm\chi' \pmod{d}$, there is a natural isomorphism $M_{d,\chi} \simeq M_{d,\chi'}$. The following theorem states that this is essentially the only case.

Theorem 1.1 [Woo, Theorem 8.1]. *For $d \geq 3$, there is an isomorphism of algebraic varieties*

$$M_{d,\chi} \simeq M_{d,\chi'}$$

if and only if $\chi \equiv \pm\chi' \pmod{d}$.

The main result of this paper is a cohomological version of Theorem 1.1:

Theorem 1.2. *For $d \geq 1$ and χ, χ' coprime to d , there is an isomorphism of graded \mathbb{C} -algebras*

$$A^*(M_{d,\chi}) \simeq A^*(M_{d,\chi'})$$

if and only if $\chi \equiv \pm\chi' \pmod{d}$.

We find this cohomological χ -dependence of ring structure interesting since the (intersection) Betti numbers of $M_{d,\chi}$ are χ -independent: In particular, we have an isomorphism

$$A^*(M_{d,\chi}) \simeq A^*(M_{d,\chi'})$$

of graded vector spaces for arbitrary χ and χ' coprime to d ; see Theorem 1.4 for the precise statement. As a corollary of Theorem 1.2, we deduce that $M_{d,\chi}$ are in general topologically different, strengthening Theorem 1.1:

Corollary 1.3. *For any χ, χ' coprime to d , the moduli spaces $M_{d,\chi}$ and $M_{d,\chi'}$ are not homeomorphic unless $\chi \equiv \pm\chi' \pmod{d}$.*

1.2. BPS invariants and χ -independence

One motivation to study the moduli spaces $M_{d,\chi}$ comes from enumerative geometry. Let $X = \text{Tot}(K_{\mathbb{P}^2})$ be the local Calabi–Yau 3-fold given by the total space of the canonical bundle on \mathbb{P}^2 . Considerations from physics [GV] predict an action of the Lie algebra $\mathfrak{sl}_2 \times \mathfrak{sl}_2$ on the cohomology of a certain moduli space of D -branes supported on degree d curves in X , yielding double-indexed integral invariants

$$n_d^{i,j} \in \mathbb{Z} \tag{1.1}$$

as the dimensions of the weight spaces of this $\mathfrak{sl}_2 \times \mathfrak{sl}_2$ -action. These are known as *refined BPS invariants* of X , named after Bogomol’nyi–Prasad–Sommerfield, which are expected to refine curve counting invariants for X defined via Gromov–Witten/Donaldson–Thomas/Pandharipande–Thomas theory [PT].

One proposal by Hosono–Saito–Takahashi [HST], Kiem–Li [KL] and Maulik–Toda [MT] suggests a mathematical definition of the invariants (1.1) by the *perverse filtration* on the cohomology of moduli spaces of one-dimensional sheaves. More precisely, the moduli space $M_{d,\chi}$ admits a Hilbert–Chow morphism

$$h : M_{d,\chi} \longrightarrow \mathbb{P}H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d)),$$

sending a sheaf to its Fitting support. This map is proper and induces an increasing filtration on the intersection cohomology

$$P_0 \text{IH}^*(M_{d,\chi}) \subset P_1 \text{IH}^*(M_{d,\chi}) \subset \dots \subset \text{IH}^*(M_{d,\chi})$$

called the perverse filtration; see [KPS, Section 1.1]. The invariants (1.1) are defined in *loc. cit.* as the dimension of the graded pieces of this filtration:

$$n_d^{i,j} := \dim \text{Gr}_i^P \text{IH}^{i+j}(M_{d,\chi}). \tag{1.2}$$

For this to be well defined, the right-hand side in Equation (1.2) should not depend on the choice of χ , which is a priori nontrivial in light of Theorem 1.1. This is a special case of Toda’s cohomological χ -independence conjecture [Tod, Conjecture 1.2]; see also [Bou, Conjecture 0.4.3] for a version on the intersection Betti numbers of $M_{d,\chi}$. The following theorem of Maulik and Shen confirms this conjecture.¹

Theorem 1.4 [MS1, Theorem 0.1]. *For any $\chi, \chi' \in \mathbb{Z}$ not necessarily coprime to d , there is a (non-canonical) isomorphism of graded vector spaces*

$$\text{IH}^*(M_{d,\chi}) \simeq \text{IH}^*(M_{d,\chi'})$$

preserving the perverse filtration on both sides.

If we restrict to the coprime case, intersection cohomology coincides with singular cohomology which admits a canonical \mathbb{Q} -algebra structure. It is then natural to ask if we can choose the isomorphism of Theorem 1.4 to be an isomorphism of \mathbb{Q} -algebras. More generally, we can ask the following:

Question 1.5 [PS, Section 0.3]. *For any χ, χ' coprime to d , is there an isomorphism*

$$H^*(M_{d,\chi}, \mathbb{Q}) \simeq H^*(M_{d,\chi'}, \mathbb{Q})$$

of graded \mathbb{Q} -algebras?

Theorem 1.2 gives a complete answer to this question in the stronger sense that we work with \mathbb{C} -coefficients instead of \mathbb{Q} .

¹The case where χ, χ' are coprime to d was previously proven by Bousseau [Bou].

1.3. Structure of the proof of Theorem 1.2

We outline the proof of our main result here, as the argument is of an elementary nature but somewhat long. It was explained in [PS] that the rings $A^*(M_{d,\chi})$ admit a minimal set of $3d - 7$ generators

$$T = \{c_0(2), c_2(0)\} \cup \bigcup_{k=2}^{d-2} \{c_{k-1}(2), c_k(1), c_{k+1}(0)\}.$$

In other words, there is a surjective graded algebra homomorphism $\mathbb{C}[T] \rightarrow A^*(M_{d,\chi})$. This homomorphism is an isomorphism up to degree $d - 1$, but in degree d there are exactly three relations, cf. Theorem 2.2. As a first step toward the proof, we show that these three relations R_1, R_2, R_3 spanning $\ker(\mathbb{C}[T]^d \rightarrow A^d(M_{d,\chi}))$ can be obtained explicitly from the tautological relations of [PS, Section 2], cf. Proposition 2.7.

We then proceed by showing that if χ and χ' are not related by the symmetries (a) and (b), that is, $\chi \not\equiv \pm\chi' \pmod{d}$, then there is no automorphism of the graded algebra $\mathbb{C}[T]$ that sends the three-dimensional subspace of relations $\ker(\mathbb{C}[T]^d \rightarrow A^d(M_{d,\chi})) \subseteq \mathbb{C}[T]^d$ to $\ker(\mathbb{C}[T]^d \rightarrow A^d(M_{d,\chi'}))$. To achieve this, we do not use the full relations R_1, R_2, R_3 , whose sizes grow considerably as d increases but only a truncated version of those. Concretely, we project the subspaces of relations onto the 18-dimensional subspace

$$c_{d-3}(2) \cdot \mathbb{C}[T]^2 \oplus c_{d-2}(1) \cdot \mathbb{C}[T]^2 \oplus c_{d-1}(0) \cdot \mathbb{C}[T]^2 \subseteq \mathbb{C}[T]^d.$$

This truncated form of the relations is completely explicit: For each of the three relations, we only need to compute the 18 coefficients that are rational functions in d and χ .

Showing that there is no automorphism of $\mathbb{C}[T]$ relating the (truncated) subspaces of relations for χ and χ' then becomes a very concrete, but still involved, linear algebra problem. For this, we employ a determinant trick in Step 2 of Section 3 that imposes strong restrictions on the possible automorphisms. The latter fall into two types, which we call *Type I* and *Type II* solutions. We treat them separately in Sections 3.1–3.3, showing that both cannot exist by analyzing certain numerical constraints (see, for example, Equation (3.7)) on the triple (d, χ, χ') .

Our proof of Theorem 1.2 involves heavy computations aided by the software Mathematica [M]. The code (together with a printed file) has been uploaded to the third author's website <https://github.com/Weite-Pi/weitepi.github.io> under the name *cohomological ring chi-dependence*, which we shall frequently refer to in the later part of this paper.

1.4. Relations to other works

The moduli space $M_{d,\chi}$ shares similar features with two types of other moduli spaces, the moduli of Higgs bundles² and the moduli of one-dimensional sheaves on K3 surfaces. The parallel of Question 1.5 for those cases has a positive answer. Cohomology rings of the moduli of Higgs bundles are proven to be χ -independent in the stronger sense that perverse filtrations are preserved, as predicted by the $P = W$ conjecture [dCHM, MS2, HMMS]. A direct proof using techniques in characteristic p is given in [dCMSZ]. The case of K3 surfaces follows from the fact that moduli of sheaves on K3 surfaces with respect to a primitive class and generic stability is birational to the Hilbert scheme of points [BM] and that any two birational projective hyperkähler manifolds are deformation equivalent [Huy].

Our main result asserts that the cohomology rings $H^*(M_{d,\chi}, \mathbb{C})$ are χ -dependent in general; on the other hand, some χ -independent multiplicative structures have been observed or speculated:

- It is conjectured and proven under certain assumptions in [BLM] that the *Virasoro constraints* hold for the moduli spaces $M_{d,\chi}$. Briefly speaking, the Virasoro constraints predict that certain intersection

²Recall that Higgs bundles on a curve C are in correspondence with one-dimensional sheaves on the surface $\text{Tot}(K_C)$ via the spectral correspondence [BNR]. This analogy provides another motivation to study $M_{d,\chi}$; see [PS, Introduction] for details.

numbers on $M_{d,\chi}$ obtained by integrating natural cohomology classes satisfy some explicit universal relations. See [BLM, Section 1.3] for the precise statement.

- The main theorem of [PS] is a uniform³ minimal generation and freeness result on $H^*(M_{d,\chi}, \mathbb{C})$ for all χ coprime to d ; see Theorem 2.2 for a precise and slightly stronger statement. The $P = C$ conjecture formulated in [KPS] seeks to identify the perverse filtration on the free part of $H^*(M_{d,\chi}, \mathbb{Q})$ with an explicit *Chern filtration* defined in terms of the generators. This prediction is also independent of χ .
- Finally, we remark that the perverse filtration on $H^*(M_{d,\chi}, \mathbb{C})$ comes from the ring structure. Indeed, denoting by L the pull-back of the hyperplane class by the Hilbert–Chow morphism, it is completely determined by the multiplication operator L according to [dCM, Proposition 5.2.4].

2. Tautological classes and relations in $A^d(M_{d,\chi})$

For the remainder of this paper, we assume χ, χ' are coprime to d . This section provides some preliminary results on the cohomology of the moduli spaces $M_{d,\chi}$. We recall the normalized tautological classes and certain tautological relations introduced in [PS] and prove a first result on the relations in $A^d(M_{d,\chi})$.

2.1. Tautological classes

Under the assumption that $\gcd(d, \chi) = 1$, there exists a universal sheaf [HL, Theorem 4.6.5] over the product $\mathbb{P}^2 \times M_{d,\chi}$ which we denote by \mathbb{F} . For a stable sheaf $[\mathcal{F}] \in M_{d,\chi}$, the restriction of \mathbb{F} to the fiber $\mathbb{P}^2 \times [\mathcal{F}]$ recovers \mathcal{F} .

Consider the two projection maps π_P and π_M from $\mathbb{P}^2 \times M_{d,\chi}$ to its first and second components. One way to obtain natural cohomology classes on $M_{d,\chi}$ is to take the Chern characters of \mathbb{F} , intersect with classes pulled back from \mathbb{P}^2 and push forward to $M_{d,\chi}$. The choice of \mathbb{F} is, however, not unique: Any two universal sheaves differ by tensoring with a line bundle pulled back from $M_{d,\chi}$. We thus conduct a normalization of \mathbb{F} as follows.

For a universal sheaf \mathbb{F} and a class

$$\delta = \pi_P^* \delta_P + \pi_M^* \delta_M \in A^1(\mathbb{P}^2 \times M_{d,\chi}), \quad \text{with } \delta_P \in A^1(\mathbb{P}^2), \quad \delta_M \in A^1(M_{d,\chi}),$$

we consider the *twisted* Chern character

$$\text{ch}^\delta(\mathbb{F}) := \text{ch}(\mathbb{F}) \cdot \exp(\delta)$$

and denote by $\text{ch}_k^\delta(\mathbb{F})$ its degree k -part. For any class $\gamma \in A^*(\mathbb{P}^2)$, we write

$$\int_\gamma \text{ch}_k^\delta(\mathbb{F}) := \pi_{M*}(\pi_P^* \gamma \cdot \text{ch}_k^\delta(\mathbb{F})) \in A^*(M_{d,\chi}).$$

It is proven in [PS, Proposition 1.2] that for a fixed \mathbb{F} , there exists a unique class $\delta_0 = \delta_0(\mathbb{F})$ as above satisfying the normalizing conditions

$$\int_H \text{ch}_2^{\delta_0}(\mathbb{F}) = 0, \quad \int_{\mathbb{1}_{\mathbb{P}^2}} \text{ch}_2^{\delta_0}(\mathbb{F}) = 0. \tag{2.1}$$

This class is computed explicitly in [KPS, Proposition 2.1]:

$$\delta_0 = \left(\frac{3}{2} - \frac{\chi}{d}\right) \cdot H - \frac{1}{d} \left(\left(\frac{3}{2} - \frac{\chi}{d}\right) \int_{H^2} \text{ch}_1(\mathbb{F}) + \int_H \text{ch}_2(\mathbb{F})\right). \tag{2.2}$$

³In the sense that the generators $c_k(j)$ in Theorem 2.2 are defined without explicit reference to χ .

Note that the twisted Chern character $\text{ch}^{\delta_0}(\mathbb{F})$ does *not* depend on the choice of \mathbb{F} anymore. With this normalizing class, we define the *tautological classes*

$$c_k(j) := \int_{H^j} \text{ch}_{k+1}^{\delta_0}(\mathbb{F}) \in A^{k+j-1}(M_{d,\chi}). \tag{2.3}$$

We collect some basic properties of the tautological classes from [PS, Section 1.2].

Proposition 2.1. *Let $c_k(j)$ be the tautological classes defined by Equation (2.3).*

- (a) *The classes $c_k(j)$ do not depend on the choice of a universal sheaf.*
- (b) *We have*

$$c_1(0) = 0 \in A^0(M_{d,\chi}), \quad c_1(1) = 0 \in A^1(M_{d,\chi}), \quad c_0(1) = d \in A^0(M_{d,\chi}).$$

- (c) *Let ψ_1 and ψ_2 be the two symmetries introduced in Section 1.1. Then we have*

$$\psi_1^* c_k(j) = c_k(j), \quad \psi_2^* c_k(j) = (-1)^k c_k(j).$$

The next theorem gives a first result on the structure of $A^*(M_{d,\chi})$ in terms of the tautological classes. Note that we change the coefficient from \mathbb{Q} to \mathbb{C} .

Theorem 2.2 [PS, Y2]. *Assume $d \geq 5$. We have:*

- (a) *$A^*(M_{d,\chi})$ is generated as a \mathbb{C} -algebra by the $3d - 7$ classes⁴ of degrees $\leq d - 2$:*

$$c_0(2), c_2(0) \in A^1(M_{d,\chi}), \quad c_k(0), c_{k-1}(1), c_{k-2}(2) \in A^{k-1}(M_{d,\chi}), \quad 3 \leq k \leq d - 1. \tag{2.4}$$

- (b) *There is no relation among these $3d - 7$ classes in degrees $\leq d - 1$.*
- (c) *There are exactly three linearly independent relations in degree d .*

We briefly recall the proof of Theorem 2.2: (i) By the results of Beauville [Bea] or Markman [Mar], the ring $A^*(M_{d,\chi})$ is generated by the tautological classes. (ii) Using the geometry of $M_{d,\chi}$, we produce tautological relations that express any tautological class in terms of the $3d - 7$ classes in Equation (2.4); see the paragraph after Proposition 2.6. (iii) Part (b) and (c) follow from comparing the Betti numbers of $M_{d,\chi}$ with those of Hilbert schemes of points on \mathbb{P}^2 ; this is explained in Yuan [Y1, Y2].

As we will see, the three relations in $A^d(M_{d,\chi})$ are the central characters in our proof of Theorem 1.2. By carefully investigating the χ -dependence of the three relations, we are able to deduce that $A^*(M_{d,\chi})$ are nonisomorphic for different choices of χ unless they are related by the symmetries in Section 1.1.

Remark 2.3. The choice of the normalization (2.1) and the shift in degrees of the Chern character in Equation (2.3) are motivated by the $P = C$ conjecture [KPS]. Roughly speaking, the $P = C$ conjecture predicts that the *Chern grading* of the tautological generators $c_k(j)$ coincides with its *perversity*; this gives a conjectural explicit description of the perverse filtration on $A^{\leq d-2}(M_{d,\chi})$ in terms of Equation (2.4). See [KPS, Conjecture 0.3 and Proposition 1.2] for details.


2.2. Tautological relations

We recall in this section certain tautological relations [PS, Section 1.2] on the cohomology of $M_{d,\chi}$. By the symmetry (a) in Section 1.1, we may assume that $0 < \chi < d$ so that any $\mathcal{F} \in M_{d,\chi}$ satisfies $0 < \mu(\mathcal{F}) < 1$.

Consider the triple product $Y := \mathbb{P}^2 \times M_{d,\chi} \times \check{\mathbb{P}}^2$, where $\check{\mathbb{P}}^2$ is the dual projective space parametrizing lines in \mathbb{P}^2 . Let $\pi_R : Y \rightarrow \check{\mathbb{P}}^2$ be the third projection. We write $p = \pi_P \times \pi_M$, $q = \pi_P \times \pi_R$ and

⁴We call these $3d - 7$ classes the *tautological generators*.

$r = \pi_M \times \pi_R$:



Let $Z \subset \mathbb{P}^2 \times \check{\mathbb{P}}^2$ be the incidence subscheme, and let \mathcal{O}_Z be its structure sheaf. For a fixed universal sheaf \mathbb{F} on $M_{d,\mathcal{X}}$, we consider the complex

$$\mathcal{H}(n) := R\mathcal{H}om(p^*\mathbb{F}, q^*\mathcal{O}_Z \otimes \pi_P^*\mathcal{O}_{\mathbb{P}^2}(-n)) \in D^b\text{Coh}(Y). \tag{2.5}$$

The projection $r : Y \rightarrow M_{d,\mathcal{X}} \times \check{\mathbb{P}}^2$ is a trivial \mathbb{P}^2 -bundle, so the derived push-forward

$$Rr_*\mathcal{H}(n) \in D^b(M_{d,\mathcal{X}} \times \check{\mathbb{P}}^2)$$

admits a three-term resolution $0 \rightarrow K^0 \rightarrow K^1 \rightarrow K^2 \rightarrow 0$ by vector bundles.

Lemma 2.4. *For $n \in \{1, 2, 3\}$, we can choose K^i with $K^0 = K^2 = 0$ and K^1 free of rank d .*

Proof. This is [PS, Lemma 2.4]; we briefly recall the proof here. Take a point

$$P = ([\mathcal{F}], [L]) \in M_{d,\mathcal{X}} \times \check{\mathbb{P}}^2,$$

where $[L] \in \check{\mathbb{P}}^2$ corresponds to the line $L \subseteq \mathbb{P}^2$. Over the point P , cohomology of the complex $K^0(P) \rightarrow K^1(P) \rightarrow K^2(P)$ computes the extension groups

$$\text{Ext}^i(\mathcal{F}, \mathcal{O}_L(-n)), \quad i = 1, 2, 3.$$

Note that $\mu(\mathcal{O}_L(-n)) = 1 - n$. By stability and Serre duality, one checks that

$$\text{Ext}^0(\mathcal{F}, \mathcal{O}_L(-n)) = 0, \quad \text{Ext}^2(\mathcal{F}, \mathcal{O}_L(-n)) = 0$$

as long as $n \in \{1, 2, 3\}$. This implies $Rr_*\mathcal{H}(n)$ can be represented by a single vector bundle K^1 concentrated in degree 1, whose rank is d by a Hirzebruch–Riemann–Roch calculation. \square

It follows that $-Rr_*\mathcal{H}(n)$ is a rank d vector bundle on $M_{d,\mathcal{X}} \times \check{\mathbb{P}}^2$, whence

Corollary 2.5. *For $\ell \geq d + 1$ and $n \in \{1, 2, 3\}$, we have*

$$c_\ell(-Rr_*\mathcal{H}(n)) = 0 \in A^\ell(M_{d,\mathcal{X}} \times \check{\mathbb{P}}^2). \tag{2.6}$$

Recall that $\mathcal{H}(n)$ is defined by Equation (2.5) in terms of a fixed universal sheaf \mathbb{F} . If we replace \mathbb{F} by

$$\mathbb{F}' := \mathbb{F} \otimes \pi_M^*L,$$

with L a fractional line bundle over $M_{d,\mathcal{X}}$, then according to Equation (2.5) and the projection formula, the vector bundle $-Rr_*\mathcal{H}(n)$ is replaced by its tensor with the dual line bundle $\pi_M^*L^\vee$. Using a formal argument involving Chern roots, one checks that the identity (2.6) also holds for \mathbb{F}' .

In particular, we choose the fractional line bundle L to have first Chern class⁵

$$c_1(L) = -\frac{1}{d} \left(\left(\frac{3}{2} - \frac{\mathcal{X}}{d} \right) c_0(2) + \int_H \text{ch}_2(\mathbb{F}) \right)$$

⁵One can check that $\int_{H^2} \text{ch}_1(\mathbb{F}) = c_0(2)$ for every universal sheaf \mathbb{F} .

so that the normalizing class (2.2) for \mathbb{F}' is given by $\delta_0(\mathbb{F}') = \left(\frac{3}{2} - \frac{\chi}{d}\right) \cdot H$. Now, we apply Equation (2.6) to \mathbb{F}' . The Chern classes in Equation (2.6) can be expressed in terms of the Chern character $\text{ch}(Rr_*\mathcal{H}(n))$, which is computed by the Grothendieck–Riemann–Roch theorem:

$$\begin{aligned} \text{ch}(Rr_*\mathcal{H}(n)) &= r_*(\text{ch}(\mathcal{H}(n)) \cdot \text{td}(\mathbb{P}^2)) \\ &= r_*(\text{ch}(p^*\mathbb{F}'^\vee) \cdot \text{ch}(q^*\mathcal{O}_Z \otimes \pi_p^*\mathcal{O}_{\mathbb{P}^2}(-n)) \cdot \text{td}(\mathbb{P}^2)). \end{aligned}$$

We denote by β the class of a line on \mathbb{P}^2 . Expanding the right-hand side of the above equation, Corollary 2.5 then gives relations in $A^*(M_{d,\chi} \times \mathbb{P}^2)$:

Proposition 2.6. *For every $\ell \geq d + 1$ and $n \in \{1, 2, 3\}$, the following identity holds:*

$$\sum_{\mathbf{m}} \prod_{s=1}^{\ell} \frac{((s-1)!)^{m_s}}{(m_s)!} \left(\pi_M^* A_s - \sum_{0 \leq i \leq 2} \frac{\pi_R^* \beta^i}{i!} (-1)^i \pi_M^* B_{s-i} \right)^{m_s} = 0. \tag{2.7}$$

Here, the first sum is over all ℓ -tuple of nonnegative integers $\mathbf{m} = (m_1, m_2, \dots, m_\ell)$ such that $m_1 + 2m_2 + \dots + \ell m_\ell = \ell$, and writing $\tilde{c}_s(j) := (-1)^{s+1} c_s(j)$, the terms A_s, B_s are given by

$$\begin{aligned} A_s &:= \tilde{c}_{s+1}(0) + \left(3 - n - \frac{\chi}{d}\right) \tilde{c}_s(1) + \frac{1}{2d^2} \left(\left(n - \frac{7}{2}\right)d + \chi \right) \left(\left(n - \frac{5}{2}\right)d + \chi \right) \tilde{c}_{s-1}(2), \\ B_s &:= \tilde{c}_{s+1}(0) + \left(2 - n - \frac{\chi}{d}\right) \tilde{c}_s(1) + \frac{1}{2d^2} \left(\left(n - \frac{5}{2}\right)d + \chi \right) \left(\left(n - \frac{3}{2}\right)d + \chi \right) \tilde{c}_{s-1}(2). \end{aligned}$$

Proof. The proof is essentially the same as [PS, Proposition 2.6], except that we use the normalized Chern character $\text{ch}^{\delta_0}(\mathbb{F}'^\vee) = \text{ch}(\mathbb{F}'^\vee) \cdot \exp(-\delta_0)$. The convoluted formula results from applying Newton’s identity to express Chern classes in the Chern characters, and the class β comes from $\text{ch}(q^*\mathcal{O}_Z)$. See also [KPS, Proposition 2.7] for an explicit version for $M_{4,1}$. □

To obtain relations in $A^*(M_{d,\chi})$, we first take the identity (2.7) for some $\ell \geq d + 1$; this gives a relation in $A^\ell(M_{d,\chi} \times \mathbb{P}^2)$. Then we multiply it by $\pi_R^*(\beta^j)$ and push forward to $M_{d,\chi}$, where $0 \leq j \leq 2$. This produces a relation in $A^{\ell+j-2}(M_{d,\chi})$ among the tautological classes (2.3). The procedure is explained in detail in [PS, Section 2.3] and the paragraph before it. Indeed, as a key step in the proof of Theorem 2.2(a), it is shown in [PS, Section 2.3] that the relations produced by Proposition 2.6 express all the tautological classes (2.3) in terms of the $3d - 7$ generators (2.4).

2.3. Three relations in $A^d(M_{d,\chi})$

The goal of this section is to prove the following result:

Proposition 2.7. *For $d \geq 5$, the three linearly independent relations in $A^d(M_{d,\chi})$, cf. Theorem 2.2(c), can be produced by Proposition 2.6.*

We first introduce some notations for the rest of the paper. Let T^k be the linear space spanned by tautological classes in degree k , that is,

$$T^1 = \text{span}_{\mathbb{C}}\{c_2(0), c_0(2)\}, \quad T^k = \text{span}_{\mathbb{C}}\{c_{k+1}(0), c_k(1), c_{k-1}(2)\}.$$

Let

$$T := \bigoplus_{k=1}^{d-2} T^k$$

be the span of the $3d - 7$ tautological generators in Equation (2.4), and let

$$T^+ := T \oplus T^{d-1} \oplus T^d.$$

We denote by $\mathbb{C}[T]$ the symmetric algebra in T (i.e., the free algebra generated by the $3d - 7$ tautological generators), and by $\mathbb{C}[T]^k$ its graded piece of degree k . Similarly, we write $\mathbb{C}[T^+]$ for the symmetric algebra on T^+ .

Furthermore, we endow the basis of T^+ with a total ordering $<$ as follows: First order the basis by cohomological degrees; within each degree, we order by the Chern grading of the basis. In a precise way, this means $c_k(j) < c_{k'}(j')$ if and only if $k + j - 1 < k' + j' - 1$, or $k + j - 1 = k' + j' - 1$ and $k < k'$. Finally, we endow monomials in each degree of $\mathbb{C}[T^+]$ the lexicographical order induced by the total ordering $<$ on T^+ . In this way, we can talk about the leading term of a polynomial in $\mathbb{C}[T^+]$, as well as in $\mathbb{C}[T]$.

We have graded algebra homomorphisms

$$\mathbb{C}[T] \hookrightarrow \mathbb{C}[T^+] \rightarrow A^*(M_{d,\chi}).$$

Theorem 2.2 states that the homomorphism $\mathbb{C}[T] \rightarrow A^*(M_{d,\chi})$ is surjective, that it is an isomorphism up to degree $d - 1$ and that $\ker(\mathbb{C}[T]^d \rightarrow A^d(M_{d,\chi}))$ is 3-dimensional. A relation is by definition an element in the kernel of $\mathbb{C}[T] \rightarrow A^*(M_{d,\chi})$ or $\mathbb{C}[T^+] \rightarrow A^*(M_{d,\chi})$.

With the above setup, Theorem 2.2 (b) and (c) are equivalent to the following.⁶

Proposition 2.8. *For $d \geq 5$, we have*

$$\dim A^k(M_{d,\chi}) = \begin{cases} \dim \mathbb{C}[T^+]^k, & k \leq d - 2, \\ \dim \mathbb{C}[T^+]^k - 3, & k = d - 1, \\ \dim \mathbb{C}[T^+]^k - 12, & k = d. \end{cases}$$

Since $\dim \mathbb{C}[T^+]^d = \dim \mathbb{C}[T]^d + 9$, to prove Proposition 2.7 it suffices to produce 12 linearly independent relations in $\mathbb{C}[T^+]^d$ from Proposition 2.6. These 12 relations in degree d are produced in the following three ways:

- (a) Take $\ell = d + 1$ in Equation (2.7), we get the vanishing of a cohomology class in $M_{d,\chi} \times \check{\mathbb{P}}^2$. Pushing forward to $M_{d,\chi}$ yields a relation $R_{(a)}^n \in \mathbb{C}[T^+]^{d-1}$ for $n = 1, 2, 3$. Each of these three relations produces two relations in degree d , namely $c_2(0)R_{(a)}^n$ and $c_0(2)R_{(a)}^n$.
- (b) Take $\ell = d + 1$ in Equation (2.7), multiply by $\pi_R^*(\beta)$ and push forward to $M_{d,\chi}$. This produces three relations $R_{(b)}^n \in \mathbb{C}[T^+]^d$ for $n = 1, 2, 3$.
- (c) Take $\ell = d + 2$ in Equation (2.7) and push forward to $M_{d,\chi}$. This produces three relations $R_{(c)}^n \in \mathbb{C}[T^+]^d$ for $n = 1, 2, 3$.

We now proceed to argue that these 12 relations are linearly independent. It can be checked, cf. det 1 in the Mathematica file and [PS, Section 2.3], that under the assumption $\gcd(d, \chi) = 1$, the 3×3 matrix of coefficients of $c_d(0), c_{d-1}(1), c_{d-2}(2)$ in the three relations $R_{(a)}^1, R_{(a)}^2, R_{(a)}^3$ is nonsingular; in fact, it has determinant

$$\det 1 = \frac{(-1)^d (d - 2)^4 (d - 1)}{4} \chi (d - \chi) (d - 2\chi).$$

In particular, equations $R_{(a)}^n$ can be used to express $c_d(0), c_{d-1}(1), c_{d-2}(2)$ in terms of the tautological generators (2.4). Multiplying these equations with $c_2(0), c_0(2)$, we obtain six relations in $\mathbb{C}[T^+]^d$. These are still linearly independent since the 6×6 matrix given by the coefficients of the monomials

⁶See [Y2, Theorem 1.1], where the relevant Betti numbers $b_{2k}(\mathbb{P}^{2[n]})$ of the Hilbert scheme equal $\dim \mathbb{C}[T^+]^k$ by Göttsche's formula [Göt].

$$\text{Mon}_1 = \{c_d(0)c_2(0), c_d(0)c_0(2), c_{d-1}(1)c_2(0), c_{d-1}(1)c_0(2), c_{d-2}(2)c_2(0), c_{d-2}(2)c_0(2)\}$$

is invertible with determinant $\det 1^2$.

We consider now the six relations $R^n_{(b)}, R^n_{(c)}$, for $n = 1, 2, 3$ and the set of monomials

$$\text{Mon}_2 = \{c_{d+1}(0), c_d(1), c_{d-1}(2), c_{d-1}(0)c_3(0), c_{d-1}(0)c_2(1), c_{d-1}(0)c_1(2)\}.$$

The 6×6 matrix of coefficients of Mon_2 in these six relations is nonsingular. Indeed, a direct computation (cf. `det2` in the Mathematica file) gives the determinant

$$\det 2 = 4(d - 2)^6(d - 1)^3 d^4 \neq 0.$$

It follows that the six relations $R^n_{(b)}$ and $R^n_{(c)}$ with $n \in \{1, 2, 3\}$ are linearly independent.

Finally, we note that the monomials in Mon_2 do not appear in the relations of the form $c_2(0)R^n_{(a)}, c_0(2)R^n_{(a)}$, so the 12×12 matrix of coefficients of monomials $\text{Mon}_1 \cup \text{Mon}_2$ in the 12 relations obtained by (a), (b), (c) is also nonsingular. We conclude that these 12 relations are all linearly independent, completing the proof of Proposition 2.7.

To explicitly obtain the three relations in $\mathbb{C}[T]^d$, we proceed as follows. We use the relations $R^n_{(a)}$ to express $c_d(0), c_{d-1}(1), c_{d-2}(2)$ in terms of lower degree generators in the image of $\mathbb{C}[T]$; we then use $R^1_{(b)}, R^1_{(c)}, R^2_{(c)}$ to write $c_{d+1}(0), c_d(1), c_{d-1}(2)$ in terms of the tautological generators. Finally, plugging these into the other three relations

$$R^2_{(b)}, R^3_{(b)}, R^3_{(c)}$$

yields the three relations in $\mathbb{C}[T]^d$. By the nonvanishing of $\det 2$, the row-echelon form (with respect to the ordering $<$ of monomials) of this system produces three relations R_1, R_2, R_3 such that

$$R_i = c_{d-1}(0)c_{4-i}(i - 1) + \left(\text{sum of monomials } < c_{d-1}(0)c_1(2)\right) \in \mathbb{C}[T]^d. \tag{2.8}$$

A truncated version (see Remark 3.2) of these relations is implemented in the Mathematica file as `TruncRelations`.

3. Truncated relations and proof of the main result

In this section, we will prove the main result of this paper, Theorem 1.2. By the symmetry (a) in Section 1.1, we may assume without loss of generality that $0 < \chi, \chi' < d$. Note that when $d < 5$ there is nothing to prove, so we assume as well that $d \geq 5$. Suppose that there is an isomorphism of graded \mathbb{C} -algebras $\phi: A^*(M_{d,\chi}) \rightarrow A^*(M_{d,\chi'})$. We first remark that ϕ can be uniquely lifted to the free algebra $\mathbb{C}[T]$, that is, there exists a map $\tilde{\phi}$ fitting into the diagram

$$\begin{array}{ccc} \mathbb{C}[T] & \overset{\tilde{\phi}}{\dashrightarrow} & \mathbb{C}[T] \\ \downarrow & & \downarrow \\ A^*(M_{d,\chi}) & \xrightarrow{\phi} & A^*(M_{d,\chi'}) \end{array}.$$

Such unique lift exists since the cohomology is freely generated up to degree $d - 2$, where the tautological generators (2.4) lie. More precisely, the lift is defined by

$$\tilde{\phi}(c_k(j)) \in A^{k+j-1}(M_{d,\chi'}) \cong \mathbb{C}[T]^{k+j-1} \quad \text{for } k + j \leq d - 1.$$

By abuse of notation, we write ϕ also for the lifted graded ring isomorphism. We will show that unless $\chi \equiv \pm\chi' \pmod{d}$, there is no such lift respecting the three relations in degree d , that is, that sends the three-dimensional subspace

$$\ker \left(\mathbb{C}[T]^d \rightarrow A^d(M_{d,\chi}) \right) \subseteq \mathbb{C}[T]^d \text{ to } \ker \left(\mathbb{C}[T]^d \rightarrow A^d(M_{d,\chi'}) \right).$$

Remark 3.1. A graded ring endomorphism of $\mathbb{C}[T]$ is defined by an element of

$$\prod_{k=1}^{d-2} \text{Hom} (T^k, \mathbb{C}[T]^k).$$

For $d = 5$, the space above has dimension $2 \times 2 + 3 \times 6 + 3 \times 13 = 61$. On the other hand, a three-dimensional subspace of $\mathbb{C}[T]^d$ defines a point in the Grassmannian $\text{Gr}(3, \mathbb{C}[T]^d)$, which has dimension $3 \times (45 - 3) = 126$ for $d = 5$. Thus, we expect no graded ring endomorphism that takes a general point in $\text{Gr}(3, \mathbb{C}[T]^d)$ to another.

Let $R_1, R_2, R_3 \in \mathbb{C}[T]^d$ be the three relations in Equation (2.8); these span $\ker \left(\mathbb{C}[T]^d \rightarrow A^d(M_{d,\chi}) \right)$ and are such that the matrix $(R_1 \ R_2 \ R_3)^\top$ is in row echelon form with respect to the total ordering introduced in Section 2.3 on the basis of $\mathbb{C}[T]^d$. Similarly, we define R'_1, R'_2, R'_3 to be the relations in $\mathbb{C}[T]^d$ which come from $M_{d,\chi'}$. Since $\phi : \mathbb{C}[T]^d \xrightarrow{\sim} \mathbb{C}[T]^d$ preserves the three-dimensional kernels, it induces an invertible matrix

$$S = (s_{ij})_{0 \leq i, j \leq 3}$$

such that

$$\phi(R_i) = \sum_{j=1}^3 s_{ij} R'_j. \tag{3.1}$$

Step 1: truncating relations

We start by truncating the relations R_i by looking only at the terms which are obtained as a product of a generator of degree $d - 2$ with a generator of degree 2. In other words, we consider the projection of R_i to $T^2 \otimes T^{d-2} \subseteq \mathbb{C}[T]^d$ and regard it as a 3×3 matrix M_i by identifying

$$M_i \in T^2 \otimes T^{d-2} \cong \text{Hom}((T^{d-2})^\vee, T^2) \cong M_{3 \times 3}.$$

This is implemented in the Mathematica file as `TopRelations[d, chi]`. The identification with a 3×3 matrix uses the ordered basis for T^2, T^{d-2} given by the generators from Section 2.3.

More concretely,

$$M_i = \left([c_{3-s}(s)c_{d-1-t}(t)] R_i \right)_{0 \leq s, t \leq 2},$$

where $[c_{3-s}(s)c_{d-1-t}(t)]$ means reading off the corresponding coefficient with respect to the monomial basis. The matrices M_1, M_2, M_3 can be explicitly calculated as follows:

$$M_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & \frac{(d-2)(d-2\chi)\chi(d-\chi)}{8d^3} \\ \frac{d-4}{8d-16} & \frac{(d-2\chi)\chi(\chi-d)}{2(d-2)d^3} & \frac{(-6\chi-1)d^5+(6\chi^2+6\chi+3)d^4+18\chi^2d^3-(48\chi^3+48\chi^2)d^2+(24\chi^4+96\chi^3)d-48\chi^4}{32(d-2)d^4} \end{bmatrix},$$

$$M_2 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & \frac{(-6\chi-1)d^2+(6\chi^2+12\chi)d-12\chi^2}{8d^2} \\ 0 & \frac{24\chi^2-24\chi d-d^3+2d^2}{8(d-2)d^2} & \frac{\chi(d^2+8d-16)(2\chi-d)(d-\chi)}{8(d-2)d^3} \end{bmatrix}, \quad M_3 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ \frac{2}{d-2} & 0 & \frac{12\chi^2-12\chi d-d^2}{8(d-2)d} \end{bmatrix}.$$

Remark 3.2. To compute these matrices and also the extended matrices that will appear later in Section 3.3, we follow the strategy described in the previous section. Note that the truncation of the relations

$R_{(a)}^n, R_{(b)}^n, R_{(c)}^n$ can be computed since only finitely many terms of Equation (2.7) contribute. For instance, for $\ell = d + 1$ the only contributing terms are the ones corresponding to the partitions

$$(d + 1), (d, 1), (d - 1, 1, 1), (d - 1, 2), (d - 2, 2, 1), (d - 2, 3), (d - 2, 1, 1, 1).$$

Similarly, we define M'_i with χ replaced by χ' . Let A, B be the invertible linear maps

$$\begin{aligned} A: T^{d-2} &\hookrightarrow \mathbb{C}[T]^{d-2} \xrightarrow{\phi} \mathbb{C}[T]^{d-2} \rightarrow T^{d-2}, \\ B: T^2 &\hookrightarrow \mathbb{C}[T]^2 \xrightarrow{\phi} \mathbb{C}[T]^2 \rightarrow T^2. \end{aligned}$$

They are implemented in the Mathematica file as `AutA` and `AutB`, respectively. By considering the standard basis for T^2, T^{d-2} we identify again A, B with 3×3 matrices. More concretely, $A = (a_{st})_{0 \leq s, t \leq 2}, B = (b_{st})_{0 \leq s, t \leq 2}$, where

$$\begin{aligned} \phi(c_{d-1-s}(s)) &= \sum_{t=0}^2 a_{st} c_{d-1-t}(t) + (\text{lower terms in } \mathbb{C}[T]^{d-2}), \\ \phi(c_{3-s}(s)) &= \sum_{t=0}^2 b_{st} c_{3-t}(t) + (\text{lower terms in } \mathbb{C}[T]^2). \end{aligned}$$

Then the truncation of Equation (3.1) to $T^2 \otimes T^{d-2}$ can be written as an equality between 3×3 matrices

$$A^T M_i B = \sum_{j=1}^3 s_{ij} M'_j, \quad i = 1, 2, 3. \tag{3.2}$$

Step 2: finding s_{ij}

We begin with solving s_{ij} for which there are matrices A, B satisfying Equation (3.2). By scaling both A, B (hence also S), we may assume that $\det(A) = \det(B) = 1$. Then taking a linear combination of Equation (3.2) and applying the determinant it follows that

$$\det\left(\sum_{i=1}^3 x_i M_i\right) = \det\left(\sum_{i,j=1}^3 x_i s_{ij} M'_j\right), \tag{3.3}$$

as an equality between two homogeneous cubic polynomials in x_1, x_2, x_3 . We let $E \subseteq \mathbb{P}^2$ be the cubic curve defined by

$$E = \left\{ [x_1 : x_2 : x_3] : \det\left(\sum_{i=1}^3 x_i M_i\right) \right\} \subseteq \mathbb{P}_{x_1, x_2, x_3}^2.$$

Similarly, define $E' \subseteq \mathbb{P}^2$. The matrix S defines an automorphism of $\mathbb{P}_{x_1, x_2, x_3}^2$ sending E to E' .

Lemma 3.3. *The cubic curves E, E' are elliptic nodal curves with a single node at $[0 : 0 : 1]$. Hence, $s_{31} = s_{32} = 0$.*

Proof. We can directly compute the equation defining E . In the chart

$$\mathbb{C}_{x_1, x_2}^2 = \{[x_1 : x_2 : 1]\} \subseteq \mathbb{P}_{x_1, x_2, x_3}^2,$$

the cubic polynomial defining E has the form

$$-\frac{\chi(d-\chi)(d-2\chi)}{4(d-2)d^2}x_1x_2 + (\text{cubic terms in } x_1, x_2).$$

It follows that $[0 : 0 : 1]$ is indeed a node and the two branches of E at $[0 : 0 : 1]$ are tangent to the lines $x_1 = 0$ and $x_2 = 0$. A cubic with a node either has exactly one node or has two nodes and is the union of a line and a conic. If the latter were the case, the line would necessarily be $x_1 = 0$ or $x_2 = 0$, but this is not possible since $\det(M_2) \neq 0$ and $\det(M_1) \neq 0$.

Finally, we note that an automorphism of $\mathbb{P}^2_{x_1, x_2, x_3}$ sending E to E' must preserve the common node $[0 : 0 : 1]$, so it follows that $s_{31} = s_{32} = 0$. □

We now use Equation (3.3) to determine the remaining entries of S . Given a triple (u, v, w) of nonnegative integers such that $u + v + w = 3$, we get an equation among the entries of S by comparing the $x_1^u x_2^v x_3^w$ coefficient of both sides of Equation (3.3); this is implemented in the Mathematica file as `coeff[u, v, w]`.⁷ The triples $(0, 2, 1)$ and $(2, 0, 1)$ give, respectively,

$$s_{21}s_{22}s_{33} = 0 \quad \text{and} \quad s_{12}s_{11}s_{33} = 0.$$

On the other hand, the equation for $(1, 1, 1)$ implies that $s_{33} \neq 0$ and $s_{12}s_{21} + s_{11}s_{22} \neq 0$. It follows that solutions must have either:

- I. $s_{11} = s_{22} = 0$. We call the solutions of this form *Type I* solutions.
- II. $s_{12} = s_{21} = 0$. We call the solutions of this form *Type II* solutions.

From now on, we divide the analysis of Equations (3.2) and (3.3) according to the type of the solutions.

3.1. Type I

Suppose that S is a solution of *Type I* to Equation (3.3) so that we have already the vanishing of the entries $s_{31} = s_{32} = s_{11} = s_{22} = 0$. We now obtain the remaining entries. By looking at the $(0, 3, 0)$ and $(3, 0, 0)$ equations, we obtain s_{21}, s_{12} up to a choice of a cubic root of unity:

$$s_{21}^3 = -d^3 \frac{2\chi(d-\chi)(d-2\chi)(d^2+8d-16)}{\chi^2(d-\chi')^2(d-2\chi')^2(d-2)},$$

$$s_{12}^3 = -\frac{1}{d^3} \frac{\chi^2(d-\chi)^2(d-2\chi)^2(d-2)}{2\chi'(d-\chi')(d-2\chi')(d^2+8d-16)}.$$

Equation (1, 1, 1) then writes s_{33} in terms of s_{12} and s_{21} and we can conclude that $s_{33}^3 = 1$. By simultaneously scaling the matrices B and S , if necessary, by a cubic root of unity⁸ we may assume without loss of generality that $s_{33} = 1$. Finally, we obtain s_{23} using $(1, 2, 0)$ and s_{13} using $(2, 1, 0)$; note that these are uniquely determined once we choose which cubic root we take for s_{21} .

After having obtained S , the next step now is to solve Equation (3.2) for the entries of A and B . The equation is quadratic in the entries of A and B ; however, by rewriting it as

$$A^T M_i = \left(\sum_{j=1}^3 s_{ij} M'_j \right) B^{-1}, \quad i = 1, 2, 3, \tag{3.4}$$

it becomes a linear (homogeneous) system in the entries a_{st} of A and in the entries \tilde{b}_{st} (written as `bb[s, t]` in the Mathematica file) of B^{-1} . It turns out that the linear system only admits the trivial

⁷In the code, we use `chi1` and `chi2` instead of χ and χ' .

⁸Note that we already scaled B so that we could assume that $\det(B) = 1$; since scaling by a cubic root of unity does not affect the determinant, we are allowed to do so.

solution $a_{st} = \tilde{b}_{st} = 0$. Indeed, we can express all a_{st}, \tilde{b}_{st} as a multiple of a_{33} , and one additional constraint, denoted by `Diff[d, chi 1, chi 2][[1]][[1, 2]]` in the Mathematica file, imposes that

$$-\frac{2\chi(d-4)(d-\chi)(d-2\chi)a_{33}}{d((d-2)d^2+24d\chi'-24\chi'^2)} = 0,$$

whence $a_{33} = 0$. Note that the denominator is always positive, thus nonzero. This gives a contradiction since A and B are invertible matrices; thus, there are no solutions to Equation (3.1) with S of Type I.

3.2. Type II

Suppose that S is a solution of Type II to Equation (3.3). We already have the vanishing of the entries $s_{31} = s_{32} = s_{12} = s_{21} = 0$. The remaining entries are obtained exactly as in the Type I case. The equations (3, 0, 0) and (0, 3, 0) determine, respectively, s_{11} and s_{22} up to a choice of a cubic root:

$$s_{11}^3 = \left(\frac{\chi(d-\chi)(d-2\chi)}{\chi'(d-\chi')(d-2\chi')}\right)^2, \quad s_{22}^3 = \frac{\chi(d-\chi)(d-2\chi)}{\chi'(d-\chi')(d-2\chi')}.$$

Equation (1, 1, 1) then writes s_{33} in terms of s_{11} and s_{22} and we can conclude that $s_{33}^3 = 1$. As we did for Type I, we may assume without loss of generality that $s_{33} = 1$. This fixes $s_{11} = s_{22}^2$. Finally, we obtain s_{13} using (2, 1, 0) and s_{23} using (1, 2, 0); note that these are uniquely determined once we choose which cubic root we take for s_{22} .

After obtaining S , Equation (3.4) is again linear in the entries a_{st} of A and in the entries \tilde{b}_{st} of B^{-1} . The linear system can be solved explicitly, and it turns out that it has a one-dimensional space of solutions. By writing all the variables in terms of a_{11} , the normalization imposed on the determinant $\det(A) = 1$ gives

$$a_{11}^3 = \frac{\chi(d-\chi)(d-2\chi)}{\chi'(d-\chi')(d-2\chi')} = s_{22}^3.$$

By further scaling A by a cubic root of unity and scaling B by its inverse hence leaving S unchanged, we may assume without loss of generality that $a_{11} = s_{22}$. This completely solves all the matrices S, A, B once we choose which cubic root we take for s_{11} . Solutions are implemented in the Mathematica file as `solS[d, chi 1, chi 2]`, `solA[d, chi 1, chi 2]`, `solB[d, chi 1, chi 2]`, respectively.

Remark 3.4. When $\chi = \chi'$, the unique solution of Type II is given by $S = A = B = \text{Id}_{3 \times 3}$, as expected. When $\chi + \chi' = d$, the solution is given by simple sign matrices

$$A = B = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \quad S = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

In either case, these truncated solutions extend to a solution of Equation (3.1) since the moduli spaces $M_{d,\chi}$ and $M_{d,\chi'}$ are isomorphic. Indeed, the solution matrices are compatible with Proposition 2.1(c) which describes how the tautological generators are mapped under the two types of symmetries.

3.3. Type II – extended matrices

Recall that we have used only a part of the full Equation (3.1) truncated via the projection $\mathbb{C}[T]^d \rightarrow T^2 \otimes T^{d-2}$. This was sufficient to determine A, B and S .

Step 3: bigger truncation

We proceed by considering a bigger truncation via the projection

$$\mathbb{C}[T]^d \twoheadrightarrow \left(T^2 \oplus \text{Sym}^2(T^1)\right) \otimes T^{d-2}.$$

We can regard the truncated relations as a block $3 \times (3 \mid 3)$ matrix by identifying

$$[M_i \ N_i] \in \left(T^2 \oplus \text{Sym}^2(T^1)\right) \otimes T^{d-2} \cong \text{Hom}((T^{d-2})^\vee, T^2 \oplus \text{Sym}^2(T^1)) \cong M_{3 \times 6},$$

where M_i is as before and N_i is implemented in the Mathematica file as `ExtRelations[d, chi]`. Similarly, we define $[M'_i \ N'_i]$ with χ replaced by χ' . As before, ϕ induces a linear map

$$\begin{bmatrix} B & U \\ 0 & V \end{bmatrix} : T^2 \oplus \text{Sym}^2(T^1) = \mathbb{C}[T]^2 \xrightarrow{\phi} \mathbb{C}[T]^2 = T^2 \oplus \text{Sym}^2(T^1),$$

where $U : T^2 \rightarrow \text{Sym}^2(T^1)$ and $V : \text{Sym}^2(T^1) \rightarrow \text{Sym}^2(T^1)$. They are implemented in the Mathematica file as `AutU` and `AutV`, respectively. Note that the lower left block is zero because ϕ is a graded ring isomorphism. Then the truncation of Equation (3.1) to $\left(T^2 \oplus \text{Sym}^2(T^1)\right) \otimes T^{d-2}$ can be written as an equality between 3×6 matrices

$$A^T [M_i \ N_i] \begin{bmatrix} B & U \\ 0 & V \end{bmatrix} = \sum_{j=1}^3 s_{ij} [M'_j \ N'_j], \quad i = 1, 2, 3.$$

Since A, B and S satisfy Equation (3.2), this reduces to

$$A^T M_i U + A^T N_i V = \sum_{j=1}^3 s_{ij} N'_j, \quad i = 1, 2, 3, \tag{3.5}$$

which is a linear system in the entries u_{st} and v_{st} of U and V , respectively.

We show that this linear system, referred to as `ExtDiff[d, chi1, chi2]` in the Mathematica file, has a solution only if $\chi = \chi'$ or $\chi + \chi' = d$. By using certain parts of the linear system (3.5), as explained in the Mathematica file, we can express $u_{11}, u_{21}, u_{31}, v_{11}, v_{21}$ in terms of v_{31} . Once this is done, we further use two of the remaining linear system (3.5), referred to as `ExtDiff[d, chi1, chi2][[1]][[1, 1]]` and `ExtDiff[d, chi1, chi2][[1]][[2, 1]]`, which take the form

$$\begin{aligned} \text{AA}[d, \text{chi1}, \text{chi2}] v_{31} + \text{BB}[d, \text{chi1}, \text{chi2}] &= 0, \\ \text{CC}[d, \text{chi1}, \text{chi2}] v_{31} + \text{DD}[d, \text{chi1}, \text{chi2}] &= 0, \end{aligned}$$

following the notations from the code. Existence of the solution v_{31} implies an equation which is purely in terms of d, χ and χ' :

$$\begin{aligned} &\text{Constraint}[d, \text{chi1}, \text{chi2}] \\ &:= \text{AA}[d, \text{chi1}, \text{chi2}] \cdot \text{DD}[d, \text{chi1}, \text{chi2}] - \text{BB}[d, \text{chi1}, \text{chi2}] \cdot \text{CC}[d, \text{chi1}, \text{chi2}] = 0. \end{aligned}$$

This expression admits a factorization

$$\text{Constraint}[d, \text{chi1}, \text{chi2}] = \text{P1}[d, \text{chi1}] \cdot \text{P2}[d, \text{chi1}, \text{chi2}] \cdot (\text{other terms})$$

with (other terms) being clearly nonzero; see the Mathematica file. Note that the roles of χ and χ' are symmetric; we conclude that

$$P_1(d, \chi) \cdot P_2(d, \chi, \chi') = 0 \quad \text{and} \quad P_1(d, \chi') \cdot P_2(d, \chi', \chi) = 0. \quad (3.6)$$

We are left to prove that Equation (3.6) implies $\chi = \chi'$ or $\chi + \chi' = d$. Suppose for the contradiction that $\chi \neq \chi'$ and $\chi + \chi' \neq d$.

We first show that $P_2(d, \chi, \chi')$ and $P_2(d, \chi', \chi)$ are nonzero. From the formula in the file, it is straightforward to check that $P_2(d, \chi, \chi') = 0$ if and only if $P_2(d, \chi', \chi) = 0$, if and only if

$$d^2(\chi - \chi')(\chi + \chi' - d) \left(216\chi^4\chi'^4 - 432\chi^3\chi'^3(\chi + \chi')d + d^2 \cdot f(d, \chi, \chi') \right) = 0 \quad (3.7)$$

for some explicit integral polynomial $f(d, \chi, \chi')$. Since we assumed that $\chi \neq \chi'$ and $\chi + \chi' \neq d$, the last term must vanish. On the other hand, $\gcd(d, \chi) = \gcd(d, \chi') = 1$ implies that d must divide 216. This further implies that d^2 divides 216 by looking at the linear term in d , so we are left with $d = 1, 2, 3, 6$. But there are no nontrivial pairs χ and χ' for such d .

Therefore, we may assume $P_1(d, \chi) = P_1(d, \chi') = 0$. We can check that $P_1(d, x)$ is a degree four polynomial in x , with the symmetry

$$P_1(d, x) = P_1(d, d - x).$$

This implies that $P_1(d, x)$ is a degree four polynomial with four distinct roots

$$x = \chi, \chi', d - \chi, d - \chi',$$

all within the interval $[1, d - 1]$. On the other hand, explicit computation shows that

$$P_1(d, 0) > 0, \quad P_1(d, 1) < 0$$

as long as $d \geq 5$. This forces $P_1(d, x)$ to have an additional root in the interval $(0, 1)$, which contradicts that it is a degree four polynomial. It follows that Equation (3.6) implies $\chi = \chi'$ or $\chi + \chi' = d$, hence completing the proof.

Acknowledgements. We thank Yakov Kononov for his help on coding and for numerous helpful conversations. The third author wishes to thank Rahul Pandharipande for inviting him to visit ETH Zürich, which eventually resulted in this collaboration, and thanks to Junliang Shen, his academic advisor, for proposing this interesting problem in the first place.

Funding statement. The first author is supported by the grant SNF-200020-182181. The second author is supported by ERC-2017-AdG-786580-MACI. The project received funding from the European Research Council (ERC) under the European Union Horizon 2020 research and innovation programme (grant agreement 786580).

Competing interest. The authors have no competing interest to declare.

Data availability statement. The code used in this article (together with a printed file of the output) has been uploaded to the third author's website <https://github.com/Weite-Pi/weitepi.github.io> under the name `cohomological ring chi-dependence`.

Ethical standards. The research meets all ethical guidelines, including adherence to the legal requirements of the study country.

References

- [BM] A. Bayer and E. Macrì, 'MMP for moduli of sheaves on K3s via wall-crossing: nef and movable cones, Lagrangian fibrations', *Invent. Math.* **198**(3) (2014), 505–590.
- [BNR] A. Beauville, M.S. Narasimhan and S. Ramanan, 'Spectral curves and the generalized theta divisor', *J. Reine Angew. Math.* **398** (1989) 169–179.
- [Bea] A. Beauville, *Sur la cohomologie de certains espaces de modules de fibrés vectoriels*, Geometry and analysis (Bombay, 1992) (Tata Inst. Fund. Res. , Bombay, 1995), 37–40.

- [BLM] A. Bojko, W. Lim and M. Moreira, ‘Virasoro constraints on moduli of sheaves and vertex algebras’, *Invent. Math.* (2024).
- [Bou] P. Bousseau, ‘Scattering diagrams, stability conditions, and coherent sheaves on \mathbb{P}^2 ’, *J. Algebraic Geom.* **31**(4) (2022), 593–686.
- [dCHM] M. A. de Cataldo, T. Hausel and L. Migliorini, ‘Topology of Hitchin systems and Hodge theory of character varieties: The case A_1 ’, *Ann. of Math. (2)* **175**(3) (2012), 1329–1407.
- [dCMSZ] M. A. de Cataldo, D. Maulik, J. Shen and S. Zhang, ‘Cohomology of the moduli of Higgs bundles via positive characteristic’, *J. Eur. Math. Soc.* (2023), **1–21**.
- [dCM] M. A. de Cataldo and L. Migliorini, ‘The Hodge theory of algebraic maps’, *Ann. Sci. École Norm. Sup. (4)* **38**(5) (2005), 693–750.
- [GV] R. Gopakumar and C. Vafa, ‘M-Theory and topological strings–II’. (1998), [arXiv:hep-th/9812127](https://arxiv.org/abs/hep-th/9812127).
- [Göt] L. Göttsche, ‘The Betti numbers of the Hilbert scheme of points on a smooth projective surface’, *Math. Ann.* **286** (1990), 193–207.
- [HMMS] T. Hausel, A. Mellit, A. Minets and O. Schiffmann. (2022), ‘ $P = W$ via H_2 ’, [arXiv: 2209.05429](https://arxiv.org/abs/2209.05429).
- [HST] S. Hosono, M.-H. Saito and A. Takahashi, ‘Relative Lefschetz action and BPS state counting’, *Internat. Math. Res. Notices* (**15**) (2001), 783–816.
- [Huy] D. Huybrechts, ‘Compact hyperkähler manifolds: Basic results’, *Invent. Math.* **135**(1) (1999), 63–113.
- [HL] D. Huybrechts and M. Lehn, *The Geometry of Moduli Spaces of Sheaves*, Aspects of Mathematics, E31 (Friedr. Vieweg and Sohn, Braunschweig, 1997), xiv+269.
- [KL] Y.-H. Kiem and J. Li, ‘Categorification of Donaldson–Thomas invariants via perverse sheaves’. (2012), [arXiv: 1212.6444v5](https://arxiv.org/abs/1212.6444v5).
- [KPS] Y. Kononov, W. Pi and J. Shen, ‘Perverse filtrations, Chern filtrations, and refined BPS invariants for local \mathbb{P}^2 ’, *Adv. Math.* **433** (2023), Paper No. 109294, 29 pp.
- [LeP] J. Le Potier, ‘Faisceaux semi-stables de dimension 1 sur le plan projectif’, *Rev. Roumaine Math. Pures Appl.* **38**(7–8): 635–678, 1993.
- [Mai] M. Maican, ‘A duality result for moduli spaces of semistable sheaves supported on projective curves’, *Rend. Semin. Mat. Univ. Padova* **123** (2010), 55–68.
- [Mar] E. Markman, ‘Integral generators for the cohomology ring of moduli spaces of sheaves over Poisson surfaces’, *Adv. Math.* **208** (2007) 622–646.
- [M] Wolfram Research, Inc., Mathematica, Version 13.2, Champaign, IL (2022).
- [MS1] D. Maulik and J. Shen, ‘Cohomological χ -independence for moduli of one-dimensional sheaves and moduli of Higgs bundles’, *Geom. Topol.* **27**(4) (2023), 1539–1586.
- [MS2] D. Maulik and J. Shen, ‘The $P = W$ conjecture for GL_n ’. (2022), [arXiv: 2209.02568](https://arxiv.org/abs/2209.02568).
- [MT] D. Maulik and Y. Toda, ‘Gopakumar–Vafa invariants via vanishing cycles’, *Invent. Math.* **213**(3) (2018), 1017–1097.
- [PT] R. Pandharipande and R. P. Thomas, ‘13/2 ways of counting curves’, in *Moduli Spaces*, London Math. Soc. Lecture Note Ser., 411 (Cambridge Univ. Press, Cambridge, 2014), 282–333.
- [PS] W. Pi and J. Shen, ‘Generators for the cohomology ring of the moduli of one-dimensional sheaves on \mathbb{P}^2 ’, *Algebr. Geom.* **10**(4) (2023), 504–520.
- [Tod] Y. Toda, ‘Gopakumar–Vafa invariants and wall-crossing’, *J. Differential Geom.* **123**(1) (2023), 141–193.
- [Woo] M. Woolf, ‘Nef and effective cones on the moduli space of torsion sheaves on the projective plane’, Preprint, 2013, [arXiv: 1305.1465](https://arxiv.org/abs/1305.1465).
- [Y1] Y. Yuan, ‘Sheaves on non-reduced curves in a projective surface’, *Sci. China Math.* **66**(2) (2023), 237–250.
- [Y2] Y. Yuan, ‘Some Betti numbers of the moduli of 1-dimensional sheaves on \mathbb{P}^2 ’, *Forum Math.* **36**(1) (2024), 215–244.