

CORRIGENDUM

Three-dimensional turbulence without vortex stretching – CORRIGENDUM

Wouter Bos and Tong Wu

doi:10.1017/jfm.2021.194, Published by Cambridge University Press, 29 March 2021

In recent work (Bos 2021) we focused on an incompressible turbulent flow governed by a modified version of the Navier–Stokes equations. The essential difference with respect to the full Navier–Stokes equations is that the curl of the modified version does not contain the vortex-stretching term and is written

$$\frac{\partial \boldsymbol{\omega}}{\partial t} + \boldsymbol{u} \cdot \nabla \boldsymbol{\omega} = -\nabla \mathcal{P}_{\omega} + \nu \Delta \boldsymbol{\omega} + \nabla \times \boldsymbol{f}, \tag{0.1}$$

with $\omega = \nabla \times u$ the vorticity, u the velocity, v the kinematic viscosity and f a solenoidal force term. The first term on the right-hand side, $\nabla \mathcal{P}_{\omega}$, represents a pressure gradient which ensures the vorticity field remains divergence free, $\nabla \cdot \omega = 0$. Its value is determined by the Poisson equation:

$$\Delta \mathcal{P}_{\omega} = -\nabla \cdot (\boldsymbol{u} \cdot \nabla \boldsymbol{\omega}). \tag{0.2}$$

In the original paper, this pressure term was omitted in the vorticity equation, and incompressibility was only imposed on the level of the velocity equation. The thereby defined set of equations was not consistent.

Denoting $\mathcal{F}[\hat{x}] = \hat{x}$ the Fourier transform and k the wave vector, we defined the velocity equation (ignoring forcing and viscous stress) as

$$\frac{\partial \hat{\boldsymbol{u}}}{\partial t} + ik^{-2}\boldsymbol{k} \times \mathcal{F}[\boldsymbol{u} \cdot \nabla \boldsymbol{\omega}] = 0. \tag{0.3}$$

It is this equation which was solved numerically in our previous investigations. Taking the curl of this equation, we find that

$$\frac{\partial \hat{\boldsymbol{\omega}}}{\partial t} - k^{-2} \boldsymbol{k} \times (\boldsymbol{k} \times \mathcal{F}[\boldsymbol{u} \cdot \nabla \boldsymbol{\omega}]) = 0, \tag{0.4}$$

or, working out the double cross-product and applying the inverse Fourier transform, $\mathcal{F}[x]^{-1}$,

$$\frac{\partial \boldsymbol{\omega}}{\partial t} + \boldsymbol{u} \cdot \nabla \boldsymbol{\omega} = \mathcal{F}^{-1} \left[\frac{\boldsymbol{k}}{k^2} \boldsymbol{k} \cdot \mathcal{F} [\boldsymbol{u} \cdot \nabla \boldsymbol{\omega}] \right]. \tag{0.5}$$

The right-hand side equals the pressure term $-\nabla \mathcal{P}_{\omega}$ and was therefore present in the numerically investigated system, but was missing in several equations in our paper.

The omission of the pressure term should be corrected in (2.3), (2.5) and (A1).

W. Bos and T. Wu

This mistake does not therefore change the numerical results, which were all obtained from the modified Navier–Stokes equation for u (equation (0.3)), which was correctly formulated in the articles. Furthermore, the presence of this term does not influence the conservation of enstrophy and helicity. The only implication is that the enstrophy is conserved as a whole but the different components $\langle \omega_x^2 \rangle$, $\langle \omega_y^2 \rangle$, $\langle \omega_z^2 \rangle$ are not conserved individually by the nonlinear interaction. The lesson here is that one cannot throw away half the nonlinear term in the vorticity equation without adding an additional mechanism to maintain incompressibility.

Acknowledgements. We thank our colleagues who pointed us to this omission of the pressure term.

REFERENCE

Bos, W.J.T. 2021 Three-dimensional turbulence without vortex stretching. J. Fluid Mech. 915.