

## Research Article

**Cite this article:** Fuller C, Mueller T, Steckel LE (2026) Managing herbicide-resistant Palmer amaranth (*Amaranthus palmeri*) from West Tennessee with sequential applications of dicamba with and without glufosinate. *Weed Technol.* **40**(e15), 1–6. doi: [10.1017/wet.2026.10103](https://doi.org/10.1017/wet.2026.10103)

Received: 24 February 2026

Revised: 15 April 2026

Accepted: 18 April 2026

**Associate Editor:**

Sandeep Singh Rana, Bayer Crop Science

**Nomenclature:**

Dicamba; glufosinate; Palmer amaranth; *Amaranthus palmeri* S. Wats AMAPA

**Keywords:**

Split-boom; tank-mixture

**Corresponding author:**

Lawrence E. Steckel; Email: [lsteckel@utk.edu](mailto:lsteckel@utk.edu)

# Managing herbicide-resistant Palmer amaranth (*Amaranthus palmeri*) from West Tennessee with sequential applications of dicamba with and without glufosinate

Colton Fuller<sup>1</sup> , Thomas Mueller<sup>1</sup>  and Lawrence E. Steckel<sup>2</sup> 

<sup>1</sup>University of Tennessee System: The University of Tennessee System, USA and <sup>2</sup>Department of Plant Sciences, University of Tennessee, Jackson, USA

**Abstract**

Palmer amaranth has become one of the most problematic weeds in row crop production in states in the mid-South. With increasing resistance to multiple herbicides, new control methods are needed. Field experiments were conducted in 2024 and 2025 at four locations in Tennessee to evaluate the efficacy of dicamba and glufosinate applied alone, tank-mixed, or sequentially using a split-boom approach for controlling auxin herbicide-resistant Palmer amaranth populations. Control varied widely among populations, reflecting differences in dicamba resistance levels. Populations with low dicamba resistance treated with glufosinate showed 99% control following sequential applications, with no differences between tank-mix and split-boom methods. In contrast, populations with high dicamba resistance demonstrated reduced control across all treatments; however, the split-boom application improved initial control at one highly resistant site compared with the tank-mixed application. Sequential applications improved control at all locations compared with the initial application. Increasing the spray carrier volume from 140 to 280 L ha<sup>-1</sup> did not improve control of Palmer amaranth. These results indicate that sequential applications of dicamba plus glufosinate are required to maximize control of auxin-resistant Palmer amaranth, but efficacy is population dependent, underscoring the need for crop scouting and integrated weed management strategies to mitigate resistance development to sustain herbicide effectiveness.

**Introduction**

In 2024, the National Cotton Council (2024) reported that 11.5 million acres of cotton (*Gossypium hirsutum* L.) were planted in the United States. Globally, the total potential loss due to pests varied in crops, but was reported to be above 80% in cotton production. Weeds were observed to have the most significant effect on cotton yield loss at 34% (Oerke 2006). Palmer amaranth is one of the most widespread, troublesome, and economically damaging weeds in agronomic fields throughout the southeastern United States. Palmer amaranth's dominance and difficulty to control are driven by its rapid growth, high seed production, genetic variability, ability to withstand undesirable environmental conditions, and the remarkable capacity to develop herbicide resistance. Palmer amaranth is recognized as the most economically significant herbicide-resistant weed in U.S. cotton production (Heap 2025; Vulchi et al. 2022).

Auxinic herbicides are synthetic auxins that have been used extensively for more than 60 yr to control broadleaf weeds (Grossmann 2000). Synthetic auxins mimic natural auxins that are found within the plant, and the effects of synthetic herbicides can be described as an auxin overdose (Gleason et al. 2011; Grossmann 2000). While weed resistance to auxinic herbicides has been confirmed in 44 species globally, the mechanism for resistance is still under study (Busi et al. 2018; Heap 2025). Busi et al. (2018) supports the theory that the mechanism for resistance to synthetic auxin herbicides can include target site and non-target site mechanisms.

In 2015, multi-gene transgenic herbicide-resistant cotton and soybean [*Glycine max* (L.) Merr.] varieties such as XtendFlex (Bayer Crop Science, St. Louis, MO) became commercially available to growers. This allowed postemergence applications of select dicamba formulations, glufosinate, and glyphosate (Bland 2024; Delong 2020). With roughly 11.4 million cotton acres planted, the U.S. Environmental Protection Agency has estimated that 75% of those acres are planted with XtendFlex (US EPA 2024; National Cotton Council 2024). The broad adoption of this technology has led to increased use of dicamba and glufosinate; likewise, increased use leads to greater selection, resulting in a higher probability of resistance evolving in Palmer amaranth. With the documentation of dicamba-resistant Palmer amaranth in Tennessee in 2020 by Foster and Steckel (2022), the need for alternative control measures is evident.

© The Author(s), 2026. Published by Cambridge University Press on behalf of Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



Use of glufosinate has increased following the introduction of XtendFlex soybean and cotton systems. A study conducted at Kansas State University evaluated herbicides applied pre-emergence, early postemergence and late postemergence for managing glyphosate-resistant Palmer amaranth (Shyam et al. 2021). Treatments that included an effective preemergence application followed by glufosinate alone or glufosinate + 2,4-D showed greater than 90% control. These results demonstrate that glufosinate remains a dependable option for managing glyphosate-resistant and multiple herbicide-resistant Palmer amaranth (Shyam et al. 2021).

The initial dicamba herbicide formulations labeled for use on XtendFlex crops did not allow glufosinate as a labeled tank-mix partner. Split-boom applications are considered an on-label application with the original over-the-top dicamba labels. This could offer another way to combat herbicide-resistant Palmer amaranth where single mode of action applications are failing (US EPA 2022).

Previous research has demonstrated that certain dicamba-based tank mixtures can result in antagonistic interactions on broadleaf weed species. Specifically, dicamba tank mixed with glyphosate and glufosinate and applied at multiple rates produced predominantly antagonistic effects on glyphosate-resistant Palmer amaranth (Constine 2021; Priess et al. 2022). Antagonism was quantified using the Colby equation by comparing expected versus observed growth reduction. Similar antagonistic responses were observed in additional broadleaf species, including glyphosate-susceptible waterhemp (*Amaranthus tuberculatus* Sauer), GR waterhemp, and glyphosate-resistant horseweed (*Erigeron canadensis* L.), indicating that the three-way mixture consistently reduced herbicide efficacy across species (Constine 2021).

The goal of this research was to evaluate whether Palmer amaranth control could be improved with a tank mixture compared with a split application of dicamba and glufosinate. Another goal was to assess whether sequential applications would mitigate resistance-related failure. An additional objective was to determine whether response differs by a known auxin-resistance level, and finally, to evaluate whether carrier volume affected Palmer amaranth control.

## Materials and Methods

In 2024 and 2025, research was conducted at four locations in western Tennessee to evaluate herbicide-resistant Palmer amaranth control. Each population was numbered and given a corresponding site name, and global positioning system coordinates and county information was recorded (Table 1). The split-boom vs. tank-mixture study was an arrangement of treatments within a randomized complete block design with four replications at the Madison location and three replications in Lauderdale 1, Lauderdale 2, and Gibson. The study examined the control of Palmer amaranth with single-ingredient, tank-mixed, and split-applied applications. The six treatments consisted of 1) dicamba, 2) glufosinate, 3) dicamba + glufosinate tank mixed, 4) dicamba and glufosinate split-boom, 5) dicamba + glufosinate tank mixed followed by a second application of that same treatment applied 21 d later, and 6) dicamba and glufosinate split-boom followed by a second application of that same treatment applied 21 d later (Table 2). All treatments, herbicide names, and rates are listed in Table 2. The treated plot size was 2 m wide by 9 m long. Split-applied applications were conducted with two applicators, both of whom used a 1.52-m boom. The first applicator began spraying a

**Table 1.** Palmer amaranth control in 2023, dicamba relative resistance factor in 2022, and Palmer amaranth control at trial site locations.

| Location                     | Coordinates                 | Control <sup>a</sup><br>% | Relative resistance factor 2022 <sup>b</sup> |
|------------------------------|-----------------------------|---------------------------|----------------------------------------------|
| Madison                      | 35.63205°N,<br>88.85838°W   | 82                        | 1.03                                         |
| Gibson                       | 35.781542°N,<br>88.851567°W | 58                        | 2.50                                         |
| Lauderdale 1                 | 35.714767°N,<br>89.919529°W | 83                        | 0.96                                         |
| Lauderdale 2                 | 35.725660°N,<br>89.873030°W | 48                        | 14.3                                         |
| Least significant difference |                             | 14                        |                                              |

<sup>a</sup>Control was evaluated after applications of dicamba (0.56 kg ha<sup>-1</sup>).

<sup>b</sup>The relative resistance factor represents herbicide resistance levels of Palmer amaranth populations (Foster and Steckel, 2022).

**Table 2.** Herbicides used in the field experiments.<sup>a,b</sup>

| Treatment No. | Herbicide treatment                                   | Rate                                                      |
|---------------|-------------------------------------------------------|-----------------------------------------------------------|
| 1             | Dicamba                                               | 0.56 kg ae ha <sup>-1</sup>                               |
| 2             | Glufosinate                                           | 0.66 kg ai ha <sup>-1</sup>                               |
| 3             | Dicamba + glufosinate TM                              | 0.56 kg ae ha <sup>-1</sup> + 0.66 kg ai ha <sup>-1</sup> |
| 4             | Dicamba and glufosinate SB                            | 0.56 kg ae ha <sup>-1</sup> + 0.66 kg ai ha <sup>-1</sup> |
| 5             | Dicamba + glufosinate fb dicamba + glufosinate TM     | 0.56 kg ae ha <sup>-1</sup> + 0.66 kg ai ha <sup>-1</sup> |
| 6             | Dicamba and glufosinate fb dicamba and glufosinate SB | 0.56 kg ae ha <sup>-1</sup> + 0.66 kg ai ha <sup>-1</sup> |

<sup>a</sup>Abbreviations: fb, followed by; SB, split-boom; TM, tank mixture.

<sup>b</sup>The trade name for dicamba is Engenia; the trade name for glufosinate is Liberty. Both herbicides are manufactured by BASF Corporation, Research Triangle Park, NC.

herbicide with a single mode of action, and the second applicator also had a bottle with a single mode-of-action herbicide. As the first applicator was 0.5 m ahead, the second applicator began spraying, following the first applicator. Each herbicide was used at the 1× rate according to the label directions (dicamba, 0.56 kg ae ha<sup>-1</sup>; glufosinate, 0.66 kg ai ha<sup>-1</sup>) (Table 2). All herbicides were applied using a CO<sub>2</sub>-pressurized backpack sprayer equipped with AIXR 11002 nozzles (TeeJet Technologies, Springfield, IL) calibrated to deliver 140 L ha<sup>-1</sup> at 4.8 kph at a pressure of 220 kPa.

## Locations and Site History

The first location was the West Tennessee Research and Education Center in Madison County. Cotton and soybean had been grown in rotation at the site from 2021 to 2024. In 2024, poor weed control (<40%) occurred with one application of 2,4-D (1.07 kg ae ha<sup>-1</sup>; data not shown) when examined 14 d after application (DAA).

The history of the Gibson field consisted of cotton grown continuously, most recently with 2,4-D-resistant cotton (Enlist Corteva AgriSciences, Indianapolis, IN) in 2020, and then dicamba-resistant cotton (Xtend Bayer Crop Science, St. Louis, MO) from 2021 to 2024. From 2021 to 2024, the grower observed escapes of Palmer amaranth after multiple applications of dicamba (0.56 kg ae ha<sup>-1</sup>).

The third (Lauderdale 1) and fourth (Lauderdale 2) locations differed by less than 2.8 km (Table 1). The history of both locations

consisted of herbicide-resistant Palmer amaranth biotypes that had been present for more than a decade. The surrounding fields had been in a cotton-soybean rotation; the locations used in the Lauderdale experiments were fallow ground.

The Madison location was equipped with lateral overhead irrigation, while all other locations received rain only. Each site consisted of a native glyphosate-resistant Palmer amaranth population that was known for escaping dicamba applications (Foster and Steckel 2022). The Madison and Gibson trials were conducted with Enlist cotton in 2024. Lauderdale 1 and 2 trials were conducted on fallow ground. In 2025, no crops were planted in any of the tested locations.

In 2024, Enlist cotton was planted on May 4 and May 10 at the Gibson and Madison locations, respectively. A burndown application of paraquat (Gramoxone, Syngenta Crop Protection Greensboro, NC) ( $0.86 \text{ kg ai ha}^{-1}$ ) plus a nonionic surfactant ( $250 \text{ mL L}^{-1}$ ) (Preference; Winfield United, Arden Hills, MN) was used to ensure the trials were free of weeds before planting. No residual herbicides were applied to the treated plots in 2024. The Lauderdale County locations were conducted on fallow ground that received no burndown or residual herbicide in 2024. In 2025, pyroxasulfone (Zidua; BASF, Research Triangle Park, NC) ( $0.12 \text{ kg ai ha}^{-1}$ ) was applied at the first application timing across all the treatments to prevent later weed flushes. The treatments were identical to those described for the other locations. Applications were initiated at all locations when Palmer amaranth reached a height of 10 cm.

Palmer amaranth control was visually assessed at 7, 14, and 21 DAA on a scale of 0% to 100%, where 0% = no injury and 100% = plant death. Following the second application, Palmer amaranth was rated again following the same rating scale. Counts were taken using one  $0.145\text{-m}^2$  quadrat per treatment, following the initial 14-d and 21-d evaluations and the subsequent 14-d and 21-d periods. Herbicides were applied a second time to evaluate sequential treatments for the tank-mix and split-boom methods. Single-ingredient treatments were not included in the second application. Ratings for the initial application were continued through 35 DAA for treatments 1 through 4 (Table 2). Treatments 5 and 6 received the sequential application and were evaluated 14 d after the second application (Table 2).

Data were analyzed using the GLIMMIX procedure in SAS software (v. 9.4; SAS Institute, Cary, NC) to analyze variance. Normal distribution of data was observed using the UNIVARIATE procedure, with both the stem-leaf and box plots showing a central peak with tails of equal length. Therefore, no data transformation was necessary to improve normality. Individual treatment differences were determined using the Fisher protected LSD test at the  $\alpha = 0.05$  level. Mean separation was performed only when the overall *F*-test was significant.

To examine whether the change in control was due to the application method as opposed to simply increased carrier volume, we conducted a study at three locations (Madison, Lauderdale 1, Gibson) in 2025 to compare a tank mix of dicamba + glufosinate applied at  $140 \text{ L ha}^{-1}$  with the same two herbicides applied at  $280 \text{ L ha}^{-1}$ . Treatments consisted of one pass and were applied as previously described for  $140 \text{ L ha}^{-1}$  and by using a  $\text{CO}_2$ -pressurized backpack sprayer equipped with TeeJet AIXR 11004 nozzles calibrated to deliver  $280 \text{ L ha}^{-1}$  at 4.8 kph using 207 kPa.

Weed control and plant density data were collected as previously described. A single degree-of-freedom contrast statement in SAS software was conducted to compare results.

## Results and Discussion

### Data Collection

Data were initially subjected to an analysis across years and populations, with each location defined as a population. In the model, locations were different but years did not differ so data were pooled across years. A significant year  $\times$  population  $\times$  treatment interaction ( $P < 0.0001$ ) was detected. Results from the mixed-model analysis indicated that location significantly influenced Palmer amaranth control, demonstrating that treatment performance varied across site conditions and population characteristics. Therefore, treatment effects were analyzed separately by population. Within each population, year was treated as a random effect and treatment was considered fixed (Carmer et al. 1989).

Previous research by Foster and Steckel (2022) supports this approach, as populations of Palmer amaranth in western Tennessee exhibited a wide range of dicamba resistance levels, with relative resistance factors of 1.03 and 0.96 for the Madison and Lauderdale 1 populations, respectively, indicating dicamba susceptibility. However, Lauderdale 2 and Gibson populations showed higher resistance, with relative resistance factor values of 14.3 and 2.5, respectively (Table 1). These documented differences in dicamba sensitivity, combined with the significant location effects observed in this study, justified analyzing each location independently to accurately capture treatment responses across resistant and susceptible Palmer amaranth populations.

### Madison

At the Madison location, when evaluated at 14 DAA, 40% of Palmer amaranth was controlled % when dicamba was applied alone at  $0.56 \text{ kg ae ha}^{-1}$  (Table 3). Glufosinate applied alone controlled Palmer amaranth by 70%, although both the tank-mix (88%) and the split-boom (91%) methods provided greater control.

Plant count data supported the visual evaluations. Plots treated with glufosinate alone had fewer plants than those treated with dicamba alone. The combination treatments (tank-mix and split-boom) contained fewer plants than plots treated with a single product (Table 3). These results demonstrate that using both herbicides, regardless of application method, improved Palmer amaranth control relative to dicamba or glufosinate applied alone when evaluated at 14 DAA.

For the treatments that did not receive a sequential application, control declined by 35 DAA (Table 4). The split-boom treatment provided similar control to that of the tank mix, but better than dicamba or glufosinate applied alone. When herbicides were applied sequentially, there was no difference between methods with  $\geq 97\%$  control. These results indicate that a sequential application is needed to achieve high levels of Palmer amaranth control at this site, regardless of whether the herbicides are applied as a tank mix or in a split-boom configuration.

### Lauderdale 1

At the Lauderdale 1 location, Palmer amaranth was controlled by 74% with dicamba, which was higher than at any other site at 14 DAA (Table 3). Glufosinate applied alone provided excellent control of Palmer amaranth, averaging 98%, and control was similar with both the tank-mix and split-boom application methods (99%). Although dicamba applied alone provided 74% control, levels were still lower than all treatments that contained glufosinate. These results suggest that using glufosinate is

**Table 3.** Palmer amaranth control and density observed 14 d after initial applications of dicamba and glufosinate.<sup>a,b</sup>

| Herbicide                            | Palmer amaranth control |            |              |            |        |            |              |            |
|--------------------------------------|-------------------------|------------|--------------|------------|--------|------------|--------------|------------|
|                                      | Madison                 |            | Lauderdale 1 |            | Gibson |            | Lauderdale 2 |            |
|                                      | %                       | No. plants | %            | No. plants | %      | No. plants | %            | No. plants |
| Dicamba                              | 40 c                    | 6 C        | 74 b         | 5 B        | 31 c   | 10 A       | 28 d         | 13 D       |
| Glufosinate                          | 70 b                    | 3 B        | 98 a         | 0 A        | 43 b   | 10 A       | 42 c         | 5 C        |
| Dicamba + glufosinate (tank mix)     | 91 a                    | 1 A        | 99 a         | 0 A        | 56 ab  | 9 A        | 61 b         | 3 B        |
| Dicamba and glufosinate (split-boom) | 88 a                    | 1 A        | 99 a         | 0 A        | 52 a   | 6 A        | 82 a         | 1 A        |
| Nontreated                           | 0 d                     | 8 C        | 0 c          | 14 C       | 0 d    | 9 A        | 0 e          | 10 D       |
| Pr > F                               | <.0001                  | <.0001     | <.0001       | <.0001     | <.0001 | 0.4384     | <.0001       | <.0001     |

<sup>a</sup>Palmer amaranth density was determined by manually counting the number of plants per 0.145 m<sup>2</sup>.

<sup>b</sup>Lowercase letters indicate mean separation for percent Palmer amaranth control; uppercase letters indicate mean separation for Palmer amaranth plant density.

**Table 4.** Palmer amaranth control and density observed 35 d after initial applications (14 d after second applications) of dicamba and glufosinate.<sup>a-d</sup>

|                                      | Palmer amaranth control |            |              |            |        |            |              |            |
|--------------------------------------|-------------------------|------------|--------------|------------|--------|------------|--------------|------------|
|                                      | Madison                 |            | Lauderdale 1 |            | Gibson |            | Lauderdale 2 |            |
|                                      | %                       | No. plants | %            | No. plants | %      | No. plants | %            | No. plants |
| Initial herbicide applications       |                         |            |              |            |        |            |              |            |
| Dicamba (35 DAA)                     | 29 d                    | –          | 73 c         | –          | 26 bc  | –          | 26 d         | –          |
| Glufosinate (35 DAA)                 | 49 c                    | –          | 83 bc        | –          | 17 c   | –          | 42 dc        | –          |
| Dicamba + glufosinate (TM, 35 DAA)   | 61 bc                   | –          | 88 ab        | –          | 35 b   | –          | 52 bc        | –          |
| Dicamba and glufosinate (SB, 35 DAA) | 63 b                    | –          | 85 ac        | –          | 35 b   | –          | 58 abc       | –          |
| Sequential herbicide applications    |                         |            |              |            |        |            |              |            |
| Dicamba + glufosinate (TM, 14 DAA)   | 97 a                    | 0 A        | 99 a         | 0 A        | 68 a   | 3 A        | 72 ab        | 3 A        |
| Dicamba and glufosinate (SB, 14 DAA) | 99 a                    | 0 A        | 99 a         | 0 A        | 65 a   | 4 A        | 78 a         | 2 A        |
| Pr > F                               | <.0001                  | 1          | <.0001       | 1          | <.0001 | 0.3774     | <.0001       | 0.3828     |

<sup>a</sup>Abbreviations: DAA, days after application; SB, split-boom; TM, tank mix.

<sup>b</sup>Palmer amaranth control was determined by manually counting the number of plants per 0.145 m<sup>2</sup>.

<sup>c</sup>35 DAA evaluations occurred after the initial postemergence application only; 14 DAA evaluations occurred after sequential applications of postemergence herbicides and were evaluated 14 d after that.

<sup>d</sup>Lowercase letters indicate mean separation for percent Palmer amaranth control; uppercase letters indicate mean separation for Palmer amaranth plant density.

necessary for achieving high levels of Palmer amaranth control at this site.

Plant count data reflected the results of the observed control. Fewer plants were counted in the plots that received dicamba alone than the plots that were used as control checks. All treatments that included glufosinate further reduced the number of plants to zero (Table 3).

For treatments that did not receive a sequential application, visual ratings declined substantially over time and were similar at 35 DAA (Table 4). Regardless of application technique, sequential treatments averaged 99% control. These results indicate that a sequential herbicide application is necessary to achieve adequate control of Palmer amaranth at this location.

### Gibson

All four herbicide applications at the Gibson location failed to provide effective control of the native Palmer amaranth population at 14 DAA. Glufosinate alone provided better control than dicamba alone (Table 3). The tank-mix and split-boom application methods achieved better control than dicamba applied alone. The poor control with glufosinate is concerning and would be consistent with grower reports and the documented glufosinate resistance in the neighboring state of Arkansas (Priess *et al.* 2022). However, resistance was not experimentally confirmed in this study. Plant counts at 14 DAA that received dicamba alone and

glufosinate alone were similar to counts in the check plots. Plant counts were similar for all treatments.

Treatments that did not receive a sequential application exhibited a further decline in control 35 DAA (Table 4). Dicamba applied alone provided 26% control, while glufosinate applied alone declined to 17%, which is due to escapes from the initial application. The initial tank-mix and split-boom treatments each averaged 35% control at 35 DAA, which was better than glufosinate alone. Sequential applications improved control and results were similar. Although this level of control remains below what would be considered acceptable in a commercial production system, the results suggest that a sequential application is necessary to improve management of this Palmer amaranth population. Plant counts for the sequential applications were also similar.

### Lauderdale 2

The Palmer amaranth at the Lauderdale 2 location had been confirmed to be resistant to dicamba (Foster and Steckel 2022), and our results confirmed this. Following the initial 14-d rating (Table 3) the level of control observed was the split boom > tankmix > glufosinate alone > dicamba alone. Plant counts supported the visual ratings. All plots except those that served as nontreated checks had fewer plants after all treatments except dicamba alone. The split-boom treatment reduced plants by 90% compared with the check.

Control of Palmer amaranth declined over time after the initial applications, except for glufosinate, for which control remained the same. A reduction in control was observed at 35 d after dicamba was applied, while the tank-mix and split-boom treatments declined to 52% and 58% control, respectively (Table 4). Sequential applications provided similar control regardless of method. These results indicate that a sequential application may be beneficial for managing Palmer amaranth at this location. Plant counts correlated to visual ratings.

### Carrier Volume

In 2025, the tank mix of dicamba + glufosinate was evaluated for its effect against Palmer amaranth populations at three locations (Madison, Lauderdale 1, Gibson) to determine whether a greater carrier volume would enhance control of dicamba-resistant Palmer amaranth. Resistance levels at these locations were previously characterized by Foster and Steckel (2022) (Table 1).

At the three locations in 2025, increasing the carrier volume from 140 to 280 L ha<sup>-1</sup> did not improve control of dicamba-resistant Palmer amaranth (Table 5). When assessed at 14 d after the initial application, visual control was 85% with an application of 140 L ha<sup>-1</sup> compared with 81% after 280 L ha<sup>-1</sup> was applied ( $P = 0.6078$ ). Likewise, plant density did not differ between the carrier volumes. Following the second application, control followed the same trend with no difference ( $P = 0.0881$ ), and plant density was also similar ( $P = 0.2962$ ). Additionally, control and plant density were similar following the second application. Overall, these results indicate that increasing the carrier volume from 140 to 280 L ha<sup>-1</sup> had no effect on Palmer amaranth control, suggesting that the standard 140 L ha<sup>-1</sup> application volume is sufficient under these conditions.

### Practical Implications

The poor Palmer amaranth control with dicamba and glufosinate applied alone at the three test locations where resistance to auxinic herbicides is high would be consistent with reports by growers and an author's personal experience. These results suggest that this confirmed dicamba-resistant Palmer amaranth population may also exhibit resistance to glufosinate. Additional research is needed to confirm these findings. Furthermore, the results indicate that the most effective strategy for managing highly resistant Palmer amaranth is to apply dicamba and glufosinate using a split application approach in combination with other management practices. These may include applying residual herbicides at planting to limit Palmer amaranth emergence, overlapping residual herbicides throughout the season, and applying a sequential application of postemergence herbicides if Palmer amaranth emerges after the initial application.

From the regulatory standpoint, a tank mix of dicamba and glufosinate in a single spray solution was not permitted with the original dicamba over-the-top herbicide labels, meaning combinations of the two products would be considered an off-label application. In contrast, applying dicamba separately (split-boom) over the top would fall in accordance with its individual label instructions and is considered an on-label application when all other label requirements are met (US EPA 2022). Though the new labels for over-the-top application of dicamba (Bayer 2026) currently allow a tank mix with glufosinate, given all the changes with the labels since 2018 this could change in the future.

**Table 5.** Contrast statement for control of dicamba-resistant Palmer amaranth comparing 140 L ha<sup>-1</sup> with 280 L ha<sup>-1</sup> carrier volume, pooled across three locations in 2025.<sup>a-d</sup>

| Application volume<br>L ha <sup>-1</sup> | Palmer amaranth control   |            |                          |            |
|------------------------------------------|---------------------------|------------|--------------------------|------------|
|                                          | 14 DA initial application |            | 14 DA second application |            |
|                                          | %                         | No. plants | %                        | No. plants |
| 140                                      | 85 a                      | 3 A        | 88 a                     | 2 A        |
| 280                                      | 81 a                      | 2 A        | 85 a                     | 1 A        |
| $P > F$                                  | 0.6078                    | 0.1163     | 0.0881                   | 0.2962     |

<sup>a</sup>Abbreviation: DA, days after.

<sup>b</sup>Palmer amaranth control was determined by manually counting the number of plants per 0.145 m<sup>2</sup>.

<sup>c</sup>The Gibson, Lauderdale 1, and Madison locations are included in the analysis.

<sup>d</sup>Lowercase letters indicate mean separation for percent Palmer amaranth control; uppercase letters indicate mean separation for Palmer amaranth plant density.

**Acknowledgments.** We thank Hayden Love, Sally Reed, Jared Buck, David Cartwright, Maddie Douglas, Ernest Merriweather, and Julie Reeves for their assistance with this research.

**Funding.** Cotton Incorporated provided partial financial support of this project.

**Competing interests.** The authors declare they have no conflicts of interest.

### References

- Bayer (2026) Stryax™ herbicide label. St. Louis, MOL Bayer CropScience LLC. <https://www.epa.gov/ingredients-used-pesticide-products/registration-dicamba-use-dicamba-tolerant-crops>. Accessed: February 16, 2026
- Bland CR (2024) Efficacy of synthetic auxin herbicides on segregating populations of waterhemp (*Amaranthus tuberculatus*) (master's thesis). West Lafayette, IN: Purdue University. [https://hammer.purdue.edu/articles/thesis/\\_b\\_Efficacy\\_of\\_Synthetic\\_Auxin\\_Herbicides\\_on\\_Segregating\\_Populations\\_of\\_Waterhemp\\_b\\_b\\_i\\_Amaranthus\\_tuberculatus\\_i\\_b\\_b\\_b\\_/25668813/1?file=45797451](https://hammer.purdue.edu/articles/thesis/_b_Efficacy_of_Synthetic_Auxin_Herbicides_on_Segregating_Populations_of_Waterhemp_b_b_i_Amaranthus_tuberculatus_i_b_b_b_/25668813/1?file=45797451). Accessed: February 16, 2026
- Busi R, Goggin DE, Heap IM, Horak MJ, Jugulam M, Masters RA, Napier RM, Riar DS, Satchivi NM, Torra J (2018) Weed resistance to synthetic auxin herbicides. *Pest Manag Sci* 74:2265–2276
- Carmer S, Nyquist W, Walker W (1989) Least significant differences for combined analyses of experiments with two-or three-factor treatment designs. *Agron J* 81:665–672
- Constine AL (2021) Weed control and tank-mix interactions in soybean resistant to dicamba, glyphosate, and glufosinate (master's thesis). Ann Arbor: Michigan State University. <https://d.lib.msu.edu/etd/49764>. Accessed: February 16, 2026
- Delong BM (2020) Economics and efficacy of weed management in Bollgard II XtendFlex cotton (master's thesis). Lubbock: Texas Tech University. <https://ttu-ir.tdl.org/server/api/core/bitstreams/e840ad38-054b-4c11-9922-685541db5c4d/content>, Accessed: February 16, 2026. 62 p
- Foster DC, Steckel LE (2022) Confirmation of dicamba-resistant Palmer amaranth in Tennessee. *Weed Technol* 36:777–780
- Gleason C, Foley RC, Singh KB (2011) Mutant analysis in Arabidopsis provides insight into the molecular mode of action of the auxinic herbicide dicamba. *PLoS One* 6:e17245
- Grossmann K (2000) Mode of action of auxin herbicides: a new ending to a long, drawn out story. *Trends Plant Sci* 5:506–508
- Heap I (2025) The International Herbicide-Resistant Weed Database. <https://www.weedscience.org/Pages/Graphs/SOAGraph.aspx>. Accessed: August 12, 2025
- National Cotton Council (2024) 2024 cotton acreage. <https://www.cotton.org/news/releases/2024/24cotacr.cfm>. Accessed: August 12, 2025
- Oerke E-C (2006) Crop losses to pests. *J Agric Sci* 144:31–43

- Priess GL, Norsworthy JK, Godara N, Mauromoustakos A, Butts TR, Roberts TL, Barber T (2022) Confirmation of glufosinate-resistant Palmer amaranth and response to other herbicides. *Weed Technol* 36:368–372
- Shyam C, Chahal PS, Jhala AJ, Jugulam M (2021) Management of glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) in 2,4-D-, glufosinate-, and glyphosate-resistant soybean. *Weed Technol* 35:136–143
- [US EPA] U.S. Environmental Protection Agency (2022) Use, Usage, and Current and Future Benefits of Enlist One and Enlist Duo Herbicides in Corn, Soybean, and Cotton. Report EPA-HQ-OPP-2021-0957-0010. Washington: EPA Office of Chemical Safety and Pollution Prevention. Accessed: February 16, 2026
- [US EPA] U.S. Environmental Protection Agency (2024) Dicamba-Tolerant Cotton Planting Estimates and Related Data. Report EPA-HQ-OPP-2024-0154. <https://www.regulations.gov/docket/EPA-HQ-OPP-2024-0154/document>. Accessed: February 16, 2026
- Vulchi R, Bagavathiannan M, Nolte SA (2022) History of herbicide-resistant traits in cotton in the US and the importance of integrated weed management for technology stewardship. *Plants* 11:1189