

Review: Automated techniques for monitoring the behaviour and welfare of broilers and laying hens: towards the goal of precision livestock farming

N. Li^a, Z. Ren[†], D. Li and L. Zeng

College of Mechanical and Electrical Engineering, Hebei Agricultural University, Baoding 071001, Hebei, P.R. China

(Received 20 December 2018; Accepted 4 July 2019; First published online 30 September 2019)

There is increasing public concern about poultry welfare; the quality of animal welfare is closely related to the quality of livestock products and the health of consumers. Good animal welfare promotes the healthy growth of poultry, which can reduce the disease rate and improve the production quality and capacity. As behaviour responses are an important expression of welfare, the study of behaviour is a simple and non-invasive method to assess animal welfare. The use of modern technology offers the possibility to monitor the behaviour of broilers and laying hens in a continuous and automated way. This paper reviews the latest technologies used for monitoring the behaviour of broilers and laying hens under both experimental conditions and commercial applications and discusses the potential of developing a precision livestock farming (PLF) system. The techniques that are presented and discussed include sound analysis, which can be an online tool to automatically monitor poultry behaviour non-invasively at the group level; wireless, wearable sensors with radio-frequency identification devices, which can automatically identify individual chickens, track the location and movement of individuals in real time and quantify some behavioural traits accordingly and image processing technology, which can be considered a direct tool for measuring behaviours, especially activity behaviours and disease early warning. All of these technologies can monitor and analyse poultry behaviour, at the group level or individual level, on commercial farms. However, the popularity and adoption of these technologies has been hampered by the logistics of applying them to thousands and tens of thousands of birds on commercial farms. This review discusses the advantages and disadvantages of these techniques in commercial applications and presents evidence that they provide potential tools to automatically monitor the behaviours of broilers and laying hens on commercial farms. However, there still has a long way to go to develop a PLF system to detect and predict abnormal situations.

Keywords: welfare assessment, behaviour detection, sound analysis, radio-frequency identification, image processing

Implications

Behaviour is one of the important indicators in animal welfare assessments and can be considered in many dimensions, including the affective state, health and needs of animals. Technological development offers automated and continuous detection of poultry behaviour, so farmers can take actions to improve welfare. This review focuses on different sensor technologies to detect behaviours of broilers or laying hens, which are applied to analyse animal welfare conditions. It has been shown that sensor-based systems can monitor animal welfare and behaviour continuously, precisely and automatically with the potential to develop precision livestock farming systems.

Introduction

Currently, there is rising concern about poultry welfare, and poultry breeding should be conducted humanely with regard to feeding and slaughtering. Simultaneously, healthy poultry and product quality rely heavily on good welfare conditions, which can improve economic efficiency (Wathes *et al.*, 2008). A good state of welfare encompasses birds having a healthy body, a positive affective state and the opportunity to express natural behaviours (Sih *et al.*, 2004). However, there are various indexes for the comprehensive assessment of animal welfare, such as behavioural indicators, physiological stress and proxies of productivity. Many of them interact together, making evaluation difficult and time-consuming, especially at the commercial level. Among these welfare indexes, behaviour is the most easily understood and commonly used

^a Present address: Hebei Agricultural University No. 289 Lingyusi Street, Baoding 071001, Hebei, P.R. China.

[†] E-mail: renzh68@163.com

welfare assessment index (De Jong *et al.*, 2016), because behaviours which may be influenced by living conditions, emotions and diseases of animals can reflect physiological conditions. Therefore, monitoring animal behaviours and formulating feeding and management measures accordingly can not only promote the growth of animals and improve the production performance, but also protect animal welfare.

Precision livestock farming (PLF) is defined as the application of process engineering principles and techniques to livestock farming to automatically monitor, model and manage animal production (Tullo *et al.*, 2017). The aim of PLF is to real-time monitor animal health and welfare automatically and continuously and prompt warning messages rapidly. Monitoring the resource use of chickens can improve the housing design and allocate resources optimally (Ringgenberg *et al.*, 2015; Li *et al.*, 2017). Through the location of chicken clustering, the quality of house facilities (e.g., feeders, drinkers) may be examined (Fraess *et al.*, 2016). Detecting the activity of poultry, such as eating, drinking, moving and perching, may estimate whether they are healthy and therefore provide early warning of diseases (Colles *et al.*, 2016; Dawkins *et al.*, 2017). Accordingly, the great potential of PLF relies on early warning, which allows farmers to take actions in the initial stages of welfare problems or diseases. However, to truly achieve PLF, it is necessary to monitor individual animals. Research on individual behaviour can control the health and production of every individual and improve the accuracy of welfare assessment. For example, detecting the movements of each chicken by a radio-frequency identification (RFID) system, the chickens can be classified into active, normal and sick level (Zhang *et al.*, 2016). Therefore, each sick chicken can be found in time. Compared with large animals (e.g., cattle and pig), poultry are not only small, but raised in hundreds, thousands or hundreds of thousands in a commercial farm, making it difficult to monitor automatically and individually. However, despite that, PLF is able to provide new opportunities to promote animal husbandry with high-efficiency, low-cost, ecological and sustainable development and offers consumers some food safety assurance (Berckmans, 2014).

The PLF system is based on sensors (e.g., RFID devices, accelerometer, camera, microphone and temperature sensor) and collects a variety of information (e.g., activity, body temperature, feed intake and weight) used for environmental monitoring, individual recognition and behaviour analysis. A range of sensor techniques are available that allow efficient and continuous data collection from multiple sources to monitor animal behaviour, health and production. Furthermore, advanced data processing techniques make it possible to analyse animal behaviour continuously, automatically, accurately and in real time (Valletta *et al.*, 2017; Van Herterem *et al.*, 2017). Ben Sassi *et al.* (2016) reviewed the sensors and technologies for poultry welfare monitoring based on the literature prior to 2015. The review emphasised that the monitoring technologies could be implemented at the commercial level but did not analyse the difficulties of commercial application.

The aim of this paper is to discuss the possibility of accomplishing PLF through review of automatic methods implemented to monitor behaviours of broilers or laying hens at the experimental or commercial level in the last 5 years. The review expounds automated techniques, including sound analysis, RFID devices and image processing, which have been applied to poultry welfare and behaviour analysis, with discussion of the detection aspects of these techniques, advantages and disadvantages on a commercial scale, and feasibility for PLF.

Sound analysis

Vocalisations of animals contain a wealth of biological information (e.g., their social interactions, alarm signals and behaviour needs) and have the possibility to be used as a welfare indicator (Manteuffel *et al.*, 2004). When applied to poultry systems, the microphone is always put low near the animals and can capture sounds continuously and non-invasively, including pecking, vocalisations and environmental noises. The sound analysis techniques usually take the duration, energy distribution, frequency and formant amplitude as the characteristic parameters to estimate the behaviour and health of birds. It can be automated, in real time, and is capable of long-term monitoring without disturbing the birds.

Growth rate detection

Monitoring feed conversion, growth rate and BW is important to improve the efficiency and economic benefits of broiler farmers. Considering the relationship between pecking behaviour and eating, 90% of the feed intake of one broiler was correctly detected with pecking sounds in an experimental pen (Aydin *et al.*, 2014). In another study, 86% of the feed intake of 10 broilers was detected accurately (Aydin *et al.*, 2015). Subsequently, short-term feeding behaviours of 10 broilers by pecking sound analysis were estimated in real time, and the accuracy of meal size reached 90%, meal duration reached 95%, the numbers of meals per day reached 84% and feeding rate reached 89% (Aydin and Berckmans, 2016). The above three papers were studied in an experimental pen of at most 10 chickens. However, the application to a commercial farm needs to be validated, including filtering out environmental noise and feasibility of individual detection with large bird numbers.

Broiler vocalisations are highly correlated with BW, and sound analysis could be a reliable method of measuring the growth rate of broiler chickens. In an intensive broiler farm, a study showed that with the growth of broiler chickens, the peak frequency of the sounds emitted by broilers decreased clearly during the entire life of the birds (Fontana *et al.*, 2015 and 2016). Furthermore, a model to predict the weight of broilers was built and validated by the peak frequency analysis of the broilers' vocalisations (Fontana *et al.*, 2017). Vocalisations could thus be used to assist in the automatic monitoring of the growth of broiler chickens.

Disease detection

There is a difference between the sound of healthy and unhealthy chickens. Analyses of the frequency domain features of 15 infected *Clostridium perfringens* type A chickens and 15 healthy chickens, classified by a supervised learning neural network, showed a classification accuracy of 66.6% 2 days after infection and reached 100% accuracy 8 days after infection (Sadeghi *et al.*, 2015). Using data-mining methods and Dempster–Shafer evidence theory, a device was made to detect whether a chicken was infected with Newcastle disease, bronchitis virus, avian influenza or healthy (Banakar *et al.*, 2016). The device could determine the type of disease in each chicken without surrounding noises.

Others

Sound analysis could identify the sex of 1-day-old chickens as well as different genetic strains (Cobb and Ross chickens) (Pereira *et al.*, 2015). Analysing four acoustic sound characteristics showed that the second spectral peak of the sound spectrum could identify the sex of chicks, and both the second spectral peak and the pitch could identify the genetic strain. Stress, such as temperature, fear and physical stress, can affect the health and production of commercial chickens. The sounds of chickens under heat or fear stress were detected in a commercial farm (Lee *et al.*, 2015). The proposed system was able to classify different stresses by support vector machine (SVM) techniques automatically and non-invasively, and both the stress detection accuracy and the average classification accuracy were greater than 96%. The vocal variations of hens anticipating different types of rewards were investigated and showed that different sound cues made different vocal parameters produced by hens (McGrath *et al.*, 2017). It was possible to use the vocal communication of hens as indicators of welfare (e.g., health, stress, growth).

In summary, sound analysis is able to be used as an early warning system by interpreting the behaviours of chickens, predicting health conditions and detecting growth rates. The group sound analysis, which was tested commercially, can isolate the sounds of each bird, but it is impossible to find the exact bird which makes the sound. The sound collection and analysis of individual chicken, which can distinguish the chicken, is still at experimental stage due to the difficulty of collecting each individual sound in a commercial farm. Like voiceprint recognition, if the voice feature model of each bird could be established, the group sound collection can also match chickens to their sounds. Meanwhile, environmental noise is also a big challenge. With the maturation of voice recognition technology and the rapid development of artificial intelligence (AI), animal sound analysis has great potential as an online detection method to improve health and welfare in the poultry industry.

Tracking location and locomotion of individuals by radio-frequency identification

Wireless wearable sensors (e.g., accelerometer, RFID microchips) are mainly used to track the location and activity of

individuals remotely. These sensors can distinguish the behaviours of individuals automatically, particularly when human observations are difficult. Siegford *et al.* (2016) reviewed the technologies, especially RFID, which was applied as a research tool for tracking individual laying hens' activity and location. However, RFID technology is rarely used in commercial poultry farms compared with farms producing larger animals (e.g., pigs and cattle), due to the small size and light weight of individual birds, the required size of the on-bird microchips and high financial investment cost. Radio-frequency identification technology is one method for automatic identification and data capture, which can be used mainly under experimental conditions to track the location of individual birds. An RFID microchip can also be injected to the leg or back of a bird produces a signal to the RFID reader when the antenna is within the magnetic field and the signal is sent to the central information system to identify and position the individual. Tracking the location of individual birds may allow observation of nesting, feeding, drinking, perching behaviour or ranging behaviour and, thus, prediction of health conditions.

Detecting nest use

Put RFID reader into nest and worn RFID tags on the leg of hens can detect a hen in and out of a nest. A smart nest box was designed to detect the laying behaviour of individual hens (Chien and Chen, 2018). A low-frequency RFID system was used to detect whether hens were in the nest. An egg detection sensor could detect whether hens laid eggs and then weigh the eggs. Design-based Internet of Things enabled all data to be recorded simultaneously in the cloud database and a local SD card at the same time. Assigning each egg to an individual can achieve product traceability and analyse the laying performance of individuals. The system had the potential to monitor the underperforming hens in egg production due to welfare problems.

Establishing hens' visits to the nest can improve the nest design. Using the Gantner Pigeon System GmbH (Schrans, Austria) RFID system, researchers found that the nest choice was more affected by the nest position, nest occupancy and age of hens than by the appearance of nest curtains (Ringgenberg *et al.*, 2015). Detecting laying behaviour could also examine the relationship between laying eggs and keel bone deviations and fractures (Gebhardt-Henrich and Frohlich, 2015).

Tracking movement and feeding behaviour of individuals

Radio-frequency identification systems can track the position of individual birds to detect the movement and duration at a particular location. With a low-frequency RFID system, an environmental preference chamber with four compartments and two tiers was designed to study the movement in different compartments of four individual birds (Sales *et al.*, 2015). Compared with video observation, successful detection rates were $62.6 \pm 11.2\%$. The detection range of the antenna, the scan interval of the antenna and the RFID tags conflicts would affect the RFID system's performance.

By monitoring the weight and moving time of individual hens using weighing sensors and RFID devices, poultry behaviours such as speed, ability to acquire food and resting time were detected, and by the K-means clustering method, chickens were classified as active, normal or sick (Zhang *et al.*, 2016). An ultra-high-frequency RFID system was developed to monitor the feeding and nesting behaviours of 60 individual hens in enriched colony housing (Li *et al.*, 2017). Using the same experimental device, feeding behaviour was assessed with various feeder spaces (12.0, 9.5, 8.5 and 6.5 cm/hen) (Oliveira *et al.*, 2018). This type of research could contribute to housing system design, animal welfare and resource allocation.

Ranging behaviour in free-range housing

Currently, the free-range system is considered to be more natural and better for poultry welfare. However, each hen within a group has a different ranging behavioural pattern. To monitor the outdoor ranging behaviour of individual chickens, RFID systems have been used by attaching tags on chickens and putting antennas at pop holes, through which chickens move inside and outside (Gebhardt-Henrich *et al.*, 2014; Campbell *et al.*, 2017; Larsen *et al.*, 2017). The frequency and duration of visiting the outdoor areas are then calculated by analysing data from chickens walking over the antennas. These systems could be used in commercial farms and potentially track thousands of birds. In addition, an RFID system based on ultra-wideband technology was developed for monitoring 42 individual chickens' locations in the 100 m × 100 m experimental field, which could calculate the position with 0.29 m of the median error and 68% of the registration rate (Stadig *et al.*, 2018a). Monitoring of outdoor range use at the individual level allows for correlation between ranging and welfare indicators such as tonic immobility test, plumage damage, foot health and spatial cognition (Campbell *et al.*, 2016, 2018a and 2018b; Hartcher *et al.*, 2016; Larsen *et al.*, 2018; Taylor *et al.*, 2018).

The impact of wearing a tag

Because the birds are small, the size, shape, weight and attachment method of tags might influence bird behaviour and the accuracy of the data. The behavioural and physiological responses of laying hens which wore backpacks containing automated monitoring devices were studied and showed that the backpacks only had a very minor effect on their behaviour after a short acclimation period (≤ 2 days) (Buijs *et al.*, 2018). Furthermore, the effects on broiler behaviour, leg health, cleanliness and growth were evaluated with a tag worn on the back of birds for tracking location in a field (Stadig *et al.*, 2018b). Six pens with 10 birds in each were observed with 5 birds fitted with a 36-g backpack and the other five with coloured markings to identify individuals. Footpad and hock dermatitis, gait and cleanliness were scored manually. These results indicated that birds wearing backpacks showed less activity early on, but no significant differences were observed in long-term behaviours between

tagged birds and coloured birds. In summary, compared to the weight of chickens, a light monitoring tag has little impact on the behaviours of chickens after a period of adjustment. Wireless sensors are a reliable tool for detecting the location and activity of individual chickens.

Image processing

Image processing technology has been applied to automatically monitor animal behaviour for the analysis of animal welfare, health conditions, weight prediction and tracking. Image processing used to detect chicken flocks can be automatic, non-invasive and continuous. Using automated computer monitoring, it is possible to record and identify different behaviours of chickens, which is an inexpensive and cost-effective approach. Typical materials and methods of image processing used in chicken behaviour detection, such as camera type, software, algorithms and application area, are shown in Table 1.

Flock activity analysis

Through the number of changed pixels between two adjacent frames divided by the overall number of pixels in the image, the flock activity level can be calculated (Youssef *et al.*, 2015; Fraess *et al.*, 2016). The method is heavily influenced by plumage differences. Group activity around the food trough was assessed by the percent pixel change (Fraess *et al.*, 2016). The study showed that the method could identify changes in activity level, identify abnormal activity away from the feeder and serve as an early health warning system.

Optical flow is a method to find the corresponding relationship between the previous frame and the current frame by analysing the change in pixels in the time domain of the image sequence and the correlation between adjacent frames, to calculate the motion of objects between adjacent frames. It is suitable for monitoring flock activity in the commercial farm. Using the optical flow patterns of chicken movements, infection by the human pathogen *Campylobacter* was detected in the early stages (Colles *et al.*, 2016). Optical flow data were collected, and the mean, variance, skew and kurtosis were calculated. Infected *Campylobacter* flocks had low mean optical flow and an apparent high kurtosis, and infection was detected 7 to 10 days earlier compared with faecal sample testing. This system has the potential for early health and welfare warnings that can benefit farmers and chickens. Using the camera-based optical flow method, early-stage footpad dermatitis and hock burn were predicted in 50 flocks from a commercial farm (Dawkins *et al.*, 2017). Optical flow analysed the movements of flocks, and lower skew and higher kurtosis values indicated heterogeneity in flock movement, which had predictive power for footpad dermatitis and hock burn.

In commercial farms, the activity and sudden variation of flock could be related to abnormal conditions, such as the equipment failure and diseases. Converting images into animal distribution index, the equipment failure in a broiler

Table 1 Typical materials and methods of automated image processing used in chicken behaviour monitoring

Reference	Camera type	Image pixel and frame rate	Main processing method	Detecting number	Main software
Zaninelli <i>et al.</i> , 2018	Thermographic camera	320 × 240; 1 fps	Threshold method and image pattern recognitions	10	LabVIEW, NI Vision Acquisition Software
Li <i>et al.</i> , 2018	IR camera	–	Threshold method	8	MATLAB
Pu <i>et al.</i> , 2018	Kinect camera	512 × 424; 30 fps	CNN	Commercial flocks	Open CV
Zhuang <i>et al.</i> , 2018	CCD camera	640 × 480	K-means clustering and the ellipse model	20	Open CV
Aydin <i>et al.</i> , 2017b	Camera	1024 × 768; 5 fps	Background subtraction and feature variables	1	MATLAB
Dawkins <i>et al.</i> , 2017	Camera	320 × 240	Optical flow	Commercial flocks	–
Fraess <i>et al.</i> , 2016	IR camera	320 × 240; 30 fps	The proportion of pixel changes	34	Etho Vision XT 10
Colles <i>et al.</i> , 2016	Web camera	320 × 240; 1 fps	Optical flow	Commercial flocks	MATLAB Python
Youssef <i>et al.</i> , 2015	CCD camera	640 × 480; 1 fps	The proportion of pixel changes	45	–
Kashiha <i>et al.</i> , 2014	Camera	1280 × 960; 0.5 fps	Binary image and count the pixel proportion	Commercial flocks	eYeNamic

CCD = charge coupled device; CNN = convolutional neural network.

house could be found out in real time by the sudden changes of the distribution index (Kashiha *et al.*, 2014). With a uniform moderate walking pace through the flock, four variables were derived from the changes in activity before and after walking, so that the gait score could be predicted automatically (Silvera *et al.*, 2017). The occupation index was used instead of the distribution index, and the activity and occupation of the flock had correlation with the welfare scores for the footpad lesions and hock burns by human experts (Fernandez *et al.*, 2018). Moreover, the precise fault location could be performed by these indices in specific areas such as feeding, drinking and resting.

By combining a computer vision system with a novel convolutional neural network (CNN) method, the flock behaviour in a commercial farm was monitored automatically with a Kinect camera (Pu *et al.*, 2018). Three categories around feeders, not crowded, slightly crowded and fairly crowded, were labelled manually among 17 135 colour images and 17 135 depth images, and a supervised learning CNN was trained. The experimental results show that our proposed CNN-based method can recognise the flock behaviour of chickens with an accuracy of 99.17%.

Behaviours of individuals

Gait score of an individual hen can be done automatically. A digital video camera was mounted above the centre of a testing corridor to record images. The image processing method could classify the number of lying events (NOL) and latency to lie (LTL) of broilers (Aydin, 2015, 2017a and 2017b). The background subtraction method was applied to segment the shape, and then an elliptical shape model was

used to define the size and orientation of the chicken (Aydin, 2015). Compared with the gait scores calculated manually from 250 broilers, the automatic monitoring system had 83% accuracy of identifying NOL. Subsequently, the image processing algorithm was improved by extracting feature variables (Aydin, 2017a). Automatically collected feature variables of broilers showed a clear correlation between feature variables and gait scores and were found to have a high level of accuracy starting from gait score three. Furthermore, instead of the 2D camera, a Kinect camera was used in the same experiment (Aydin, 2017b). Additional distance information between the broiler and the 3D camera was collected as broilers walked along a test corridor, and data were extracted to detect NOL and assess LTL with the back posture. The accuracy of NOL classification reached 93% compared to manual scoring.

In addition to RFID, image processing is another technology that has been considered to detect nest use. It can detect double nest occupation (Zaninelli *et al.*, 2015 and 2018), while RFID cannot. Photographs taken in a nest that was painted black, with the number of colour pixels counted and compared with the threshold double nest occupation, could detect whether there were two or more hens in a nest (Zaninelli *et al.*, 2015). Later, with a thermographic camera to identify laying hens in a closed room (Zaninelli *et al.*, 2016 and 2017), this team developed a better 'nest usage sensor', which could count the number of hens in the nest (Zaninelli *et al.*, 2018). The thermographic images were processed by binarisation and coloured, and the number of colour pixels was counted to detect whether there were more than one hen in the nest. If there had a multiple nest occupation,

pattern recognition was performed to make sure how many hens were in the nest.

The image processing algorithm could calculate the number of hens using an IR camera in an automatic lighting preference test system (Li *et al.*, 2018). Original colour images were first cropped, converted into grey scale and then converted into binary images. The number of hens could be concluded by the number of white pixels. The accuracy of the image algorithm was 71.23%, due to the number of pixels changing significantly for one hen when feeding, wing flapping and preening.

The area required by each hen for the behaviours of standing, lying, perching, wing flapping and dust bathing was studied using ImageJ software and MATLAB (Riddle *et al.*, 2018). However, it required manual selection of perfect images. Diseased broilers were predicted and classified automatically by the SVM model (Zhuang *et al.*, 2018). To obtain the posture features, images were segmented by K-means clustering, and the skeleton structure was obtained by the thinning method. Superior to other machine learning (ML) algorithms, the SVM model obtained an accuracy rate of 99.47%.

Weight prediction

An automatic weight prediction system based on a 3D camera was developed and assessed on a commercial farm (Mortensen *et al.*, 2016). The algorithm consisted of a range-based watershed algorithm and was predicted by a Bayesian artificial neural network. The system was able to weigh several chickens simultaneously. Due to the overlapping of broilers in the image, better prediction could be improved by a better segmentation method. Moreover, to estimate the weight of live broiler chickens, a machine vision

method with an artificial neural network (ANN) was proposed (Amraei *et al.*, 2017). Compared to Levenberg–Marquardt, scaled conjugate gradient and gradient descent, Bayesian regulation was the best network for the prediction of broiler weight. The maximum error of the system was less than 50 g. The aforementioned systems weighed individual broilers.

Discussion

Over the last few years, the prominent emergence of sensor technology, AI and big data analysis has made it possible to implement PLF. In this review, there are a total of 50 papers using automated techniques to detect poultry welfare. Of these, 14 papers including 3 of sound analysis, 4 of RFID and 7 of image processing were tested on a commercial level (even if just experimentally in a commercial flock). These techniques all have the ability to monitor the behaviour of birds under commercial conditions. Figure 1 shows the number of papers classified by detection contents and sensors.

The use of sound and image processing technologies offers a comparatively non-invasive way to monitor the health and behaviour of animals over the long term. Due to the increasing size in commercial production systems for broilers and laying hens, it is common to observe and assess welfare at the group level. At the group level, sound analysis technology can be used to detect weight and stress of chickens on a commercial farm (Lee *et al.*, 2015; Fontana *et al.*, 2017). Other detections based on sound analysis are at the experimental stage with a small number of birds, but they have a great potential to be applied in commercial chicken houses. Due to the environmental noise and simultaneous




Figure 1 (Colour online) The articles, which monitored behaviour and welfare of broilers and laying hens automatically, are classified by detection contents and sensors. The height of the bars denotes the number of articles. Each sensor, including microphone, RFID and camera, has two bars. The left bar shows all the articles used in the sensor which are divided into five categories. The right bar shows the commercially tested articles which also have five categories. Activities refer to basic behaviours, such as feeding, drinking, perching, walking ability (including gait score) and laying behaviours (multiple nest occupation). Preference test includes environment preference, lighting preference, nest choice and free-range use. Others mean not belonging to the other four categories (e.g., equipment failure, welfare assessment). RFID, radio-frequency identification.

vocalisations, the accuracy of algorithms may be affected in complex environments. Using image processing, the activity of chicken flocks can be monitored on commercial farms and in real time (Colles *et al.*, 2016; Dawkins *et al.*, 2017). Mainly defined as the pixels change in the whole image, the activity means the flock is active in general but the specific behaviours of the flock are not classified. Moreover, some cameras have blind spots that cannot cover the entire chicken house. Another problem is that cameras, except IR cameras, cannot monitor the activity of chickens when lights are off.

However, PLF systems should be based on individual welfare assessments, because the average group welfare level may not accurately predict specific individual welfare. Monitoring individual behaviour, especially monitoring multiple individuals in a large group, requires the recognition and tracking of animals. Radio-frequency identification technology has been utilised in PLF systems to identify, track and register animals to detect behaviour and welfare in a reliable, automated and inexpensive way. While the RFID system can collect a large amount of data, it estimates behaviour by indirectly tracking the position of individuals. Additionally, the accuracy and tracking of this system depends on the number of antennas installed and the number of chickens wearing transponders. Because of the huge number of birds on commercial farms, it is currently unrealistic for each bird to wear an RFID tag. Typically, a random sample of birds is selected to wear a tag. Although RFID systems can be applied to hundreds of chickens, fitting and recycling tags is a time-consuming and laborious problem. Image processing technology can be used for target detection and tracking. With image target tracking and identification algorithms, an individual bird was tracked and identified in a group of six birds (Wang *et al.*, 2016). Due to the high breeding density on commercial farms, it is difficult to automatically identify and track individual birds in a long time based on image processing with currently technology. Occlusion is another consideration. By combining image processing with RFID, individual birds can be tracked and behaviour can be detected more accurately (Nakarmi *et al.*, 2014). The combination of multiple methods is such a way to implement PLF.

Since the advent of cheaper sensors and advanced sensor technology used for the observation of animal behaviour, huge data sets have been collected. These data sets are large, complex and diverse; thus, it is difficult to analyse by classical statistical methods. Unlike classical mathematical statistics, ML is a branch of AI that can analyse complex model data sets with unknown relationships between measured variables. ML attempts to mine the hidden variables from large collections of historical data for future prediction or classification. Many ML methods, such as decision trees, SVMs (Banakar *et al.*, 2016; Zhuang *et al.*, 2018) and ANNs (Pu *et al.*, 2018), have been used in welfare assessments. Therefore, ML can provide strong support in developing an early warning system for PLF systems. On the other hand, multi-sensor integration system has a trend made for PLF (Van Hertem *et al.*, 2017). It is a hard work to perform a meta-analysis of the collected data of control units and

behaviour detect devices such as climate control units, feed distribution units, camera-based system and microphone-based system. For farmers, only systems which are simple, easy to understand and easy to operate are been willing to use, for example, a visualisation tool. The stability of the system and the accuracy of measurement and prediction are also other factors that affect the use of farmers.

Conclusion

The development of modern technology offers powerful tools to automatically detect poultry health and welfare. However, most applications have the limitations of only being used in experimental conditions, although they have the potential to be used in commercial systems. The assessment of animal welfare is being transformed from the group level to the individual level. Consequently, the full implementation of the PLF system remains a great challenge. Further research is required to improve the accuracy of data processing and modelling, as well as to improve device detection before commercial validation. Once available, PLF systems will create added value to improve animal welfare, health and efficiency for farmers.

Acknowledgements

The authors would like to thank American Journal Experts for editing the English language in this paper. This work was also supported by the Key Science and Technology Research and Development Program of Hebei Province (16236605D-2 (2018)). The authors also thank the anonymous reviewers whose critiques and comments greatly improved the manuscript.

Declaration of interest

The authors declare no conflicts of interest.

Ethics statement

Conducting this review involved no animal handling or procedures.

Software and data repository resources

No new software or original data resources were generated as part of this study.

References

- Amraei S, Abdanan Mehdizadeh S and Salari S 2017. Broiler weight estimation based on machine vision and artificial neural network. *British Poultry Science* 58, 200–205.
- Aydin A 2015. Automatic classification of measures of lying to assess the lameness of broilers. *Animal Welfare* 24, 335–343.
- Aydin A 2017a. Using 3D vision camera system to automatically assess the level of inactivity in broiler chickens. *Computers and Electronics in Agriculture* 135, 4–10.

- Aydin A 2017b. Development of an early detection system for lameness of broilers using computer vision. *Computers and Electronics in Agriculture* 136, 140–146.
- Aydin A, Bahr C and Berckmans D 2015. A real-time monitoring tool to automatically measure the feed intakes of multiple broiler chickens by sound analysis. *Computers and Electronics in Agriculture* 114, 1–6.
- Aydin A, Bahr C, Viazzi S, Exadaktylos V, Buysse J and Berckmans D 2014. A novel method to automatically measure the feed intake of broiler chickens by sound technology. *Computers and Electronics in Agriculture* 101, 17–23.
- Aydin A and Berckmans D 2016. Using sound technology to automatically detect the short-term feeding behaviours of broiler chickens. *Computers and Electronics in Agriculture* 121, 25–31.
- Banakar A, Sadeghi M and Shushtari A 2016. An intelligent device for diagnosing avian diseases: Newcastle, infectious bronchitis, avian influenza. *Computers and Electronics in Agriculture* 127, 744–753.
- Ben Sassi N, Averós X and Estevez I 2016. Technology and poultry welfare. *Animals* 6, 62. doi: [10.3390/ani6100062](https://doi.org/10.3390/ani6100062).
- Berckmans D 2014. Precision livestock farming technologies for welfare management in intensive livestock systems. *Revue Scientifique et Technique – Office International des Epizooties* 33, 189–196.
- Buijs S, Booth F, Richards G, McGaughey L, Nicol CJ, Edgar J and Tarlton JF 2018. Behavioural and physiological responses of laying hens to automated monitoring equipment. *Applied Animal Behaviour Science* 199, 17–23.
- Campbell DLM, Hinch GN, Downing JA and Lee C 2016. Fear and coping styles of outdoor-preferring, moderate-outdoor and indoor-preferring free-range laying hens. *Applied Animal Behaviour Science* 185, 73–77.
- Campbell DLM, Hinch GN, Downing JA and Lee C 2018a. Early enrichment in free-range laying hens: effects on ranging behaviour, welfare and response to stressors. *Animal* 12, 575–584.
- Campbell DLM, Hinch GN, Dyall TR, Warin L, Little BA and Lee C 2017. Outdoor stocking density in free-range laying hens: radio-frequency identification of impacts on range use. *Animal* 11, 121–130.
- Campbell DLM, Talk AC, Loh ZA, Dyall TR and Lee C 2018b. Spatial cognition and range use in free-range laying hens. *Animals* 8, 26. doi: [10.3390/ani8020026](https://doi.org/10.3390/ani8020026).
- Chien YR and Chen YX 2018. An RFID-based smart nest box: an experimental study of laying performance and behavior of individual hens. *Sensors* 18, 859. doi: [10.3390/s18030859](https://doi.org/10.3390/s18030859).
- Colles FM, Cain RJ, Nickson T, Smith AL, Roberts SJ, Maiden MCJ, Lunn D and Dawkins MS 2016. Monitoring chicken flock behaviour provides early warning of infection by human pathogen *Campylobacter*. *Proceedings of the Royal Society B-Biological Sciences* 283, 20152323. doi: [10.1098/rspb.2015.2323](https://doi.org/10.1098/rspb.2015.2323).
- Dawkins MS, Roberts SJ, Cain RJ, Nickson T and Donnelly CA 2017. Early warning of footpad dermatitis and hockburn in broiler chicken flocks using optical flow, bodyweight and water consumption. *Veterinary Record* 180, 499–U60.
- De Jong IC, Hindle VA, Butterworth A, Engel B, Ferrari P, Gunnink H, Moya TP, Tuytens FAM and van Reenen CG 2016. Simplifying the Welfare Quality (R) assessment protocol for broiler chicken welfare. *Animal* 10, 117–127.
- Fernandez AP, Norton T, Tullo E, van Hertem T, Youssef A, Exadaktylos V, Vranken E, Guarino M and Berckmans D 2018. Real-time monitoring of broiler flock's welfare status using camera-based technology. *Biosystems Engineering* 173, 103–114.
- Fontana I, Tullo E, Butterworth A and Guarino M 2015. An innovative approach to predict the growth in intensive poultry farming. *Computers and Electronics in Agriculture* 119, 178–183.
- Fontana I, Tullo E, Carpentier L, Berckmans D, Butterworth A, Vranken E, Norton T, Berckmans D and Guarino M 2017. Sound analysis to model weight of broiler chickens. *Poultry Science* 96, 3938–3943.
- Fontana I, Tullo E, Scrase A and Butterworth A 2016. Vocalisation sound pattern identification in young broiler chickens. *Animal* 10, 1567–1574.
- Fraess GA, Bench CJ and Tierney KB 2016. Automated behavioural response assessment to a feeding event in two heritage chicken breeds. *Applied Animal Behaviour Science* 179, 74–81.
- Gebhardt-Henrich SG and Frohlich EKF 2015. Early onset of laying and hockbone favor keel bone fractures. *Animals* 5, 1192–1206.
- Gebhardt-Henrich SG, Toscano MJ and Frohlich EKF 2014. Use of outdoor ranges by laying hens in different sized flocks. *Applied Animal Behaviour Science* 155, 74–81.
- Hartcher KM, Hickey KA, Hemsworth PH, Cronin GM, Wilkinson SJ and Singh M 2016. Relationships between range access as monitored by radio frequency identification technology, fearfulness, and plumage damage in free-range laying hens. *Animal* 10, 847–853.
- Kashiha M, Bahr C, Vranken E, Hong SW and Berckmans D 2014. Monitoring system to detect problems in broiler houses based on image processing. In *Proceedings International Conference of Agricultural Engineering*, 6–10 July 2014, Zurich, Switzerland, pp. 6–10.
- Larsen H, Cronin GM, Gebhardt-Henrich SG, Smith CL, Hemsworth PH and Rault JL 2017. Individual ranging behaviour patterns in commercial free-range layers as observed through RFID tracking. *Animals* 7, 21. doi: [10.3390/ani7030021](https://doi.org/10.3390/ani7030021).
- Larsen H, Hemsworth PH, Cronin GM, Gebhardt-Henrich SG, Smith CL and Rault JL 2018. Relationship between welfare and individual ranging behaviour in commercial free-range laying hens. *Animal* 12, 2356–2364.
- Lee J, Noh B, Jang S, Park D, Chung Y and Chang HH 2015. Stress detection and classification of laying hens by sound analysis. *Asian-Australasian Journal of Animal Sciences* 28, 592–598.
- Li GM, Li BM, Shi ZX, Zhao Y and Ma H 2018. Design and evaluation of a lighting preference test system for laying hens. *Computers and Electronics in Agriculture* 147, 118–125.
- Li L, Zhao Y, Oliveira J, Verhoijens W, Liu K and Xin H 2017. A UHF RFID system for studying individual feeding and nesting behaviors of group-housed laying hens. *Transactions of the ASABE* 60, 1337–1347.
- Manteuffel G, Puppe B and Schon PC 2004. Vocalization of farm animals as a measure of welfare. *Applied Animal Behaviour Science* 88, 163–182.
- McGrath N, Dunlop R, Dwyer C, Burman O and Phillips CJC 2017. Hens vary their vocal repertoire and structure when anticipating different types of reward. *Animal Behaviour* 130, 79–96.
- Mortensen AK, Lisouski P and Ahrendt P 2016. Weight prediction of broiler chickens using 3D computer vision. *Computers and Electronics in Agriculture* 123, 319–326.
- Nakarmi AD, Tang L and Xin H 2014. Automated tracking and behavior quantification of laying hens using 3D computer vision and radio frequency identification technologies. *Transactions of the ASABE* 57, 1455–1472.
- Oliveira JL, Xin H and Wu H 2018. Impact of feeder space on laying hen feeding behavior and production performance in enriched colony housing. *Animal* 13, 374–383.
- Pereira EM, Naas ID and Garcia RG 2015. Vocalization of broilers can be used to identify their sex and genetic strain. *Engenharia Agrícola* 35, 192–196.
- Pu HT, Lian J and Fan MQ 2018. Automatic recognition of flock behavior of chickens with convolutional neural network and Kinect sensor. *International Journal of Pattern Recognition and Artificial Intelligence* 32, 1850023. doi: [10.1142/S0218001418500234](https://doi.org/10.1142/S0218001418500234).
- Riddle ER, Ali ABA, Campbell DLM and Siegford JM 2018. Space use by 4 strains of laying hens to perch, wing flap, dust bathe, stand and lie down. *PLoS ONE* 13, e190532.
- Ringgenberg N, Frohlich EKF, Harlander-Matuschek A, Toscano MJ, Wurbel H and Roth BA 2015. Effects of variation in nest curtain design on pre-laying behaviour of domestic hens. *Applied Animal Behaviour Science* 170, 34–43.
- Sadeghi M, Banakar A, Khazaei M and Soleimani MR 2015. An intelligent procedure for the detection and classification of chickens infected by *Clostridium perfringens* based on their vocalization. *Brazilian Journal of Poultry Science* 17, 537–544.
- Sales GT, Green AR, Gates RS, Brown-Brandt TM and Eigenberg RA 2015. Quantifying detection performance of a passive low-frequency RFID system in an environmental preference chamber for laying hens. *Computers and Electronics in Agriculture* 114, 261–268.
- Siegford JM, Berezowski J, Biswas SK, Daigle CL and Gebhardt-Henrich SG 2016. Assessing activity and location of individual laying hens in large groups using modern technology. *Animals* 6, 10. doi: [10.3390/ani6020010](https://doi.org/10.3390/ani6020010).
- Sih A, Bell AM, Johnson JC and Ziemba RE 2004. Behavioral syndromes: an integrative overview. *Quarterly Review of Biology* 79, 241–277.
- Silvera AM, Knowles TG, Butterworth A, Berckmans D, Vranken E and Blokhuis HJ 2017. Lameness assessment with automatic monitoring of activity in commercial broiler flocks. *Poultry Science* 96, 2013–2017.
- Stadig LM, Ampe B, Rodenburg TB, Reubens B, Maselyne J, Zhuang SJ, Criel J and Tuytens FAM 2018a. An automated positioning system for monitoring chickens' location: accuracy and registration success in a free-range area. *Applied Animal Behaviour Science* 201, 31–39.

- Stadig LM, Rodenburg TB, Ampe B, Reubens B and Tuytens FAM 2018b. An automated positioning system for monitoring chickens' location: effects of wearing a backpack on behaviour, leg health and production. *Applied Animal Behaviour Science* 198, 83–88.
- Taylor PS, Hemsworth PH, Groves PJ, Gebhardt-Henrich SG and Rault JL 2018. Ranging behavior relates to welfare indicators pre- and post-range access in commercial free-range broilers. *Poultry Science* 97, 1861–1871.
- Tullo E, Fontana I, Diana A, Norton T, Berckmans D and Guarino M 2017. Application note: labelling, a methodology to develop reliable algorithm in PLF. *Computers and Electronics in Agriculture* 142, 424–428.
- Valletta JJ, Torney C, Kings M, Thornton A and Madden J 2017. Applications of machine learning in animal behaviour studies. *Animal Behaviour* 124, 203–220.
- Van Hertem T, Rooijackers L, Berckmans D, Fernandez AP, Norton T, Berckmans D and Vranken E 2017. Appropriate data visualisation is key to Precision Livestock Farming acceptance. *Computers and Electronics in Agriculture* 138, 1–10.
- Wang C, Chen H, Zhang X and Meng C 2016. Evaluation of a laying-hen tracking algorithm based on a hybrid support vector machine. *Journal of Animal Science and Biotechnology* 7, 60. doi: [10.1186/s40104-016-0119-3](https://doi.org/10.1186/s40104-016-0119-3).
- Wathes CM, Kristensen HH, Aerts JM and Berckmans D 2008. Is precision livestock farming an engineer's daydream or nightmare, an animal's friend or foe, and a farmer's panacea or pitfall? *Computers and Electronics in Agriculture* 64, 2–10.
- Youssef A, Exadaktylos V and Berckmans DA 2015. Towards real-time control of chicken activity in a ventilated chamber. *Biosystems Engineering* 135, 31–43.
- Zaninelli M, Costa A, Tangorra FM, Rossi L, Agazzi A and Savoini G 2015. Preliminary evaluation of a nest usage sensor to detect double nest occupations of laying hens. *Sensors* 15, 2680–2693.
- Zaninelli M, Redaelli V, Luzi F, Bontempo V, Dell'Orto V and Savoini G 2017. A monitoring system for laying hens that uses a detection sensor based on infrared technology and image pattern recognition. *Sensors* 17, 1195. doi: [10.3390/s17061195](https://doi.org/10.3390/s17061195).
- Zaninelli M, Redaelli V, Luzi F, Mitchell M, Bontempo V, Cattaneo D, Dell'Orto V and Savoini G 2018. Development of a machine vision method for the monitoring of laying hens and detection of multiple nest occupations. *Sensors* 18, 132. doi: [10.3390/s18010132](https://doi.org/10.3390/s18010132).
- Zaninelli M, Redaelli V, Tirloni E, Bernardi C, Dell'Orto V and Savoini G 2016. First results of a detection sensor for the monitoring of laying hens reared in a commercial organic egg production farm based on the use of infrared technology. *Sensors* 16, 1757. doi: [10.3390/s16101757](https://doi.org/10.3390/s16101757).
- Zhang FY, Hu YM, Chen LC, Guo LH, Duan WJ and Wang L 2016. Monitoring behavior of poultry based on RFID radio frequency network. *International Journal of Agricultural and Biological Engineering* 9, 139–147.
- Zhuang XL, Bi MN, Guo JL, Wu SY and Zhang TM 2018. Development of an early warning algorithm to detect sick broilers. *Computers and Electronics in Agriculture* 144, 102–113.