

NOTES FOR AUTHORS

The *Bulletin of Entomological Research* publishes original research papers concerning insects, mites, ticks or other arthropods of economic importance in agriculture, forestry, stored products, biological control, medicine, animal health and natural resource management. The geographical scope of the *Bulletin* is worldwide but with emphasis on the tropics. Taxonomic papers are accepted if relevant. Short review papers, although normally by invitation, will also be considered for publication.

Page Format. The *Bulletin* is printed in a two-column format (column width of 80 mm) with a text area of 170×225 mm.

Text. Papers should be typed, on one side of the paper only, with double line spacing and ample margins (at least 1.5 cm) on each side and with no underlining or bold in text except for scientific names. Draft quality print from a word-processor is not acceptable. Standard abbreviations (e.g. fig. and figs) and metric units must be used. Guidelines for taxonomic papers are available.

When the paper has been accepted word-processed text stored on floppy disk is encouraged, providing the software is IBM/DOS compatible, but floppy discs must be accompanied by a hard copy. This will enable papers to be handled rapidly, and with fewer type-setting errors.

Abstract. Each paper must commence with a carefully prepared, accurate, informative abstract, in one paragraph, that is complete in itself and intelligible without reference to text or figures. It should not exceed 250 words. A short title should be provided as a running head.

Tables. Tables should be reduced to the simplest form, and should not be used where text or illustrations give the same information. They should be submitted on separate sheets at the end of the article and must fit conveniently into single column, full width or landscape (if absolutely necessary) format. Table captions should be typed on a separate sheet.

Illustrations. Copies only of artwork should be submitted. The original illustrations should accompany the paper after acceptance and revision. Text figures, line drawings, computer-generated figures and graphs should be of sufficient size and quality to allow for reduction by half or two-thirds. Half-tone photographs are acceptable where they are a real contribution to the text. Figure and captions should be typed on a separate sheet in

the following format:

Figs 23–26. Figs 23–24, Urophora eggs: 23, U. hispanica; 24, U. stigma. Figs 25–26, spermathecae: 25, U. maura; 26, U. stigma; scale lines = 0.05 mm.

Voucher specimens. The deposition of voucher specimens should be considered where appropriate.

References. References must be based on the name and year system, give full journal titles and conform to the following styles:

Powell, W. (1986) Enhancing parasitoid activity in crops. pp. 319–340 in Waage, J. & Greathead, D. (Eds) *Insect parasitoids*. London, Academic Press (Symposium, Royal Entomological Society of London No. 13).

Southwood, T.R.E. (1978) *Ecological methods with particular reference to the study of insect populations*. 2nd edn. 524 pp. London, Chapman & Hall.

Zhou, X., Carter, N. & Mumford, J. (1989) A simulation model describing the population dynamics and damage potential of the rose grain aphid, Metopolophium dirhodum (Walker) (Hemiptera: Aphididae), in the UK. *Bulletin of Entomological Research* 79, 373–380.

Citation of authors in the text should appear in the form: Polaszek (1990) or (Polaszek, 1990). More than one author should be cited in chronological order as: (Holloway *et al.*, 1987; Walker & Huddleston, 1988).

Offprints. 50 copies of each paper are provided free to the author (or major author) of each paper. Further copies may be obtained on payment, and the number required should be specified and ordered at proof stage.

Manuscripts. Three copies of the manuscript and artwork should be submitted to:

The Editors
Bulletin of Entomological Research
International Institute of Entomology
56, Queen's Gate
London
SW7 5JR, UK.

Bulletin of Entomological Research

Baker, P. S., Ley, C., Balbuena, R. & Barrera, J. F. Factors affecting the emergence of <i>Hypothenemus hampei</i> (Coleoptera: Scolytidae) from coffee berries	145
Black, W. C. IV., DuTeau, N. M., Puterka, G. J., Nechols, J. R. & Pettorini, J. M. Use of the random amplified polymorphic DNA polymerase chain reaction (RAPD-PCR) to detect DNA polymorphisms in aphids	151
Blackman, R. L. & Spence, J. M. Electrophoretic distinction between the peach-potato aphid, <i>Myzus persicae</i> , and the tobacco aphid, <i>M. nicotianae</i> (Homoptera: Aphididae)	161
Brightwell, R., Dransfield, R. D. & Williams, B. G. Factors affecting seasonal dispersal of the tsetse flies <i>Glossina pallidipes</i> and <i>G. longipennis</i> (Diptera: Glossinidae) at Nguruman, south-west Kenya	167
Busch-Petersen, E. & Baumgartner, H. Simulation of instability in genetic sexing strains: effects of wild-type contamination in association with other biological parameters	183
Disney, R. H. L., Khochare, N. N. & Godase, S. K. Jr. A new parasitoid (Diptera: Phoridae) of the mango shoot borer, <i>Chlumetia transversa</i> (Lepidoptera: Noctuidae), in India	191
Donnelly, D., Calitz, F. J. & Van Aarde, I. M. R. Insecticidal control of <i>Melanterius servulus</i> (Coleoptera: Curculionidae), a potential biocontrol agent of <i>Paraserianthes lophantha</i> (Leguminosae), in commercial seed orchards of black wattle, <i>Acacia mearnsii</i> (Leguminosae)	197
Guglielmone, A. A. The effect of temperature and humidity on development and longevity of <i>Amblyomma triguttatum triguttatum</i> (Acarina: Ixodidae)	203
Kamhawi, S., Lane, R. P., Cameron, M., Phillips, A., Milligan, P. & Molyneux, D. H. The cuticular hydrocarbons of <i>Phlebotomus argentipes</i> (Diptera: Phlebotominae) from field populations in northern India and Sri Lanka, and their change with laboratory colonization	209
Löhr, B. The pugnacious ant, <i>Anoplolepis custodiens</i> (Hymenoptera: Formicidae), and its beneficial effect on coconut production in Tanzania	213
Millest, A. L., Cheke, R. A., Howe, M. A., Lehane, M. J. & Garms, R. Determining the ages of adult females of different members of the <i>Simulium damnosum</i> complex (Diptera: Simuliidae) by the pteridine accumulation method	219
Natochin, Yu. V., Parnova, R. G., Sundukov, O. V. & Shakhmatova, E. I. The effect of insecticide treatment and injection of sodium chloride and sucrose solutions on the water-salt homeostasis of lepidopterous larvae	227
Price, R. E. & Brown, H. D. Incubation and overwintering in the egg stage of the African migratory locust, <i>Locusta migratoria migratorioides</i> (Orthoptera: Acrididae), on the highveld of South Africa	235
Riley, J. R., Armes, N. J., Reynolds, D. R. & Smith, A. D. Nocturnal observations on the emergence and flight behaviour of <i>Helicoverpa armigera</i> (Lepidoptera: Noctuidae) in the post-rainy season in central India	243
Sommer, C., Steffansen, B., Overgaard Nielsen, B., Grønvold, J., Vagn Jensen, K.-M., Brøchner Jespersen, J., Springborg, J. & Nansen, P. Ivermectin excreted in cattle dung after subcutaneous injection pour-on treatment: concentrations and impact on dung fauna	257
Strong, L. Avermectins: a review of their impact on insects of cattle dung	265
White, I. M. & Wang, X. j. Taxonomic notes on some dacine (Diptera: Tephritidae) fruit flies associated with citrus, olives and cucurbits	275
Witz, J. A., Lopez, J. D. Jr. & Latheef, M. A. Field density estimates of <i>Heliothis virescens</i> (Lepidoptera: Noctuidae) from catches in sex pheromone-baited traps	281
Book Reviews	287

© C.A.B International, 1992

All rights reserved. No part of this publication may be reproduced, in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without prior permission of the copyright owner.