



Chapter III: Astronomical Observations

About the atomic and molecular databases in the planetary community – A contribution in the Laboratory Astrophysics Data WG IAU 2022 GA session

M. Rengel^{ID}

Max-Planck-Institut für Sonnensystem Forschung, Justus-von-Liebig-Weg 3, 37077 Göttingen,
Germany,
email: rengel@mps.mpg.de

Abstract. This paper corresponds to an invited oral contribution to the session 5A organised by the IAU inter-commission B2-B5 working group (WG) “Laboratory Astrophysics Data Compilation, Validation and Standardization: from the Laboratory to FAIR usage in the Astronomical Community” at the IAU 2022 General Assembly (GA) Rengel (2022). This WG provides a platform where to discuss the Findability, Accessibility, Interoperability, Reuse (FAIR) usage of laboratory Atomic and Molecular (A&M) data in astronomy and astrophysics.

A&M data play a key role in the understanding of the physics and chemistry of processes in several research topics, including planetary science and interdisciplinary research in particular the atmospheres of planets and planetary explorations, etc. Databases, compilation of spectroscopic parameters, and facility tools are used by computer codes to interpret spectroscopic observations and simulate them. In this talk I presented existing A&M databases of interest to the planetary community focusing on access, organisation, infrastructures, limitations and issues, etc.

Keywords. planets, atmospheres, exoplanets, atomic data, molecular data, laboratory astrophysics, experiment, databases, data network, data analysis

1. Introduction

The talk represented a tour on A&M databases used in the planetary and exoplanetary communities not from the perspective of a developer, but of an user. A&M databases compile and provide detailed spectral information for A&M to feed codes that predict and simulate radiation in gaseous media. Between the applications, we find atmospheric physics: (exo) planetary atmospheres, comets and small bodies. These databases are a critical input for the codes which predict and interpret spectra of planetary atmospheres (hydrostatic equilibrium atmospheres and expanding comas), and space and ground-based telescopes facilities depend on the quality and extent of reference A&M parameters.

Line lists typically contain hundreds to billions of individual transitions. A&M databases contains detailed A&M spectroscopic parameters of A&M like pressure-broadening (shapes), collision-induced absorption (CIA), transition intensity or cross sections, line shape parameters, rotation-vibration transition position (wavelength, frequency), parameters to describe how these vary with temperature and pressure, aerosol indices of refraction, microphysical and optical properties of atmospheric aerosols, for example.

© The Author(s), 2024. Published by Cambridge University Press on behalf of International Astronomical Union. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

2. Some existing A&M databases - Data and file formats

Several groups worldwide generate and compile A&M data through measurement and/or calculation (e.g. HITRAN[†] [Gordon *et al.* (2022)], GEISA[‡] [Jacquinot-Husson *et al.* (2016)], JPL Molecular Spectroscopy[§], CDMS[¶] [Endres *et al.* (2016)], VAMDC || [Dubernet *et al.* (2010), Dubernet *et al.* (2016), Albert *et al.* (2020)], ExoMol** [Tennyson *et al.* (2020)], HITEMP^{††} [Rothman *et al.* (2010)], VALD <http://vald.astro.uu.se>, MoLLIST <http://bernath.uwaterloo.ca/molecularlists.php>, Ames Molecular Spectroscopic Data for Astrophysical and Atmospheric Studies (<http://huang.seti.org>, TheoReTs <https://theorets.univ-reims.fr/>, etc.). Several secondary databases and information services are fed with data from such sources in a fragmented manner. A variety of data formats (cross sections, K-tables, line-by-line, super-lines) and file formats (e.g. .hdf5, .pickle, .mp4, .txt, .npy) are generated. There are online tools such as HAPI to extend functionalities <https://hitran.org/hapi/> [Kochanov *et al.* (2016)], and exo-k library to handle radiative opacities <https://pypi.org/project/exo-k/> [Leconte (2021)], that enable conversion between different formats. As part of the spectroscopic input to atmospheric codes, the HITRAN molecular spectroscopic database is already internationally recognised as standard in the planetary community, and the ExoMol database, valid over extended temperature ranges, is widely used in the exoplanetary community.

3. Some (exo)planet atmospheric radiative transfer and inversion codes

One commonly used way to interpret the measured spectra is calculating a synthetic spectrum for comparison with that measured by solving the radiative transfer (forward model) -i.e., computation of the outgoing radiation from the planetary surface for a given set of free parameters-, and inferring parameters like temperature and abundance. This last step is called inversion and consists in comparing the measured and best modelled spectra adjusting the atmospheric parameters in such a way as to minimise any discrepancy. A number of radiative transfer codes or forward models and inversion algorithms (retrieval technique) are already generally available and used by the planetary and exoplanetary characterisation communities (Table 1). When calculating a synthetic atmospheric spectrum, lines are read from A&M databases. The retrieval or parameter fitting techniques commonly used are Optimal Estimation algorithm, nested sampling, Markov chain Monte Carlo method, and Grid search. Some examples of applications of such algorithms are given in [Rengel *et al.* (2008); Hartogh *et al.* (2010); Rengel *et al.* (2014); Shulyak *et al.* (2019); Shulyak *et al.* (2020); Rengel *et al.* (2022); Villanueva *et al.* (2022)]

4. Discussion: Needs and wish-list

In spite of the tremendous advances and current efforts in the generation of databases, there are still improvements in progress. A growing demand for spectroscopic data for (exo) planetary studies and other atmospheres is being driven by scientists who are interested in modelling as well as observing diverse bodies. Line lists are generated from experiments and/or ab initio calculations and may be incomplete or contain errors.

[†] <https://hitran.org>
[‡] <https://geisa.aeris-data.fr/>
[§] <https://spec.jpl.nasa.gov>
[¶] <https://cdms.astro.uni-koeln.de>
|| <https://vamdc.org>
** <https://www.exomol.com>
^{††} <https://hitran.org/hitemp>

Table 1. Some radiative transfer and inversion codes used in the (exo) planetary community.

Code name	Link	Reference
ARCiS	http://www.exoclouds.com	Min <i>et al.</i> (2020)
TauReX	https://taurex3-public.readthedocs.io/en/latest/	Al-Refaie <i>et al.</i> (2021)
NEMESIS	https://users.ox.ac.uk/atmp0035/nemesis.html	Irwin <i>et al.</i> (2008)
petitRADTRANS	https://petitradtrans.readthedocs.io/en/latest/	Mollière <i>et al.</i> (2019)
PSG	https://psg.gsfc.nasa.gov/	Villanueva <i>et al.</i> (2018)
CHIMERA	https://github.com/mrline/CHIMERA	Line <i>et al.</i> (2013)
PLATON	https://github.com/ideasrule/platon	Zhang <i>et al.</i> (2020)
ATMO	https://www.erc-atmo.eu	Tremblin <i>et al.</i> (2015)
MOLIERE	Urban <i>et al.</i> (2012)	Urban (2004)
ARTS	https://radiativetransfer.org/	Buehler <i>et al.</i> (2018)
Home made (MPS)	-	Jarchow, C. (1998)
Helios-r2	https://github.com/exoclime/Helios-r2	Kitzmann <i>et al.</i> (2020)
SCARLET	-	Benneke (2015)
BART	https://github.com/exosports/BART	Blecic (2016)
INARA	https://gitlab.com/frontierdevelopmentlab/astrobiology/inara	Soboczenski <i>et al.</i> (2018)

Databases differ in completeness, and some ones do not accurately characterise high-frequency spectral regions. Sometimes there are no datasets for a specific problem at hand. Atmospheric codes used by the planetary and exoplanetary characterisation communities, that are designed to solve the radiative transfer equation by looking at the propagation of radiation through a medium and simulate observations and infer parameters, have their own methods for the computation of opacities and there are no community standards. Furthermore, there are also no community standards in the selection of atmospheric codes in mission planning. There are needs to increase accessibility of opacities (computation, access, visualisation, manipulation), laboratory measurements of molecular cross-sections, and pressure broadening description for some species, among many other aspects.

Furthermore, going into details, the community identifies needs in the following aspects: isotopologues: CIA (more lists of N₂O, CH₄, SO), expansion of CIA of secondary species for wider temperature and wavelength ranges, additional experiments for CO intensities, improvement of the CH₄ quality of the line shape parameters, continuum absorption by water vapour, partition functions at higher temperatures for all species (around 5000 K), more friendly ways implementing new data in the RT codes, more aerosol refractive indexes above 500-600 K, more saturation vapour pressure data, kinetic data at high temperature, Rayleigh scattering for non-H₂, collisional xsecs (in particular: mid-sized organics, H₂O⁺, diatom-H₂, CH₃OH, HCN, SO₂, CH₂) and high-resolution xsecs (R=1E6+) atmospheres, organic sulphide gases for the line lists (biomarkers).

Acknowledgments

I thank Sergey Yurchenko and Iouli Gordon for the input and discussions. I thank also the members of the SOC of the session, and the members of the inter-commission B2-B5 working group in special to Marie-Lise Dubernet. I acknowledge the support by the DFG priority program SPP 1992 “Exploring the Diversity of Extrasolar Planets” (DFG PR 36 24602/41).

For the purpose of Open Access, a CC-BY-SA 4.0 public copyright licence has been applied by the author to the present document and will be applied to all subsequent versions up to the Author Accepted Manuscript arising from this submission.

References

- Albert, D., Anthony, B.L., Ba, Y.A. *et al.* 2020, *Atoms*, 8, 76 10.3390/atoms8040076
- Al-Refaie, A. F., Changeat, Q., Waldmann, I. P., *et al.* 2021, *Astrophysical Journal*, 917, 37. doi:10.3847/1538-4357/ac0252
- Benneke, B. 2015, arXiv:1504.07655
- Blecic, J. 2016, arXiv:1604.02692
- Buehler, S. A., Mendrok, J., Eriksson, P., *et al.* 2018, *Geoscientific Model Development*, 11, 1537. doi:10.5194/gmd-11-1537-2018
- Dubernet, M.-L., Boudon, V., Cullhane, J. L., *et al.* 2010, *JQRST*, 111, 2151-2159; 10.1016/j.jqsrt.2010.05.004
- Dubernet, M.-L., Antony, B. K., Ba, Y.-A., *et al.* 2016, *J. of Physics B*, 49, 074003 10.1088/0953-4075/49/7/074003
- Endres, C. P., Schlemmer, S., Schilke, P., *et al.* 2016, *Journal of Molecular Spectroscopy*, 327, 95. doi:10.1016/j.jms.2016.03.005
- Gordon, I. E., Rothman, L. S., Hargreaves, R. J., *et al.* 2022, *JQSRT*, 277, 107949 10.1016/j.jqsrt.2021.107949
- Hartogh, P., Blecka, M. I., Jarchow, C., *et al.* 2010, *A&A*, 521, L48. doi:10.1051/0004-6361/201015159
- Irwin, P. G. J., Teanby, N. A., de Kok, R., *et al.* 2008, *J. Quant. Spectrosc. and Rad. Trans*, 109, 1136. doi:10.1016/j.jqsrt.2007.11.006
- Jacquinet-Husson, N., Armante, R., Scott, N. A., *et al.* 2016, *Journal of Molecular Spectroscopy*, 327, 31. doi:10.1016/j.jms.2016.06.007
- Jarchow, C., 1998. Bestimmung atmosphärischer Wasserdampf- und Ozone profile mittels bodengebundener Millimeterwellen- Fernerkundung. Ph.D. thesis.
- Kitzmann, D., Heng, K., Oreshenko, M., *et al.* 2020, *The Astrophysical Journal*, 890, 174. doi:10.3847/1538-4357/ab6d71
- Kochanov, R. V., Gordon, I. E., Rothman, L. S., *et al.* 2016, *JQSRT*, 177, 15. doi:10.1016/j.jqsrt.2016.03.005
- Leconte, J. 2021, *A&A*, 645, A20. doi:10.1051/0004-6361/202039040
- Line, M. R., Wolf, A., Zhang, X., *et al.* 2013, *The Astrophysical Journal*
- Min, M., Ormel, C. W., Chubb, K., *et al.* 2020, *A&A*, 642, A28. doi:10.1051/0004-6361/201937377
- Mollière, P., Wardenier, J. P., van Boekel, R., *et al.* 2019, *A&A*, 627, A67. doi:10.1051/0004-6361/201935470
- Rengel, M., Hartogh, P., & Jarchow, C. 2008, *Planet. Space Sci*, 56, 1368. doi:10.1016/j.pss.2008.07.004
- Rengel, M., Sagawa, H., Hartogh, P., *et al.* 2014, *A&A*, 561, A4. doi:10.1051/0004-6361/201321945
- Rengel, M., Shulyak, D., Hartogh, P., *et al.* 2022, *A&A*, 658, A88. doi:10.1051/0004-6361/202141422
- Rengel Miriam. (2022, September 1). About the atomic and molecular databases in the planetary community. Zenodo. <https://doi.org/10.5281/zenodo.7040446>
- Rothman, L. S., Gordon, I. E., Barber, R. J., *et al.* 2010, *JQSRT*, 111, 2139. doi:10.1016/j.jqsrt.2010.05.001
- Shulyak, D., Rengel, M., Reiners, A., *et al.* 2019, *A&A*, 629, A109. doi:10.1051/0004-6361/201935691
- Shulyak, D., Lara, L. M., Rengel, M., *et al.* 2020, *A&A*, 639, A48. doi:10.1051/0004-6361/201937210
- Soboczenski, F., Himes, M. D., O'Beirne, M. D., *et al.* 2018, arXiv:1811.03390
- Tennyson, J., Yurchenko, S. N., Al-Refaie, A. F., *et al.* 2020, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 255, 107228 10.1016/j.jqsrt.2020.107228
- Tremblin, P., Amundsen, D. S., Mourier, P., *et al.* 2015, *The Astrophysical Journal*, 804, L17. doi:10.1088/2041-8205/804/1/L17

- Urban, J. 2004, Journal of Quantitative Spectroscopy and Radiative Transfer, 83, 529.
doi:10.1016/S0022-4073(03)00104-3
- Urban, J., Baron, P., & Lautie, N. 2012, Astrophysics Source Code Library. ascl:1212.004
- Villanueva, G. L., Smith, M. D., Protopapa, S., *et al.* 2018, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 217, 86. doi:10.1016/j.jqsrt.2018.05.023
- Villanueva, G. L., Liuzzi, G., Faggi, S., *et al.* 2022, Fundamentals of the Planetary Spectrum Generator. 2022 edition of the handbook by G.L. Villanueva *et al.* ISBN 978-0-578-36143-7, 2022
- Zhang, M., Chachan, Y., Kempton, E. M.-R., *et al.* 2020, The Astrophysical Journal, 899, 27.
doi:10.3847/1538-4357/aba1e6