

Dietary patterns and breast-feeding in Australian children

Jessica A Grieger*, Jane Scott and Lynne Cobiac

Department of Nutrition and Dietetics, Flinders Clinical and Molecular Medicine, Level 7E, Flinders Medical Centre, Flinders Drive, Bedford Park, SA 5042, Australia

Submitted 18 May 2010; Accepted 4 April 2011; First published online 23 June 2011

Abstract

Objective: To determine the dietary patterns of a national sample of 2–8-year-old Australian children and to establish whether breast-feeding is associated with dietary patterns in this age group.

Design: Cross-sectional study using 24 h recall data from the 2007 Australian National Children's Nutrition and Physical Activity Survey.

Setting: Australia.

Subjects: A total of 2287 children aged 2–8 years.

Results: Principal component factor analysis identified three distinct patterns. The 'Non-core food groups' pattern included food groups such as whole-fat dairy products, cheese, medium–high sugar-sweetened breakfast cereals and sweet biscuits, no fruit, reduced/low-fat dairy products and wholegrain bread/rolls. The 'Healthy, meat and vegetable' pattern included vegetables, red meat, fruit and wholegrain bread/rolls and was inversely associated with take-away foods and carbonated sugar-sweetened beverages. The 'Combination' pattern contained many food groups including candy (not chocolate based), pasta/rice products, nuts/seeds, cakes and chocolate, but no fruit or vegetables. Of the 2287 children, 2064 (89.3%) had been breast-fed. A positive association was found between breast-feeding and the healthy, meat and vegetable pattern ($r=0.267$) but not with the other two patterns. Higher scores on this pattern were also associated with younger age, lower BMI, higher birth weight, high likelihood of being in the less-disadvantaged Socio-economic Indexes for Areas category and less likelihood of the child's parents having a lower educational level.

Conclusions: These results provide suggestive evidence that breast-feeding during infancy is associated with a healthy dietary pattern in childhood and offers a likely pathway to explain the previously reported association between breast-feeding and chronic disease.

Keywords
Factor analysis
Children
Breast-feeding
Eating patterns

Children's food choices and preferences develop from a young age and may be influenced by genetic predisposition towards certain tastes, by food availability and by cultural and parental influences^(1–3). There is evidence that behaviours and attitudes inculcated at a young age track through childhood and into adulthood^(4–6). Therefore, the importance of establishing healthy eating habits in early life is crucial to reducing the risk for longer-term diet-related diseases in later years.

Experimental studies have revealed that infants who are breast-fed experience flavours in their mother's milk before they are exposed to the same flavours in solid foods^(7,8) and that breast-fed infants have a greater acceptability of new foods and flavours at 2–8 months of age compared with formula-fed infants^(9–12). For example, breast-fed infants have been reported to have a greater diversity in foods and to consume less cordial and juice at 9 months of age⁽¹³⁾. Furthermore, there is convincing evidence that, compared with formula-fed

infants, breast-fed infants have a reduced risk of being overweight or obese and of developing type 2 diabetes during child- and adulthood^(14,15). This suggests that postnatal programming in infancy has an effect on feeding preferences and practices, which may subsequently affect health outcomes during later life.

A healthy diet is important for the prevention and management of a number of chronic diseases⁽¹⁶⁾. However, it is of concern that, in young children, nutrient intakes, particularly Ca, folate, K and Zn^(17–19), and intakes of fruit and vegetables^(20,21) have been reported to be low, whereas intakes of take-away foods and soft drinks have been reported to be high⁽²²⁾. Some studies have reported associations between unhealthier dietary patterns of children and various factors such as fat mass⁽²³⁾, sociodemographic status^(24–26) and lower levels of education of the main food provider⁽²⁵⁾.

There appears to be a link between breast-feeding and diet in infants; however, it is not known whether

*Corresponding author: Email jessica.grieger@flinders.edu.au

breast-feeding is associated with food choices and dietary patterns during later childhood. Therefore, the aims of the present study were to determine the dietary patterns of a national sample of 2–8-year-old Australian children and to establish whether breast-feeding is associated with dietary patterns in this age group.

Methods

Study population and procedures

The 2007 Australian National Children's Nutrition and Physical Activity Survey was a national survey involving 4487 children aged between 2 and 16 years. Children and their parents were interviewed in their home during which time the children's food and beverage intakes and use of supplements were assessed using a 24 h food recall and a food habits survey. At this time, their height, weight and waist circumference were measured and further information was collected on their level of physical activity and demographic characteristics.

Food and nutrient intake data were collected on two occasions using a 24 h food recall. A computer-assisted personal interview (CAPI) was conducted in the child's home and was followed 7–21 d later by a computer-assisted telephonic interview (CATI). All field interviewers were trained in recruitment procedures and in dietary assessment, physical activity and anthropometry measures. Age- and sex-specific BMI cut-offs for normal weight, overweight and obese children and for adolescents were applied to the data⁽²⁷⁾. For underweight participants, grade-3 thinness (corresponding to an adult BMI of 18.5 kg/m²) was used as the cut-off point⁽²⁸⁾.

Sample selection

The survey was conducted using a quota-sampling scheme stratified by state/territory and by capital city statistical division or by the rest of the state/territory. Postcodes were used as the primary sampling units in each state, allocated to a stratum using the Australian Bureau of Statistics (ABS) postal area-to-statistical local area concordance. Random digit dialling was used to recruit households (private dwellings) from selected postcodes to the survey. The telephone number prefix acted as a 'geographical indicator' that corresponded to the postcode. Households with children aged between 2 and 16 years were identified and were asked to participate. One child within the household was selected as the 'study child' using a defined method for children's selection. For example, using a predefined Kish table (refer to user guide for additional details), the parent or caregiver was asked for the name, gender and age in years of each child in the household aged 2–16 years, arranging the children by age (from the oldest to the youngest). As the number of children required in the sample for each of the age cohorts was not proportional to the number of children in each of the age cohorts in the population, children aged

2–3 or 14–16 years had a higher probability of selection compared with children aged 4–13 years in any one household. Therefore, the Kish table showed a bias towards selecting children who were aged 14–16 years, followed by children aged 2–3 years and finally children aged 4–13 years. In cases where the age and gender quota for a particular location had already been filled, recruitment of the study child did not proceed. Some postcodes were excluded if they were in very remote areas, having few in-scope children according to the 2001 ABS Census.

Further details of the survey methodology and sample design can be obtained from the 2007 Australian National Children's Nutrition and Physical Activity Survey Users Guide available at [http://www.health.gov.au/internet/main/publishing.nsf/Content/AC3F256C715674D5CA2574D6000237D/\\$File/user-guide-v2.pdf](http://www.health.gov.au/internet/main/publishing.nsf/Content/AC3F256C715674D5CA2574D6000237D/$File/user-guide-v2.pdf).

The current analysis was focused on 2–8-year-old children using the 24 h dietary food recall data collected through the CAPI from the primary caregiver of the child.

Food group consumption

Dietary intake information collected from the CAPI 24 h food recall generated a large number of foods. In order to process and clean the data and identify meaningful dietary patterns, a number of foods were eliminated that were considered unlikely to contribute to a dietary pattern according to the following criteria. Food items that were not included in the analysis as they contributed to a cumulative <10% frequency of all individual food items consumed were: dietary/multivitamin and oil supplements; gelatin; meat substitutes; vegetable-based chutneys/dressings/relishes; energy drinks; essences/baking powder; flour; yeast; stuffing; alcohol; chewing gum; soya beverages; infant drinks/formulae; pretzels; unspecified fats; mature legumes and pulses; chives; garlic; ginger; electrolyte and fortified drinks; extruded or reformed snacks; and batter-based products. A frequency of the 'gram intake' for each food was calculated. Individual foods were eliminated from the analysis when the amount consumed was <3rd percentile of the study population (e.g. <20 g for meat/poultry or <5 g for vegetables). It was believed that intakes in these amounts were unlikely to make a significant contribution to a dietary pattern.

A total of forty-five food groups were created that were mutually exclusive and were included in the analysis (Table 1). Frequency of consumption was binary coded to 0 (not consumed) or 1 (consumed).

Factor analysis

Dietary patterns were derived using factor analysis with factor loadings extracted using the principal component method and Varimax/orthogonal rotation. Factor loadings for binary variables are usually underestimated and therefore tetrachoric correlation coefficients were calculated to estimate the correlation between the continuous versions of the variables. The number of dietary patterns

Table 1 Food groups and the description of foods used in the factor analysis

Food group	Description of food
Non-carbonated/non-nutritive drinks	Non-carbonated drinks containing minimal kilojoules (i.e. intense sweetened/low-joule fruit juice/cordial)
Fruit juice	Juice (100 % juice; NFS, homemade)
Cordial/fruit drinks	Cordials (<25 % fruit juice, 25 % fruit juice, 40 % juice, NFS), fruit drink (25 % juice, 35 % juice), mineral water with fruit juice (5 %)
Carbonated, non-nutritive drinks	Intense sweetened drinks (e.g. diet coke, carbonated water with flavouring)
Carbonated sugar-sweetened drinks	Cola, flavoured soft drinks
Mineral water and water	All water (non-carbonated, unflavoured)
Tea/coffee	Tea/coffee base or beverage (prepared mainly with water)
Milk: whole fat	Whole-fat milk (cow, sheep, goat, flavoured)
Milk: non-whole fat	Skimmed, low-fat, reduced fat milk
Yoghurt: whole fat	Whole-fat yoghurt
Yoghurt: non-whole fat	Low-fat, reduced-fat or intense/artificially sweetened yoghurt
Cheese	Cheese, cream cheese, cream cheese dips
Other dairy products	Ice cream, ice confection (milk based), ice cream desserts (with nuts, toppings, etc.), condensed milk, evaporated milk
Vegetables: orange	Pumpkin, carrot, sweet potato
Vegetables: others	All other vegetables, mixed vegetables (frozen or fresh)
Potatoes: take-away	Potatoes bought as take-away (e.g. fries/hash brown from McDonalds, KFC)
Potatoes: home baked, no oil	Potatoes that were cooked at home without oil/fat (e.g. dry, baked, frozen wedges/chips, boiled potatoes, roast potatoes)
Potatoes: home baked, with oil	Potatoes that were cooked at home using oil/fat (e.g. frozen wedges/chips, roast potatoes, mashed potatoes with butter/margarine)
Fruit	Fresh, canned, dried, pureed
Savoury snacks	Potato crisps, rice/wheat biscuits, rice cakes, crisp-bread
Muesli bars and cereal bars	Muesli bars and cereal bars, specified as muesli/breakfast bars. May contain chocolate chips or are coated in yoghurt but not coated in chocolate
Bread/bread rolls: white	White-flour bread/bread rolls, cheese/ham-topped rolls, foccacia, pita bread
Bread/bread rolls: non-white	Rye, wholemeal, wholegrain bread/bread rolls
Breakfast cereals: low-medium sugar sweetened (<15 g sugar/100 g)	Cereals (including oats) containing <15 g sugar/100 g
Breakfast cereals: medium-high sugar sweetened (≥15 g sugar/100 g)	Cereals containing ≥15 g sugar/100 g
Soups	Vegetable, meat, chicken: prepared with water/milk, homemade
Pasta and pasta products/rice and rice products	Pasta dishes, rice dishes, plain/boiled rice, plain/boiled pasta, noodles, lasagne, risotto, cannelloni
Cereal-based dishes	Pancakes, pizza, quiche, sandwiches, crumpets, pikelets, savoury scones, burrito, meat pie, sausage roll, pastie
Eggs and egg-based dishes	Eggs, whole, fried, omelette
Seafood and seafood dishes	Fresh/frozen fish, seafood, crustaceans, fish pie
Take-away/restaurant foods	Any main food item bought from a fast-food outlet (but with a minor component – e.g. lettuce, listed as a vegetable), ready-to-eat foods eaten away from home (e.g. meat pie, pastie, sausage roll)
Nuts and seeds	Nuts, seeds
Sweets: cakes	Cake, muffins, sweet scones, buns, fruit/sweet pies, doughnuts
Sweets: candy (not chocolate based)	Jelly confectionary (hard/soft), caramel, liquorice, marshmallow, chocolate-coated confectionary, nougat, cones (wafer)
Sweets: chocolate	Chocolate, chocolate bars, chocolate-coated bars
Sweets: biscuits	Biscuits, slices
Edible oils and spreads	Butter, margarine, edible oil and spreads
Other spreads	Peanut butter, vegemite, honey, jam, Nutella
Sauces	Tomato sauce, tomato paste, cranberry sauce, pasta sauces
Added sugar	When sugar was listed in quantities >10 g
Beverage bases	Milo, Ovaltine, Nesquick
White meat: chicken/turkey	Mince Sausage Patty/burger Cut: breast, thigh, wing, drumstick
Pork	Mince Sausages Roast/fillet/leg Chop/spare ribs
Red meat: beef/lamb/veal	Mince Sausages Cut/steak/rump Roast Burger/patty
Luncheon and deli-type meats	Frankfurts (not bought at take-away outlets), luncheon meats, deli-meats, ham, kabana, bacon

NFS, not further specified.

identified was based on eigenvalues > 1.5 on identification of a break point in the scree plot and on interpretability⁽²⁹⁾. Items were loaded on a factor if they had a correlation > 0.25 with that factor. These items represent the foods most highly related to the identified factor^(30,31). Foods that cross-loaded on several factors were also retained. Inter-item reliability for each factor was assessed using Cronbach's α coefficients.

Initially, a factor analysis was conducted separately for the 2–3-year-olds and 4–8-year-olds; however, food groups that loaded on the patterns were similar, and therefore the age groups were combined and the analysis re-run. Three distinct dietary patterns emerged and all children received a dietary pattern score for each of the identified patterns. A confirmatory factor analysis was also conducted using a second 24 h recall from the CATI data in which similar food groups loaded on a three-factor solution (data not shown).

Food habits survey

As part of the interview process, a 'food habits' survey was completed by the child (if aged > 9 years) or by a parent/caregiver during the CAPI. The survey included questions related to diet and food habits (e.g. type of milk used, servings of fruit/vegetables normally consumed, who prepares meals, is salt added to meals) with a series of answers pertaining to each question that the parent/caregiver could choose from. Four questions were included on breast-feeding, such as whether the child had ever been breast-fed; duration of breast-feeding; age of introduction of solids; and if, and at what age, the child received formula. For the current data analysis, breast-feeding was categorised into a dichotomous variable ('ever' or 'never' breast-fed).

Socio-economic Indexes for Areas

The Socio-economic Indexes for Areas (SEIFA) ranks geographical areas across Australia according to a 'score' that is created for the area on the basis of the characteristics of people, families and dwellings within that area⁽³²⁾. Every geographical area in Australia is given a SEIFA score for each index, which shows how disadvantaged that area is compared with other areas in Australia. Lower scores

indicate greater disadvantage⁽³²⁾. The SEIFA index was categorised into tertiles.

Statistical analyses

All analyses were performed using the STATA statistical software package for Windows version 10.1 (StataCorp., College Station, TX, USA) with population weights applied. Continuous data were assessed for normality and, if required, normalised with natural log transformation. Back transformations from the logarithmic scale to the original scale were undertaken in the multivariate regression models. Data are reported as OR and 95% CI. Only significant covariates in the final regression models are reported. The α level for significance was set at $P < 0.05$.

Results

A total of 2287 children were included in the analysis (51% male). Selected characteristics (mean and SEM) of the children are reported in Table 2. The median duration of breast-feeding and the interquartile range were 30.0 and 8.2–51.6 weeks, respectively; 223 children (10.7%) were not breast-fed.

Factor analysis

The dietary pattern analysis revealed three distinct patterns that explained 4.2%, 4.1% and 3.7% of the variance for each respective pattern. The first pattern was labelled 'Non-core food groups', with food groups such as whole-fat dairy products, cheese, luncheon and deli-type meats, medium-high sugar-sweetened breakfast cereals and sweet biscuits loading on this pattern (Table 3). Some food groups that were inversely associated with this pattern were reduced/low-fat dairy products, wholegrain/wholemeal breads and non-carbonated, non-nutritive beverages. The second pattern was labelled 'Healthy, meat and vegetable' as it included two types of vegetables, red meat, fruit and wholegrain/wholemeal breads. Food groups that were inversely associated with this pattern were potatoes from take-away outlets, carbonated sugar-sweetened beverages, take-away foods and added sugar. The final dietary pattern

Table 2 Characteristics of the children

	2–3-year-old children (n 1071)		4–8-year-old children (n 1216)		P value
	Mean or n	SEM or %	Mean or n	SEM or %	
Age (years)	2.5	0.0	6.0	0.0	< 0.001
Body weight (kg)	15.5	0.1	24.2	0.2	< 0.001
BMI (kg/m ²)	16.7	0.0	16.6	0.1	0.320
Obese*	39	4	65	5	0.504
Overweight*	174	1	170	14	0.487
Normal weight*	812	7	931	77	0.552
Underweight/very underweight*	46	4	50	4	0.690
Duration of breast-feeding (weeks)	35.3	1.4	36.5	0.9	0.479

*Data are presented as n and %.

Table 3 Whole-day dietary patterns and food group loadings for 2–8-year-old children

Non-core food groups		Healthy, meat and vegetable		Combination	
Food group	Loading	Food group	Loading	Food group	Loading
Milk: whole fat	0.5689	Vegetables: other	0.6532	Candy (not chocolate based)	0.3792
Luncheon and deli-type meats	0.4290	Vegetables: orange	0.5660	Other dairy	0.3615
Cheese	0.4008	Fruit	0.4870	Pasta/rice products	0.3525
Beef/lamb/veal	0.3851	Bread/rolls: non-white	0.4209	Milk: non-whole fat	0.3521
Bread/rolls: white	0.3320	Non-carbonated, non-sugar-sweetened beverages	0.3839	Nuts/seeds	0.3498
Yoghurt: whole fat	0.3318	Cheese	0.3265	Cakes	0.3467
Sauces	0.3224	Cereal: low–medium sugar sweetened	0.3227	White meat	0.3392
Muesli and cereal bars	0.2709	Edible oils and spreads	0.2883	Sauces	0.3294
Soups	0.2963	Beef/lamb/veal	0.2771	Egg dishes	0.3272
Potatoes: home cooked, no oil	0.2915	Water	0.2719	Seafood	0.3252
Cereal: medium–high sugar sweetened	0.2659	Pasta/rice products	0.2578	Take-away	0.3245
Beverage bases	0.2598			Cereal: medium–high sugar sweetened	0.3184
Juice	0.2594	Potatoes: take-away	–0.7334	Potatoes: home cooked, with oil	0.3184
Sweet biscuits	0.2570	Carbonated sugar-sweetened beverages	–0.6446	Yoghurt: non-whole fat	0.3147
		Take-away	–0.6048	Cereal-based dishes	0.3146
Non-carbonated, non-sugar-sweetened beverages	–0.9055	Added sugar	–0.3590	Beverage bases	0.3054
Milk: non-whole fat	–0.5036			Pork	0.2901
White meat	–0.3229			Added sugar	0.2887
Yoghurt: non-whole fat	–0.3213			Cordial/fruit drinks	0.2866
Bread/rolls: non-white	–0.2779			Chocolate	0.2791
Savoury snacks	–0.2599				
				Carbonated, non-sugar-sweetened beverages	–0.9865
				Edible oils and spreads	–0.4254
				Bread/rolls: white	–0.4243

Table 4 Multivariate linear regression coefficients (β) associating factor scores with breast-feeding (not breast-fed *v.* ever breast-fed)

	β Coefficient	SEM	P	OR	95% CI
Non-core food groups					
Ever breast-fed	0.079	0.093	0.395	1.08	0.901, 1.299
Age at which solids were introduced	-0.400	0.195	0.041	0.67	0.456, 0.984
Number of children in the household	0.079	0.032	0.015	1.08	1.015, 1.153
SEIFA tertile 1 (more disadvantaged)	0.633	0.201	0.002	1.88	1.269, 2.797
SEIFA tertile 2	0.527	0.203	0.010	1.69	1.136, 2.525
SEIFA tertile 3 (least disadvantaged)	0.522	0.202	0.010	1.69	1.132, 2.507
Energy intake (MJ)	0.069	0.016	<0.001	1.07	1.037, 1.106
Healthy, meat and vegetable					
Ever breast-fed	0.230	0.106	0.030	1.26	1.022, 1.550
Age (years)	-0.073	0.107	<0.001	0.93	0.910, 0.949
BMI (kg/m ²)	-0.022	0.106	0.040	0.98	0.959, 0.999
Birth weight	0.023	0.007	0.002	1.02	1.008, 1.038
Primary parent's educational level*	-0.062	0.011	<0.001	0.94	0.920, 0.960
SEIFA tertile 1 (more disadvantaged)	0.185	0.132	0.163	1.20	0.928, 1.560
SEIFA tertile 2	0.155	0.129	0.230	1.17	0.906, 1.505
SEIFA tertile 3 (least disadvantaged)	0.271	0.129	0.035	1.31	1.019, 1.689
Energy intake (MJ)	0.044	0.011	<0.001	1.04	1.023, 1.068
Combination					
Ever breast-fed	-0.168	0.107	0.118	0.85	0.685, 1.044
Age (years)	0.034	0.016	0.032	1.03	1.003, 1.068
Female	0.211	0.074	0.005	1.22	1.057, 1.419
Primary parent's educational level*	-0.047	0.019	0.014	0.95	0.919, 0.991
SEIFA tertile 1 (more disadvantaged)	-0.598	0.166	<0.001	0.55	0.397, 0.762
SEIFA tertile 2	-0.454	0.163	0.005	0.64	0.461, 0.874
SEIFA tertile 3 (least disadvantaged)	-0.481	0.158	0.002	0.62	0.453, 0.843
Energy intake (MJ)	0.042	0.019	0.022	1.04	1.006, 1.082

SEIFA, Socio-economic Indexes for Areas.

Final models with significant covariates.

*Higher values represent lower educational level.

was loaded with the greatest number of food groups with similar food group loadings and was labelled 'Combination'. This pattern did not include any fruit or vegetables but included candy (not chocolate based), other dairy products (e.g. ice cream, custard), pasta/rice products, reduced/low-fat milk, nuts/seeds, cakes and chocolate.

Cronbach's α for inter-item reliability was 0.78 for the non-core food groups pattern, 0.74 for the healthy, meat and vegetable pattern and 0.71 for the combination pattern.

Breast-feeding and dietary patterns

Children were classified as 'breast-fed' (n 2064, 89.3%) or 'never breast-fed' (n 223, 10.7%). After adjusting for a number of potential confounders, i.e. age, gender, energy intake, BMI, SEIFA, number of adults in the household, number of children in the household, birth weight, waist girth, age when the child started solids (if formula fed), age (weeks) at which the child received infant formula for the first time and the highest completed level of education/qualification by the primary caregiver and their partner, the results of the final models for breast-fed children compared with children who were 'never breast-fed' are shown in Table 4.

There was no association between breast-feeding and the non-core food groups pattern or the combination pattern (Table 4). Higher scores on the non-core food groups pattern were associated with the following: earlier introduction to solid foods; more children in the household; and

greater likelihood of being in the disadvantaged SEIFA category. Higher scores on the combination pattern were associated with: being older; greater likelihood of being a girl; less likelihood of having a lower educational level; and lower SEIFA category (indicating more disadvantaged).

There was a positive association between breast-feeding and the healthy, meat and vegetable pattern (Table 4). Scores on the healthy, meat and vegetable pattern were 26% higher for children who were breast-fed compared with children who were not breast-fed. Higher scores on this pattern were also associated with younger age, lower BMI, higher birth weight, greater likelihood of being in the less-disadvantaged SEIFA category and less likelihood of the children's parents having lower educational level.

Discussion

The present study is the first in Australia to conduct a factor analysis on a national sample of 2-8-year-old children and to associate the resulting dietary patterns with prior breast-feeding.

A positive association was found between the healthy, meat and vegetable dietary pattern and breast-feeding. Higher factor scores indicate more food groups loading on this pattern. Therefore, compared with children who were not breast-fed, children who were breast-fed were more likely to consume healthy foods such as vegetables,

fruit, wholegrain bread and red meat, which were retained in this pattern. Only one study has examined the relationship between breast-feeding and dietary patterns; however, this was conducted in infants at 6 months and 1 year of age⁽³³⁾. In that study, a dietary pattern characterised by a higher frequency of consumption of vegetables, fruit, meat and fish, in addition to high consumption of breast milk, emerged in infants at 6 months of age and again at 1 year of age⁽³³⁾. The present study extends these findings in infants to an older age group of Australian children, indicating that breast-feeding appears to be associated with an overall healthy eating pattern, and tracks throughout childhood.

The mechanism behind this relationship may relate to findings from earlier studies in animals^(34–37) and to subsequent human studies that revealed that breast-feeding was associated with greater acceptability to new foods and flavours^(7,11,38). Moreover, it has been shown that flavours from the mothers' diet are transmitted through amniotic fluid⁽³⁹⁾ and later through breast milk^(40,41). Therefore, flavours and components consumed in the mothers' diet are associated with, and may have a significant effect on, feeding preferences and practices in later life.

There is also some, albeit limited, evidence linking breast-feeding to improved cardiovascular risk factors such as blood pressure and lipids^(42,43), type 2 diabetes⁽⁴⁴⁾ and risk for obesity⁽⁴⁵⁾. All these outcomes are also affected by dietary intakes. Therefore, if at the onset of life positive breast-feeding practices influence childhood and adolescent food choices and dietary intakes, it is imperative that the benefits of breast-feeding be promoted and clearly understood by new mothers.

The WHO recommends that all infants be exclusively breast-fed for the first 6 months of life and that breast-feeding continues into the second year of life⁽⁴⁶⁾. This is not currently achieved, as data from the 2004–2005 Australian National Health Survey reported that, although 88% of mothers initiated breast-feeding, only 50% of infants were still breast-fed at 6 months and 23% were breast-fed at 12 months⁽⁴⁷⁾. This further suggests the need to educate and promote breast-feeding and inform mothers on possible health benefits and its associated outcomes in young children.

There was no association between the non-core food groups pattern and breast-feeding. However, this pattern was associated with earlier introduction to solids, to the presence of more children in the household, to a greater likelihood of being in the more disadvantaged SEIFA categories and to higher energy intakes. This finding is supported by the results of another Australian study which reported that infants were more likely to have received non-core foods if they received solids for the first time before 17 weeks and if they had two or more older siblings⁽⁴⁸⁾. No association was found between the combination dietary pattern and breast-feeding. However, the direction of breast-feeding was negative, suggesting that

being breast-fed was less likely to be associated with higher scores on this pattern. This pattern was characterised by a large number of food groups, with many being non-core foods. In the study by Robinson *et al.*⁽³³⁾ conducted in infants, the second dietary pattern identified was a pattern containing 'adult'-type foods such as bread, savoury snacks, breakfast cereals and chips. This pattern was inversely associated with consumption of breast milk and infant-type foods⁽³³⁾. These food groups were loaded on either the combination or non-core food groups pattern, supporting the fact that poorer food choices and dietary patterns are associated with little or no breast-feeding.

The role of the primary parent's educational level (typically that of the mother) also needs to be highlighted. It is of interest that the association between breast-feeding and the healthy, meat and vegetable pattern was independent of education, which was associated with this pattern as well. Other studies have reported positive associations between breast-feeding and education^(49,50), and women who breast-fed for >16 weeks were more often highly educated⁽⁵¹⁾. In the latter study, breast-fed children were less likely to have a low consumption of fruit and vegetables and a high consumption of snacks or an unhealthy diet, independent of maternal education⁽⁵¹⁾. This earlier study and our data indicate that the association between breast-feeding and diet is independent and that breast-feeding is not just a proxy measure of educational level.

There are some limitations to the present study. This was a cross-sectional study in which only a snapshot of the foods normally consumed by these children was captured. However, the large sample size reduced the estimated variance. Factor analysis is a validated method for analysis of dietary patterns^(52,53); however, rather than assigning an individual to a specific dietary pattern, this method produced continuous dietary factor scores for each individual for each pattern. Therefore, it was not possible to determine statistically significant differences between each dietary pattern, and only associations and relationships between continuous variables and dietary pattern scores could be tested. Factor analysis is also sample specific; therefore, the results may not be generalisable to other populations. However, we performed the same factor analysis methodology in this population using the 24h CATI data, in which three similar factors emerged.

The total variance explained by each factor was also low, compared with previous factor analysis studies conducted in older age groups^(27,31,54,55). However, Cronbach's α coefficients were all >0.7, indicating acceptable inter-item reliability. In addition, the food groups loading on the factors were varied and many were greater than the 0.25 cut-off value. This suggests that our population had a varied diet that was, nevertheless, still specific to the identified factors. As the present study is the first of its kind in this age group, further studies are required to refute or support our findings.

In conclusion, our results provide suggestive evidence that breast-feeding during infancy is associated with a

healthy dietary pattern in childhood and offers a likely mechanistic pathway to explain the previously reported association of breast-feeding with chronic disease. A number of health benefits have been associated with breast-feeding and these have also been associated with health outcomes in adulthood. Although we did not find any association between breast-feeding and the unhealthier patterns, it is of concern that these patterns were identified at such a young age, as these may track through adulthood. It is important to inform parents of a healthy diet during pregnancy and of the significance of breast-feeding and how food choices in childhood may track into adulthood. It is also essential to inform young children of the importance of healthy eating.

Further studies investigating dietary patterns and duration of breast-feeding are warranted to supplement the limited existing literature on the significance of breast-feeding and how it may relate to children's diet and to later health outcomes.

Acknowledgements

The present study was supported by the Meat and Live-stock Association but the Association had no input into the results presented in the current report. The authors have no conflict of interest to declare. J.A.G. was responsible for analysing and interpreting the data and for writing the manuscript; L.C. and J.S. were responsible for the study design and for reading and editing the drafts. The authors acknowledge the assistance of Dr Richard Woodman who provided invaluable statistical assistance.

References

- Birch LL & Fisher JO (1998) Development of eating behaviors among children and adolescents. *Pediatrics* **101**, 539–549.
- Oliveria SA, Ellison RC, Moore LL *et al.* (1992) Parent-child relationships in nutrient intake: the Framingham Children's Study. *Am J Clin Nutr* **56**, 593–598.
- Vereecken CA, Keukelier E & Maes L (2004) Influence of mother's educational level on food parenting practices and food habits of young children. *Appetite* **43**, 93–103.
- Mannino ML, Lee Y, Mitchell DC *et al.* (2004) The quality of girls' diets declines and tracks across middle childhood. *Int J Behav Nutr Phys Act* **1**, 5–15.
- Nicklas TA, Webber LS & Berenson GS (1991) Studies of consistency of dietary intake during the first four years of life in a prospective analysis: Bogalusa Heart Study. *J Am Coll Nutr* **10**, 234–241.
- Nicklaus S, Boggio V, Chabanet C *et al.* (2005) A prospective study of food variety seeking in childhood, adolescence and early adult life. *Appetite* **44**, 289–297.
- Mennella JA, Forestell CA, Morgan LK *et al.* (2009) Early milk feeding influences taste acceptance and liking during infancy. *Am J Clin Nutr* **90**, 780S–788S.
- Mennella JA, Jagnow CP & Beauchamp GK (2001) Prenatal and postnatal flavor learning by human infants. *Pediatrics* **107**, E88.
- Hausner H, Nicklaus S, Issanchou S *et al.* (2010) Breast-feeding facilitates acceptance of a novel dietary flavour compound. *Clin Nutr* **29**, 141–148.
- Maier AS, Chabanet C, Schaal B *et al.* (2008) Breastfeeding and experience with variety early in weaning increase infants' acceptance of new foods for up to two months. *Clin Nutr* **27**, 849–857.
- Mennella JA & Beauchamp GK (1997) Mothers' milk enhances the acceptance of cereal during weaning. *Pediatr Res* **41**, 188–192.
- Sullivan SA & Birch LL (1994) Infant dietary experience and acceptance of solid foods. *Pediatrics* **93**, 271–277.
- Conn JA, Davies MJ, Walker RB *et al.* (2009) Food and nutrient intakes of 9-month-old infants in Adelaide, Australia. *Public Health Nutr* **12**, 2448–2456.
- Ip S, Chung M, Raman G *et al.* (2007) *Breastfeeding and Maternal and Child Health Outcomes in Developed Countries*. AHRQ Publication no. 07-E007. Rockville, MD: Agency for Healthcare Research and Quality.
- Horta BL, Bahl R, Martines JC *et al.* (2007) *Evidence on the Long-term Effects of Breastfeeding. Systematic Reviews and Meta-analysis*. Geneva: WHO.
- Hu FB & Willett WC (2002) Optimal diets for prevention of coronary heart disease. *JAMA* **288**, 2569–2578.
- Coppinger T, Jeanes YM, Dobinett J *et al.* (2010) Physical activity and dietary intake of children aged 9–11 years and the influence of peers on these behaviours: a 1-year follow-up. *Eur J Clin Nutr* **64**, 776–781.
- Hennessy-Priest KA, Mustard JL, Keller HH *et al.* (2008) Zinc-fortified foods do not improve intake of total dietary zinc for Ontario preschoolers. *J Am Coll Nutr* **27**, 561–568.
- Jennings A, Davies GJ, Costarelli V *et al.* (2010) Micro-nutrient intakes of pre-adolescent children living in London. *Int J Food Sci Nutr* **61**, 68–77.
- Jones LR, Steer CD, Rogers IS *et al.* (2010) Influences on child fruit and vegetable intake: sociodemographic, parental and child factors in a longitudinal cohort study. *Public Health Nutr* **13**, 1122–1130.
- Magarey A, Daniels LA & Smith A (2001) Fruit and vegetable intakes of Australians aged 2–18 years: an evaluation of the 1995 National Nutrition Survey data. *Aust N Z J Public Health* **25**, 155–161.
- Babey SH, Jones M, Yu H *et al.* (2009) Bubbling over: soda consumption and its link to obesity in California. *Policy Brief UCLA Cent Health Policy Res* 1–8.
- Wosje KS, Khoury PR, Claytor RP *et al.* (2010) Dietary patterns associated with fat and bone mass in young children. *Am J Clin Nutr* **92**, 294–303.
- Aranceta J, Perez-Rodrigo C, Ribas L *et al.* (2003) Socio-demographic and lifestyle determinants of food patterns in Spanish children and adolescents: the enKid study. *Eur J Clin Nutr* **57**, Suppl. 1, S40–S44.
- Craig LC, McNeill G, Macdiarmid JI *et al.* (2010) Dietary patterns of school-age children in Scotland: association with socio-economic indicators, physical activity and obesity. *Br J Nutr* **103**, 319–334.
- North K & Emmett P (2000) Multivariate analysis of diet among three-year-old children and associations with socio-demographic characteristics. The Avon Longitudinal Study of Pregnancy and Childhood (ALSPAC) Study Team. *Eur J Clin Nutr* **54**, 73–80.
- Cole TJ, Bellizzi MC, Flegal KM *et al.* (2000) Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* **320**, 1240–1243.
- Cole TJ, Flegal KM, Nicholls D *et al.* (2007) Body mass index cut offs to define thinness in children and adolescents: international survey. *BMJ* **335**, 194.
- Schulze MB, Hoffmann K, Kroke A *et al.* (2003) An approach to construct simplified measures of dietary patterns from exploratory factor analysis. *Br J Nutr* **89**, 409–419.
- McCann SE, Marshall JR, Brasure JR *et al.* (2001) Analysis of patterns of food intake in nutritional epidemiology: food classification in principal components analysis and the

- subsequent impact on estimates for endometrial cancer. *Public Health Nutr* **4**, 989–997.
31. Slattery ML, Boucher KM, Caan BJ *et al.* (1998) Eating patterns and risk of colon cancer. *Am J Epidemiol* **148**, 4–16.
 32. Australian Bureau of Statistics (2008) *Information Paper: An Introduction to Socio-Economic Indexes for Areas (SEIFA)*. Catalogue no. 2039.0. Canberra: ABS; available at <http://www.abs.gov.au/ausstats/abs@.nsf/mf/2033.0.55.001/2006>
 33. Robinson S, Marriott L, Poole J *et al.* (2007) Dietary patterns in infancy: the importance of maternal and family influences on feeding practice. *Br J Nutr* **98**, 1029–1037.
 34. Galef BG Jr & Henderson PW (1972) Mother's milk: a determinant of the feeding preferences of weaning rat pups. *J Comp Physiol Psychol* **78**, 213–219.
 35. Galef BG Jr & Sherry DF (1973) Mother's milk: a medium for transmission of cues reflecting the flavor of mother's diet. *J Comp Physiol Psychol* **83**, 374–378.
 36. Morrill J & Dayton AD (1978) Effect of feed flavor in milk and calf starter on feed consumption and growth. *J Dairy Sci* **61**, 229–232.
 37. Campbell R (1976) A note on the use of feed flavor to stimulate the feed intake of weaner pigs. *Anim Prod* **23**, 417–419.
 38. Mennella JA, Griffin CE & Beauchamp GK (2004) Flavor programming during infancy. *Pediatrics* **113**, 840–845.
 39. Mennella JA, Johnson A & Beauchamp GK (1995) Garlic ingestion by pregnant women alters the odor of amniotic fluid. *Chem Senses* **20**, 207–209.
 40. Desage M, Schaal B, Soubeyrand J *et al.* (1996) Gas chromatographic–mass spectrometric method to characterise the transfer of dietary odorous compounds into plasma and milk. *J Chromatogr B Biomed Appl* **678**, 205–210.
 41. Mennella JA & Beauchamp GK (1991) Maternal diet alters the sensory qualities of human milk and the nursing's behavior. *Pediatrics* **88**, 737–744.
 42. Singhal A, Cole TJ, Fewtrell M *et al.* (2004) Breastmilk feeding and lipoprotein profile in adolescents born preterm: follow-up of a prospective randomised study. *Lancet* **363**, 1571–1578.
 43. Singhal A, Cole TJ & Lucas A (2001) Early nutrition in preterm infants and later blood pressure: two cohorts after randomised trials. *Lancet* **357**, 413–419.
 44. Owen CG, Martin RM, Whincup PH *et al.* (2006) Does breastfeeding influence risk of type 2 diabetes in later life? A quantitative analysis of published evidence. *Am J Clin Nutr* **84**, 1043–1054.
 45. Owen CG, Martin RM, Whincup PH *et al.* (2005) Effect of infant feeding on the risk of obesity across the life course: a quantitative review of published evidence. *Pediatrics* **115**, 1367–1377.
 46. World Health Organization (2002) *Global Strategy for Infant and Young Child Feeding*. Report by the Secretariat (WHA55 A55/15, paragraph 10, page 5). Geneva: WHO; available at http://apps.who.int/gb/archive/pdf_files/IUHA55/ea5515.pdf
 47. Amir LH & Donath SM (2008) Socioeconomic status and rates of breastfeeding in Australia: evidence from three recent national health surveys. *Med J Aust* **189**, 254–256.
 48. Koh G, Scott JA, Oddy WH *et al.* (2010) Exposure to non-core foods and beverages in the first year of life: results from a cohort study. *Nutr Diet* **67**, 137–142.
 49. Rebhan B, Kohlhuber M, Schwegler U *et al.* (2009) Infant feeding practices and associated factors through the first 9 months of life in Bavaria, Germany. *J Pediatr Gastroenterol Nutr* **49**, 467–473.
 50. Senarath U, Dibley MJ & Agho KE (2007) Breastfeeding practices and associated factors among children under 24 months of age in Timor-Leste. *Eur J Clin Nutr* **61**, 387–397.
 51. Scholtens S, Brunekreef B, Smit HA *et al.* (2008) Do differences in childhood diet explain the reduced overweight risk in breastfed children? *Obesity (Silver Spring)* **16**, 2498–2503.
 52. Hu FB, Rimm E, Smith-Warner SA *et al.* (1999) Reproducibility and validity of dietary patterns assessed with a food-frequency questionnaire. *Am J Clin Nutr* **69**, 243–249.
 53. Khani BR, Ye W, Terry P *et al.* (2004) Reproducibility and validity of major dietary patterns among Swedish women assessed with a food-frequency questionnaire. *J Nutr* **134**, 1541–1545.
 54. Ambrosini GL, Oddy WH, Robinson M *et al.* (2009) Adolescent dietary patterns are associated with lifestyle and family psychosocial factors. *Public Health Nutr* **12**, 1807–1815.
 55. Hu FB, Rimm EB, Stampfer MJ *et al.* (2000) Prospective study of major dietary patterns and risk of coronary heart disease in men. *Am J Clin Nutr* **72**, 912–921.