

Review Article

Food-chain selenium and human health: spotlight on speciation

Margaret P. Rayman^{1*}, Heidi Goenaga Infante² and Mike Sargent²

¹Nutritional Sciences Division, Faculty of Health and Medical Sciences, University of Surrey, Guildford, Surrey GU2 7XH, UK

²LGC Limited, Queens Road, Teddington, Middlesex TW11 OLY, UK

(Received 14 March 2007 – Revised 16 November 2007 – Accepted 21 November 2007 – First published online 18 March 2008)

There is a growing appreciation that it is not just the total intake of dietary Se that is important to health but that the species of Se ingested may also be important. The present review attempts to catalogue what is known about Se species in foods and supplements and the health effects in which they are implicated. The biosynthetic pathways involved in Se assimilation by plants and the way in which Se species are metabolised in animals are presented in order to give an insight into the species likely to be present in plant and animal foods. Known data on the species of Se in the food chain and in food supplements are tabulated along with their concentrations and the analytical methodology used. The latter is important, since identification that is only based on retention-time matching with authentic standards must be considered as tentative: for evidence of structural confirmation, fragmentation of the molecular ion in addition to MS data is required. Bioavailability, as normally defined, is higher for organic Se species. Health effects, both beneficial and toxic, thought to be associated with specific Se species are described. Potent anti-tumour effects have been attributed to the low-molecular-weight species, Se-methyl-selenocysteine and its γ -glutamyl-derivative, found in a number of edible plants of the *Allium* and *Brassica* families. There remain considerable gaps in our knowledge of the forms of Se that naturally occur in foods. Without adequate knowledge of Se speciation, false conclusions may be drawn when assessing Se requirements for optimal health.

Selenium: Speciation: Selenium in foods: Human health

The extent of the literature on the essential trace element Se appears to have increased exponentially over the last decade reflecting the tremendous growth of interest in this nutrient since it was shown by Clark *et al.* to reduce cancer risk in their landmark trial⁽¹⁾. Though the form of Se used in that trial was high-Se yeast, when large-scale funding was obtained from the National Cancer Institute for a follow-up randomised trial of the effect of supplemental Se on prostate cancer risk (SELECT), the decision was taken to use selenomethionine (SeMet) owing to the perceived importance of being able to define the specific form of Se that might be associated with an important health effect⁽²⁾. Thus we are no longer satisfied with knowing simply the amount of Se that may be associated with benefit but seek to know the species of Se to which that alleged benefit may be attributed. Furthermore, we have come to realise that different species of an element (for example, arsenic) can have very different health effects. The present review therefore attempts to pull together what is known about the species of Se in foods and supplements, the pathways by which they are synthesised, their apparent bioavailability as found in different food sources as this has implications for Se requirements, and the health effects that can be ascribed to specific Se species.

Biosynthesis and metabolism of dietary selenium species

A consideration of Se speciation in plant and animal food sources requires some understanding of the biosynthetic pathways involved in Se assimilation by plants and how these species are metabolised in animals. Such knowledge enables us to predict to some extent the Se species likely to be contained in foods. The biosynthetic pathways for Se in plants, some of which are assumed by analogy with S pathways, are shown in Fig. 1 (adapted from Ellis & Salt⁽³⁾, Whanger^(4,5), Terry *et al.*⁽⁶⁾, Tagmount *et al.*⁽⁷⁾ and Sors *et al.*⁽⁸⁾). The relative dominance of the pathways differs for Se-accumulators and non-accumulators.

The major species in plant sources of Se are: selenate (translocated directly from the soil and less readily bound to soil components than selenite); SeMet (biosynthesised) and a smaller amount of selenocysteine (SeCys; biosynthesised); Se-containing proteins (where SeMet and SeCys have been incorporated non-specifically in place of methionine and cysteine); Se-methyl-selenocysteine and γ -glutamyl-Se-methyl-selenocysteine (considered as detoxification products, notably formed in Se-accumulators and plants of the *Brassica* and *Allium* families). Plants can volatilise significant amounts

Abbreviations: SeCys, selenocysteine; SeMet, selenomethionine.

*Corresponding author: Professor Margaret Rayman, fax +44 1483 686401, email m.rayman@surrey.ac.uk




Fig. 1. Biosynthetic pathways elucidated for Se in higher plants (some by analogy with S pathways) (adapted from Ellis & Salt⁽³⁾, Whanger^(4,5), Terry *et al.*⁽⁶⁾, Tagmount *et al.*⁽⁷⁾ and Sors *et al.*⁽⁸⁾). It should be noted that reactions vary from species to species so that compounds formed and their relative quantities differ between species and strains. APSe, adenosine-5'-phosphoselenate; GSSe $_{\text{O}_3}^{2-}$, glutathione-S-selenite; GSH, glutathione; DMDSe, dimethyl-diselenide (volatile); GSSe $_{\text{SG}}$, selenodiglutathione; MeSeH, methyl selenol; GSSeH, glutathione-selenopersulfide; MeSeCysSeO, Se-methyl-selenocysteine selenoxide; GSSe $^{-}$, glutathione-conjugated selenide; γ -GMeSeCys, γ -glutamyl-Se-methyl-selenocysteine; MeSeCys, Se-methyl-selenocysteine; SeCys MeTransferase, selenocysteine methyltransferase; SeCys, selenocysteine; DMSe, dimethyl selenide (volatile); DMSeP, dimethyl-selenonio-propionate ($\text{CH}_3\text{Se}^+(\text{CH}_3)_2\text{CH}_2\text{CH}_2\text{COO}^-$); MeSeMet $^+$, Se-methyl-selenomethionine; SeCysth, selenocystathione; γ -GSeCysth, γ -glutamyl-selenocystathione; SAM, S-adenosyl methionine; Se-ASeMet, Se-adenosyl-selenomethionine; SeMet, selenomethionine; MeTHF, methyl-tetrahydrofolate; SeHomocysteine, selenohomocysteine.

of Se as dimethylselenide (non-accumulators) and dimethyl-diselenide (accumulators)⁽⁶⁾. To avoid an over-complicated Fig. 1, the enzymes implicated in these pathways are not shown, with the exception of SeCys methyltransferase, the enzyme notably present in Se-accumulators and responsible for the methylation of SeCys to the characteristic methylated metabolites that are believed to have anti-cancer properties.

While a study of these pathways suggests Se species that may be expected in foods from plant sources, it should be noted that compounds formed and their relative quantities

differ not only between Se-accumulators and non-accumulators but also between species.

There is much less information on the species of Se in dietary sources of animal origin⁽⁹⁾. When inorganic Se is given to animals, SeCys is the main seleno-compound formed but when animals eat Se-containing foods of plant origin, protein-bound SeMet will also be formed from the non-specific incorporation of plant-derived SeMet in place of methionine. Selenotrisulfide, glutathione selenopersulfide and metallic selenides have also been reported in tissues⁽¹⁰⁾. The presence

of some of these compounds can be explained by the metabolic pathway of dietary Se in animals which resembles that in humans as described below.

Most of what we know about the metabolism of dietary (or supplement) Se in humans is inferred from studies in rats and mice. A simplified version of the metabolic pathway is shown in Fig. 2 and clearly illustrates the central role of hydrogen selenide (H_2Se) (adapted from Combs⁽¹¹⁾ and Rayman⁽¹²⁾)^(13,14). SeMet catabolised from proteins can be trans-selenated to SeCys (by analogy with the trans-sulfuration pathway). SeCys, either from this source or directly from the diet, is then converted to H_2Se by SeCys β -lyase. Alternatively, SeMet can undergo α,γ -elimination catalysed by a γ -lyase to yield CH_3SeH , though the relative importance of this route in humans is not known^(13,15,16). CH_3SeH is also produced by a β -lyase from plant sources containing *Se*-methyl-selenocysteine and γ -glutamyl-*Se*-methyl-selenocysteine. Utilisation of selenate or selenite (plant sources or supplements) for selenoprotein synthesis, or excretion of excess, first requires reduction to the central Se metabolite, H_2Se , via interaction with the tripeptide, glutathione. The H_2Se so formed may be converted to selenophosphate ($HSePO_3^{2-}$) which then reacts with tRNA-bound serinyl residues to give SeCys-bound tRNA from which SeCys is inserted co-translationally, at loci encoded by specific UGA codons, to give selenoproteins^(17,18). As CH_3SeH can be demethylated to H_2Se in an equilibrium reaction, both it and its precursors can also act as Se sources for selenoprotein synthesis⁽¹³⁾. Oxidation of excess H_2Se can lead to the production of superoxide and other reactive oxygen species with associated toxic effects⁽¹¹⁾.

Surplus Se is transformed to methylated metabolites mostly for excretion into urine. Excretion of Se is either from H_2Se through a methylated selenosugar (1 *β* -methylseleno-N-acetyl-D-galactosamine) in urine or by further methylation of CH_3SeH to dimethyl selenide ((CH_3)₂Se) which is exhaled in breath, and trimethyl selenonium ion ((CH_3)₃Se⁺) excreted

in urine⁽¹⁹⁻²¹⁾. Though 1 *β* -methylseleno-N-acetyl-D-galactosamine is the most significant urinary metabolite in most individuals, (CH_3)₃Se⁺ is a major product from Se-methyl-selenocysteine^(13,21,22).

Selenium in food sources and dietary supplements: speciation and concentration

Table 1 shows the Se species apparently identified in foods and dietary supplements and their concentrations or relative concentrations in some cases^(4,5,9,10,12,23-72) (H Goenaga Infante, G O'Connor and MP Rayman, unpublished results). In terms of identification, it must be borne in mind that many of these studies were carried out when the available analytical strategies that combined both elemental and molecular MS were less well developed than is currently the case. In the case of most foods, however, they are the only data we have and can help focus the direction of further studies. Column 5 shows the methodology used for Se species identification. Readers should be aware, however, that identification that is only based on retention-time matching with authentic standards by HPLC-inductively coupled plasma MS is tentative and that electrospray ionisation MS data alone do not provide enough evidence of structural confirmation. To obtain this, fragmentation of the molecular ion has to be performed⁽²⁸⁾. Table 1 contains some speciation data that have been obtained in this way, for example, by inductively coupled plasma MS combined with MS/MS data obtained by matrix-assisted laser desorption/ionisation (MALDI) or electrospray ionisation MS with fragmentation of the precursor/molecular ion (electrospray ionisation MS/MS)^(27,30-33,37,47,48,56-58,61,62). Those wishing to understand more about speciation-analysis methodology are referred to critical reviews of recent analytical developments for the Se speciation analysis of foods, supplements and biosamples^(28,73).

Most quantitative data in Table 1 have been calculated from the peak area for a particular Se species expressed as a per-




Fig. 2. Metabolic pathway of dietary Se in humans (adapted from Combs⁽¹¹⁾ and Rayman⁽¹²⁾)^(13,14). SeMet, selenomethionine; SeCys, selenocysteine; GSSeSG, selenodiglutathione; γ -glutamyl-CH₃SeCys, γ -glutamyl-Se-methyl-selenocysteine; H_2Se , hydrogen selenide; $HSePO_3^{2-}$, selenophosphate; CH₃SeCys, Se-methyl-selenocysteine; CH₃SeH, methyl selenol; (CH_3)₂Se, dimethyl selenide; SeO_2 , selenium dioxide; (CH_3)₃Se⁺, trimethyl selenonium ion.

Table 1. Species and concentrations of selenium in foods and supplements (concentrations of selenium species are expressed in terms of concentration of elemental selenium) and the methodology used for species identification

Food or source	Growth conditions	Species	Typical concentration	Identification methodology	References
Supplements (also see Se-yeast)	Not applicable	Selenite/selenate Selenomethionine Se-methyl-selenocysteine –	30–100 µg/tablet 60–200 µg/tablet 100–200 µg/tablet Total Se: 1200–2200 µg/g		Whanger (1989, 2004) ^(4,5)
Se-yeast (in supplements; as a food additive; in functional foods)	Medium enriched with inorganic Se	Selenomethionine Selenocysteine Selenite Se-methyl-selenocysteine γ-Glutamyl-Se-methyl-selenocysteine Se-adenosyl-selenohomocysteine Se-cystathione Se-lanthionine Selenocystine Selenomethionine–selenoxide/hydrate SeMet–Asn–Ala–Gly–Arg Selenodiglutathione and the mixed seleno- notrisulfide of glutathione and cysteinylglycine Se-containing proteins (SIP18 and HSP12) S-selenomethyl-glutathione Glutathione-S-selenoglutathione – – –	60–84 % 3–5 % <1 % total extracted Se 0.5 % 0.5 % 0.5–5 % 0.5–1 % 0.5 % 0.5 % 0.5 % 0.5 % – –	ESI-MS/MS Retention time Retention time ESI-MS/MS ESI-MS/MS ESI-MS/MS Retention time Retention time Retention time ESI-MS ESI-MS/MS ESI-MS/MS	Whanger (2002) ⁽⁹⁾ Rayman (2004) ⁽¹²⁾ Kotrebai (2000) ⁽²³⁾ , Uden (2003) ⁽²⁴⁾ , Larsen (2004) ⁽²⁵⁾ , Goenaga Infante (2004) ⁽²⁶⁾ , Goenaga Infante (2005) ^(27,28) , Mester (2006) ⁽²⁹⁾
Cereals			0.1–30 µg/g (Western USA) 0.016–0.021 µg/g (Denmark) 0.007–0.011 µg/g (Finland, Sweden, Norway) ≤ 30 µg/g		Encinar (2003) ⁽³⁰⁾ Lindemann (2002) ⁽³¹⁾
Wheat, maize, barley, oats, rye	North and South Dakota, Nebraska, Kansas, Colorado Other parts of USA	–	0.1 µg/g 12–19 % total eluted Se	ESI-MS/MS Retention time	Encinar (2003) ⁽³²⁾ Goenaga Infante (2006) ⁽³³⁾
Wheat grain	Other parts of USA	Selenate (includes selenite, and selenocysteic acid) Selenomethionine Selenocysteine Se-methyl-selenocysteine Selenate	56–83 % total eluted Se 4–12 % total eluted Se 1–4 % total eluted Se ≤ 40–50 % total Se at ≤ 100 µg/g dry weight	Retention time Retention time Retention time Retention time	Whanger (2002) ⁽⁹⁾
Wheat sprouts					Lintschinger (2000) ⁽³⁵⁾
Wheat grain Wheat flour	China USA	Selenomethionine – Selenomethionine Selenate	50.4–81.4 % of total Se Total: 439 ng/g 256 ng/g 5.3 ng/g	Retention time ESI-MS/MS Retention time	Yang (1997) ⁽³⁶⁾ Warburton (2007) ⁽³⁷⁾
Maize	Seleniferous soil High-Se area of China	– Selenomethionine Selenocysteine Selenomethionine	8.7 (range 0.6–44.0) µg/g (selenosis) 18 µg/g 61–64 % of total eluted Se 15–16 % of total eluted Se 0.01–19.01 µg/g representing 46–82 % of the total Se	Retention time Retention time Retention time	Whanger (1996) ⁽³⁸⁾ Beilstein (1991) ⁽³⁹⁾
Barley Rice	Denmark Seleniferous soil	– –	≤ 0.11 µg/g 4.0 (0.3–20.2) µg/g (selenosis)		Whanger (1989) ⁽⁴⁾ Whanger (1996) ⁽³⁸⁾

Selenium speciation in food and health

Table 1. Continued

Food or source	Growth conditions	Species	Typical concentration	Identification methodology	References
	High-Se area of China	–	3.6 µg/g Se		Whanger (1996) ⁽³⁸⁾
Soya beans	China, grain	Selenomethionine	68–81 % of total eluted Se	Retention time	
	–	Selenocysteine	6–10 % of total eluted Se	Retention time	
	China	Selenite	5–13 % of total eluted Se	Retention time	
	–	Selenate	1–3 % of total eluted Se	Retention time	
Lima beans	China	Selenomethionine	54.9–86.5 % of total Se	Retention time	Yang (1997) ⁽³⁶⁾
	–	Selenomethionine	>80 % of eluted Se	Retention time	Whanger (1989) ⁽⁴⁾
Brazil nuts	Natural	Se-methyl-selenocysteine	62.9–71.8 % of total Se	Retention time	Yang (1997) ⁽³⁶⁾
	Acre – Rondonia, Brazil (shelled)	γ-Glutamyl-Se-methyl-selenocysteine	–	Retention time	Whanger (1989) ⁽⁴⁾
	Manaus – Belém, Brazil (unshelled)	Se-cystathione	–	Retention time	
	Natural (UK purchased)	–	2.54 µg/g		Barceloux (1999) ⁽⁴⁰⁾
	Natural (unshelled)	–	3.06 (range 0.03–31.7) µg/g fresh weight		Chang (1995) ⁽⁴¹⁾
	Natural	–	36.0 (range 1.25–512.0) µg/g fresh weight		
	Purchased with shells	–	2.54 (range 0.85–6.86) µg/g		Barclay (1995) ⁽⁴²⁾
	Purchased without shells	Selenomethionine	22 µg/g		Lisk (1988) ⁽⁴³⁾
		Weakly protein-bound Se	≤100 µg/g	Retention time	Palmer (1982) ⁽⁴⁴⁾
		Low-molecular-weight compounds	–	Retention time	Wrobel (2003) ⁽⁴⁵⁾
		–	Total Se: 35.1 µg/g		Kannankumarath (2002) ⁽⁴⁶⁾
	Natural	Selenomethionine	Major Se species, 25 % total Se	Retention time	
		Weakly protein-bound Se	12 % total Se		
		Low-molecular-weight compounds	3.1 % total Se		
		–	Total Se: 8.3 µg/g		
Walnuts	Black and white	Selenomethionine	Major Se species, 21 % total Se	Retention time	
		Weakly protein-bound Se	12.0 % total Se		
		Low-molecular-weight compounds	5.0 % total Se		
		Fifteen Se-containing peptides (fourteen selenomethionine-containing peptides + one Se-cysteine-containing peptide)	Total Se: 82.9 µg/g	ESI-MS/MS	Dernovics (2007) ⁽⁴⁷⁾
Cashew nuts	Unspecified	–	Total Se: 0.38 and 0.20 µg/g respectively		Kannankumarath (2002) ⁽⁴⁶⁾
	Natural	Selenomethionine	19 % and 23 % respectively of total Se	Retention time	
		Low-molecular-weight compounds	15 % and 10 % respectively of total Se		
		–	Total Se: 0.031 (0.012–0.060) µg/g		Barclay (1995) ⁽⁴²⁾
		Selenomethionine	Total Se: 0.27 µg/g		Kannankumarath (2002) ⁽⁴⁶⁾
		Low-molecular-weight compounds	22 % of total Se	Retention time	
	Natural	–	12 % of total Se		
			Total Se: 0.27 (range 0.17–0.39) µg/g		Barclay (1995) ⁽⁴²⁾
Pecan nuts	Natural	–	Total Se: 0.10 µg/g		Kannankumarath (2002) ⁽⁴⁶⁾
		Selenomethionine	25 % of total Se	Retention time	
Monkeypot nuts	Natural	Low-molecular-weight compounds	Not detected		Dernovics (2007) ⁽⁴⁸⁾
		–	Total Se: 4.5 mg/g		

M. P. Rayman *et al.*

Table 1. Continued

Food or source	Growth conditions	Species	Typical concentration	Identification methodology	References
Pumpkin seeds	Enriched by leaf spraying	Selenocystathione Isoforms of γ -glutamyl-selenocystathione	– –	ESI-MS/MS	
Vegetables	Natural	Selenomethionine	81 ± 8 % total Se	Retention time	Smrkolj (2005) ⁽⁴⁹⁾
Vegetables: rutabagas, cabbage, peas, beans, carrots, tomatoes, beets, potatoes, cucumbers	Seleniferous soil	– Selenate	<0.01 µg/g ≤6 µg/g total Se	Retention time	Barceloux (1999) ⁽⁴⁰⁾ Whanger (1989) ⁽⁴⁾
Vegetables	Sludge-amended soil	– Selenate	Up to 50 % of total Se 0.001–0.064 µg/g		Whanger (1989) ⁽⁴⁾
Asparagus	UK	–	11 µg/g		Barclay (1995) ⁽⁴²⁾
Broccoli	Seleniferous soil	–	Total Se: 345 µg/g		Whanger (1989) ⁽⁴⁾
	Enriched	Se-methyl-selenocysteine: selenocysteine	Ratio 2:1	Retention time	Cai (1995) ⁽⁵⁰⁾
Broccoli florets and leaves	Enriched	Selenomethionine Selenate	Minor amounts ≤44 % and ≤38 % total Se respectively	Retention time Retention time	Whanger (2002) ⁽⁹⁾
Broccoli sprouts	Enriched	– Se-methyl-selenocysteine Selenate Selenomethionine Adenosyl-selenohomocysteine	Total Se: 62.3 ± 0.6 µg/g dry weight 45 % 20 % 12 % 3 %	Retention time Retention time Retention time Retention time	Finley (2001) ⁽⁵¹⁾
Broccoli roots	Grown in hydroponic culture with Na ₂ SeO ₃	Selenomethionine	Major species	Retention time	Pedrero (2007) ⁽⁵²⁾
Broccoli florets	Enriched	Se-methyl-selenocysteine Se-methyl-selenocysteine-selenoxide	Major species ≤21.5 % of total extractable Se	Retention time Retention time	Hamilton (1975) ⁽⁵³⁾
Cabbage (<i>Brassica oleracea capitata</i>)	Sludge-amended soil	Selenate	≤40 % of total Se	Retention time	Whanger (2002) ⁽⁹⁾
Cabbage	Enriched	–	Total: 94 µg/g		H Goenaga Infante, G O'Connor and MP Rayman (unpublished results)
Radish (<i>Raphanus sativus</i>)	Hydroponic culture with (i) Na ₂ SeO ₃ (ii) Na ₂ SeO ₄	Se-methyl-selenocysteine Selenomethionine Selenate Secystine	30 ng/g 302 ng/g 38 µg/g (i) Total 112 ± 7 µg/g (ii) Total 120 ± 6 µg/g µg/g fresh weight (i) 6 ± 1 (ii) 19 ± 2 83 ± 7 7 ± 1 18 ± 1 20 ± 1 1 ± 0 68 ± 5	Retention time Retention time Retention time Retention times	Pedrero (2006) ⁽⁵⁴⁾
Mushrooms (<i>Agaricus bisporus</i> and <i>Lentinula edodes</i>)	Enriched	Se-methyl-selenocysteine Selenomethionine Selenate Selenocystine	Total Se dry weight: <i>Agaricus bisporus</i> 0.77 µg/g; <i>Lentinula edodes</i> 0.043 µg/g		Gergely (2006) ⁽⁵⁵⁾
			–	Retention times	

Selenium speciation in food and health

Table 1. Continued

Food or source	Growth conditions	Species	Typical concentration	Identification methodology	References
<i>Lentinula edodes</i> (shiitake mushrooms)	Enriched	Selenomethionine Se-methyl-selenocysteine Inorganic Se Selenomethionine	— — — 40.6 % of the total water-soluble Se	ESI-MS/MS	Ogra (2004) ⁽⁵⁶⁾
	Enriched	—	Total Se: 69 µg/g		H Goenaga Infante, G O'Connor and MP Rayman (unpublished results), Rayman (2007) ⁽⁵⁷⁾
Onions (<i>Allium cepa</i>)	Natural (USA)	Se-methyl-selenocysteine Selenomethionine Selenate	114 ng/g 17 µg/g 61 parts per billion	ESI-MS/MS ESI-MS/MS Retention time	Kotrebai (2000) ⁽²³⁾
	—	Selenate	Total Se: <0.5 µg/g	Retention time	Whanger (1989) ⁽⁴⁾
Onions (<i>Allium cepa</i>)	Seleniferous soil	—	100% of total enzymatic extract of Se	Retention time	Cai (1995) ⁽⁵⁰⁾
	Enriched	Selenocysteine	17 µg/g	Retention time	
Onions (<i>Allium cepa</i>)	Enriched	Se-methyl-selenocysteine	Total Se: 96 µg/g; equal amounts of both compounds	Retention time	Kotrebai (2000) ⁽²³⁾
	Enriched with 15 µg/g Se IV added to growth medium	—	Total Se 96 140 µg/g of which % distribution as follows:		
Garlic (<i>Allium sativum</i>)	Natural	γ-Glutamyl-Se-methyl-selenocysteine Selenomethionine Se-methyl-selenocysteine Selenate Se-cystathione Selenocystine	35 63 % 10 5 % 5 1 % 33 10 % 4 0.5 % — 1 % Sum 92 96 %	Retention times	
	Natural	—	30.3 µg Se/g plant tissue		Shah (2004) ⁽⁵⁸⁾
Garlic (<i>Allium sativum</i>)	Supplement	Selenocystine Se-methyl-selenocysteine SeMet γ-Glutamyl-Se-methyl-selenocysteine Inorganic Se	— — — — —	Retention time Retention time Retention time ESI-MS/MS Retention time	Cai (1995) ⁽⁵⁰⁾ Kotrebai (2000) ⁽²³⁾
	—	Selenocysteine — γ-Glutamyl-Se-methyl-selenocysteine Selenomethionine Se-methyl-selenocysteine Selenate	Total Se: 0.02 µg/g Total Se: <0.5 µg/g; % distribution: 31 % 53 % 12 % 4 %	Retention times	
Garlic (<i>Allium sativum</i>)	Enriched	—	Total Se: <0.5 µg/g; % distribution: 48 % 28 % 14 % 10 %	ESI-MS ESI-MS ESI-MS Retention time	Kotrebai (2000) ⁽²³⁾
	Enriched	γ-Glutamyl-Se-methyl-selenocysteine Selenomethionine Se-methyl-selenocysteine Selenate Selenocysteine —	Total Se: 68 µg/g Total Se: 1355 µg/g	Retention times	Cai (1995) ⁽⁵⁰⁾ Cai (1995) ⁽⁵⁰⁾
Garlic (<i>Allium sativum</i>)	Enriched	—	— — —		
	Enriched	Selenomethionine	Total Se: 1355 and 235 µg/g		Bird (1997) ⁽⁵⁹⁾ , Whanger (2002) ⁽⁹⁾

M. P. Rayman *et al.*

Table 1. Continued

Food or source	Growth conditions	Species	Typical concentration	Identification methodology	References
Enriched	Enriched	Se-methyl-selenocysteine (γ -glutamyl-Se-methyl-selenocysteine)	Predominant form	Retention times	
		Selenocystine	—		
		Selenomethionine	—		
		Selenoethionine	—		
		Se-propyl-selenocysteine	—		
		Selenate	—		
		Selenite	—		
		—	296 µg/g Se		Ip (2000) ⁽⁶⁰⁾
		γ -Glutamyl-Se-methyl-selenocysteine	73% of total eluted Se	ESI-MS	
		Selenomethionine	13% of total eluted Se	ESI-MS	
Enriched	Enriched	γ -Glutamyl-selenomethionine	4% of total eluted Se	Retention time	
		Se-methyl-selenocysteine	3% of total eluted Se	ESI-MS	
		Selenate	2% of total eluted Se	Retention time	
		Selenocystine	0.5% of total eluted Se	Retention time	
		Selenocystathione	0.5% of total eluted Se	Retention time	
		—	28.3 µg/g wet weight; 96 µg/g lyophilised		Dumont (2006) ⁽⁶¹⁾
		γ -Glutamyl-Se-methyl-selenocysteine	49.7% Se in hot water extract	ESI-MS/MS	
		Se-methyl-selenocysteine	28.8% Se in hot water extract	ESI-MS/MS	
		Selenomethionine	15.5% Se in hot water extract	ESI-MS/MS	
		Selenocystine	6.0% Se in hot water extract	Retention time	
Enriched	Enriched	—	Total Se, 68 112 235 296 1355 µg/g, of which % distribution as follows:		Kotrebai (2000) ⁽²³⁾
		γ -Glutamyl-Se-methyl-selenocysteine	68 52 70 73 8%	ESI-MS/MS	
		Selenomethionine	18 28 17 13 13%	ESI-MS/MS	
		Se-methyl-selenocysteine	2.5 3.0 3.0 3.0 60%	ESI-MS/MS	
		Selenate	1.0 0.1 1.5 2.0 4.0%	Retention time	
		Se-cystathione	0.5 1.5 0.5 0.5 1.5%	Retention time	
		Selenocystine	0.5 1.0 0.5 0.5 — %	Retention time	
		Sum	93 93 95 96 87%		
		—	Total: 205 µg/g		McSheehy (2000) ⁽⁶²⁾
		γ -Glutamyl-Se-methyl-selenocysteine	>78% of total Se	ESI-MS/MS	
Ramps (wild leeks, <i>Allium tricoccum</i>)	Enriched	—	Total: 1.980 µg/g		H Goenaga Infante, G O'Connor and MP Rayman (unpublished results)
		γ -Glutamyl-Se-methyl-selenocysteine	496 ng/g	ESI-MS/MS	
		Se-methyl-selenocysteine	35 ng/g	ESI-MS/MS	
		—	Total Se, 48 77* 230 252 405 524 µg/g, of which % distribution as follows:	Retention times	Kotrebai (2000) ⁽²³⁾
		γ -Glutamyl-Se-methyl-selenocysteine	— 3 1 1 1.5 1.5%		
		Selenomethionine	10 21 8 8 5 5%		
		Se-methyl-selenocysteine	35 34 50 50 44 44%		
		Selenate	25 1 15 15 25 22%		
		Se-cystathione	3 1 2 2 0.5 1.5%		
		Sum	73 60 76 76 76 74%		
Enriched	Enriched	Se-methyl-selenocysteine	35–50% total Se (at 48–524 µg/g)	Retention times	Whanger (2000) ⁽⁶³⁾
		Selenate	15–25%		
		Se-cystathione	0.5–3%		
		γ -Glutamyl-Se-methyl-selenocysteine	1–2%		

Selenium speciation in food and health

Table 1. Continued

Food or source	Growth conditions	Species	Typical concentration	Identification methodology	References
Chives (<i>Allium schoenoprasum</i>) leaves	Enriched with Se IV	–	Total: 222 µg/g	Retention times	Kápolna (2007) ⁽⁶⁴⁾
	Enriched with Se VI	Selenocystine	49 % of total Se		
		Se-methyl-selenocysteine	38 % of total Se		
		Selenomethionine	5 % of total Se		
	Enriched with SeMet	Inorganic Se (Se(IV) + Se(VI))	8 % of total Se		
		–	Total: 613 µg/g		
		Selenocystine	24 % of total Se		
		Se-methyl-selenocysteine	22 % of total Se		
		Se(VI)	>50 % of total Se		
Horseradish Meat and poultry	Enriched	–	Total: 265 µg/g	Unknown Retention times	Stewart (1974) ⁽⁶⁵⁾ Burk (1976) ⁽¹⁰⁾ , Behne (1998) ⁽⁶⁶⁾ , Whanger (2002) ⁽⁹⁾
		Selenocystine	44 % of total Se		
Meats Beef, pork, lamb Turkey (raw) Liver, kidney (raw) Fish (whole) Fish: black marlin blue marlin Fish: mackerel, octopus Fish: sardine, swordfish, tuna Fish: tuna	USA UK UK UK 109 US stations USA Spain/Portugal Spain/Portugal Canned	Se-methyl-selenocysteine	52 % of total Se	– Selenocysteine is predominant	Barceloux (1999) ⁽⁴⁰⁾ Barclay (1995) ⁽⁴²⁾
		Selenomethionine	3 % of total Se		
		Selenosinigrin (a glucosinolate)	–		
		Selenocysteine, selenomethionine, seleno-trisulfide, selenopersulfide, metallic selenide	Selenocysteine is predominant		
		–	–		
		–	0.3 µg/g		
		–	0.03–0.15 µg/g		
		–	0.1–0.2 µg/g		
		–	0.2–2.0 µg/g		
		–	0.42 µg/g		
Several fish, mollusks, crustaceans and pods Several species of marine and freshwater fish Cod Cod muscle	Spain/Portugal Spain/Portugal Canned, edible tissue Muscle Cooked Dried and powdered	0.4–4.3 µg/g	–	Barceloux (1999) ⁽⁴⁰⁾ Barceloux (1999) ⁽⁴⁰⁾	Cabanero (2005) ⁽⁶⁷⁾ Cabanero (2005) ⁽⁶⁷⁾
		2.5–4.2 µg/g	–		
		0.26 and 0.13 µg/g respectively	–		
		Total Se: 0.43, 0.47 and 0.92 µg/g respectively	Retention time		
		4–47 % of total Se (at 0.15– 4.15 µg/g)	Retention time		
		14–36 % of total Se (at 0.14– 0.83 µg/g)	Retention time		
		12 % of total Se at 1.5 µg/g	Retention time		
		70 % of the total Se	Retention time		
		Canned	7.6–44.8 % of total Se (at 0.36– 1.33 µg/g)	Retention time	Cappon (1982) ⁽⁶⁸⁾
		Selenate Se(-II), Se(IV)	4–47 % of total Se (at 0.15– 4.15 µg/g)	Retention time	Cappon (1982) ⁽⁶⁹⁾

ESI-MS, electrospray ionisation MS; ESI-MS/MS, electrospray ionisation MS with fragmentation of the precursor/molecular ion.

*For ramps, sample from 2nd year of growth.

tage of the total area of eluted Se peaks. However, accurate measurements by isotope-dilution MS or standard additions are also reported for methylated Se compounds such as SeMet and γ -glutamyl-Se-methyl-selenocysteine^{(25,27,29,72). Ideally, full mass balance data (i.e. total Se, total extracted Se, Se species, sum of species, extraction efficiency) should be considered together with recovery results from spiking experiments or analysis of ‘speciated’ certified reference materials for validation of speciation methodologies.}

The total Se concentration has been reported in Table 1 where possible, as it can affect the distribution of Se between species, as in the case of Se-enriched garlic and yeast⁽²³⁾. As the concentration of Se in Se-enriched foods is considerably higher than in the corresponding natural foods, such foods must be treated with caution, though the amounts in which they are eaten (for example, garlic) may reduce the risk of toxicity.

It is noteworthy that while wheat, other grains and soya contain predominantly SeMet with lesser amounts of SeCys and selenate, the major seleno-amino acids found in *Allium* and *Broccoli* species are Se-methyl-selenocysteine and γ -glutamyl-Se-methyl-selenocysteine. The latter two compounds are characteristic of the Se species produced by Se-accumulator plants which avoid the toxic effects of incorporation of excessive amounts of SeCys and SeMet into their proteins by accumulating non-protein seleno-amino acids or their γ -glutamyl derivatives⁽⁶⁾. Other non-protein seleno-amino acids that have been identified in Se-accumulator plants are selenocystathionine, Se-methyl-selenomethionine, γ -glutamyl-selenocystathionine, selenopeptides and selenohomocysteine⁽⁹⁾, though, of these, only selenocystathionine has been fully identified in foods (Table 1).

Given that Brazil nuts are potentially the richest food source of Se, and the tree that produces them, *Bertholletia excelsa*, is regarded as an Se-accumulator, it might be expected that the major Se species would be Se-methyl-selenocysteine or γ -glutamyl-Se-methyl-selenocysteine, as described above. Instead the major species in Brazil nuts appears to be SeMet^(44–46). This may to some extent be an illustration of the differences in concentration and speciation found between different plant tissues, Brazil nuts being seeds rather than fleshy leaves or florets as in the case of garlic or broccoli^(3,6). However, it may also be due to more general differences in Se metabolism between plant species (Dr Martin Broadley (2007), personal communication).

Considerably less information is available on Se species in animal foods than is available for plant foods. Although the Se content of fish and other seafoods has been reviewed by Reilly⁽⁷⁴⁾, normally ranging from 0·1 to 1·0 µg/g fresh weight, there is little information on specific Se species in fish. Several studies have found that seafood Se appears to be less bioavailable than that from other dietary sources, the implication being that the molecular form of at least some of the fish Se is such that it is not utilisable for selenoprotein synthesis^(40,75,76). Though it has been suggested that an explanation for this lower bioavailability may be interaction with Hg in seafood, the molar concentration of Se exceeds that of Hg by one or two orders of magnitude except in the case of sea mammals (cetaceans), suggesting that this is an unlikely explanation^(77–79). While Se and Hg undoubtedly have very high affinity for one another⁽⁸⁰⁾, there are as yet no

published data identifying Se–Hg species in seafood. However, according to Dr Nick Ralston (2007, personal communication) it appears that inorganic HgSe is present in the muscle meat of blue marlin as has already been shown in organs of mammals⁽⁸¹⁾. SeMet was the only compound identified in fish samples of high Se content in a speciation study⁽⁶⁷⁾ though other studies found from 4 to 47 % of total fish Se in the form of selenate^(68–70). This is an area ripe for further speciation studies.

Recently, new Se-containing glutathione species, S-selenomethyl-glutathione and glutathione-S-selenogluthathione have been identified in aqueous extracts of Se-yeast⁽³³⁾. As shown in Fig. 1, bonding of Se to glutathione via a non-enzymic reaction occurs in metabolism at the point where selenite enters the pathway to SeCys⁽⁶⁾. Alternatively, as glutathione is a tripeptide of γ -glutamine, cysteine and glycine, it seems possible that the formation of these Se-containing glutathione species could result from the incorporation of SeCys (or methylated SeCys) in place of cysteine in the biosynthetic pathway to glutathione.

While on the subject of Se-yeast, we should make it clear that it is not a defined form of Se. There is considerable variability in products described as Se-yeast which is reflected in the species composition. Se-yeast is produced by fermenting yeast in an Se-enriched medium when the Se becomes organically bound to yeast components. With reputable manufacturers, the percentage of Se that is organically bound should be greater than 90 % and more than 80 % should be bound to yeast proteins, including cell-wall proteins⁽¹²⁾. However, in some products, the percentage of sodium selenite is such that most of the Se is clearly not bound to the yeast; at worst, there may merely be a mixture of sodium selenite and yeast, the Se not being bound to the yeast⁽²⁴⁾. Such products dupe the consumer, as they do not conform to the normal understanding of Se-yeast as containing Se in an organic form. While they may be capable of increasing the production of selenoproteins, they will be less good at increasing plasma Se and acting as a storage form of Se in the body (see below), thereby maintaining Se status⁽⁸²⁾.

Selenium in food sources and dietary supplements: bioavailability

Bioavailability of a nutrient is conventionally defined as that fraction of ingested nutrient that is utilised for normal physiological functions⁽⁸³⁾; absorption and retention of the nutrient are taken as indirect measures of bioavailability as these are measurable⁽⁸³⁾ though they cannot address functional bioavailability which is that most likely to be relevant to health.

Absorption of Se is not homeostatically regulated and is not believed to be affected by nutritional status. Absorption of dietary Se is generally believed to be good – about 80 % from food⁽⁷⁴⁾. Guar gum is thought to reduce its absorption in humans⁽⁸⁴⁾, as is high dietary sulfur, probably because of competition between chemically similar sulfur and Se species^(74,85). Absorption of SeMet is active and uses the same enzyme transport system as does methionine⁽⁷⁴⁾. Absorption and retention of a commercially produced Se-yeast, in which 66 % of the Se present was in the form of SeMet (SelenoPrecise™), were measured as 90 and 75 % respectively (see Rayman⁽¹²⁾⁽⁸⁶⁾).

A number of supplementation studies have compared the bioavailability of different forms of Se to humans, i.e. Se-rich wheat, Se-enriched yeast, SeMet, sodium selenate and sodium selenite (for a review, see Rayman⁽¹²⁾). Organic forms of Se (wheat Se, SeMet and high-Se-yeast) were found to be more bioavailable than selenate and selenite in that they were more effective in raising blood Se concentrations (suggesting better absorption and retention), though all forms were able to increase selenoenzyme (glutathione peroxidase) activity. This difference is undoubtedly due to the ability of SeMet from digested organic Se sources to be incorporated in place of methionine into tissue proteins such as skeletal muscle, erythrocytes and plasma albumin where it can act as a Se store though it becomes available to the body only upon turnover of tissue proteins⁽⁸⁷⁾. Organic Se (Se-yeast) was also more effective than inorganic forms in its ability to transfer Se to breast-fed infants or suckling animals, thereby reducing the risk of deficiency in the offspring⁽¹²⁾. Foods that contain high proportions of SeMet, such as Brazil nuts and wheat, are good bioavailable sources of the element^(88,89). Though the Se content of mushrooms is higher than that of most other vegetables⁽⁷⁴⁾, its bioavailability is said to be very low⁽⁹⁰⁾. However, our own recent work on Se-enriched mushrooms shows SeMet to be the major Se species and bioavailability to be good⁽⁵⁷⁾. A speciation effect may be responsible for the bioavailability of Se from fish being inconsistent⁽⁹¹⁾; one study has shown a daily intake of 115 µg Se from fish to be unable to increase Se status⁽⁷⁶⁾.

There is good evidence that the increased Se status attained after supplementation with organic forms of Se is retained for a longer period after supplementation has ceased than is the case with selenite or selenate⁽¹²⁾. Reported whole-body half-lives of SeMet and selenite in humans were 252 and 102 d respectively, implying that Se administered as SeMet is retained 2.5 times longer in the body than is selenite⁽⁸⁵⁾. Accordingly, foods or supplements containing SeMet can maintain the activities of selenoenzymes during Se depletion for longer periods of time than those containing inorganic Se owing to the recycling of SeMet catabolised from protein stores⁽⁸⁵⁾.

No bioavailability data exist for *Se*-methyl-selenocysteine or γ -glutamyl-*Se*-methyl-selenocysteine.

Health effects associated with specific selenium species in foods and supplements

While the nutritionally essential functions of Se are understood to be fulfilled by the selenoproteins, dietary Se can be metabolised to small-molecular-weight species that have more recently generated interest because of putative anti-cancer effects. In contrast to such beneficial effects, at a sufficiently high dose level, Se metabolites can also cause toxicity.

Species-related beneficial effects

Though supplementation with Se or a good Se intake or status has been associated with health benefits, there is little or no evidence to connect such benefits with particular Se species. We know from studies in transgenic mice that selenoproteins are important for the cancer-protective effects of Se⁽⁹²⁾ and it

seems likely that antioxidant selenoproteins may be of benefit in counteracting diseases of oxidative stress. However, selenoproteins can be synthesised more or less efficiently from many different Se species, though if consumed in foods, they are digested and must be resynthesised as shown in Fig. 2.

In mice with genetically impaired selenoprotein expression, the presence of low-molecular-weight selenocompounds has been shown to reduce colon cancer risk⁽⁹²⁾. Such low-molecular-weight selenocompounds may be an *in vivo* source of the methylated metabolite, CH₃SeH, which is believed to be responsible for the potent anti-carcinogenic and anti-angiogenic effects of Se shown in the rat mammary tumour model and in cells in culture^(5,60,93–97). As shown in Fig. 2 and explained above, CH₃SeH can be formed directly from the low-molecular-weight selenocompounds *Se*-methyl-selenocysteine, by the action of a β -lyase⁽¹¹⁾, and SeMet by the action of a γ -lyase, also known as methioninase^(13,15,16,97–99).

Se-methyl-selenocysteine and its γ -glutamyl-derivative are found in a number of edible plants, including garlic, onions and broccoli and others of the *Allium* and *Brassica* families, particularly when grown in Se-enriched conditions^(5,23,60). Se-enriched plants such as broccoli and garlic have been shown to have potent anti-tumour effects in animals that are attributed to the presence of these species^(60,96). Though these species have not yet been tested in human interventions, a number of groups are planning pharmacokinetic studies as a prelude to human trials (Dr C Ip (2006), personal communication). Small amounts of both *Se*-methyl-selenocysteine and γ -glutamyl-*Se*-methyl-selenocysteine have also been identified in *Se*-yeast which may possibly be relevant to the anti-cancer effects seen in human trials with *Se*-yeast^(26,27). *Se*-methyl-selenocysteine has been commercially available for some time and can be bought over the counter as a supplement.

Though there is as yet no evidence of it, it appears possible that Se analogues of anti-cancer sulfur compounds such as diallyl disulfide and ajoene may also be isolable from Se-enriched garlic or onions. As diallylselenide was found to be more than 300 times more effective than diallylsulfide in protecting against carcinogen-induced mammary adenocarcinoma in rats⁽⁹⁷⁾, attempts to find such species may be worthwhile.

Species-related toxic effects

More is known about species-related toxic effects of Se than about species-related beneficial effects. The toxicity of Se and the mechanisms by which it exerts its toxic effects depend on its form, though there are few species-specific data on the toxicity of Se in humans and none relating to dose nor safe upper limits of particular species.

It is likely that a number of different mechanisms are involved in Se toxicity. According to Spallholz *et al.*^(97,98), Se compounds that can easily form the anion, RSe[–], generate superoxide in the presence of thiols such as glutathione, resulting in redox cycling, cell-cycle arrest and apoptosis. Spallholz ascribes the toxic (and indeed the carcinostatic) effects of Se to this oxidative-stress mechanism. Superoxide has been shown to be generated from selenite and diselenides such as selenocystamine in the presence of reduced glutathione *in vitro*, though not from selenate, SeMet or *Se*-methyl-selenocysteine⁽⁹⁷⁾. Neither SeMet nor *Se*-methyl-selenocysteine is

very toxic to cells in culture nor to animals or humans in line with their inability to generate superoxide, although both are capable of conversion to CH₃SeH by enzymic systems either *in vitro* or *in vivo*⁽⁹⁷⁾.

Selenodiglutathione, an intermediate in the formation of superoxide from selenite and glutathione, has been found to be even more toxic than selenite itself^(98,99). However, in contradiction to Spallholz's belief, Harrison *et al.*⁽¹⁰⁰⁾ did not find that the growth inhibition observed with this compound resulted from induction of an oxidative-stress mechanism, at least not of the type observed with oxidants such as H₂O₂. Supporting an oxidative-stress mechanism, selenite-induced redox cycles have been suggested to be responsible for oxygen-dependent DNA fragmentation in Se toxicity to hepatocyte model systems⁽¹⁰¹⁾ and high levels of selenite have been shown to induce the formation of 8-hydroxy-2-deoxyguanosine in rat liver DNA⁽¹⁰²⁾.

Other suggested mechanisms of Se toxicity include inhibition of Se methylation, the major detoxification pathway for Se, allowing the accumulation of hepato-toxic selenides, notably H₂Se. For instance, in mice, high doses of SeCys have been shown to cause hepatic toxicity by depressing Se methylation through the inactivation of methionine adenosyltransferase, the enzyme responsible for *S*-adenosyl methionine synthesis⁽¹⁰³⁾.

Although it has been suggested that organic forms of Se may be more toxic than inorganic forms during long-term consumption as they can be incorporated into tissue proteins rather than be excreted rapidly⁽¹⁰⁴⁾, there is no evidence that this is the case⁽⁴⁰⁾. Long-term supplementation studies with Se-yeast (60–80% of which is SeMet) at doses of 200, 300, 400 and even 800 µg Se/d for lengthy periods (up to 12 years in the case of the 200 µg dose) have been carried out by a number of research groups without any indication of toxic effects (for references, see Rayman⁽¹²⁾). Furthermore men with prostate cancer tolerated doses of 1600 and 3200 µg Se/d, as Se-yeast, for almost 12 months 'without any obvious Se-related serious toxicity'⁽¹⁰⁵⁾. Thus these results imply that uncontrolled accumulation of tissue Se does not occur.

Though there is no direct evidence in humans, it is generally accepted on the basis of animal studies that inorganic forms of Se are more acutely toxic than organic forms, selenite being slightly more toxic than selenate⁽⁴⁰⁾. Though of equivalent toxicity to SeCys in animals, sodium selenite is considerably more acutely toxic than SeMet, dimethyl selenide, trimethyl selenonium ion, selenoethers, selenobetaine or Se-yeast, the major Se component of which is SeMet⁽⁴⁰⁾. From lethal dose 50% (LD₅₀) determinations, selenite was found to be four-fold more toxic than SeMet when administered to mice intravenously⁽¹⁰⁶⁾ and three-fold more toxic than Se-yeast when given orally to rats⁽¹⁰⁷⁾.

Chronic toxicity of SeCys is equivalent to that of selenite and both are more toxic than SeMet (the L-isomer of which is more toxic than the D-isomer) and other organic Se compounds in animal studies⁽⁴⁰⁾. Comparison of selenite and Se-yeast diets in rats showed that Se-yeast was much less toxic than selenite; although the livers of animals fed Se-yeast showed up to 50% greater deposition of Se, there was no corresponding toxicity, as evidenced by histological examination⁽¹⁰⁸⁾. Se-yeast also seems to be less toxic than L-SeMet;

after 2 weeks of feeding 30 µg Se/g diet, survival in mallard ducklings was 36% for L-SeMet and 88% for Se-yeast⁽¹⁰⁹⁾. Human studies have also shown a lower chronic toxicity of organically bound Se, though there are limited data on the toxicity of individual compounds⁽⁴⁰⁾. However, SeMet is known to be the main Se species present in the diet of Chinese who developed chronic selenosis from consumption of maize and rice grown in the Enshi area of China⁽³⁹⁾.

The toxicity of the Se-accumulators to livestock has been linked to the high levels of *Se*-methyl-selenocysteine found in these species⁽¹¹⁰⁾. Se-accumulator plants are able to circumvent the toxicity that would result from the non-specific integration of the seleno-amino acids SeCys and SeMet into proteins by converting the precursor, SeCys, into the non-protein amino acids *Se*-methyl-selenocysteine, γ-glutamyl-*Se*-methyl-selenocysteine and selenocystathionine⁽⁸⁾. The potent toxicity of Se-accumulator plants to grazing animals is probably more a reflection of the extremely high concentrations of Se that can build up in these plants – up to 10–15 mg Se/g dry weight even on non-seleniferous soils⁽⁸⁾ – rather than the toxicity of *Se*-methyl-selenocysteine *per se*. According to Dr C Ip (2006, personal communication) who has worked with *Se*-methyl-selenocysteine for many years, it should be a safer compound than SeMet based on its biochemistry; though both compounds are equally well absorbed, *Se*-methyl-selenocysteine is converted to excretable metabolites more rapidly resulting in lower tissue retention of Se. Comparison of the no observable adverse effect level (NOAEL) in male and female rats for *Se*-methyl-selenocysteine (1.0 and 0.5 mg/kg per d, respectively) with that for selenite (0.14 and 0.2 mg/kg per d, respectively) suggests that *Se*-methyl-selenocysteine is less toxic at least than selenite⁽¹¹¹⁾ (C Ip (2006), personal communication). Results from Hasegawa *et al.*⁽¹⁰³⁾ similarly suggest that methylated forms of Se are generally less toxic than non-methylated compounds. This postulated lower toxicity may be highly relevant to the potential for use of *Se*-methyl-selenocysteine in human cancer prevention studies.

Conclusion

The development of state-of-the-art analytical methods that combine elemental and molecular mass spectrometric detection to investigate different chemical forms of Se in food has made possible the identification of a variety of Se species in foods and supplements. However, this is such a difficult and exacting area of research that, to date, we have only scratched the surface. It is difficult to maintain the integrity of species through the extraction process. Though we may know the identity of some Se species present in foods, there is no case where we know the identity of all the Se species; only where we have mass balance can we ensure that all species have been captured. We need to take food processing and preparation into account so that we are actually investigating the species that will be consumed (for example, Japanese soup stock made from shiitake mushrooms⁽⁵⁶⁾).

There remain considerable gaps in our knowledge of the forms of Se that naturally occur in foods. For instance, we know little about species of Se, other than SeMet, in fish, normally considered a good source of the element, or indeed what Se–Hg species may be present; we need to know full

speciation of Se in Se-yeast because of its frequent use in human intervention studies; and perhaps most importantly, there is a need to know to which Se species beneficial or detrimental health effects can be attributed.

We need to continue to develop speciation methodology, and to further investigate biosynthetic and metabolic pathways in order to have a steer on what species we should be searching for. Where we do suspect we know the identity of an active species (for example, *Se*-methyl-selenocysteine), we need single-species trials to prove efficacy or relative efficacy to help us towards a better understanding of how dietary Se should be supplemented.

Finally, there is a clear need for analytical chemists to present the data in a form that is understandable to and usable by consumers, nutritionists and legislators. Without adequate knowledge of Se speciation, false conclusions may be drawn when assessing Se requirements for optimal health. Furthermore, the ability to identify and accurately quantify Se species with powerful anti-cancer or other valuable effects will be essential for the development of plant-breeding programmes to optimise the biosynthesis of such species if clear proof of their health effects should be forthcoming.

Acknowledgements

M. P. R. wrote the manuscript, prepared the figures and compiled the first draft of the table. H. G. I. contributed to sections of the manuscript and to the table. M. S. initiated the writing of the review and advised on its content. The authors have no conflict of interest to declare.

References

- Clark LC, Combs GF Jr, Turnbull BW, *et al.* (1996) Effects of selenium supplementation for cancer prevention in patients with carcinoma of the skin. A randomized controlled trial. Nutritional Prevention of Cancer Study Group. *JAMA* **276**, 1957–1963.
- Lippman SM, Goodman PJ, Klein EA, *et al.* (2005) Designing the Selenium and Vitamin E Cancer Prevention Trial (SELECT). *J Natl Cancer Inst* **97**, 94–102.
- Ellis DR & Salt DE (2003) Plants, selenium and human health. *Curr Opin Plant Biol* **6**, 273–279.
- Whanger PD (1989) Selenocompounds in plants and their effects on animals. In *Toxicants of Plant Origin, Vol. III, Proteins and Amino Acids*, pp. 141–167 [PR Cheeke, editor]. Boca Raton, FL: CRC Press.
- Whanger PD (2004) Selenium and its relationship to cancer: an update. *Br J Nutr* **91**, 11–28.
- Terry N, Zayed AM, De Souza MP & Tarun AS (2000) Selenium in higher plants. *Annu Rev Plant Physiol Plant Mol Biol* **51**, 401–432.
- Tagmount A, Berken A & Terry N (2002) An essential role of S-adenosyl-L-methionine:L-methionine *S*-methyltransferase in selenium volatilization by plants. Methylation of selenomethionine to selenium-methyl-L-selenomethionine, the precursor of volatile selenium. *Plant Physiol* **130**, 847–856.
- Sors TG, Ellis DR & Salt DE (2005) Selenium uptake, translocation, assimilation and metabolic fate in plants. *Photosynth Res* **86**, 373–389.
- Whanger PD (2002) Selenocompounds in plants and animals and their biological significance. *J Am Coll Nutr* **21**, 223–232.
- Burk RF (1976) Selenium in man. In *Trace elements in Human Health and Disease*, Vol. 2, pp. 105–133 [AS Prasad, editor]. New York: Academic press.
- Combs GF Jr (2001) Selenium in global food systems. *Br J Nutr* **85**, 517–547.
- Rayman MP (2004) The use of high-selenium yeast to raise selenium status: how does it measure up? *Br J Nutr* **92**, 557–573.
- Suzuki K, Doi C & Suzuki N (2006) Metabolism of 76Se-methyl-selenocysteine compared with that of 77Se-methylselenomethionine and 82-selenite. *Toxicol Appl Pharmacol* **217**, 185–195.
- Suzuki KT, Somekawa L & Suzuki N (2006) Distribution and reuse of ⁷⁶Se-selenosugar in selenium-deficient rats. *Toxicol Appl Pharmacol* **216**, 303–308.
- Okuno T, Kubota T, Kuroda T, Ueno H & Nakamuro K (2001) Contribution of enzymic α , γ -elimination reaction in detoxification pathway of selenomethionine in mouse liver. *Toxicol Appl Pharmacol* **176**, 18–23.
- Okuno T, Ueno H & Nakamuro K (2006) Cystathione γ -lyase contributes to selenomethionine detoxification and cytosolic glutathione peroxidase biosynthesis in mouse liver. *Biol Trace Elem Res* **109**, 155–171.
- Berry MJ, Banu L, Chen YY, Mandel SJ, Kieffer JD, Harney JW & Larsen PR (1991) Recognition of UGA as a selenocysteine codon in type I deiodinase requires sequences in the 3' untranslated region. *Nature* **353**, 273–276.
- Berry MJ, Banu L, Harney JW & Larsen PR (1993) Functional characterization of the eukaryotic SECIS elements which direct selenocysteine insertion at UGA codons. *Embo J* **12**, 3315–3322.
- Francesconi KA & Pannier F (2004) Selenium metabolites in urine: a critical overview of past work and current status. *Clin Chem* **50**, 2240–2253.
- Kobayashi Y, Ogra Y, Ishiwata K, Takayama H, Aimi N & Suzuki KT (2002) Selenosugars are key and urinary metabolites for selenium excretion within the required to low-toxic range. *Proc Natl Acad Sci U S A* **99**, 15932–15936.
- Kuehnelt D, Kienzl N, Traar P, Le NH, Francesconi KA & Ochi T (2005) Selenium metabolites in human urine after ingestion of selenite, L-selenomethionine, or DL-selenomethionine: a quantitative case study by HPLC/ICPMS. *Anal Bioanal Chem* **383**, 235–246.
- Kuehnelt D, Juresa D, Kienzl N & Francesconi KA (2006) Marked individual variability in the levels of trimethylselenonium ion in human urine determined by HPLC/ICPMS and HPLC/vapor generation/ICPMS. *Anal Bioanal Chem* **386**, 2207–2212.
- Kotrebai M, Birringer M, Tyson JF, Block E & Uden PC (2000) Selenium speciation in enriched and natural samples by HPLC-ICP-MS and HPLC-ESI-MS with perfluorinated carboxylic acid ion-pairing agents. *Analyst* **125**, 71–78.
- Uden PC, Totoe Boakye H, Kahakachchi C, Hafezi R, Nolibos P, Block E, Johnson S & Tyson JF (2003) Element selective characterization of stability and reactivity of selenium species in selenized yeast. *J Anal At Spectrom* **18**, 1–10.
- Larsen EH, Hansen M, Paulin H, Moesgaard S, Reid M & Rayman M (2004) Speciation and bioavailability of selenium in yeast-based intervention agents used in cancer chemoprevention studies. *J AOAC Int* **87**, 225–232.
- Goenaga Infante H, O'Connor G, Rayman M, Wahlen R, Entwistle J, Norris P, Hearn R & Catterick T (2004) Selenium speciation analysis of selenium-enriched supplements by HPLC with ultrasonic nebulisation ICP-MS and electrospray MS/MS detection. *J Anal At Spectrom* **19**, 1529–1538.
- Goenaga Infante H, O'Connor G, Rayman M, Wahlen R, Spalholz J, Norris P, Hearn R & Catterick T (2005) Identification of water-soluble γ -glutamyl-CH₃SeCys in yeast-based Se supplements by HPLC with ICP-MS and electrospray tandem MS. *J Anal At Spectrom* **20**, 864–870.

28. Goenaga Infante H, Hearn R & Catterick T (2005) Current mass spectrometry strategies for selenium speciation in dietary sources of high-selenium. *Anal Bioanal Chem* **382**, 957–967.
29. Mester Z, Willie S, Lu Y, et al. (2006) Certification of a new selenised yeast reference material (SELM-1) for methionine, selenomethionine and total selenium content and its use in an intercomparison exercise for quantifying these analytes. *Anal Bioanal Chem* **385**, 168–180.
30. Encinar JR, Ruzik R, Buchmann W, Tortajada J, Lobinski R & Szpunar J (2003) Detection of selenocompounds in a tryptic digest of yeast selenoprotein by MALDI time-of-flight MS prior to their structural analysis by electrospray ionization triple quadrupole MS. *Analyst* **128**, 220–224.
31. Lindemann T & Hintelmann H (2002) Identification of selenium-containing glutathione S-conjugates in a yeast extract by two-dimensional liquid chromatography with inductively coupled plasma MS and nanoelectrospray MS/MS detection. *Anal Chem* **74**, 4602–4610.
32. Encinar JR, Ouerdane L, Buchmann W, Tortajada J, Lobinski R & Szpunar J (2003) Identification of water-soluble selenium-containing proteins in selenized yeast by size-exclusion-reversed-phase HPLC/ICPMS followed by MALDI-TOF and electrospray Q-TOF mass spectrometry. *Anal Chem* **75**, 3765–3774.
33. Goenaga Infante H, O'Connor G, Rayman M, Hearn R & Cook K (2006) Simultaneous identification of selenium-containing glutathione species in selenised yeast by on-line HPLC with ICP-MS and electrospray ionisation quadrupole time of flight (QTOF)-MS/MS. *J Anal At Spectrom* **21**, 1256–1263.
34. Gissel Nielsen G, Gupta UC, Lamand M & Westermark T (1984) Selenium in soils and plants and its importance in livestock and human nutrition. *Adv Agron* **37**, 397–460.
35. Lintscherger J, Fuchs N, Moser J, Kuehnelt D & Goessler W (2000) Selenium-enriched sprouts. A raw material for fortified cereal-based diets. *J Agric Food Chem* **48**, 5362–5368.
36. Yang X, Tian Y, Ha P, Gu L & Xia Y (1997) Determination of the selenomethionine content in grain and human blood (article in Chinese). *Weisheng Yanjiu* **26**, 113–116.
37. Warburton E & Goenaga Infante H (2007) Methane mixed plasma-improved sensitivity of inductively coupled plasma mass spectrometry detection for selenium speciation analysis of wheat-based food. *J Anal At Spectrom* **22**, 370–376.
38. Whanger P, Vendeland S, Park YC & Xia Y (1996) Metabolism of subtoxic levels of selenium in animals and humans. *Ann Clin Lab Sci* **26**, 99–113.
39. Beilstein MA, Whanger PD & Yang GQ (1991) Chemical forms of selenium in corn and rice grown in a high selenium area of China. *Biomed Environ Sci* **4**, 392–398.
40. Barceloux DG (1999) Selenium. *J Toxicol Clin Toxicol* **37**, 145–172.
41. Chang JC, Gutenmann WH, Reid CM & Lisk DJ (1995) Selenium content of Brazil nuts from two geographic locations in Brazil. *Chemosphere* **30**, 801–802.
42. Barclay MNI, MacPherson A & Dixon J (1995) Selenium content of a range of UK foods. *J Food Compost Anal* **8**, 307–318.
43. Lisk DJ, Bache CA, Essick LA, Reid CM, Rutzke M & Crown K (1988) Absorption and excretion of selenium and barium in humans from consumption of Brazil nuts. *Nutr Rep Int* **38**, 183–191.
44. Palmer I, Herr A & Nelson T (1982) Toxicity of selenium in Brazil nuts to rats. *J Food Sci* **47**, 1595–1597.
45. Wrobel K, Kannamkumarath SS, Wrobel K & Caruso JA (2003) Hydrolysis of proteins with methanesulfonic acid for improved HPLC–ICP-MS determination of seleno-methionine in yeast and nuts. *Anal Bioanal Chem* **375**, 133–138.
46. Kannamkumarath SS, Wrobel K, Wrobel K, Vonderheide A & Caruso JA (2002) HPLC–ICP-MS determination of selenium distribution and speciation in different types of nut. *Anal Bioanal Chem* **373**, 454–460.
47. Dernovics M, Giusti P & Lobinski R (2007) ICP-MS-assisted nanoHPLC-electrospray Q/time-of-flight MS/MS selenopeptide mapping in Brazil nuts. *J Anal At Spectrom* **22**, 41–50.
48. Dernovics M, García-Barrera T, Bierla K, Preud'homme H & Lobinski R (2007) Standardless identification of selenocystathione and its γ -glutamyl derivatives in monkeypot nuts by 3D liquid chromatography with ICP-MS detection followed by nano HPLC-Q-TOF-MS/MS. *Analyst* **132**, 439–449.
49. Smrkolj P, Stibilj V, Kreft I & Kapolna E (2005) Selenium species determination in selenium-enriched pumpkin (*Cucurbita pepo L.*) seeds by HPLC-UV-HG-AFS. *Anal Sci* **21**, 1501–1504.
50. Cai X-J, Block E, Uden PC, Zhang X, Quimby BD & Sullivan JJ (1995) Allium chemistry: identification of selenoamino acids in ordinary and selenium-enriched garlic, onion and broccoli using gas chromatography with atomic emission detection. *J Agric Food Chem* **43**, 1754–1757.
51. Finley JW, Ip C, Lisk DJ, Davis CD, Hintze KJ & Whanger PD (2001) Cancer-protective properties of high-selenium broccoli. *J Agric Food Chem* **49**, 2679–2683.
52. Pedrero Z, Elvira D, Cámaras C & Madrid Y (2007) Selenium transformation studies during Broccoli (*Brassica oleracea*) growing process by liquid chromatography-inductively coupled plasma mass spectrometry (LC-ICP-MS). *Anal Chim Acta* **596**, 251–256.
53. Hamilton JW (1975) Chemical examination of seleniferous cabbage *Brassica oleracea capitata*. *J Agric Food Chem* **23**, 1150–1152.
54. Pedrero Z, Madrid Y & Cámaras C (2006) Selenium species bioaccessibility in enriched radish (*Raphanus sativus*): a potential dietary source of selenium. *J Agric Food Chem* **54**, 2412–2417.
55. Gergely V, Kubachka KM, Mounicou S, Fodor P & Caruso JA (2006) Selenium speciation in *Agaricus bisporus* and *Lentinula edodes* mushroom proteins using multi-dimensional chromatography coupled to inductively coupled plasma mass spectrometry. *J Chromatogr A* **1101**, 94–102.
56. Ogra Y, Ishiwata K, Ruiz-Encinar J, Lobinski R & Suzuki KT (2004) Speciation of selenium in selenium-enriched shiitake mushroom, *Lentinula edodes*. *Anal Bioanal Chem* **379**, 861–866.
57. Rayman MP, Angus F & Goenaga-Infante H (2007) Selenium-enriched mushrooms as a functional food: bioavailability and speciation of selenium. *Proc Nutr Soc* **66**, 55A.
58. Shah M, Kannamkumarath SS, Wuilloud JCA, Wuilloud RG & Caruso JA (2004) Identification and characterization of selenium species in enriched green onion (*Allium fistulosum*) by HPLC–ICP-MS and ESI-ITMS. *J Anal At Spectrom* **19**, 381–386.
59. Bird SM, Ge H, Uden PC, Tyson JF, Block E & Denoyer E (1997) High-performance liquid chromatography of selenoamino acids and organo selenium compounds. Speciation by inductively coupled plasma mass spectrometry. *J Chromatogr A* **789**, 349–359.
60. Ip C, Birringer M, Block E, Kotrebai M, Tyson JF, Uden PC & Lisk DJ (2000) Chemical speciation influences comparative activity of selenium-enriched garlic and yeast in mammary cancer prevention. *J Agric Food Chem* **48**, 2062–2070.
61. Dumont E, Ogra Y, Vanhaecke F, Suzuki KT & Cornelis R (2006) Liquid chromatography-mass spectrometry (LC-MS): a powerful combination for selenium speciation in garlic (*Allium sativum*). *Anal Bioanal Chem* **384**, 1196–1206.
62. McSheehy S, Yang W, Pannier F, Szpunar J, Lobinski R, Auger J & Potin-Gautier M (2000) Speciation analysis of selenium in garlic by two-dimensional high-performance liquid

- chromatography with parallel inductively coupled plasma mass spectrometric and electrospray tandem mass spectrometric detection. *Anal Chim Acta* **421**, 147–153.
63. Whanger PD, Ip C, Polan CE, Uden PC & Welbaum G (2000) Tumorigenesis, metabolism, speciation, bioavailability, and tissue deposition of selenium in selenium-enriched ramps (*Allium tricoccum*). *J Agric Food Chem* **48**, 5723–5730.
 64. Kápolna E, Shah M, Caruso J & Fodor P (2007) Selenium speciation studies in Se-enriched chives (*Allium schoenoprasum*) by HPLC–ICP-MS. *Food Chem* **101**, 1398–1406.
 65. Stewart JM, Nigam SN & McConnell WB (1974) Metabolism of Na⁷⁵SeO₄ in horseradish: formation of selenosinigrin. *Can J Biochem* **52**, 144–145.
 66. Behre D, Hammel C, Pfeifer H, Rothlein D, Gessner H & Kyriakopoulos A (1998) Speciation of selenium in the mammalian organism. *Analyst* **123**, 871–873.
 67. Cabanero AI, Carvalho C, Madrid Y, Batoreu C & Camara C (2005) Quantification and speciation of mercury and selenium in fish samples of high consumption in Spain and Portugal. *Biol Trace Elem Res* **103**, 17–35.
 68. Cappon CJ & Smith JC (1982) Chemical form and distribution of mercury and selenium in canned tuna. *J Appl Toxicol* **2**, 181–189.
 69. Cappon CJ & Smith JC (1982) Chemical form and distribution of mercury and selenium in edible seafood. *J Anal Toxicol* **6**, 10–21.
 70. Cappon CJ & Smith JC (1981) Mercury and selenium content and chemical form in fish muscle. *Arch Environ Contam Toxicol* **10**, 305–319.
 71. Crews H, Clarke PA, Lewis DL, Owen LM, Strutt PR & Izquierdo A (1996) Investigation of selenium speciation in *in vitro* gastrointestinal extracts of cooked cod by high-performance liquid chromatography-inductively coupled plasma mass spectrometry and electrospray mass spectrometry. *J Anal At Spectrom* **11**, 1177–1182.
 72. Diaz Huerta V, Fernandez Sanchez ML & Sanz-Medel A (2004) Quantitative selenium speciation in cod muscle by isotope dilution ICP-MS with a reaction cell: comparison of different reported extraction procedures. *J Anal At Spectrom* **19**, 644–648.
 73. Dumont E, Vanhaecke F & Cornelis R (2006) Selenium speciation from food source to metabolites: a critical overview. *Anal Bioanal Chem* **385**, 1304–1323.
 74. Reilly C (2006) *Selenium in Food and Health*, 2nd ed. New York: Springer.
 75. Huang W, Akesson B, Svensson BG, Schutz A, Burk RF & Skerfving S (1995) Selenoprotein P and glutathione peroxidase (EC 1.11.1.9) in plasma as indices of selenium status in relation to the intake of fish. *Br J Nutr* **73**, 455–461.
 76. Meltzer HM, Bibow K, Paulsen IT, Mundal HH, Norheim G & Holm H (1993) Different bioavailability in humans of wheat and fish selenium as measured by blood platelet response to increased dietary selenium. *Biol Trace Elem Res* **36**, 229–241.
 77. Endo T, Haraguchi K, Hotta Y, Hisamichi Y, Lavery S, Dalebout ML & Baker CS (2005) Total mercury, methyl mercury, and selenium levels in the red meat of small cetaceans sold for human consumption in Japan. *Environ Sci Technol* **39**, 5703–5708.
 78. Bates CJ, Prentice A, Birch MC, Delves HT & Sinclair KA (2006) Blood indices of selenium and mercury, and their correlations with fish intake, in young people living in Britain. *Br J Nutr* **96**, 523–531.
 79. Bates CJ, Prentice A, Birch MC & Delves HT (2007) Dependence of blood indices of selenium and mercury on estimated fish intake in a national survey of British adults. *Public Health Nutr* **10**, 508–517.
 80. Dyrssen D & Wedborg M (1991) The sulfur-mercury(II) system in natural waters. *Water Air Soil Pollut* **56**, 507–519.
 81. Wageman R, Trebacz E, Boila G & Lockhart WL (1999) Mercury species in the liver of ringed seals. *Sci Total Environ* **261**, 21–32.
 82. Burk RF, Norsworthy BK, Hill KE, Motley AK & Byrne DW (2006) Effects of chemical form of selenium on plasma biomarkers in a high-dose human supplementation trial. *Cancer Epidemiol Biomarkers Prev* **15**, 804–810.
 83. Fox TE, Van den Heuvel EG, Atherton CA, *et al.* (2004) Bioavailability of selenium from fish, yeast and selenate: a comparative study in humans using stable isotopes. *Eur J Clin Nutr* **58**, 343–349.
 84. Fairweather-Tait SJ (1997) Bioavailability of selenium. *Eur J Clin Nutr* **51**, Suppl. 1, S20–S23.
 85. Combs GF (1988) Selenium in foods. In *Advances in Food Research*, vol. 32, pp. 85–113 [C Chichester and B Schweiger, editors]. San Diego: Academic Press.
 86. Sloth JJ, Larsen EH, Bugel SH & Moesgaard S (2003) Determination of total selenium and ⁷⁷Se in isotopically enriched human samples by ICP-dynamic reaction cell-MS. *J Anal At Spectrom* **18**, 317–322.
 87. Schrauzer GN (2000) Selenomethionine: a review of its nutritional significance, metabolism and toxicity. *J Nutr* **130**, 1653–1656.
 88. Ip C & Lisk DJ (1994) Bioactivity of selenium from Brazil nut for cancer prevention and selenoenzyme maintenance. *Nutr Cancer* **21**, 203–212.
 89. World Health Organization (1987) *Selenium. A Report of the International Programme on Chemical Safety*. Environmental Health Criteria 58. Geneva: WHO.
 90. Fordyce FM (2005) Selenium deficiency and toxicity in the environment. In *Essentials of Medical Geology*, pp. 373–415 [O Selinus, B Alloway, JA Centeno, RB Finkelman, R Fuge, U Lindh and P Smedley, editors]. London: Elsevier.
 91. Rayman MP (2000) The importance of selenium to human health. *Lancet* **356**, 233–241.
 92. Irons R, Carlson BA, Hatfield DL & Davis CD (2006) Both selenoproteins and low molecular weight selenocompounds reduce colon cancer risk in mice with genetically impaired selenoprotein expression. *J Nutr* **136**, 1311–1317.
 93. Ip C (1998) Lessons from basic research in selenium and cancer prevention. *J Nutr* **128**, 1845–1854.
 94. Ip C, Dong Y & Ganther HE (2002) New concepts in selenium chemoprevention. *Cancer Metastasis Rev* **21**, 281–289.
 95. Jiang C, Jiang W, Ip C, Ganther H & Lu J (1999) Selenium-induced inhibition of angiogenesis in mammary cancer at chemopreventive levels of intake. *Mol Carcinog* **26**, 213–225.
 96. Davis CD & Finley JW (2003) Chemical versus food forms of selenium in cancer prevention. In *Functional Foods and Nutraceuticals in Cancer Prevention*, pp. 55–85 [RR Watson, editor]. Ames, IA: Iowa State Press.
 97. Spallholz JE, Palace VP & Reid TW (2004) Methioninase and selenomethionine but not Se-methylselenocysteine generate methylselenol and superoxide in an *in vitro* chemiluminescent assay: implications for the nutritional carcinostatic activity of selenoamino acids. *Biochem Pharmacol* **67**, 547–554.
 98. Nakamuro K, Nakanishi K, Okuno T, Hasegawa T & Sayato Y (1997) Comparison of methylated selenium metabolites in rats after oral administration of various selenium compounds. *Jpn J Toxicol Environ Health* **43**, 1482–1489.
 99. Wang Z, Jiang C & Lu J (2002) Induction of caspase-mediated apoptosis and cell-cycle G1 arrest by selenium metabolite methylselenol. *Mol Carcinog* **34**, 113–120.
 100. Harrison PR, Lanfear J, Wu L, Fleming J, McGarry L & Blower L (1997) Chemopreventive and growth inhibitory effects of selenium. *Biomed Environ Sci* **10**, 235–245.

101. Garberg P, Stahl A, Warholm M & Hogberg J (1988) Studies of the role of DNA fragmentation in selenium toxicity. *Biochem Pharmacol* **37**, 3401–3406.
102. Wycherly BJ, Moak MA & Christensen MJ (2004) High dietary intake of sodium selenite induces oxidative DNA damage in rat liver. *Nutr Cancer* **48**, 78–83.
103. Hasegawa T, Mihara M, Nakamuro K & Sayato Y (1996) Mechanisms of selenium methylation and toxicity in mice treated with selenocystine. *Arch Toxicol* **71**, 31–38.
104. Patterson B & Levander O (1997) Naturally occurring selenium compounds in cancer chemoprevention trials: a workshop summary. *Cancer Epidemiol Biomarkers Prev* **6**, 63–69.
105. Reid ME, Stratton MS, Lillico AJ, Fakih M, Natarajan R, Clark LC & Marshall JR (2004) A report of high-dose selenium supplementation: response and toxicities. *J Trace Elem Med Biol* **18**, 69–74.
106. Ammar EM & Couri D (1981) Acute toxicity of sodium selenite and selenomethionine in mice after ICV or IV administration. *Neurotoxicity* **2**, 383–386.
107. Vinson JA & Bose P (1987) Comparison of the toxicity of inorganic and natural selenium. In *Selenium in Biology and Medicine (Part A)*, pp. 513–515 [JE Combs, JE Spallholz, OA Levander and JE Oldfield, editors]. New York: Van Nostrand Reinhold Co.
108. Spallholz JE & Raftery A (1987) Nutritional, chemical, and toxicological evaluation of a high-selenium yeast. In *Selenium in Biology and Medicine (Part A)*, pp. 516–529 [JE Combs, JE Spallholz, OA Levander and JE Oldfield, editors]. New York: Van Nostrand Reinhold Co.
109. Heinz GH, Hoffman DJ & LeCaptain LJ (1996) Toxicity of seleno-L-methionine, seleno-DL-methionine, high selenium wheat, and selenized yeast to mallard ducklings. *Arch Environ Contam Toxicol* **30**, 93–99.
110. Spallholz JE (1994) On the nature of selenium toxicity and carcinostatic activity. *Free Radic Biol Med* **17**, 45–64.
111. Jia X, Li N & Chen J (2005) A subchronic toxicity study of elemental Nano-Se in Sprague–Dawley rats. *Life Sci* **76**, 1989–2003.