

switching. Think-aloud interviews with 4 psychiatrists and pharmacists have been conducted so far to refine iSWITCHED.

Conclusion: The iSWITCHED switching intervention combines lived experiences, clinical expertise and integrates seamlessly into existing MH record systems. Before wider-implementation, it will be refined using insights from think-aloud interviews and role-play activities and piloted in a larger feasibility study.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard BJPsych Open peer review process and should not be quoted as peer-reviewed by BJPsych Open in any subsequent publication.

Perinatal Factors Associated With Mental Health Problems Among Postnatal Women Referred to the University Psychiatry Unit, National Hospital of Sri Lanka

Dr Amanda Kasturi Mudali^{1,2}, Prof Chathurie Suraweera³, Dr TVM Pieris⁴, Dr MRIF Ariyachandra⁵ and Dr AGUS Gunathilaka⁶

¹University Psychiatry Unit, National Hospital of Sri Lanka, Colombo, Sri Lanka; ²Kent and Medway NHS Social Care and Partnership Trust, Maidstone, United Kingdom; ³Department of Psychiatry, Faculty of Medicine, University of Colombo, Colombo, Sri Lanka; ⁴National Institute of Mental Health, Colombo, Sri Lanka; ⁵Lady Ridgeway Hospital for Children, Colombo, Sri Lanka and ⁶Colombo North Teaching Hospital, Colombo, Sri Lanka

doi: [10.1192/bjo.2025.10158](https://doi.org/10.1192/bjo.2025.10158)

Aims: The postpartum period exerts a profound influence on maternal mental health. This study aimed to describe mental health problems and identify associated perinatal factors in women referred to the University Psychiatry Unit, National Hospital of Sri Lanka (NHSL) during the postnatal period.

Methods: A retrospective descriptive cross-sectional study with analytical component was conducted. Data was collected from records of 201 postnatal women randomly selected between 2021–2023 referred from De Soysa Maternity Hospital (DMH) and Castle Street Hospital for Women (CSHW). Women who experienced a perinatal death in their most recent pregnancy were excluded.

Results: The mean age of the study population was 30 years. Most were unemployed (73.4%), completed Advanced Level education (39.3%), lived in Colombo district (71.6%), and were married (89.1%) and referred before 5 days postpartum (52%). Among referred patients, 7% had history of depression. Affective symptoms (33.7%) and anxiety (11.2%) were common reasons for referral. The proportion of patients referred for suicidal thoughts was 4.3% and for heroin use was 8%. During assessment 67.2% were diagnosed with a mental illness. Majority had maternal blues (30.4%), depression in 11.8% and postpartum psychosis in 7.4%. Outpatient treatment was offered to 98% of those diagnosed with 81.3% offered pharmacotherapy. It was found that 34.6% defaulted and 21.6% were not sent for review during admission. Postpartum mental illness was significantly associated with gestational age at delivery ($\chi^2=8.347$, $p=0.004$), whether baby was admitted to Neonatal Intensive Care Unit or High Dependency Unit ($\chi^2=5.424$, $p=0.02$), and spousal support ($\chi^2=5.580$, $p=0.018$). No association was found with age, education, past psychiatric history, parity, mode of delivery, or breastfeeding difficulty.

Conclusion: The high rate of treatment defaulting indicates a need for improved follow-up and support strategies for women with postpartum mental illness. The identified associated factors of

gestational age, neonatal admission status and spousal support can be utilized to design targeted preventative interventions.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard BJPsych Open peer review process and should not be quoted as peer-reviewed by BJPsych Open in any subsequent publication.

A Study of the Relationship Between Mental Health and Online Platforms for Education

Dr Nastaran Kazemi Rad, Dr Atefeh Mohammad Jafari, Dr Zahra Mirsepassi and Dr Pedram Sarikhani

Tehran University of Medical Sciences, Tehran, Iran, Islamic Republic of

doi: [10.1192/bjo.2025.10159](https://doi.org/10.1192/bjo.2025.10159)

Aims: 1. Providing practical solutions for richer and broader use of online platforms for education among students. 2. Finding factors related to online education that are involved in causing mental disorders in students and preventing and correcting them.

Methods: This cross-sectional study was conducted after obtaining approval from the Research Council and Ethics Committee of Tehran University of Medical Sciences. Medical students were selected through simple random sampling, and those who met the inclusion criteria completed an online questionnaire that included the PHQ-9 and PSS-10 questionnaires developed by the researcher. Informed consent was obtained from all participants, and the data was analysed using SPSS software.

Results: A total of 330 medical students participated in the study, of whom 137 (41.5%) were female and 193 (58.5%) were male. The results revealed that 73.5% of participants had mild depression, 17.3% had moderate depression, and 9.4% had severe depression. Additionally, 4.2% had mild stress, 91.5% had moderate stress, and 4.2% had severe stress.

Conclusion: The study findings revealed that virtual education quality during the Covid-19 era, accessibility to virtual learning resources, financial and health worries due to Covid-19, and communication level among students with their family and friends significantly influence their mental health. The study suggests that while designing virtual classrooms, it is essential to consider the availability of resources, facilitate peer-group communication, and provide financial support to enhance students' mental health.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard BJPsych Open peer review process and should not be quoted as peer-reviewed by BJPsych Open in any subsequent publication.

A Study on Online Gaming and Its Effect on Emotional Regulation – Online Survey

Dr Poornima Khadanga¹, Dr Madhavi Rairikar², Dr Gaurav Pawar³ and Dr Gomati Nadadgalli⁴

¹Northamptonshire Healthcare Foundation NHS Trust, Northampton, United Kingdom; ²Salvation Church Medical Centre, Mumbai, India; ³Symbiosis Medical College for Women, Pune, India and ⁴KLE's JGMMC, Hubballi, India

doi: [10.1192/bjo.2025.10160](https://doi.org/10.1192/bjo.2025.10160)

Aims: The study aimed to study the prevalence of problematic online gaming and find out association between problematic gaming with domains of emotional regulation.

Methods: An online survey questionnaire was created by the researchers after approval and validation from experts and sent through various social media platforms. Those who did not consent for the study were excluded. The questionnaire included basic sociodemographic details and questions taken from two scales – (i) Problematic Online Gaming Questionnaire SF for identifying problematic online gaming and (ii) Difficulties in Emotional Regulation Scale (DERS-18) to identify domains of emotional regulation. Descriptive statistics, Pearson correlation, ANOVA were used for statistical analysis.

Results: 108 participants reverted with the completed forms, out of which, 69.4% were regularly engaged in online gaming, 19.4% fell under the domain of problematic online gaming. 4.6% of the participants were engaged in online gaming for more than 8 hours in a day. With regards to various domains of emotional regulation, the mean scores in each subscale of emotional regulation (Nonacceptance of Emotional Responses, Difficulties Engaging in Goal-Directed Behaviour, Impulse Control Difficulties, Lack of Emotional Awareness, Limited Access to Emotion Regulation Strategies and Lack of Emotional Clarity) were more in those with problematic online gaming as compared with non-problematic individuals and was statistically significant ($p < 0.05$). There was also a positive correlation between problematic online gaming and sub-domains of emotional regulation ($p < 0.01$).

Conclusion: The study clearly depicts the rise in online gaming that has been clearly demonstrated in the recent studies worldwide. The results clearly illustrate the association of online gaming with difficulty in handling emotions but also a means to attenuate negative emotions. It could be potentially being used as an escape mechanism. Though details of the causal link between the two parameters were beyond the scope of the study, it would be worth looking at it in future research.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Structured Literature Review of the Impact of Using Cognitive-behavioural Therapy and Interpersonal Therapy On Depressive Symptoms in People Living With HIV/AIDS.

Dr June Kilonzo

Cardiff University, Wales, United Kingdom

doi: [10.1192/bjo.2025.10161](https://doi.org/10.1192/bjo.2025.10161)

Aims: A diagnosis of depression is three times more likely to occur in people living with HIV/AIDS. Management is imperative, as it not only ensures reduced depressive symptoms but also improves quality of life. Interpersonal psychotherapy (IPT) and cognitive-behavioural therapy (CBT) have been linked to reduced depressive symptoms. This paper will investigate to what degree both treatment options can help reduce depressive symptoms in people living with HIV/AIDS.

Methods: Randomized controlled trial articles from the Cardiff University repository, PsycARTICLES, PsycINFO, Cinahl, and PubMed databases were collected and dated not later than ten years from 2024. The participants were individuals with an HIV/AIDS diagnosis and depression as a comorbidity. They also had been exposed to either IPT or CBT psychotherapy. The articles were screened, and data from 19 articles with 4805 participants were extracted.

Results: Findings showed that CBT reduced depressive symptoms, and its effects were visible in the long term, which involves more than six months. Similarly, IPT was found to have long-term effects, but the difference in the symptoms in IPT was minimal compared with CBT. No significant difference in the efficiency of either therapy was observed.

Conclusion: Both the psychotherapy modalities were found to have positive impact by reducing depressive symptoms. There is a need for more research into improving the psychotherapy intervention time and mode of delivery.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Exploring the Interplay Between Inflammatory and Metabolic Markers in Depression Treatment Outcomes: A Focus on Interleukin-6 and Ghrelin

Professor Jae-Min Kim

Chonnam National University Hospital, Gwangju, Korea, Republic of

doi: [10.1192/bjo.2025.10162](https://doi.org/10.1192/bjo.2025.10162)

Aims: This study aimed to elucidate the modulating effects of serum ghrelin on the relationships between interleukin-6 (IL-6) and antidepressant treatment outcomes, particularly focusing on 12-week remission and 24-month relapse in patients with depressive disorders.

Methods: We analysed baseline serum levels of ghrelin and IL-6 in 1,086 patients engaged in a naturalistic, stepwise antidepressant treatment protocol. Remission was assessed using the Hamilton Depression Rating Scale (HAMD), with a score of 7 or less defining remission at 12 weeks. Patients achieving a response ($\text{HAMD} \leq 14$) at this interval were tracked for relapse ($\text{HAMD} > 14$) quarterly up to 24 months. Logistic regression models, adjusting for sociodemographic and clinical variables, explored the interactive effects of these biomarkers on treatment outcomes.

Results: Our analysis indicated that while serum ghrelin levels did not directly impact treatment outcomes, they significantly modulated the relationship between high IL-6 levels and the likelihood of non-remission at 12 weeks as well as relapse at 24 months. Notably, elevated IL-6 was strongly associated with these negative outcomes primarily in the context of lower ghrelin levels. The modulatory effects of ghrelin were statistically significant in the context of relapse after controlling for relevant covariates.

Conclusion: The findings from this study underscore the critical interplay between inflammatory and metabolic markers in determining the trajectory of depression treatment outcomes. By demonstrating the significant roles of IL-6 and ghrelin, particularly their interactive effects, this research highlights the potential to enhance personalized antidepressant strategies through the integration of biomarker profiles. Future investigations should focus on unravelling the dynamic mechanisms behind these interactions, which could pave the way for refining prediction models for treatment responsiveness and developing targeted interventions that more effectively address the complexities of managing depression.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.