

However, shorter periods were not associated with worse concordance. The NHSN should evaluate whether the burden of a 3-month validation period is justified.

Funding: None

Disclosures: None

Doi:10.1017/ice.2020.973

Presentation Type:

Poster Presentation

Predicting Community-Onset Candidemia in an Academic Medical Center Using Machine Learning

Mohammed ALZunitan, University of Iowa Health Care; Alexandre Marra, Hospital Israelita Albert Einstein; Michael Edmond, University of Iowa Hospitals and Clinics; Nick Street, Department of Business Analytics, Henry B. Tippie College of Business, University of Iowa; John Cromwell; Daniel Diekema, University of Iowa Carver College of Medicine; Jorge Salinas, University of Iowa

Background: Candidemia is a leading cause of bloodstream infections (BSIs), and community-onset candidemia is being recognized as a public health problem. In the era of electronic health records (EHRs), we can use machine learning to detect patterns in patient data that may predict infections. **Objective:** We aimed to predict community-onset candidemia in patients admitted to the University of Iowa Hospital & Clinics (UIHC) using machine-learning algorithms. **Methods:** We retrospectively reviewed data for patients admitted to UIHC during 2015–2018. All adult inpatients who had a requested blood culture were included. Candidemia was defined as a blood culture positive for *Candida* within 48 hours after admission. Variables of interest were extracted from the EHR: age, sex, body mass index, and month of admission. We also included comorbidities upon admission defined by the *International Classification of Diseases, 10th Revision, Clinical Modification* (ICD-10-CM): cardiovascular

diseases, neurological disorders, chronic pulmonary disease, dementia, rheumatoid disease, peptic ulcer disease, liver disease, diabetes mellitus, hypothyroidism, renal failure, coagulopathy, obesity, weight loss, fluid and electrolyte disorders, anemia, alcohol abuse, drug abuse, psychiatric diseases, malignancy, and HIV/AIDS. We calculated Charlson and Elixhauser scores based on ICD-10-CM codes. We also included prehospitalization conditions (90 days before admission): *Candida*-positive cultures from sites other than blood, antibiotics/antifungals, hemodialysis, central lines, corticosteroids, surgeries, and intensive care unit (ICU) admissions. Mode and median imputation were used for missing information. Random forests with resampled training sets were used for prediction, and results were evaluated using 10-fold cross validation. **Results:** In total, 30,528 adult admissions were extracted; 73 admissions had an episode of candidemia (<1%). Median admission age was 61 years, and nearly half of admissions were female patients (44.7%). Mean BMI was 27.67. The most admissions occurred during the months of March, August, and November. The 3 most common ICD-10-CM codes were diabetes mellitus, hypertension, and cancer. Median Charlson and Elixhauser scores were 1 and 2, respectively. The model used 103 variables. The 3 most predictive variables were Elixhauser score on admission, and characteristics in the 90 days prior to admission were *Candida* from sites other than blood, use of a central line, and recent use of antibiotics/antifungals. The model's area under the receiver operating characteristic curve was 0.72. **Conclusions:** Preadmission patient characteristics predicted community-onset candidemia. Machine-learning models may help detect patients eligible for screening for candidemia and prompt empiric antifungal therapy in high-risk patients in the first 48 hours of their admission.

Funding: None

Disclosures: None

Doi:10.1017/ice.2020.974

Presentation Type:

Poster Presentation

Prediction of Surgical Risk in General Surgeries: Process Optimization Through Support Vector Machine (SVM) Algorithm

Flávio Souza, Centro Universitário de Belo Horizonte (UNIBH); Braulio Couto, Centro Universitário de Belo Horizonte – UniBH; Felipe Leandro Andrade da Conceição, Centro Universitário de Belo Horizonte; Gabriel Henrique Silvestre da Silva, Centro Universitário de Belo Horizonte; Igor Gonçalves Dias, Centro Universitário de Belo Horizonte; Rafael Vieira Magno Rigueira, Centro Universitário de Belo Horizonte; Gustavo Maciel Pimenta, Centro Universitário de Belo Horizonte; Maurilio Martins, Centro Universitário de Belo Horizonte; Julio Cesar Mendes, Centro Universitário de Belo Horizonte; Gabriele Maria Braga, Centro Universitário de Belo Horizonte; Jéssica Angelina Teixeira, Centro Universitário de Belo Horizonte; Renata Carvalho Santos, Centro Universitário de Belo Horizonte; Julia Maria Campos Martins, Centro Universitário de Belo Horizonte; Karla Silvia de Sousa, Centro Universitário de Belo Horizonte; Douglas Nascimento de Souza, Centro Universitário de Belo Horizonte; Gustavo Barros Alves, Centro Universitário de Belo Horizonte; Vladimir Alexei Rodrigues Rocha, Universidade Federal de Minas Gerais

Background: In 5 hospitals in Belo Horizonte (population, 3 million) between July 2016 and June 2018, a survey was performed regarding surgical site infection (SSI). We statistically evaluated




Figure 1: ROC curve

Fig. 1.

SSI incidents and optimized the power to predict SSI through pattern recognition algorithms based on support vector machines (SVMs). **Methods:** Data were collected on SSIs at 5 different hospitals. The hospital infection control committees (CCHs) of the hospitals collected all data used in the analysis during their routine SSI surveillance procedures; these data were sent to the NOIS (Nosocomial Infection Study) Project. NOIS uses SACIH software (an automated hospital infection control system) to collect data from hospitals that participate voluntarily in the project. In the NOIS, 3 procedures were performed: (1) a treatment of the database collected for use of intact samples; (2) a statistical analysis on the profile of the hospitals collected; and (3) an assessment of the predictive power of SVM with a nonlinear separation process varying in configurations including kernel function (Laplace, Radial Basis, Hyperbolic Tangent and Bessel) and the k-fold cross-validation-based resampling process (ie, the use of data varied according to the amount of folders that cross and combine the evaluated data, being $k = 3, 5, 6, 7,$ and 10). The data were compared by measuring the area under the curve (AUC; range, 0–1) for each of the configurations. **Results:** From 13,383 records, 7,565 were usable, and SSI incidence was 2.0%. Most patients were aged 35–62 years; the average duration of surgery was 101 minutes, but 76% of surgeries lasted >2 hours. The mean hospital length of stay without SSI was 4 days versus 17 days for the SSI cases. The survey data showed that even with a low number of SSI cases, the prediction rate for this specific surgery was 0.74, which was 14% higher than the rate reported in the literature. **Conclusions:** Despite the high noise index of the database, it was possible to sample relevant data for the evaluation of general surgery patients. For the predictive process, our results were >0.50 and were 14% better than those reported in the literature. However, the database requires more SSI case samples because only 2% of positive samples unbalanced the database. To optimize data collection and to enable other hospitals to use the SSI prediction tool, a mobile application was developed (available at www.sacihweb.com).

Funding: None

Disclosures: None

Doi:[10.1017/ice.2020.975](https://doi.org/10.1017/ice.2020.975)

Presentation Type:

Poster Presentation

Prescribing Pattern of Antibiotics among Children in a Tertiary-Care Hospital, Bangladesh

Sukanta Chowdhury, ICCDR,B; Rajib Sarker, Patuakhali Science and Technology University, Bangladesh; Md. Shafiqul Islam Khan, Patuakhali Science and Technology University, Bangladesh; Probir Kumar Ghosh, ICCDR,B; Md. Abu Tareq, Patuakhali Science and Technology University, Bangladesh; Musammet Rasheda Begum, Chattogram Veterinary and Animal Sciences University, Bangladesh

Background: The inappropriate and irrational use of antibiotics both in humans and animals causes bacterial resistance. Bacterial resistance is common in low- and middle-income countries, including Bangladesh. Bangladesh has very limited information on antibiotic use and associated resistance. We sought to better understand antibiotic use in low-resource settings for the development of effective strategies to address inappropriate antibiotic use. **Methods:** We conducted a cross-sectional study among hospitalized children <5 years of age in a tertiary-care hospital in Barishal, Bangladesh, to collect data on antibiotic use. We collected data from 400 children during February–April 2019. **Results:** Among these 400 children, >50% were aged <1 year, and >60%

of these children were boys. The average hospital stay was 3 days (range, 1–14). Most of the children had history of diarrhea and 18% had pneumonia. Most children (82%) were prescribed antibiotics. A combined form of antibiotics was prescribed for 17% of these children. In total, 14 different antibiotics were used. The most commonly used antibiotic was ceftriaxone (57%) followed by azithromycin (14%). The parental route was mostly preferred (75%) for antibiotic administration. **Conclusions:** Antibiotic prescription was common in children aged <5 years visiting a tertiary-care hospital. Most of the prescribed antibiotics were broad spectrum, which can promote bacterial resistance. Further studies are needed to identify the factors associated with overuse of antibiotics and bacterial resistance in low-resource settings.

Funding: None

Disclosures: None

Doi:[10.1017/ice.2020.976](https://doi.org/10.1017/ice.2020.976)

Presentation Type:

Poster Presentation

Prevalence and Carbapenem Resistance of *Acinetobacter baumannii* and Other Than *A. baumannii* Isolates From Intensive Care Units (ICUs) and non-ICUs

Tomasz Kasperski, Biophage Pharma S.A., Kraków; Agnieszka Chmielarczyk, Jagiellonian University Collegium Medicum; Monika Pomorska-Wesołowska, Analytical and Microbiological Laboratory of Ruda Slaska KORLAB; Dorota Romaniszyn, Jagiellonian University Medical School; Jadwiga Wojkowska-Mach, Jagiellonian University Medical School

Background: *Acinetobacter* spp are gram-negative bacteria that have emerged as a leading cause of hospital-associated infections, most often in the intensive care unit (ICU) setting. This is particularly important in Poland, where the prevalence of *A. baumannii* in various types of infections, including bloodstream infection (BSI), pneumonia, skin and soft-tissue infection (SSTI), and urinary tract infection (UTI) is higher than in neighboring countries. Recently, other *Acinetobacter* spp, including *A. lwoffii* or *A. ursingii*, have been found to be clinically relevant. In Poland, we have also observed a very rapid increase in antimicrobial resistance, significantly faster for *A. baumannii* than for other nosocomial pathogens. **Methods:** A study was conducted in 12 southern Polish hospitals, including 3 ICUs, from January 1 to December 31, 2018. Only adult hospitalized patients were included. Strains were identified using the MALDI-TOF method. Carbapenem resistance was determined using the minimum inhibitory concentration (MIC). **Results:** During the study, 194 strains belonging to the *Acinetobacter* genus were isolated. *A. baumannii* was the dominant species, 88.1% ($n = 171$), and 23 isolates (11.9%) were other *Acinetobacter* spp: *A. ursingii* ($n = 5$), *A. lwoffii* ($n = 4$), *A. haemolyticus* ($n = 4$), *A. junii* ($n = 3$), *A. radioresistens* ($n = 2$), *A. bereziniae* ($n = 2$), and *A. johnsonii* ($n = 2$). Moreover, 15 *Acinetobacter* strains were collected from ICUs. The most *Acinetobacter* strains were isolated from SSTIs ($n = 115$) from non-ICU settings. Non-*A. baumannii* strains were also most frequently isolated from SSTIs; they constituted 11.3% of all *Acinetobacter* strains from this type of infection ($n = 13$). The total *Acinetobacter* prevalence was 2.6%, whereas the prevalence in the ICU setting was 7%. *Acinetobacter* prevalence in SSTIs was 10.4%. In pneumonia, *Acinetobacter* prevalence was 18.6% for ICUs ($n = 13$) and 2.7% for non-ICUs ($n = 46$). Strains from UTIs were isolated only with the non-ICU setting, and their prevalence was 0.7% ($n = 14$). More