

THE SEMI-BALAYABILITY OF REAL CONVOLUTION KERNELS

MASAYUKI ITÔ AND NORIAKI SUZUKI

Dedicated to Professor Yukio Kusunoki on his 60th birthday

§1.

Let X be a locally compact, σ -compact and non-compact abelian group. Throughout this paper, we shall denote by ξ a fixed Haar measure on X and by δ the Alexandroff point of X .

A real convolution kernel (i.e., a real Radon measure) N on X is said to be semi-balayable if N satisfies the semi-balayage principle on all open sets (see Definition 6). We know that every convolution kernel N of logarithmic type is semi-balayable (see [8]). Here N is said to be of logarithmic type if, with a vaguely continuous, markovian, semi-transient and recurrent convolution semi-group $(\alpha_t)_{t \geq 0}$ of non-negative Radon measures on X ,

$$N * \mu = \int_0^\infty \alpha_t * \mu dt \left(= \lim_{t \rightarrow \infty} \int_0^t \alpha_s * \mu ds \right)^{1)}$$

for all real Radon measure μ on X with compact support and $\int d\mu = 0$.

In this paper, we shall show that the semi-balayability is an essential property to characterize convolution kernels of logarithmic type. More precisely, we shall establish the following theorems.

THEOREM 1. *Let N be a real convolution kernel on X . If $X \approx R \times F$ or $X \approx Z \times F$, we suppose an additional condition: $N = o(|x|)$ at the infinity²⁾. Then N is of logarithmic type if and only if N is semi-balayable, non-periodic and satisfies $\inf_{x \in X} N * f(x) \leq 0$ for any finite continuous function f on X with compact support and $\int f d\xi = 0$.*

Received May 28, 1984.

¹⁾ For a net $(\mu_\alpha)_{\alpha \in A}$ of real Radon measures and a real Radon measure μ , we write $\mu = \lim_{\alpha \in A} \mu_\alpha$ if $(\mu_\alpha)_{\alpha \in A}$ converges vaguely to μ along A .

²⁾ If $X = R \times F$ or $X = Z \times F$, $N = o(|x|)$ at the infinity means that for any $f \in C_K^+(X)$, $N * f((x, y)) = o(|x|)$ as $|x| \rightarrow \infty$, where $(x, y) \in R \times F$ or $\in Z \times F$. In the case of $X \approx R \times F$ or $X \approx Z \times F$, the definition $N = o(|x|)$ at the infinity follows naturally from the above definition.

Here R , Z and F denote the additive group of real numbers, the additive group of integers and a certain compact abelian group, respectively.

By virtue of the main theorems in [8] (see Théorèmes 52 and 52'), Theorem 1 follows immediately from the following

THEOREM 2. *Let N be a non-periodic real convolution kernel on X satisfying $\inf_{x \in X} N * f(x) \leq 0$ for any finite continuous function f on X with compact support and $\int f d\xi = 0$. Then N is semi-balayable if and only if N satisfies the semi-complete maximum principle and $\eta_{N,\delta} = -\infty$, i.e., for any exhaustion $(K_n)_{n=1}^\infty$ of X ³⁾ and any non-negative continuous function $f \neq 0$ on X with compact support, $\lim_{n \rightarrow \infty} \int f d\eta_{N,CK_n} = -\infty$, where η_{N,CK_n} is the N -reduced measure of N on CK_n .*

The “if” part is already known (see Proposition 28 in [8]), so that this paper will be devoted principally to the proof of the “only if” part.

It is interesting to compare Theorem 1 with the Choquet-Deny theorem for Hunt convolution kernels⁴⁾ (see [3]).

Contrary to a conjecture in [8] (see Problème 29), Theorem 1 shows that, under some additional conditions, non-periodic and semi-balayable real convolution kernels are of logarithmic type.

§ 2.

We denote by:

$C(X)$ the usual Fréchet space of finite continuous functions on X ;

$C_K(X)$ the usual topological vector space of finite continuous functions on X with compact support;

$M(X) = C_K(X)^*$ the topological vector space of real Radon measures on X with the vague (weak*) topology;

$M_K(X) = C(X)^*$ the usual topological vector space of real Radon measures on X with compact support;

$C^+(X)$, $C_K^+(X)$, $M^+(X)$ and $M_K^+(X)$ their subsets of non-negative elements.

Furthermore, we put

$$C_K^0(X) = \left\{ f \in C_K(X); \int f d\xi = 0 \right\} \quad \text{and} \quad M_K^0(X) = \left\{ \mu \in M_K(X); \int d\mu = 0 \right\}.$$

³⁾ An exhaustion $(K_n)_{n=1}^\infty$ of X means a sequence of compact sets satisfying $K_n \subset$ the interior of K_{n+1} and $\bigcup_{n=1}^\infty K_n = X$.

⁴⁾ A non-negative convolution kernel N_0 on X is a Hunt convolution kernel if and only if N_0 is balayable (see Remark 14 (3)) and not pseudo-periodic.

DEFINITION 3. A real convolution kernel N on X is said to satisfy the semi-complete maximum principle (denoted by $N \in (\text{SMP})$) if for any $f, g \in C_K^+(X)$ with $\int f d\xi = \int g d\xi$ and any $a \in R$, we have the implication:

$$N * f(x) \leq N * g(x) + a \quad \text{on } \text{supp}(f) \implies N * f(x) \leq N * g(x) + a \quad \text{on } X,$$

where $\text{supp}(f)$ denotes the support of f .

DEFINITION 4. A real convolution kernel N on X is said to satisfy the transitive semi-complete maximum principle with respect to ξ (denoted by $(N, \xi) \in (\text{TSMP})$) if for any $f, g \in C_K^+(X)$ with $\int f d\xi = \int g d\xi$ and any $a \in R$, we have the implication:

$$N * f(x) \leq N * g(x) + a \quad \text{on } \text{supp}(f) \implies a \geq 0.$$

We can describe the above principles by the term of non-negative Radon measures.

Remark 5. $N \in (\text{SMP})$ (resp. $(N, \xi) \in (\text{TSMP})$) if and only if for any $\mu, \nu \in M_K^+(X)$ with $\int d\mu = \int d\nu$ and any $a \in R$, we have the implication:

$$\begin{aligned} N * \mu &\leq N * \nu + a\xi && \text{in a certain open set } \supset \text{supp}(\mu) \\ \implies N * \mu &\leq N * \nu + a\xi && \text{on } X \text{ (resp. } \implies a \geq 0), \end{aligned}$$

where $\text{supp}(\mu)$ denotes also the support of μ .

For a real convolution kernel N on X , we put

$$D^+(N) = \{\mu \in M^+(X); N * \mu \text{ is defined in } M(X)\}.$$

Let $\mu \in M^+(X)$. Evidently $\mu \in D^+(N)$ if and only if for any $f \in C_K^+(X)$, $\int |\check{N} * f| d\mu < \infty$. Here \check{N} denotes the real convolution kernel on X defined by $\int f d\check{N} = \int \check{f} dN$ for all $f \in C_K(X)$, where $\check{f}(x) = f(-x)$.

DEFINITION 6. A real convolution kernel N on X is said to satisfy the semi-balayage principle (resp. the semi-balayage principle on all open sets) (denoted by $N \in (\text{SBP})$ (resp. denoted by $N \in (\text{SBP}_g)$)) if for any $\mu \in M_K^+(X)$, any $a \in R$ and any relatively compact open set (resp. any open set) $\omega \neq \emptyset$ in X , there exists an element $(\mu', a') \in M^+(X) \times R$ such that:

$$(B.1) \quad \int d\mu' = \int d\mu.$$

$$(B.2) \quad \text{supp}(\mu') \subset \bar{\omega}.$$

(B.3) $\mu' \in D^+(N)$ and $N * \mu' + a'\xi = N * \mu + a\xi$ in ω .

(B.4) $N * \mu' + a'\xi \leq N * \mu + a\xi$ on X .

In this case, we call (μ', a') a semi-balayaged couple of (μ, a) on ω with respect to N and denote by $SB_N((\mu, a); \omega)$ the totality of such couples. If $N \in (SBP_g)$, we say that N is semi-balayable.

We set

$$\begin{aligned} \underline{SB}_N((\mu, a); \omega) &= \{(\mu', a') \in SB_N((\mu, a); \omega); N * \mu + a'\xi \\ &= \inf \{N * \mu'' + a''\xi; (\mu'', a'') \in SB_N((\mu, a); \omega)\}^5\}. \end{aligned}$$

When $\bar{\omega}$ is non-compact, it is not easy to examine directly whether $\underline{SB}_N((\mu, a); \omega) \neq \phi$ or $= \phi$.

Let $N \in (SBP)$ (resp. $N \in (SBP_g)$). For $\mu \in D^+(N)$ with $\int d\mu < \infty$, $a \in R$ and a relatively compact open set (resp. an open set) $\omega \neq \phi$ in X , we can define $SB_N((\mu, a); \omega)$ and $\underline{SB}_N((\mu, a); \omega)$ analogously.

We shall use known results concerning potential theoretic principles for a real convolution kernel N on X (see Remarques 2, 7, Proposition 11 and Corollaire 14 in [8]).

Remark 7. (1) $N \in (SMP)$ and $N \in (SBP)$ are equivalent.

(2) Assume that $N \in (SMP)$. Then $(N, \xi) \in (TSMP)$ is equivalent to $\inf_{x \in X} N * f(x) \leq 0$ for any $f \in C_K^0(X)$.

(3) Assume that $(N, \xi) \in (TSMP)$. Then N and \check{N} satisfy the maximum principle, that is, for any $f \in C_K^+(X)$, we have $N * f(x) \leq \sup_{y \in \text{supp}(f)} N * f(y)$ on X and $\check{N} * f(x) \leq \sup_{y \in \text{supp}(f)} \check{N} * f(y)$ on X .

LEMMA 8. *Let $N \in (SMP)$ and $\omega \neq \phi$ be a relatively compact open set in X . Then we have:*

(1) *For any $\mu \in D^+(N)$ with $\int d\mu < \infty$ and any $a \in R$, we have $\underline{SB}_N((\mu, a); \omega) \neq \phi$, and for any $(\mu', a') \in \underline{SB}_N((\mu, a); \omega)$, there exist nets $(\mu_\alpha)_{\alpha \in A}$ in $M_K^+(X)$ and $(a_\alpha)_{\alpha \in A}$ in R such that $\text{supp}(\mu_\alpha) \subset \omega$ and $(N * \mu_\alpha + a_\alpha \xi)_{\alpha \in A}$ converges increasingly to $N * \mu' + a'\xi$ on X along A .*

(2) *For $0 < c \in R$, we denote by $SP_c(N)$ the vague closure of*

$$\left\{ N * \nu + a\xi; \nu \in M_K^+(X), \int d\nu = c, a \in R \right\}.$$

For any $\eta \in SP_c(N)$, there exists an element $(\mu', a') \in M_K^+(X) \times R$ such that

⁵⁾ This means that $\inf \{N * \mu'' + a''\xi; (\mu'', a'') \in SB_N((\mu, a); \omega)\}$ exists as a real Radon measure on X and it is equal to $N * \mu' + a'\xi$.

$\int d\mu' = c$, $\text{supp}(\mu') \subset \bar{\omega}$, $N * \mu' + a'\xi = \eta$ in ω and $N * \mu' + a'\xi \leq \eta$ on X .

Proof. The assertion (1) is proved in the same manner as in [8] (see Corollaire 12). We shall show the assertion (2). We choose nets $(\mu_\alpha)_{\alpha \in A}$ in $M_K^+(X)$ with $\int d\mu_\alpha = c$ and $(a_\alpha)_{\alpha \in A}$ in R such that $\lim_{\alpha \in A} (N * \mu_\alpha + a_\alpha \xi) = \eta$. Let $(\mu'_\alpha, a'_\alpha) \in \text{SB}_N((\mu_\alpha, a_\alpha); \omega)$. Since $\int d\mu'_\alpha = \int d\mu_\alpha = c$, we may assume that $(\mu'_\alpha)_{\alpha \in A}$ converges vaguely. Put $\mu' = \lim_{\alpha \in A} \mu'_\alpha$. All μ'_α being supported by the compact set $\bar{\omega}$, we have $N * \mu' = \lim_{\alpha \in A} N * \mu'_\alpha$. This implies that $(a'_\alpha)_{\alpha \in A}$ converges. Putting $a' = \lim_{\alpha \in A} a'_\alpha$, we see that (μ', a') is a required element.

We shall use a more general form of the semi-complete maximum principle.

PROPOSITION 9. *Let $N \in (\text{SMP})$, $(N, \xi) \in (\text{TSMP})$, $\mu \in D^+(N)$ with $c = \int d\mu < \infty$, $a \in R$ and let $\eta \in \text{SP}_c(N)$. If $N * \mu + a\xi \leq \eta$ in a certain open set containing $\text{supp}(\mu)$, then the same inequality holds on X .*

For the proof of this proposition, we shall use the following known lemma.

LEMMA 10 (see Lemme 21 in [8]). *Let $N \in (\text{SMP})$ and $(\mu_\alpha)_{\alpha \in A}$ be a net in $M_K^+(X)$. If $\lim_{\alpha \in A} \int d\mu_\alpha = 0$ and $(N * \mu_\alpha)_{\alpha \in A}$ converges vaguely, then there exists $b \in R$ such that $\lim_{\alpha \in A} N * \mu_\alpha = b\xi$. Furthermore, if $(N, \xi) \in (\text{TSMP})$, then $b \leq 0$.*

Proof of Proposition 9. If $\mu \in M_K^+(X)$, then our assertion follows from Remark 5 and Lemma 8. In general case, we choose an open exhaustion $(\omega_n)_{n=1}^\infty$ of X ⁶⁾. Let ω be an open set in X satisfying $\omega \supset \text{supp}(\mu)$ and $N * \mu + a\xi \leq \eta$ in ω . We may assume that $\omega \cap \omega_1 \neq \emptyset$. Put $\mu_n = \mu|_{\omega_n}$ ⁷⁾ and $\lambda_n = \mu - \mu_n$. Let $(\lambda'_n, a'_n) \in \text{SB}_N((\lambda_n, a); \omega \cap \omega_n)$. Then $(\mu_n + \lambda'_n, a'_n) \in \text{SB}_N((\mu, a); \omega \cap \omega_n)$, and Lemma 8 (1) gives

$$N * (\mu_n + \lambda'_n) + a'_n \xi \leq \eta \quad \text{on } X.$$

Hence it suffices to show that $\lim_{n \rightarrow \infty} (N * (\mu_n + \lambda'_n) + a'_n \xi) = N * \mu + a\xi$.

⁶⁾ An open exhaustion $(\omega_n)_{n=1}^\infty$ of X means a sequence of relatively compact open sets $\neq \emptyset$ in X satisfying $\omega_{n+1} \supset \bar{\omega}_n$ and $\cup_{n=1}^\infty \omega_n = X$.

⁷⁾ For $\mu \in M(X)$ and a universally measurable set E in X , $\mu|_E$ denotes the real Radon measure on X defined by $\mu|_E = \mu$ on E and $\mu|_E = 0$ on CE .

From $(N, \xi) \in (\text{TSMP})$, we see that $a'_n \leq a'_{n+1} \leq a$ for all $n \geq 1$, so that $(N * \lambda'_n)_{n=1}^\infty$ converges vaguely. By Lemma 10 and $\lim_{n \rightarrow \infty} \int d\lambda'_n = 0$, there exists $0 \leq b \in R$ such that $\lim_{n \rightarrow \infty} N * \lambda'_n = b\xi$. Since

$$\lim_{n \rightarrow \infty} N * (\mu_n + \lambda'_n) + (\lim_{n \rightarrow \infty} a'_n)\xi = N * \mu + a\xi \quad \text{in } \omega ,$$

$\lim_{n \rightarrow \infty} a'_n = a$ and $b = 0$. Thus $N * (\mu_n + \lambda'_n) + a'_n\xi$ converges increasingly to $N * \mu + a\xi$ as $n \uparrow \infty$, which completes the proof.

Similarly we obtain the following

PROPOSITION 11. *Let $N \in (\text{SBP}_g)$ and $(N, \xi) \in (\text{TSMP})$. Then, for any $\mu \in M_K^+(X)$, any $a \in R$, any open set $\omega \neq \phi$ in X and any $(\mu', a') \in \text{SB}_N((\mu, a); \omega)$, we have $a' \leq a$. Furthermore, if $C\omega$ is compact, $a' = a$.*

Proof. Let $(\omega_n)_{n=1}^\infty$ be an open exhaustion of X . Put $\mu'_n = \mu'|_{\omega_n}$ and $\lambda_n = \mu' - \mu'_n$. Choose $(\lambda'_n, a'_n) \in \text{SB}_N((\lambda_n, a'); \omega_n)$; then $(\mu'_n + \lambda'_n, a'_n) \in \text{SB}_N((\mu', a'); \omega_n)$. Then $(N, \xi) \in (\text{TSMP})$ gives $a'_n \leq a$. From the above proof, we see that $\lim_{n \rightarrow \infty} a'_n = a'$, that is, $a' \leq a$.

The latter part is shown in the same manner as in Proposition 28 (2) in [8].

It is a question when $a' = a$ holds.

§ 3.

In this paragraph, we shall prepare some potential theoretic results concerning shift-bounded Hunt convolution kernels.

DEFINITION 12. A non-negative convolution kernel N_0 on X is said to be a Hunt convolution kernel if it is of form

$$(3.1) \quad N_0 = \int_0^\infty \alpha_t dt \quad \left(\text{i.e., for any } f \in C_K(X), \int f dN_0 = \int_0^\infty dt \int f d\alpha_t \right),$$

where $(\alpha_t)_{t \geq 0}$ is a vaguely continuous convolution semi-group (of positive Radon measures) on X , i.e., $\alpha_0 =$ the unit measure ε at the origin 0 , $\alpha_t * \alpha_s = \alpha_{t+s}$ for all $t \geq 0, s \geq 0$ and $t \rightarrow \alpha_t$ is vaguely continuous.

In this case, $(\alpha_t)_{t \geq 0}$ is uniquely determined (see [5]) and called the convolution semi-group of N_0 .

A vaguely continuous convolution semi-group $(\alpha_t)_{t \geq 0}$ is said to be sub-markovian (resp. markovian) if $\int d\alpha_t \leq 1$ (resp. $\int d\alpha_t = 1$) for all $t \geq 0$.

DEFINITION 13. A family $(N_p)_{p>0}$ of non-negative convolution kernels on X is said to be a resolvent if for any $p > 0$ and $q > 0$,

$$(3.2) \quad N_p - N_q = (q - p)N_p * N_q \text{ (The resolvent equation).}$$

A non-negative convolution kernel N_0 on X possesses the resolvent if there exists a resolvent $(N_p)_{p>0}$ with $N_0 = \lim_{p \downarrow 0} N_p$.

In this case, $N_0 - N_p = pN_0 * N_p$ and $\text{supp}(N_0) = \text{supp}(N_p)$ ($p > 0$) hold, and $(N_p)_{p>0}$ is uniquely determined (see [5]). We call it the resolvent of N_0 .

A resolvent $(N_p)_{p>0}$ is said to be sub-markovian (resp. markovian) if for any $p > 0$, $p \int dN_p \leq 1$ (resp. $p \int dN_p = 1$).

The following results are fundamental for Hunt convolution kernels (see [1], [3], [5], [6] and [7]).

Remark 14. (1) A non-negative convolution kernel N_0 on X is a Hunt convolution kernel if and only if its resolvent exists and N_0 is non-periodic, i.e., for any $x \in X$, $N_0 \neq N_0 * \epsilon_x$ provided with $x \neq 0$, where ϵ_x denotes the unit measure at x .

(2) Let N_0 be a Hunt convolution kernel on X . Then the equivalences (a) \Leftrightarrow (b) \Leftrightarrow (c) hold:

(a) The convolution semi-group of N_0 is sub-markovian (resp. markovian).

(b) The resolvent of N_0 is sub-markovian (resp. markovian).

(c) N_0 is shift-bounded, i.e., for any $f \in C_K(X)$, $N * f$ is bounded on X (resp. shift-bounded and $\int dN_0 = \infty$).

(3) Let N_0 be a shift-bounded Hunt convolution kernel on X . Then we have:

(a) (The balayability). For any $\mu \in M_K^+(X)$ and any open set ω in X , there exists $\mu' \in D^+(N_0)$ such that $\text{supp}(\mu') \subset \bar{\omega}$, $N_0 * \mu' = N_0 * \mu$ in ω and $N_0 * \mu' \leq N_0 * \mu$ on X .

In this case, μ' is called an N_0 -balayaged measure of μ on ω , and $\int d\mu' \leq d\mu$ holds. We have $\int dN_0 = \infty$ if and only if, for any $\mu \in M_K^+(X)$, any open set ω in X whose complement is compact and any N_0 -balayaged measure μ' of μ on ω , $\int d\mu' = \int d\mu$.

(b) (The complete maximum principle). For any $\mu, \nu \in M_K^+(X)$ and any $0 \leq c \in R$, $N_0 * \mu \leq N_0 * \nu + c\epsilon$ in a certain neighborhood of $\text{supp}(\mu)$

implies that the same inequality holds on X .

(c) (The equilibrium principle). For any relatively compact open set ω in X , there exists $\gamma \in M_K^+(X)$ such that $\text{supp}(\gamma) \subset \bar{\omega}$, $N_0 * \gamma = \xi$ in ω and $N_0 * \gamma \leq \xi$ on X .

In this case, γ is called an N_0 -equilibrium measure of ω .

(d) (The positive mass principle). For any $\mu, \nu \in M_K^+(X)$, $N_0 * \mu \leq N_0 * \nu$ on X implies $\int d\mu \leq \int d\nu$.

(e) (The dominated convergence property). Let $(\mu_\alpha)_{\alpha \in A}$ be a net in $D^+(N_0)$ and $\mu \in M^+(X)$. If $\lim_{\alpha \in A} \mu_\alpha = \mu$ and there exists $\nu \in D^+(N_0)$ satisfying $N_0 * \mu_\alpha \leq N_0 * \nu$ on X for all $\alpha \in A$, then $\lim_{\alpha \in A} N_0 * \mu_\alpha = N_0 * \mu$.

(f) (The injectivity). For any $\mu, \nu \in D^+(N_0)$, $N_0 * \mu = N_0 * \nu$ on X implies $\mu = \nu$.

For $\mu \in D^+(N_0)$ and an open set ω in X , we can define analogously N_0 -balayaged measures of μ on ω and denote by $B_{N_0}(\mu; \omega)$ their totality. It is well-known that $B_{N_0}(\mu; \omega) \neq \phi$. Put

$$B_{N_0}(\mu; \omega) = \{\mu' \in B_{N_0}(\mu; \omega); N_0 * \mu' = \inf \{N_0 * \mu''; \mu'' \in B_{N_0}(\mu; \omega)\}\} \quad (\text{see}^5)$$

and

$$\bar{B}_{N_0}(\mu; \omega) = \{\mu' \in B_{N_0}(\mu; \omega); N_0 * \mu' = \sup \{N_0 * \mu''; \mu'' \in B_{N_0}(\mu; \omega)\}\} \quad (\text{see}^5).$$

For an open set ω in X , we can define analogously N_0 -equilibrium measures of ω and denote by $E_{N_0}(\omega)$ their totality. Put

$$\underline{E}_{N_0}(\omega) = \{\gamma \in E_{N_0}(\omega); N_0 * \gamma = \inf \{N_0 * \gamma'; \gamma' \in E_{N_0}(\omega)\}\} \quad (\text{see}^5)$$

provided with $E_{N_0}(\omega) \neq \phi$.

LEMMA 15. *Let N_0 be a shift-bounded Hunt convolution kernel on X . Then we have:*

(1) *For any $\mu \in D^+(N_0)$ and any open set ω in X , $B_{N_0}(\mu; \omega) \neq \phi$ and $\bar{B}_{N_0}(\mu; \omega) \neq \phi$. Moreover, $B_{N_0}(\mu; \omega)$ and $\bar{B}_{N_0}(\mu; \omega)$ form only one element.*

(2) *For any $\mu \in D^+(N_0)$ and any two open sets ω_1, ω_2 in X with $\omega_1 \subset \omega_2$, we have $N_0 * \mu'_1 \leq N_0 * \mu'_2$ and $N_0 * \mu''_1 \leq N_0 * \mu''_2$ on X , where $\mu'_i \in B_{N_0}(\mu; \omega_i)$ and $\mu''_i \in \bar{B}_{N_0}(\mu; \omega_i)$ ($i = 1, 2$).*

(3) *Put $P(N_0) = \overline{\{N_0 * \mu; \mu \in D^+(N_0)\}}$, where the closure is in the sense of the vague topology. For any $\mu \in D^+(N_0)$ and any $\eta \in P(N_0)$, $N_0 * \mu \leq \eta$ in a certain open set $\supset \text{supp}(\mu)$ implies that the same inequality holds on X .*

(4) For an open set ω in X , $E_{N_0}(\mu) \neq \phi$ implies $\underline{E}_{N_0}(\mu) \neq \phi$. In this case, $\underline{E}_{N_0}(\mu)$ forms only one element.

(5) For $0 < c \in R$, we put $P_c(N_0) = \overline{\left\{ N_0 * \mu; \mu \in D^+(N_0), \int d\mu \leq c \right\}}$. For any $\mu \in D^+(N_0)$ and any $\eta \in P_c(N_0)$, $N_0 * \mu \leq \eta$ on X implies $\int d\mu \leq c$.

(6) Let $(\mu_\alpha)_{\alpha \in A}$ be a net in $D^+(N_0)$ and $0 \neq \lambda_1, 0 \neq \lambda_2 \in M_K^+(X)$. If there exist $\nu \in D^+(N_0)$ and a relatively compact net $(x_\alpha)_{\alpha \in A}$ in X such that $N_0 * \mu_\alpha * \lambda_1 \leq N_0 * \nu * \varepsilon_{x_\alpha} * \lambda_2$ on X , then $(\mu_\alpha)_{\alpha \in A}$ is vaguely bounded. If $\mu_\alpha \rightarrow \mu \in M^+(X)$, then $\lim_{\alpha \in A} N_0 * \mu_\alpha = N_0 * \mu$.

Proof. (1) Let $(\omega_\alpha)_{\alpha \in A}$ be a net of open sets in X with $\bar{\omega}_\alpha \subset \omega_\beta (\alpha \preceq \beta)$ and $\bigcup_{\alpha \in A} \omega_\alpha = \omega$. We choose $\mu'_\alpha \in B_{N_0}(\mu; \omega_\alpha)$. Then the complete maximum principle of N_0 implies that for any $\mu'' \in B_{N_0}(\mu; \omega)$, $N_0 * \mu'_\alpha \leq N_0 * \mu''$ on X . This and the dominated convergence property of N_0 show that $(\mu'_\alpha)_{\alpha \in A}$ is vaguely bounded and every vaguely accumulation point of $(\mu'_\alpha)_{\alpha \in A}$ as $\omega_\alpha \uparrow \omega$ is contained in $\underline{B}_{N_0}(\mu; \omega)$, which gives $\underline{B}_{N_0}(\mu; \omega) \neq \phi$. Let $(\omega'_\alpha)_{\alpha' \in A'}$ be a net of open sets in X with $\omega'_\alpha \supset \bar{\omega}'_{\beta'} (\alpha' \preceq \beta')$ and $\bigcap_{\alpha' \in A'} \omega'_\alpha = \bar{\omega}$. We choose $\mu''_{\alpha'} \in B_{N_0}(\mu; \omega'_\alpha)$. Similarly as above, $(\mu''_{\alpha'})_{\alpha' \in A'}$ as $\omega'_\alpha \downarrow \bar{\omega}$ is contained in $\bar{B}_{N_0}(\mu; \omega)$, that is, $\bar{B}_{N_0}(\mu; \omega) \neq \phi$. The injectivity of N_0 shows that $\underline{B}_{N_0}(\mu; \omega)$ and $\bar{B}_{N_0}(\mu; \omega)$ form only one element.

Consequently, let $\mu'_\alpha \in \underline{B}_{N_0}(\mu; \omega_\alpha)$, $\mu' \in \underline{B}_{N_0}(\mu; \omega)$, $\mu''_{\alpha'} \in \bar{B}_{N_0}(\mu; \omega'_\alpha)$ and $\mu'' \in \bar{B}_{N_0}(\mu; \omega)$; then $\lim_{\alpha \in A} \mu'_\alpha = \mu'$ and $\lim_{\alpha' \in A'} \mu''_{\alpha'} = \mu''$.

(2) Using the complete maximum principle of N_0 and noting the above proof, we see easily (2).

(3) Let $\nu \in M_K^+(X)$ with $\nu \leq \mu$. We choose a relatively compact open set ω in X such that $\omega \supset \text{supp}(\nu)$ and $N_0 * \nu \leq \eta$ in ω . By virtue of the balayability of N_0 , we can choose $\lambda \in M_K^+(X)$ such that $\text{supp}(\lambda) \subset \bar{\omega}$, $N_0 * \lambda = \eta$ in ω and $N_0 * \lambda \leq \eta$ on X . This shows that $N_0 * \nu \leq N_0 * \lambda \leq \eta$ on X , and ν being arbitrary, we have $N_0 * \mu \leq \eta$ on X .

(4) In the same manner as in the proof of $\underline{B}_{N_0}(\mu; \omega) \neq \phi$ in (1), we see that $E_{N_0}(\omega) \neq \phi$ implies $\underline{E}_{N_0}(\omega) \neq \phi$. For any $\gamma \in E_{N_0}(\omega)$, $\underline{E}_{N_0}(\omega) = \underline{B}_{N_0}(\gamma; \omega)$. If $\underline{E}_{N_0}(\omega) \neq \phi$, the injectivity of N_0 shows that $\underline{E}_{N_0}(\omega)$ forms only one element.

(5) By using the positive mass principle of N_0 and the similar method to (3), we obtain (5).

(6) Evidently $(\mu_\alpha)_{\alpha \in A}$ is vaguely bounded. We shall show only the latter half part. Let $(K_n)_{n=1}^\infty$ be an exhaustion of X . We choose $\varepsilon'_n \in B_{N_0}(\varepsilon, CK_n)$. The dominated convergence property of N_0 gives $\lim_{n \rightarrow \infty} N_0 * \varepsilon'_n = 0$. Let

$f \in C_K^+(X)$. Since $(x_\alpha)_{\alpha \in A}$ is relatively compact, $\int fdN_0 * \varepsilon'_n * \varepsilon_{x_\alpha} * \nu * \lambda_2$ converges uniformly to 0 on $(x_\alpha)_{\alpha \in A}$ as $n \rightarrow \infty$. Hence

$$\overline{\lim}_{\alpha \in A} \int fdN_0 * \mu_\alpha * \lambda_1 \leq \int fdN_0 * \mu * \lambda_1.$$

Using the lower semi-continuity of convolutions of non-negative Radon measures, we have $\underline{\lim}_{\alpha \in A} \int fdN_0 * \mu_\alpha \geq \int fdN_0 * \mu$. Thus $\mu \in D^+(N_0)$ and $\lim_{\alpha \in A} N_0 * \mu_\alpha = N_0 * \mu$.

From Lemma 15 and its proof, we see the following

LEMMA 16. *Let N_0 be a shift-bounded Hunt convolution kernel on X , $(\Omega_j)_{j=1}^m$ and $(\omega_k)_{k=1}^n$ two finite families of open sets in X and let $(\mu_j)_{j=1}^m \subset D^+(N_0)$. Assume that $E_{N_0}(\omega_k) \neq \phi$ ($k = 1, 2, \dots, n$). Let $\mu'_j \in \underline{B}_{N_0}(\mu_j; \Omega_j)$, $\gamma_k \in \underline{B}_{N_0}(\omega_k)$ ($j = 1, 2, \dots, m; k = 1, 2, \dots, n$) and let $\eta \in P(N_0)$. If $\sum_{j=1}^m \sum_{k=1}^n N_0 * (\mu'_j + \gamma_k) \leq \eta$ in $(\cup_{j=1}^m \Omega_j) \cup (\cup_{k=1}^n \omega_k)$, then the same inequality holds on X .*

LEMMA 17. *Let N_0 be the same as above, $\mu \in D^+(N_0)$ and let ω be an open set in X . For $x \in X$, we denote by μ'_x and μ''_x the unique element in $\underline{B}_{N_0}(\mu * \varepsilon_x; \omega)$ and that in $\overline{B}_{N_0}(\mu * \varepsilon_x; \omega)$, respectively. Then we have:*

(1) *The mapping $x \rightarrow \mu'_x$ and $x \rightarrow \mu''_x$ are universally measurable, that is, for any $f \in C_K(X)$, the functions $\int fd\mu'_x$ and $\int fd\mu''_x$ of x are universally measurable on X .*

(2) *For any $\nu \in M_K^+(X)$, $(\mu * \nu)' \in \underline{B}_{N_0}(\mu * \nu; \omega)$ and $(\mu * \nu)'' \in \overline{B}_{N_0}(\mu * \nu; \omega)$ are of form*

$$(3.3) \quad (\mu * \nu)' = \int \mu'_x d\nu(x)^{8)} \quad \text{and} \quad (\mu * \nu)'' = \int \mu''_x d\nu(x).$$

Proof. Let $x \in X$ and $(x_\alpha)_{\alpha \in A}$ be a net in X with $x_\alpha \rightarrow x$. Then Lemma 15 (6) shows that $(\mu'_{x_\alpha})_{\alpha \in A}$ and $(\mu''_{x_\alpha})_{\alpha \in A}$ are vaguely bounded and that every vaguely accumulation point of $(\mu'_{x_\alpha})_{\alpha \in A}$ and that of $(\mu''_{x_\alpha})_{\alpha \in A}$ as $x_\alpha \rightarrow x$ belong to $\underline{B}_{N_0}(\mu * \varepsilon_x; \omega)$. This implies that the mapping $x \rightarrow N_0 * \mu'_x$ is lower semi-continuous (i.e., for any $f \in C_K^+(X)$, the function $\int fdN_0 * \mu'_x$ is lower semi-continuous) and the mapping $x \rightarrow N_0 * \mu''_x$ is upper semi-continuous. Let $(N_p)_{p>0}$ be the resolvent of N_0 . Then, for any $p > 0$, $x \rightarrow N_0 * N_p * \mu'_x$

⁸⁾ This means that for any $f \in C_K(X)$, $\int fd(\mu * \nu)' = \int \int fd\mu'_x d\nu(x)$

is also lower semi-continuous and $x \rightarrow N_0 * N_p * \mu'_x$ is also upper semi-continuous, because N_p is also a Hunt convolution kernel on X , so that N_p possesses the dominated convergence property. Hence, for any $f \in C_K(X)$ and any $p > 0$, the resolvent equation shows that $\int f dN_p * \mu'_x$ and $\int f dN_p * \mu''_x$ are universally measurable functions of x on X . Since $\lim_{p \rightarrow \infty} pN_p = \varepsilon$ and there exists $g \in C_K^+(X)$ such that $|p\check{N}_p * f| \leq \check{N}_0 * g$ on X for all $p > 0$, the Lebesgue dominated convergence theorem gives $\int f d\mu'_x = \lim_{p \rightarrow \infty} p \int f dN_p * \mu'_x$ and $\int f d\mu''_x = \lim_{p \rightarrow \infty} p \int f dN_p * \mu''_x$, which show that $x \rightarrow \mu'_x$ and $x \rightarrow \mu''_x$ are universally measurable.

We shall show the assertion (2). For any $f \in C_K^+(X)$, $\iint f d\mu'_x d\nu(X)$ and $\iint f d\mu''_x d\nu(x)$ are defined and

$$\iint \check{N}_0 * f d\mu'_x d\nu(x) \leq \iint \check{N}_0 * f d\mu''_x d\nu(x) \leq \int f dN_0 * (\mu * \nu),$$

so that $\int \mu'_x d\nu(x)$ and $\int \mu''_x d\nu(x)$ belong to $D^+(N_0)$. We see easily that $\int \mu'_x d\nu(x), \int \mu''_x d\nu(x) \in B_{N_0}(\mu * \nu; \omega)$. Let $(\omega_\alpha)_{\alpha \in A}$ be a net of open sets in X satisfying $\bar{\omega}_\alpha \subset \omega_\beta (\alpha \preceq \beta)$ and $\bigcup_{\alpha \in A} \omega_\alpha = \omega$. We choose $\mu'_{x,\alpha} \in \underline{B}_{N_0}(\mu * \varepsilon_x; \omega_\alpha)$. Then Lemma 15 (1), (3) show that $N_0 * \mu'_{x,\alpha} \uparrow N_0 * \mu'_x$ as $\omega_\alpha \uparrow \omega$, that is,

$$N_0 * \left(\int \mu'_{x,\alpha} d\nu(x) \right) \uparrow N_0 * \left(\int \mu'_x d\nu(x) \right) \quad \text{as } \omega_\alpha \uparrow \omega.$$

This shows that $\int \mu'_x d\nu(x) \in \underline{B}_{N_0}(\mu * \nu; \omega)$, and Lemma 15 (1) gives the first equality in (3.3). Let $(\omega'_{\alpha'})_{\alpha' \in A'}$ be a net of open sets in X satisfying $\omega'_{\alpha'} \supset \bar{\omega}'_{\beta'} (\alpha' \preceq \beta')$ and $\bigcap_{\alpha' \in A'} \omega'_{\alpha'} = \bar{\omega}$. We choose $\mu''_{x,\alpha'} \in \bar{B}_{N_0}(\mu * \varepsilon_x; \omega'_{\alpha'})$. Similarly as above, we have

$$N_0 * \left(\int \mu''_{x,\alpha'} d\nu(x) \right) \downarrow N_0 * \left(\int \mu''_x d\nu(x) \right) \quad \text{as } \omega'_{\alpha'} \downarrow \bar{\omega},$$

and hence the second equality in (3.3) holds. Thus Lemma 17 is shown.

The following proposition will play an important role to prove our main theorem.

PROPOSITION 18. *Let N_0 be a shift-bounded Hunt convolution kernel on X and assume that the closed subgroup generated by $\text{supp}(N_0)$ is equal to X . Then, for any $0 \neq \mu \in M_K^+(X)$, there exist an open set $\omega \neq \phi$ in X*

and an open neighborhood V of the origin such that:

- (1) For any $\mu' \in B_{N_0}(\mu, \omega + V)^{9)}$, $\int d\mu' < \int d\mu$.
- (2) N_0 -equilibrium measures of ω with finite total mass do not exist.

For the poof of this proposition, we use the following result:

LEMMA 19 (see [2], [4]). Let $\sigma \in M^+(X)$ with $\int d\sigma = 1$. If a shift-bounded real Radon measure μ on X satisfies $\mu = \mu * \sigma$, then, for any x in the closed subgroup Γ generated by $\text{supp}(\sigma)$, we have $\mu = \mu * \varepsilon_x$, that is, each x in Γ is a period of μ .

Proof of Proposition 18. It suffices to show the following assertion:

Let $0 \neq f \in C_K^+(X)$. Then there exist an open set $\omega \neq \phi$ in X and open neighborhood V of the origin such that:

- (1') For $(f\xi)'' \in \bar{B}_{N_0}(f\xi; \omega + V)$, $\int d(f\xi)'' < \int fd\xi$.
- (2') $E_{N_0}(\omega) = \phi$, or $E_{N_0}(\omega) \neq \phi$ and for $\gamma \in \underline{E}_{N_0}(\omega)$, $\int d\gamma = \infty$.

In fact, admit this assertion and let $0 \neq \mu \in M_K^+(X)$. Choose $\varphi \in C_K^+(X)$ with $\int \varphi d\xi = 1$. Then there exist an open set $\omega \neq \phi$ in X and an open neighborhood V of the origin such that, for $f = \mu * \varphi$, (1') and (2') are verified. Since $\int d\mu = \int \mu * \varphi d\xi$, Lemma 17 (2) shows that there exists $x \in \text{supp}(\varphi)$ such that for $(\mu * \varepsilon_x)'' \in \bar{B}_{N_0}(\mu * \varepsilon_x; \omega + V)$, $\int d(\mu * \varepsilon_x)'' < \int d\mu * \varepsilon_x = \int d\mu$. We remark here that $(\mu * \varphi)\xi = \int \mu * \varepsilon_x \varphi(x) d\xi(x)$ and for any $y \in X$, $\int d(\mu * \varepsilon_y)'' \leq \int d\mu * \varepsilon_y$. Put $\omega_x = \omega - \{x\}$ and $\mu_x'' \in \bar{B}_{N_0}(\mu; \omega_x + V)$. Then we see easily that $(\mu * \varepsilon_x)'' = \mu_x'' * \varepsilon_x$, which implies $\int d\mu_x'' < \int d\mu$. We remark that $E_{N_0}(\omega) = \phi$ and $E_{N_0}(\omega_x) = \phi$ are equivalent and if $E_{N_0}(\omega) \neq \phi$, then, for $\gamma \in \underline{E}_{N_0}(\omega)$ and $\gamma_x \in \underline{E}_{N_0}(\omega_x)$, $\gamma = \gamma_x * \varepsilon_x$. By the positive mass principle and Lemma 15 (5), we see that ω_x and V are our required open set and open neighborhood of the origin.

Dividing into the following two cases, we shall show our required assertion.

- (a) Assume that there exists $0 \neq g \in C_X^+(X)$ with $\overline{\lim}_{x \rightarrow \delta} N_0 * g(x) > 0$. Then $\int dN_0 = \infty$. Noting that $(N_0 * \varepsilon_x)_{x \in X}$ is vaguely bounded, we can

⁹⁾ For subsets A, B of X , $A + B = \{x + y; x \in A, y \in B\}$, $-B = \{-x; x \in B\}$.

choose a net $(x_\alpha)_{\alpha \in A}$ in X with $x_\alpha \rightarrow \delta$ such that $(N_0 * \varepsilon_{x_\alpha})_{\alpha \in A}$ converges vaguely and $\lim_{\alpha \in A} N_0 * \varepsilon_{x_\alpha} * g(0) = \overline{\lim}_{x \rightarrow \delta} N_0 * g(x)$. Put $\eta = \lim_{\alpha \in A} N_0 * \varepsilon_{x_\alpha}$; then $\eta \neq 0$. Let $(N_p)_{p > 0}$ be the resolvent of N_0 . By the resolvent equation and $p \int dN_p = 1$ ($p > 0$), we have

$$\eta = pN_p * \eta \quad (p > 0).$$

Since $\text{supp}(N_p) = \text{supp}(N_0)$ ($p > 0$) and η is shift-bounded, Lemma 19 gives $\eta = c\xi$ with some constant $c > 0$. We may assume that $\int fd\xi = 1$. Let Ω be a relatively compact open set with $\Omega \supset \text{supp}(f)$. Since $(N_0 * \varepsilon_x * f)_{x \in X}$ converges uniformly to $N_0 * f$ on $\bar{\Omega}$ as $x \rightarrow 0$, there exists an open neighborhood V of the origin such that $V = -V$, $\text{supp}(f) + \bar{V} \subset \Omega$ and for any $x \in \bar{V}$, $|N_0 * \varepsilon_x * f - N_0 * f| < \frac{1}{3}c$ on $\bar{\Omega}$. By virtue of the complete maximum principle of N_0 , we have $|N_0 * \varepsilon_x * f - N_0 * f| < \frac{1}{3}c$ on X for all $x \in \bar{V}$. Put $\omega = \{x \in X; N_0 * f(x) < \frac{1}{3}c\}$ and $\omega' = \{x \in X; N_0 * f(x) < \frac{2}{3}c\}$. Then $\bar{\omega} + \bar{V} \subset \omega'$. We shall show that ω and V are our required open set and open neighborhood of the origin. First we see that $E_{N_0}(\omega) = \phi$, because, if there exists $\gamma \in E_{N_0}(\omega)$, then $N_0 * (\frac{1}{3}c\gamma + f\xi) \geq \frac{1}{3}c\xi$ on X , which contradicts $p \int dN_p = 1$ for all $p > 0$ and $pN_p * N_0 \downarrow 0$ as $p \downarrow 0$. It remains to prove that (1') is verified. By Lemma 15 (2), it suffices to show that for any $(f\xi)' \in B_{N_0}(f\xi; \omega')$, $\int d(f\xi)' < \int fd\xi = 1$. For any integer $m \geq 1$, $N_0 * (f\xi)' \leq (\frac{2}{3} + 1/m)\eta$ in a certain open set $\supset \text{supp}((f\xi)')$, so that Lemma 15 (3) gives $N_0 * (f\xi)' \leq (\frac{2}{3} + 1/m)\eta$ on X . Letting $m \uparrow \infty$ and using Lemma 15 (5), we obtain $\int d(f\xi)' \leq \frac{2}{3}$. Thus ω and V are our required open set and open neighborhood of the origin.

(b) Assume that N_0 vanishes at the infinity (i.e., for any $g \in C_K(X)$, $\lim_{x \rightarrow \delta} N_0 * g(x) = 0$). Let U_0 be a relatively compact open set $\neq \phi$ in X with $\bar{U}_0 \subset \{x \in X; f(x) > 0\}$. Since $\text{supp}(N_0) \ni 0$, we may assume that $N_0 * f(x) > 1$ on \bar{U}_0 . We choose an open set $\omega_0 \neq \phi$ and an open neighborhood V of the origin such that $\bar{\omega}_0 + \bar{V} \subset U_0$. Since $\lim_{x \rightarrow \delta} N_0 * \varepsilon_x = 0$, we can choose inductively a sequence $(x_n)_{n=0}^\infty$ in X with $x_0 = 0$ and $x_n \rightarrow \delta$ ($n \rightarrow \infty$) such that, for any $n \geq 0$ and $m \geq 0$ with $n \neq m$,

$$N_0 * \varepsilon_{x_n} * f \leq \frac{1}{2^{|n-m|+1}} \quad \text{on } \{x_m\} + U_0.$$

Put $U_n = \{x_n\} + U_0$ ($n = 1, 2, \dots$) and $U = \bigcup_{n=1}^\infty U_n$. Evidently $\bar{U}_m \cap \bar{U}_n = \phi$

if $n \neq m$. Put $\omega_n = \{x_n\} + \omega_0$ ($n = 1, 2, \dots$) and $\omega = \bigcup_{n=1}^\infty \omega_n$. Then $\omega + V \subset U$. For any $(f\xi)' \in B_{N_0}(f\xi; U)$, we set $(f\xi)'_n = (f\xi)'|_{\bar{\omega}_n}$ ($n \geq 1$). Then, by virtue of the complete maximum principle of N_0 ,

$$N_0 * (f\xi)'_n \leq \frac{1}{2^{n+1}}(N_0 * \varepsilon_{x_n} * f) \quad \text{on } X,$$

and hence $\int d(f\xi)'_n \leq (1/2^{n+1}) \int f d\xi$. Consequently, $\int d(f\xi)' \leq \frac{1}{2} \int f d\xi$. From Lemma 15 (2), it follows that for $(f\xi)'' \in \bar{B}_{N_0}(f\xi; \omega + V)$, $\int d(f\xi)'' \leq \frac{1}{2} \int f d\xi$.

Let $\gamma'_n \in \underline{E}_{N_0}(\omega_n)$. Then $N_0 * \gamma'_n \leq (N_0 * \varepsilon_{x_n} * f)\xi$ on X . For any $n \geq 1$ and any k with $1 \leq k \leq n$, we have, in ω_k ,

$$N_0 * \left(\sum_{j=1}^n \gamma'_j\right) \leq \xi + \sum_{j=1}^{k-1} (N_0 * \varepsilon_{x_j} * f)\xi + \sum_{j=k+1}^n (N_0 * \varepsilon_{x_j} * f)\xi \leq 2\xi,$$

that is, $N_0 * (\sum_{j=1}^n \gamma'_j) \leq 2\xi$ in $\bigcup_{j=1}^n \omega_j$. This and Lemma 16 show that the same inequality holds on X . Thus $\sum_{n=1}^\infty \gamma'_n$ converges vaguely. Put $\gamma' = \sum_{n=1}^\infty \gamma'_n$; then $N_0 * \gamma' \geq \xi$ in ω and $N_0 * \gamma' \leq 2\xi$ on X . Let $\gamma_n \in \underline{E}_{N_0}(\bigcup_{k=1}^n \omega_k)$. Then $N_0 * \gamma' \geq N_0 * \gamma_n$ and $\sum_{k=1}^n N_0 * \gamma'_k \leq 2N_0 * \gamma_n$ on X . By virtue of the dominated convergence property of N_0 , we have $\underline{E}_{N_0}(\omega) \neq \phi$. Let $\gamma \in \underline{E}_{N_0}(\omega)$; then $\lim_{n \rightarrow \infty} \gamma_n = \gamma$. This implies that

$$N_0 * \gamma \leq N_0 * \gamma' \leq 2N_0 * \gamma \quad \text{on } X.$$

Evidently $\int d\gamma'_n = \int d\gamma'_m$ for all $n \geq 1, m \geq 1$ and $\gamma' \neq 0$, so that $\int d\gamma' = \infty$.

The positive mass principle of N_0 gives $\int d\gamma = \infty$. Thus ω and V are our required open set and open neighborhood of the origin.

It is a question if there exist an open set $\omega \neq \phi$ in X and an open neighborhood V of the origin such that for any $0 \neq \mu \in M^+(X)$ with $\text{supp}(\mu) \subset C(\overline{\omega + V})$ and any $\mu' \in B_{N_0}(\mu; \omega + V)$, $\int d\mu' < \int d\mu$ and N_0 -equilibrium measures γ of ω with $\int d\gamma < \infty$ do not exist.

§4.

We return to the argument of real convolution kernels. We begin with the following

DEFINITION 20. For a real convolution kernel N on X and an open set $\omega \neq \phi$ in X , we denote by $\text{SP}_1(N; \omega)$ the vague closure of

$$\left\{ N * \mu + a\xi; \mu \in M_K^+(X), \int d\mu = 1, \text{supp}(\mu) \subset \omega, a \in R \right\}$$

and put

$$\eta_{N,\omega} = \sup \{ \eta \in SP_1(N; \omega); \eta \leq N \text{ on } X \}$$

provided that the right hand exists in $M(X)$. If $\eta_{N,\omega}$ exists, we call it the N -reduced measure of N on ω .

Assume that $N \in (\text{SMP})$. Then $\eta_{N,\omega}$ always exists and satisfies $\eta_{N,\omega} = N$ in ω , $\eta_{N,\omega} \leq N$ on X (see Remarque 19 in [8]). Let $(K_n)_{n=1}^\infty$ be an exhaustion of X . Then $(\eta_{N,CK_n})_{n=1}^\infty$ is decreasing and $\lim_{n \rightarrow \infty} \eta_{N,CK_n}$ is independent of the choice of $(K_n)_{n=1}^\infty$ (see § 3 in [8]). Put $\eta_{N,\delta} = \lim_{n \rightarrow \infty} \eta_{N,CK_n}$. Then $\eta_{N,\delta} = -\infty$, i.e., for any $0 \neq f \in C_K^+(X)$, $\lim_{n \rightarrow \infty} \int f d\eta_{N,CK_n} = -\infty$, or $\eta_{N,\delta} \in M(X)$ (see Remarque 19 in [8]).

Proposition 9 gives immediately the following

Remark 21. Let $N \in (\text{SMP}_g)$, $(N, \xi) \in (\text{TSMP})$, $(K_n)_{n=1}^\infty$ be an exhaustion of X and let $(\epsilon'_{CK_n}, 0) \in \text{SB}_N((\epsilon, 0); CK_n)$ (see Proposition 11). Then, for any $n \geq 2$,

$$\eta_{N,CK_n} \leq N * \epsilon'_{CK_n} \leq \eta_{N,CK_{n-1}} \quad \text{on } X.$$

The following proposition is shown in [8] (see Théorème 20).

PROPOSITION 22. *Let $N \in (\text{SMP})$, $(N, \xi) \in (\text{TSMP})$ and let $(\omega_n)_{n=1}^\infty$ be an open exhaustion of X . Then we have:*

(1) *For any $0 < p \in R$ and any $n \geq 1$, there exists a uniquely determined $(\epsilon'_{p,n}, a_{p,n}) \in M_K^+(X) \times R$ such that $\int d\xi'_{p,n} = 1$, $\text{supp}(\epsilon'_{p,n}) \subset \bar{\omega}_n$, $(N + (1/p)\epsilon) * \epsilon'_{p,n} + a_{p,n}\xi = N$ in ω_n , $(N + (1/p)\epsilon) * \epsilon'_{p,n} + a_{p,n}\xi \leq N$ on X and for any $\nu \in M_K^+(X)$ with $\int d\nu = 1$ and any $a \in R$, $(N + (1/p)\epsilon) * \nu + a\xi \geq (N + (1/p)\epsilon) * \epsilon'_{p,n} + a_{p,n}\xi$ on X whenever $(N + (1/p)\epsilon) * \nu + a\xi \geq N$ in ω_n .*

(2) *Put $V_{p,\omega_n,\epsilon} = (1/p)\epsilon'_{p,n}$. Then $V_{p,\omega_n,\epsilon} \geq V_{p,\omega_{n+1},\epsilon}$ in ω_n and $\lim_{n \rightarrow \infty} V_{p,\omega_n,\epsilon}$ exists.*

(3) *Put*

$$(4.1) \quad N_p = \lim_{n \rightarrow \infty} V_{p,\omega_n,\epsilon} (\in M^+(X)),$$

then $(N_p)_{p>0}$ is a sub-markovian resolvent and independent of the choice of $(\omega_n)_{n=1}^\infty$.

By using Proposition 22, we have the following

LEMMA 23. *Let $N \in (\text{SBP}_g)$, $(N, \xi) \in (\text{TSMP})$ and assume that N is non-periodic. Then there exists a uniquely determined resolvent $(N_p)_{p>0}$ such that*

$$(4.2) \quad N = pN * N_p + N_p .$$

Proof. First we remark that $N \in (\text{SBP})$ and $N \in (\text{SMP})$ are equivalent. Let $V_{p, \omega_n \xi}, N_p$ and $a_{p, n}$ be the same as in Proposition 22. Then, for any $p > 0$,

$$(4.3) \quad \lim_{n \rightarrow \infty} ((pN + \varepsilon) * V_{p, \omega_n \xi} + a_{p, n} \xi) = N .$$

Let $(K_m)_{m=1}^\infty$ be an exhaustion of X with $K_1 \ni 0$. We shall show that for any $m \geq 2$, $N \neq \eta_{N, CK_m}$. Assume contrary that for an $m \geq 2, N = \eta_{N, CK_m}$. Then Remark 21 gives $N = N * \varepsilon'_{CK_m}$, where $(\varepsilon'_{CK_m}, 0) \in \text{SB}_N((\varepsilon, 0); CK_m)$. Let Γ be the closed subgroup generated by $\text{supp}(\xi'_{CK_m})$; then $\Gamma \neq \{0\}$. For any $x \in X$, $N * (\varepsilon - \varepsilon_x)$ is shift-bounded (see Remarque 4 in [8]), and Lemma 19 shows that for any $y \in \Gamma$, $N * (\varepsilon - \varepsilon_x) * \varepsilon_y = N * (\varepsilon - \varepsilon_x)$. This implies that for any $x \in \Gamma$ and any integer $n \geq 1$, $N - N * \varepsilon_{nx} = n(N - N * \varepsilon_x)$. Since for any $f \in C_K^+(X)$, $\check{N} * f$ is upper bounded (see Remark 7 (3)), we have $\int f d(N - N * \varepsilon_x) \geq 0$, and Γ being a subgroup of X , we see that $N = N * \varepsilon_x$ for all $x \in \Gamma$. This contradicts the non-periodicity of N . Thus $N \neq \eta_{N, CK_m}$ for all $m \geq 2$. Next we shall show that $(N_p)_{p>0}$ is markovian. From (4.1), $\int d\varepsilon'_{CK_m} = 1$ and $(pN + \varepsilon) * V_{p, \omega_n \xi} + a_{p, n} \xi \uparrow N$ as $n \uparrow \infty$, it follows that

$$(4.4) \quad N - N * \varepsilon'_{CK_m} = p(N - N * \varepsilon'_{CK_m}) * N_p + N_p * (\varepsilon - \varepsilon'_{CK_m}) .$$

Assume that $(N_p)_{p>0}$ is not markovian. Then, for any $p > 0, p \int dN_p < 1$. From (4.4), it follows that for any $p > 0$, any $n \geq 1$ and any $m \geq 1$,

$$N - N * \varepsilon'_{CK_m} = (N - N * \varepsilon'_{CK_m}) * (pN_p)^n + \frac{1}{p} \sum_{k=1}^n (pN_p)^k * (\varepsilon - \varepsilon'_{CK_m}) ,$$

where $(pN_p)^1 = pN_p$ and $(pN_p)^n = (pN_p)^{n-1} * (pN_p)$ ($n \geq 2$). Letting $n \uparrow \infty$, we have

$$N - N * \varepsilon'_{CK_m} = \frac{1}{p} \sum_{k=1}^\infty (pN_p)^k * (\varepsilon - \varepsilon'_{CK_m}) .$$

Since $\int d(\sum_{k=1}^{\infty} (pN_p)^k) < \infty$ and $\int d\varepsilon'_{CK_m} = 1$, we have $\int d(N - N * \varepsilon'_{CK_m}) = 0$, so that $N = N * \varepsilon'_{CK_m}$. This contradicts $N \neq \eta_{N,CK_m}$ and $\eta_{N,CK_{m-1}} \geq N * \varepsilon'_{CK_m}$ ($m \geq 2$). Thus $(N_p)_{p>0}$ is markovian. In the same manner as in [8] (see Théorème 20 and Remarque 24), we see the rest of the proof.

DEFINITION 24. Let $N \in (\text{SMP})$. If a sub-markovian resolvent $(N_p)_{p>0}$ satisfying (4.2) exists, then $(N_p)_{p>0}$ is called the resolvent associated with N .

The resolvent associated with N is uniquely determined if it exists (see Remarque 24 in [8]).

LEMMA 25. Let $N \in (\text{SMP})$ and $(N, \xi) \in (\text{TSMP})$. Assume that $\eta_{N,\delta} \neq -\infty$, N is non-periodic and that the resolvent $(N_p)_{p>0}$ associated with N exists and is markovian. Put $N' = \eta_{N,\delta}$ and $N_0 = N - N'$. Then N_0 is a shift-bounded Hunt convolution kernel on X , $N_0 = \lim_{p \rightarrow 0} N_p$ and every point in the closed subgroup generated by $\text{supp}(N_0)$ is a period on N' .

Proof. Let $(K_n)_{n=1}^{\infty}$ and $(\omega_m)_{m=1}^{\infty}$ be an exhaustion of X and an open exhaustion of X , respectively. We choose $(\varepsilon'_{n,m}, a_{n,m}) \in \text{SB}_N((\varepsilon, 0); CK_n \cap \omega_m)$ whenever $CK_n \cap \omega_m \neq \emptyset$. Then $N * \varepsilon'_{n,m} + a_{n,m} \xi \uparrow \eta_{N,CK_n}$ as $m \uparrow \infty$ (see Remarque 19 in [8]). Here we may assume that $(\varepsilon'_{n,m})_{m=1}^{\infty}$ converges vaguely as $m \rightarrow \infty$. Put $\varepsilon'_n = \lim_{m \rightarrow \infty} \varepsilon'_{n,m}$; then $\int d\varepsilon'_n \leq 1$. Since $\int dN_p = 1/p$ ($p > 0$), we have, for any $p > 0$ and any $n \geq 1$,

$$\begin{aligned}
 (4.5) \quad p(N - \eta_{N,CK_n}) * N_p &= \lim_{m \rightarrow \infty} p(N - N * \varepsilon'_{n,m} - a_{n,m} \xi) * N_p \\
 &= \lim_{m \rightarrow \infty} (N - N * \varepsilon'_{n,m} - N_p + N_p * \varepsilon'_{n,m} - a_{n,m} \xi) \\
 &= N - \eta_{N,CK_n} - N_p + N_p * \varepsilon'_n.
 \end{aligned}$$

Letting $n \uparrow \infty$, we have $pN_0 * N_p = N_0 - N_p$. Letting $p \downarrow 0$ in (4.5), we have $\lim_{p \downarrow 0} N_p \geq N - \eta_{N,CK_n}$. Thus we see $\lim_{p \rightarrow 0} N_p = N_0$, that is, $(N_p)_{p>0}$ is the resolvent of N_0 . Since N is non-periodic, (4.2) shows that N_p is also non-periodic ($p > 0$), which implies that N_0 is also non-periodic. Remark 14 (1), (2) show that N_0 is a shift-bounded Hunt convolution kernel. On the other hand, we have $pN' * N_p = N'$ for all $p > 0$. Let Γ be the closed subgroup generated by $\text{supp}(N_0)$. For any $x \in X$, $N - N * \varepsilon_x$ is shift-bounded (see Remarque 4 in [8]), and $N' \in \text{SP}_1(N)$ gives the shift-boundedness of $N' - N' * \varepsilon_x$. Lemma 19 and $\text{supp}(N_0) = \text{supp}(N_p)$ ($p > 0$) show that for any $y \in \Gamma$, $(N' - N' * \varepsilon_x) * \varepsilon_y = N' - N' * \varepsilon_x$. This implies

that for any $x \in \Gamma$ and any integer $n \geq 1$, $N' - N' * \varepsilon_{nx} = n(N' - N' * \varepsilon_x)$. For any $f \in C_K^+(X)$, we have $\check{N}' * f(x) \leq \check{N} * f(x) \leq \sup_{y \in \text{supp}(f)} \check{N} * f(y)$ on X . Similarly as in Lemma 23, we have $N' = N' * \varepsilon_x$ for all $x \in \Gamma$. Thus every point in Γ is a period of N' .

We shall give the proof of the “only if” part in Theorem 2. By Remark 7, it suffices to show the following

PROPOSITION 26. *If a real convolution kernel N on X is semi-balayable, non-periodic and satisfies $(N, \xi) \in (\text{TSMP})$, then $\eta_{N,\delta} = -\infty$.*

Proof. Assume contrary that $\eta_{N,\delta} \neq -\infty$. Then $\eta_{N,\delta} \in M(X)$. Put $N' = \eta_{N,\delta}$ and $N_0 = N - N'$. We denote by Γ the closed subgroup generated by $\text{supp}(N_0)$. First we shall show that $N' \in (\text{SMP})$. Let $\mu, \nu \in M_K^+(X)$ with $\int d\mu = \int d\nu \neq 0$ and $a \in R$. Assume that $N' * \mu \leq N' * \nu + a\xi$ in a certain open set $\omega' \supset \text{supp}(\mu)$. By Lemma 23 and Lemma 25, we have $N' * \mu \leq N' * \nu + a\xi$ in $\omega' + \Gamma$. We choose a relatively compact open set ω in X such that $\omega' \supset \bar{\omega} \supset \omega \supset \text{supp}(\mu)$. Let $(\mu', a') \in \text{SB}_N(\mu, 0); C(\bar{\omega} + \Gamma)$. Then $N * \mu' + a'\xi \leq N' * \mu$ in $C(\text{supp}(\mu) + \Gamma)$. Put $c = \int d\mu$. Then $N * \mu \in \text{SP}_c(N)$. Hence Proposition 9 gives $N * \mu' + a'\xi \leq N' * \mu$ on X . Evidently $N * \mu' + a'\xi = N' * \mu$ in $C(\bar{\omega} + \Gamma)$. For an exhaustion $(K_n)_{n=1}^\infty$ of X , we choose $\varepsilon'_{CK_n} \in B_{N_0}(\varepsilon; CK_n)$. Then $\text{supp}(\varepsilon'_{CK_n}) \subset \Gamma$ and $\int d\varepsilon'_{CK_n} = 1$ (see Remark 14 (2) and Lemmas 23, 25), so that

$$N * \mu' * \varepsilon'_{CK_n} + a'\xi \leq N' * \mu * \varepsilon'_{CK_n} = N' * \mu \quad \text{on } X$$

and

$$N * \mu' * \varepsilon'_{CK_n} + a'\xi = N' * \mu \quad \text{in } C(\bar{\omega} + \Gamma).$$

Letting $n \uparrow \infty$, we obtain that

$$N' * \mu' + a'\xi \leq N' * \mu \quad \text{on } X \quad \text{and} \quad N' * \mu' + a'\xi = N' * \mu \quad \text{in } C(\bar{\omega} + \Gamma),$$

because $\lim_{n \rightarrow \infty} N_0 * \varepsilon'_{CK_n} = 0$. Hence $N' * \mu' = N * \mu'$ in $C(\bar{\omega} + \Gamma)$, which shows that $\text{supp}(N_0 * \mu') \subset \bar{\omega} + \Gamma$. This implies $\text{supp}(\mu') \subset \bar{\omega} + \Gamma$. On the other hand, $\text{supp}(\mu') \subset \overline{C(\bar{\omega} + \Gamma)}$, that is, $\text{supp}(\mu')$ is contained in the boundary $\partial(\bar{\omega} + \Gamma)$ of $\bar{\omega} + \Gamma$. Thus $N * \mu' + a'\xi \leq N' * \mu \leq N' * \nu + a\xi$ in $\omega' + \Gamma \supset \text{supp}(\mu')$, and Proposition 9 gives $N * \mu' + a'\xi \leq N' * \nu + a\xi$ on X . This implies $N' * \mu \leq N' * \nu + a\xi$ in $C(\bar{\omega} + \Gamma)$, that is, $N' * \mu \leq N' * \nu + a\xi$ on X , which shows that $N' \in (\text{SMP})$. From $(N, \xi) \in (\text{TSMP})$ and $N' \in (\text{SMP})$, we see also $(N', \xi) \in (\text{TSMP})$.

Evidently N_0 may be considered as a shift-bounded Hunt convolution kernel on Γ . We denote by ξ_r a fixed Haar measure on Γ . Proposition 18 shows that, for any positive Radon measure $\mu \neq 0$ on Γ with compact support (i.e., $\mu \in M^+_k(\Gamma)$), there exist an open set $\omega_r \neq \phi$ in Γ and a relatively compact open neighborhood V_r of the origin in Γ such that:

- (A) For any $\mu'' \in B_{N_0,r}(\mu; \omega_r + V_r)$, $\int d\mu'' < \int d\mu$.
- (B) $E_{N_0,r}(\omega_r) = \phi$, or $E_{N_0,r}(\omega_r) \neq \phi$ and for any $\gamma \in E_{N_0,r}(\omega_r)$, $\int d\gamma = \infty$,

where N_0 being considered as a shift-bounded Hunt convolution kernel on Γ , $B_{N_0,r}(\mu; \omega_r + V_r)$ denotes the totality of N_0 -balayaged measures of μ on $\omega_r + V_r$ and $E_{N_0,r}(\omega_r)$ denotes the totality of N_0 -equilibrium measures of ω_r ¹⁰. Let V be a relatively compact open neighborhood of the origin in X with $\bar{V} \cap \Gamma = \bar{V}_r$. Put $\omega_v = \omega_r + V$; then ω_v is open in X . We choose another open neighborhood U of the origin in X such that $U = -U$ and $U + U \subset V$. We may consider $M^+_k(\Gamma)$ as a subset of $M^+_k(X)$. Choose $(\mu', a') \in SB_N((\mu, 0); \omega_v)$. Then $N * \mu \geq N * \mu' + a'\xi$ on X implies $N' * \mu \geq N' * \mu' + a'\xi$ on X . Assume that $N' * \mu - N' * \mu' - a'\xi = 0$. Then $N_0 * \mu' = N_0 * \mu$ in ω_v and $N_0 * \mu' \leq N_0 * \mu$ on X . Hence $\text{supp}(\mu') = \bar{\omega}_v \cap \Gamma = (\bar{\omega}_r + \bar{V}) \cap \Gamma = \bar{\omega}_r + \bar{V}_r$. Thus we may consider μ' as in $M^+(\Gamma)$. This shows that $\mu' \in B_{N_0,r}(\mu; \omega_r + V_r)$ and $\int d\mu' = \int d\mu$, which contradicts (A). Therefore $N' * \mu - N' * \mu' - a'\xi \neq 0$. By $N' \in (\text{SMP})$ and Proposition 9, we have $\text{supp}(N' * \mu - N' * \mu' - a'\xi) \cap \text{supp}(\mu) \neq \phi$, which implies $\text{supp}(N' * \mu - N' * \mu' - a'\xi) \supset \Gamma$. Let $f \in C^+_k(X)$ with $\text{supp}(f) \subset U$ and $f(0) > 0$. Then there exists $g \in C^+_k(X)$ such that $g \leq f$, $g(0) > 0$ and

$$(4.6) \quad (N' * \mu - N' * \mu' - a'\xi) * f \geq \xi_r * g \quad \text{on } X.$$

Since $N_0 * \mu' = N_0 * \mu + (N' * \mu - N' * \mu' - a'\xi)$ in ω_v , we obtain that

$$(4.7) \quad N_0 * \mu' * f = N_0 * \mu * f + (N' * \mu - N' * \mu' - a'\xi) * f \quad \text{in } \omega_r + U.$$

Let $(\omega_{r,\alpha})_{\alpha \in A}$ be a net of relatively compact open sets in Γ with $\bar{\omega}_{r,\alpha} \subset \omega_{r,\beta}$ ($\alpha \leq \beta$) and $\bigcup_{\alpha \in A} \omega_{r,\alpha} = \omega_r$, $\gamma_\alpha \in E_{N_0,r}(\omega_{r,\alpha})$ ($\alpha \in A$) and let $\mu''_\alpha \in B_{N_0,r}(\mu; \omega_r)$ ¹¹. Then, by (4.6) and (4.7), we have

¹⁰ In the case of $E_{N_0,r}(\omega_r) \neq \phi$, each $\gamma \in E_{N_0,r}(\omega_r)$ satisfies $\text{supp}(\gamma) \subset \bar{\omega}_r$, $N_0 * \gamma \leq \xi_r$ and $N_0 * \gamma = \xi_r$ on ω_r .

¹¹ Similarly as in the definition of $B_{N_0}(\mu; \omega)$, we define $B_{N_0,r}(\mu; \omega_r)$ from $B_{N_0,r}(\mu; \omega_r)$.

$$\begin{aligned} N_0 * (\mu''_{\omega_r} + \gamma_a) * g &\leq N_0 * \mu * g + \xi_r * g \\ &\leq N_0 * \mu * f + (N' * \mu - N' * \mu' - a'\xi) * f \\ &= N_0 * \mu' * f \quad \text{in } \omega_r + U. \end{aligned}$$

Since $\text{supp}((\mu''_{\omega_r} + \gamma_a) * g) \subset \omega_r + U$, the complete maximum principle of N_0 gives $N_0 * (\mu''_{\omega_r} + \gamma_a) * g \leq N_0 * \mu' * f$ on X . Letting $\omega_{r,\alpha} \uparrow \omega_r$, we see, from the dominated convergence property of N_0 , that there exists $\gamma \in E_{N_0, r}(\omega_r)$ such that

$$N_0 * (\mu''_{\omega_r} + \gamma) * g \leq N_0 * \mu' * f \quad \text{on } X$$

(see also Lemma 15 (6)). By the positive mass principle of N_0 (see also Lemma 15 (5)), we have $(\int d\mu''_{\omega_r} + \int d\gamma) \cdot \int g d\xi \leq (\int d\mu') \cdot \int f d\xi$, which implies $\int d\gamma < \infty$. This contradicts (B). The assumption $\eta_{N,\delta} \neq -\infty$ leads to this contradiction. Consequently, $\eta_{N,\delta} = -\infty$. This completes the proof.

Let $(\alpha_t)_{t \geq 0}$ be a vaguely continuous convolution semi-group on X . It is said to be recurrent if there exists $0 \neq f \in C^+_K(X)$ with $\lim_{t \rightarrow \infty} \int_0^t \int f d\alpha_s ds = \infty$, and it is said to be semi-transient if $\lim_{t \rightarrow \infty} \alpha_t = 0$ and $\mu \in M^0_K(X)$, $(\int_0^t \alpha_s * \mu ds)_{t > 0}$ is vaguely bounded.

As we mentioned in Section 1, Theorem 2 and main theorems in [8] (Théorèmes 52 and 52') imply Theorem 1. By Theorem 2 and a result in [8] (see Théorème 25), it can be also stated as follows:

THEOREM 27. *If a real convolution kernel N on X is semi-balayable, non-periodic and satisfies $\inf_{x \in X} N * f(x) \leq 0$ for all $f \in C^0_K(X)$, then there exists a uniquely determined vaguely continuous, markovian, semi-transient and recurrent convolution semi-group $(\alpha_t)_{t \geq 0}$ on X such that for any $t > 0$, $N \geq N * \alpha_t$ and*

$$\lim_{t \rightarrow 0} \frac{N - N * \alpha_t}{t} = \varepsilon.$$

In Theorem 2, it is a question if the condition $\inf_{x \in X} N * f(x) \leq 0$ for all $f \in C^0_K(X)$ can be removed. By Theorem 2 and Proposition 28 in [8], we have the following

Remark 28. Assume that a real convolution kernel N on X satisfies the same conditions as in Theorem 27. Then, for any $\mu \in D^+(N)$ with

$\int d\mu < \infty$ and any open set $\omega \neq \phi$ in X , $\underline{\text{SB}}_N((\mu, 0); \omega) \neq \phi$ and it forms only one element.

In fact, it is known that if $\mu \in M_K^+(X)$, $\underline{\text{SB}}_N((\mu, 0); \omega) \neq \phi$ (see Proposition 28 in [8]). Assume that $\text{supp}(\mu)$ is non-compact. Then we write $\mu = \sum_{n=1}^{\infty} \mu_n$, where $\mu_n \in M_K^+(X)$. Let $(\mu'_n, a'_n) \in \underline{\text{SB}}_N((\mu_n, 0); \omega)$. Then $a'_n \leq 0$. Let ω' be a relatively compact open set $\neq \phi$ in X with $\bar{\omega}' \subset \omega$ and $(\nu, b) \in \underline{\text{SB}}_N((\mu, 0); \omega')$ (see Lemma 8). Then $\sum_{n=1}^{\infty} a'_n \geq b$, that is, $\sum_{n=1}^{\infty} a'_n > -\infty$. This implies that $\sum_{n=1}^{\infty} \mu'_n \in D^+(N)$. Hence we see easily that $(\sum_{n=1}^{\infty} \mu'_n, \sum_{n=1}^{\infty} a'_n) \in \underline{\text{SB}}_N((\mu, 0); \omega)$, that is, $\underline{\text{SB}}_N((\mu, 0); \omega) \neq \phi$. Let (μ', a') and (μ'', a'') be in $\underline{\text{SB}}_N((\mu, 0); \omega)$. Then $N * \mu' + a'\xi = N * \mu'' + a''\xi$. Let $(N_p)_{p>0}$ be the resolvent associated with N and $x \in X$. Since $N * \mu' * (\varepsilon - \varepsilon_x)$ and $N * \mu'' * (\varepsilon - \varepsilon_x)$ are shift-bounded, the above equality and (4.2) give

$$N_p * (\mu' * (\varepsilon - \varepsilon_x)) = N_p * (\mu'' * (\varepsilon - \varepsilon_x)) \quad \text{for all } p > 0,$$

which implies $\mu' - \mu' * \varepsilon_x = \mu'' - \mu'' * \varepsilon_x$. Letting $x \rightarrow \delta$, we have $\mu' = \mu''$, because $\int d\mu' = \int d\mu'' = \int d\mu < \infty$, so that $a' = a''$. Thus $\underline{\text{SB}}_N((\mu, 0); \omega)$ forms only one element.

BIBLIOGRAPHY

- [1] C. Berg and G. Forst, Potential theory on locally compact abelian groups, Springer-Verlag, 1975.
- [2] G. Choquet and J. Deny, Sur l'équation de convolution $\mu = \mu * \sigma$, C. R. Acad. Sc. Paris, **250** (1960), 799–801.
- [3] —, Aspects linéaires de la théorie du potentiel III. Noyaux de composition satisfaisant au principe du balayage sur tout ouvert, C. R. Acad. Sc. Paris, **250** (1960), 4260–4262.
- [4] J. Deny, Sur l'équation de convolution $\mu = \mu * \sigma$, Sém. de la théorie du potentiel, 4ème année, n°5, 1959/60.
- [5] —, Noyaux de convolution de Hunt et les noyaux associés à une famille fondamentale, Ann. Inst. Fourier (Grenoble), **12** (1962), 643–667.
- [6] M. Itô, Sur la famille sous-ordonnée au noyau de convolution de Hunt II, Nagoya Math. J., **53** (1974), 115–126.
- [7] —, Sur le principe relatif de domination pour les noyaux de convolution, Hiroshima Math. J., **5**, 2 (1975), 239–350.
- [8] —, Une caractérisation de noyaux de convolution réels de type logarithmique, Nagoya Math. J., **97** (1985), 1–49.

Masayuki Itô
 Department of Mathematics
 Faculty of Sciences
 Nagoya University
 Chikusa-ku, Nagoya 464
 Japan

Noriaki Suzuki
Department of Mathematics
Faculty of Sciences
Hiroshima University
Naka-ku, Hiroshima 730
Japan