

## Short Note

# On the age and relation between metamorphic gneisses and the Trinity Peninsula Group, Bowman Coast, Graham Land, Antarctica

MICHAEL J. FLOWERDEW

*British Antarctic Survey, NERC, High Cross, Madingley Road, Cambridge CB3 0ET, UK  
mf@bas.ac.uk*

Received 5 March 2008, accepted 21 April 2008

### Introduction

The Trinity Peninsula Group (TPG) of northern Graham Land, a weakly metamorphosed thick sequence of predominantly quartz- and feldspar-rich greywacke, has tentatively been correlated with metasedimentary rocks exposed along the Bowman Coast of Graham Land (Stubbs 1968). The base of the TPG is not observed but the Bowman Coast rocks, here newly defined as the Bowman Coast Succession (BCS), is proximal to high-grade gneisses, which may represent the local basement. Recent geological mapping along the Bowman Coast has allowed a revision of the local geology (Fig. 1) and this note focuses on the relationship of the BCS with the adjacent gneisses.

### Bowman Coast Succession


The BCS varies from being relatively undeformed to intensely folded slates and all of the sequence was later hornfelsed. These factors combined with intermittent exposure and poor rock accessibility hinder detailed stratigraphy. The base of the sequence was not observed. Stratigraphically, the lowest beds consist of slates, originally mudstones and siltstones with rare, thin and persistent quartz-rich sandstone beds. Up-stratigraphy the quartz-sandstone beds become thicker and more frequent. The sequence abruptly becomes dominated by buff-weathering, moderately thickly bedded, laminated, graded and cross-laminated southerly-derived quartz-sandstones and quartzites. This unit also displays soft sediment dewatering and lode structures. Water-lain tuffs conformably overly these sandstones and isolated exposures of intermediate and silicic lavas are thought to occupy the upper-most part of the BCS. The succession is interpreted as initial marine distal turbidites that give way to more proximal, possibly non-marine, deposits with thick volcanic units. Lithologically, the BCS resembles TPG and an association of intermediate volcanic rocks with soft sediment is also recorded within the main TPG outcrop at Mount Cardinal. Assuming a tectonic setting comparable to the TPG, sedimentation may record arc migration through accretionary sediments and the emergence of volcanism in the area. The BCS may be more extensive as similar lithologies outcrop on the Wilkins Coast (Fraser & Grimley 1972).

### Age of the succession and contact relationships

A contact is not observed between the BCS and the gneisses. The BCS is extensively intruded by unfoliated granitoids which have hornfelsed the already deformed sedimentary rocks. None of the cross cutting granitoids is reliably dated. Recalculations from Hole (1986) and Hole *et al.* (1991) suggest the pink granite at Scharer Bluff is  $169 \pm 11$  Ma (whole-rock Rb–Sr isochron). When considered together with adamellite from the same locality an errorchron of  $176 \pm 5$  Ma results. Whilst this age may be erroneous through isotopic disequilibrium between the two rock types, it is in agreement with the 164–169 Ma ages for more reliably dated plutons reported further north on the east coast (Pankhurst *et al.* 2000).

Muscovite and K feldspar from a foliated granite yields a Rb–Sr mineral age of  $173 \pm 3$  Ma. This intrusion does not cut the Curran Bluff metasediments as claimed by Hole (1986). Unfoliated granite does, itself cutting the dated foliated intrusion and obscuring, over a distance of several hundred metres, the relationship between foliated granite and metasediment. The  $173 \pm 3$  Ma age is valid in that it either records the thermal effect of the later granite intrusion or cooling of the whole assembly. The BCS is therefore likely to be older than  $173 \pm 3$  Ma and correlate with the TPG as the TPG has yielded Triassic metamorphic ages (Smellie & Millar 1995), rather than the Late Jurassic back-arc metasedimentary rocks of north-east Palmer Land (Meneilly *et al.* 1987). Gneisses do occur in close proximity to the BCS. Granitic gneisses from Solberg Inlet yield a protolith age of  $238 \pm 3$  Ma, an age widely recorded in Palmer Land (Flowerdew *et al.* 2006). This fieldwork identifies that much of the orthogneiss, including the dated 238 Ma rock, are variably foliated and compositionally banded coarse-grained basic to acidic intrusions with extensive, late, or unrelated injection and veining in places. The term igneous complex is therefore given for these rocks.

Paragneisses are associated with extensive late leucocratic granite and aplite veining. The paragneisses contain garnet- or amphibole-bearing leucosomes, are extensively intruded by granitoids and the whole assembly, here termed metamorphic complex, was folded and foliated during which amphibolite-facies metamorphic minerals grew. The



**Fig. 1.** Geological sketch map for the Bowman Coast showing the location and age of previously dated rocks.

metamorphic complex in many ways resembles that at Auriga Nunataks, north-west Palmer Land (Vaughan *et al.* 1999), with which they may correlate. The age of the metamorphism affecting the Bowman Coast metamorphic complex is not directly dated. However, the pre-metamorphic granitoids cutting the paragneiss are texturally and compositionally similar to the  $238 \pm 3$  Ma igneous complex and are likely to be co-magmatic. The sedimentary protolith has alternating quartz-rich psammite and pelite bands, in many ways resembling a high-grade BCS. Indeed, paragneiss within the metamorphic complex between Stubbs Pass and Pylon Point (Fig. 1) becomes more schistose, contains fewer (mostly not garnet-bearing) leucosomes, is mylonitised by south-east dipping high strain zones and gives the impression of decreasing metamorphic grade eastwards. Rocks of a similar nature are mapped along small parts of Solberg Inlet and on Tonkin Island.

Paragneiss within the metamorphic complex and the BCS may in fact represent different parts of the same sedimentary protolith, different crustal levels of an accretionary prism through which a Triassic magmatic arc has developed. Granitoid - BCS contacts represent the highest crustal levels while the igneous and metamorphic complexes document the interaction of voluminous arc plutonism with the sedimentary protolith host at a lower crustal level. To

some extent, therefore, an oblique cross section through the middle crust of a Triassic arc is exposed at the Bowman Coast. Uplift from lower crustal levels is evident from decompression textures such as plagioclase corona around garnet in some paragneisses, and was further facilitated by ductile shear zones and by ascent within the late plutons. Future work will test the validity of this model by determining whether or not the metamorphic complex is older and unrelated to the BCS and Triassic arc and whether or not the BCS is a correlative of the TPG.

### Acknowledgements

This work could not have been achieved without the tireless work of the operations and air unit staff at Rothera research station and field assistant Catrin Thomas. Discussions with Teal Riley and insightful reviews by John Smellie and Alan Vaughan are greatly appreciated. This is a contribution to the east coast mapping project of LTMS-G.

### References

- FLOWERDEW, M.J., MILLAR, I.L., VAUGHAN, A.P.M., HORSTWOOD, M.S.A. & FANNING, C.M. 2006. The source of granitic gneisses and migmatites in the Antarctic Peninsula: a combined U-Pb SHRIMP and laser ablation Hf isotope study of complex zircons. *Contributions to Mineralogy and Petrology*, **151**, 751–768.
- FRASER, A.G. & GRIMLEY, P.H. 1972. The geology of parts of the Bowman and Wilkins coasts, Antarctic Peninsula. *Falkland Island Dependencies Survey Scientific Reports*, No. 67, 59 pp.
- HOLE, M.J., PANKHURST, R.J. & SAUNDERS, A.D. 1991. Geochemical evolution of the Antarctic Peninsula magmatic arc: the importance of mantle-crust interactions during granitoid genesis. In THOMSON, M.R.A., CRAME, J.A. & THOMSON, J.W., eds. *Geological evolution of Antarctica*. Cambridge: Cambridge University Press, 369–374.
- HOLE, M.J. 1986. *Time controlled geochemistry of igneous rocks of the Antarctic Peninsula*. PhD thesis, University of London, 396 pp. [Unpublished].
- MENEILLY, A.W., HARRISON, S.M., PIERCY, B.A. & STOREY, B.C. 1987. Structural evolution of the magmatic arc in northern Palmer Land, Antarctic Peninsula. In MCKENZIE, G.D., ed. *Gondwana Six: structure, tectonics and geophysics*. American Geophysical Union, *Geophysical Monograph*, **40**, 209–219.
- PANKHURST, R.J., RILEY, T.R., FANNING, C.M. & KELLEY, S.P. 2000. Episodic silicic volcanism in Patagonia and the Antarctic Peninsula: chronology of magmatism associated with the break-up of Gondwana. *Journal of Petrology*, **41**, 605–625.
- SMELLIE, J.L. & MILLAR, I.L. 1995. New K-Ar isotopic ages of schists from the Nordenskjöld Coast, Antarctic Peninsula: oldest part of the Trinity Peninsula Group? *Antarctic Science*, **7**, 191–196.
- STUBBS, G.M. 1968. *Geology of parts of the Foyen and Bowman Coasts, Graham Land*. PhD thesis, University of Birmingham, 245 pp. [Unpublished].
- VAUGHAN, A.P.M., MILLAR, I.L. & THISTLEWOOD, L. 1999. The Auriga Nunataks Shear zone: Mesozoic transfer faulting and arc deformation in northwest Palmer Land, Antarctica. *Tectonics*, **18**, 911–928.