# Complete Culture Media

## **WO ENTIRELY NEW FORMS\***

### \* TABLETS

\* GRANULES

—for preparation of single of media. Each tablet when dissolved in water produces a convenient quantity.

-for larger scale preparatubes and small quantities tions. Hard and of even size, the granules are less hygroscopic and easier to handle than spray-dried powder.

Both tablets and granules retain all the properties of the original medium, from which they are indistinguishable when reconstituted.

#### ADVANTAGES ARE APPARENT-

Big time-saving in preparation • No specialised equipment needed Variability reduced to minimum . Easily portable for field work Results reproducible from one laboratory to another when using media from central source . Economical, as only sufficient for immediate requirements need be used.



#### THE FOLLOWING ARE AMONG THE MEDIA AVAILABLE AS TABLETS AND GRANULES

Nutrient Broth Nutrient Agar . McConkey Broth McConkey Agar . Peptone Water . Glucose Peptone Broth Glucose Peptone Agar · Agar-Agar



OXO LIMITED (Medical Dept.)

Thames House, Queen St. Place, London, E.C.4. Tel. CENtral 9781

#### CONTENTS

(All rights reserved)

	PAGE
LANCASTER, H. O. The theory of amoebic surveys. (With 1 Figure in the Text) .	257
HOYLE, L. The multiplication of influenza viruses in the fertile egg. (With Plates 3 and 4, and 5 Figures in the Text)	277
ARMITAGE, P. and Allen, IRENE. Methods of estimating the LD 50 in quantal	
response data	298
ROWNTREE, G. Accidents among children under two years of age in Great Britain .	323
Bruce, H. M. Feeding and breeding of laboratory animals. XII. A note on the breeding of guinea-pigs without fresh green food. (With 1 Figure in the Text).	338
LAUGHTON, NANCY. Vaginal corynebacteria. (With Plates 5 and 6)	346
STEWART, F. S. and McKeever, J. D. The anti-globulin technique applied to the detection of non-agglutinating antibody against Salmonella typhi O in human	
sers	357
Springert, V. H. A comparative study of tuberculosis mortality rates. (With 20	
Figures in the Text)	361

The Journal of Hygiene is issued four times a year.

Quotations can be given for back volumes on application.

Papers for publication should be sent to Professor E. T. C. Spooner, London School of Hygiene and Tropical Medicine, Keppel Street, Gower Street, London, W.C. 1. Other communications should be addressed to the University Press, Cambridge.

Papers forwarded to the Editor for publication are understood to be offered

to The Journal of Hygiene alone, unless the contrary is stated.

Contributors receive twenty copies of their papers free. Twenty-five additional copies may be had at cost price: these should be ordered when the final proof is returned.

The subscription price is £3. 15s. 0d. per volume (post-free), payable in advance; single numbers 20s. net. Subscriptions may be sent to any Bookseller, or to The Cambridge University Press, Bentley House, 200 Euston Road, London, N.W. 1.

Enquiries from the U.S.A. should be addressed to the American Branch of the Press at 51 Madison Avenue, New York, 10.