BRITISH MUSEUM NATURAL RADIOCARBON MEASUREMENTS VI

HAROLD BARKER, RICHARD BURLEIGH, and NIGEL MEEKS

Research Laboratory, British Museum, London W.C. 1

Dates listed below are based on measurements made up to May 1968, and cover a period during which the technique of gas proportional counting using CO₂ was gradually replaced by liquid scintillation counting using benzene. The gas counting measurements were carried out by the method and techniques previously described (Barker and Mackey, 1968) the only modifications being the replacement of some old electronic units by more stable solid-state equipment; proportional counting results are indicated in the text by (P) at the end of the relevant sample descriptions. Liquid scintillation counting, which is now the preferred method in this laboratory, is carried out using a Packard Tri-Carb liquid scintillation spectrometer model 3315/AES fitted with selected low-noise quartz-faced photomultipliers. Normally 3 ml of benzene is prepared from each sample. This is dissolved in 12 ml of scintillation grade toluene containing 5 gm/liter of scintillator (PPO) and the solution is measured in a standard low-potassium glass vial at a temperature of 0°C. Photomultiplier E.H.T., amplifier, and channel width settings are optimized for C¹⁴, and measurements are carried out at ca. 65% efficiency of detection for C¹⁴ to eliminate interference from any tritium which may be present in the benzene. Under these circumstances the background is approx. 8.6 cpm and the modern (95% Aox) is approx. 24.0 cpm. Samples are counted in groups of 3 to 5 together with background and modern reference samples and are measured for at least one week, the instrument being set to cycle at 100 min intervals. In this period, the counts accumulated are such that the background is always measured to a statistical accuracy of better than 1% and most other samples to a higher accuracy than this. Background and modern counts used in the calculation of each result are only those relevant to the period of measurement of that particular sample. Statistical analysis of groups of replicate measurements made under these conditions over a very long period of time has demonstrated the excellent long-term stability of the equipment and indicates that the technique is quite capable of achieving results of very high statistical accuracy when required.

The synthesis of benzene follows broadly the procedure described by Noakes, Kim, and Stipp (1965), *i.e.*, acetylene is prepared from CO_2 via lithium carbide as described by Barker (1953; Barker and Mackey, 1959) and this is converted into benzene using a silica-alumina catalyst activated with vanadium pentoxide. The yield at each stage of the synthesis is very high and the over-all conversion efficiency from CO_2 to benzene ranges from 92 to 95%.

Adoption of the liquid scintillation technique has also coincided with the development of a new approach to combustion of samples (Barker *et al.*, 1969) in which samples are burnt in oxygen at high pressure within a thick-walled stainless steel pressure vessel. This procedure, which is semi-automatic, has several advantages over the conventional system using a combustion tube furnace; in particular, it saves considerable operator time and yields, without further purification CO_2 which is low in oxides of nitrogen and sulphur even when prepared from materials rich in protein such as the organic fraction of bones and antler.

Ages are calculated on the basis of a half-life of 5568 years and unless otherwise stated, the error terms also include a contribution of \pm 80 years to allow for possible isotopic fractionation effects. The practice adopted in previous date lists of widening the error term to include possible errors due to fluctuations in the C¹⁴ content of the carbon exchange reservoir in the past (de Vries effects) has been discontinued, thus leaving the user of the dates free to apply corrections for such deviations as they become available (e.g., Stuiver and Suess, 1966).

ACKNOWLEDGMENT

Thanks are due to Miss G. I. Hassall of the National Physical Laboratory, for isotopic fractionation measurements made in connection with the investigation of age discrepancies in Egyptian materials.

SAMPLE DESCRIPTIONS

ARCHAEOLOGIC SAMPLES

A. Crete

Knossos series

Five samples from Neolithic settlement of Knossos, Crete $(35^{\circ} 31'$ N Lat, 25° 20' E Long), from sounding below central court of Minoan Palace (Evans, 1964a,b). Coll. 1960 and subm. by J. D. Evans, Inst. of Archaeol., Univ. of London, for comparison with unexpectedly early dates previously obtained for Knossos Neolithic, BM-124, 8050 \pm 180 and BM-126, 7000 \pm 180 (Radiocarbon, 1963, v. 5, p. 104).

7570 ± 150 5620 в.с.

6140 ± 150 4190 в.с.

Sample 2, ref. Area AC, Level 24. Charcoal from Occupation layer assoc. with 1st brick houses and immediately overlying earliest camp occupation above bedrock (Stratum IX, Evans, 1964b). Expected age early 6th millennium B.C. based on BM-124 and BM-126. (P)

6210 ± 150 4260 в.с.

Sample 3, ref. Area AC, Level 17. Charcoal from habitation deposit in Stratum VI (Evans, 1964b). Expected age late 6th millennium B.C. based on BM-124 and BM-126. (P)

BM-274. Knossos 4

BM-273. Knossos 3

BM-272. Knossos 2

Sample 4, ref. Area A, Level 15. Charcoal from habitation level in Stratum V (Evans, 1964b). Expected age end of 6th millennium B.C. based on BM-124 and BM-126. (P)

5680 ± 150 3730 в.с.

BM-279. Knossos 6

Charcoal sample from Upper Neolithic level, Stratum IV (Evans, 1964b). (P)

General Comment (J.D.E.): new series of dates in conjunction with Stratum X date (BM-124), but without BM-126, gives consistent and acceptable picture of development of Knossos Neolithic through 3 millennia.

Knossos, the cereal problem

Carbonized grain containing high proportion of naked hexaploid wheat (bread wheat, id. by H. Helbaek) found in Stratum X assoc. with carbonized wooden stake previously dated at 8050 ± 180 (BM-124). Date was unexpectedly early by comparison with other evidence for westward spread of domesticated cereals from earliest centers of cultivation. However, date for BM-124 was confirmed by remeasurement, using liquid scintillation technique (BM-278) and measurement on carbonized grain provided direct confirmation of early occurrence of grain of this type at Knossos (ref. Evans, Athens, Ann. of British School, in press).

BM-278.Knossos 1 7910 ± 140 5960 B.C.

Charcoal, remains of carbonized wooden stake found in Pit F, Area AC, Level 27, in central court of Minoan Palace (ref. BM-124).

BM-436. Knossos 7

7740 ± 140 5790 в.с.

Carbonized grain, assoc. with carbonized stake described above (ref. BM-124).

B. Egypt

Tarkhan II

Linen from Mastaba 2050 at Tarkhan near Cairo, Egypt (29° 40' N Lat, 31° 13' E Long). Coll. 1912 by W. M. Flinders Petrie and considered by him to date to First Dynasty (Petrie, 1914). Subm. by I. E. S. Edwards, British Mus. Also dated by UCLA and Arizona labs; UCLA-739, 4265 ± 80 ; A569, 4295 ± 90 . Expected age 5000 B.P. (ca. 3050 B.C.).

BM-203. Tarkhan II	4150 ± 110
Pretreated with dilute acid only. (P)	2200 b.c.
BM-248. Tarkhan II	4160 ± 110 2210 в с

Pretreated with dilute acid followed by boiling in 10% aqueous solution of 0.880 ammonia until fabric was almost pure white, followed by thorough washing in distilled water. *Comment*: 2nd measurement after more rigorous pretreatment was made because of suggestion that dark color of linen might be due to non-contemporaneous contaminant such as bitumen. Evidently this was not so. (P)

Egyptian chronology series

Radiocarbon measurements of Egyptian material of known age from late 2nd and 3rd millennia B.C. have already indicated that there are serious discrepancies between radiocarbon and calendar ages in this period (see, e.g., Smith, 1964; Damon *et al.*, 1963; Barker and Mackey, 1959, 1961; Berger *et al.*, 1965). More recently, Suess produced detailed plot of discrepancies from ca. 4000 to 1500 B.C. based on radiocarbon measurements of bristlecone pine wood from E central California (Suess, 1967). Following series of freshly coll. samples obtained specifically to investigate this problem are securely dated archaeologically and of exactly known provenance (see Emery, 1938, 1939, 1954, 1958, 1961, 1965). Where possible material such as reed or fabric, representing one or only a few seasons' growth was obtained.

Samples have also been measured by Arizona and UCLA labs (Berger and Libby, 1967). Samples 1-4, 6, 7, 10-13 are from Egypt Exploration Society's excavations at Sakkara Archaic Cemetery (29° 51' N Lat, 31° 14' E Long). Sample 15 is from El Lahun (Fayûm) (29° 13' N Lat, 30° 59' E Long). Coll. 1966 by G. T. Martin; subm. by I .E. S. Edwards, British Mus., who initiated and arranged project. More detailed analysis of results will be publ. elsewhere. Samples were pretreated with dilute acid.

		4300 ± 65
BM-228.	Sakkara 10/66	2350 в.с.
	,	$\delta C^{13} = -23.2 + 1.0\%$

Reed used as bonding between mud-brick courses, W side of superstructure, Tomb 3357 (Hor-Aha). Beginning of 1st Dynasty, reign of Hor-Aha, ca. 3100 B.C. (P) Age based on 5730 yr half-life 4430 \pm 65, 2480 B.C.

		4520 ± 65
BM-229.	Sakkara 1/66	2570 в.с.
		$\delta C^{_{13}} = -21.8 \pm 1.0\%$

Reed used as bonding between mud-brick courses, N side of superstructure, Tomb 3503 (Mer-Neit). Mid 1st Dynasty, reign of Den, ca. 3000 B.c. (P) Age based on 5730 yr half-life 4660 ± 65 , 2710 B.c.

			4380 ± 65
BM-230.	Sakkara	11/66	2430 в.с.
		,	$\delta C^{_{13}}=-23.7\pm1.0\%$

Reed used as bonding between mud-brick courses, S side of superstructure, Tomb 3035 (Hemaka). Mid 1st Dynasty, end of reign of Den, ca. 2950 B.C. (P) Age based on 5730 yr half-life 4510 ± 65 , 2560 B.C.

			4270 ± 65
BM-231.	Sakkara	2/66	2320 в.с.
		'	$\delta C^{_{13}} = -23.9 \pm 1.0\%$

Reed used as bonding between mud-brick courses of inner enclosure wall, W side, Tomb 3505 (Ka'a). End of 1st Dynasty, reign of Qaa, ca. 2900 B.C. (P) Age based on 5730 yr half-life 4400 \pm 65, 2450 B.C.

BM-232.Sakkara 12/66 4230 ± 65
2280 B.C.
 $\delta C^{13} = -23.1 \pm 1.0\%$

Reed used as bonding between mud-brick courses, W side of superstructure, Tomb 3046. Second Dynasty, ca. 2750 B.C. (P) Age based on 5730 yr half-life 4360 ± 65 , 2410 B.C.

DM 000		4000 ± 65
BM-233.	Sakkara 13/66	2050 в.с.
		$\delta C^{_{13}} = -23.0 \pm 1.0\%$

Reed used as bonding between mud-brick courses, W side of entrance stairway, Tomb 3030. Early 3rd Dynasty, ca. 2675 B.C. (P) Age based on 5730 yr half-life 4120 ± 65 , 2170 B.C.

BM-234.	Sakkara 4/66	3790 ± 65 1840 в.с.
		$\delta C^{_{13}} = -25.1 \pm 1.0\%$

Wood (Acacia sp. prob A. nilotica L.) from floor of S chapel or offering niche on E side of superstructure, Tomb 3510. Third Dynasty, ca. 2650 B.C. (P) Age based on 5730 yr half-life 3900 \pm 65, 1950 B.C.

BM-235.	Sakkara 3/66	4070 ± 65 2120 B.C.
		$\delta C^{13} = -23.7 \pm 1.0\%$

Reed used as bonding between mud-brick courses of S side of superstructure, Tomb 3073-5 (Kha'-bau-Sokar). End of 3rd Dynasty to beginning of 4th Dynasty, ca. 2600 B.C. (P) Age based on 5730 yr half-life 4190 ± 65 , 2240 B.C.

BM-236.	Sakkara 6/66	3840 ± 65 1890 в.с.
		$\delta C^{_{13}} = -24.8 \pm 1.0\%$

Cloth (linen) from undisturbed burial between 3rd Dynasty Tombs 3508 and 3510 (Emery, 1965) tentatively dated to 4th Dynasty, ca. 2550 B.C. (P) Age based on 5730 yr half-life 3960 \pm 65, 2010 B.C.

BM-237. Sakkara 7/66

3720 ± 110 1770 в.с.

Human bone collagen, from undisturbed burial between Tomb 3508 and 3510 adjacent to tomb containing Sample 6 (BM-236, above). Fourth Dynasty (?), ca. 2550 B.C. (P) Age based on 5730 yr half-life 3830 \pm 110, 1880 B.C.

BM-238. El Lahun

 3580 ± 65 1630 b.c. $\delta C^{13} = -12.0 \pm 1.0\%$

Sample 15/66. Reed used as bonding between mud-brick courses, N boundary wall of pyramid of Senusret II (Sesostris II). Twelfth Dynasty, ca. 1880 B.C. (P) Age based on 5730 yr half-life 3690 ± 65 , 1740 B.C.

BM-280. El Lahun

3550 ± 65 1600 в.с.

 $\delta C^{13} = -11.1 \pm 1.0\%$

Sample 15/66. Remeasurement using fresh portion of sample (BM-238, above) pretreated with dilute alkali in addition to standard acid pretreatment. (P) Age based on 5730 yr half-life 3660 ± 65 , 1710 B.C.

C. France

Trou des Forges, Bruniquel

Antler biserial harpoon fragments and antler waste from Trou des Forges, Bruniquel, Tarn-et-Garonne, France (44° 03' N Lat, 1° 40' E Long). Coll. 1864 by Vicomte de Lastic; subm. by G. de G. Sieveking, British Mus., from British Mus. collection. Antler biserial harpoons define industry termed by Breuil Magdalenian VI; by comparison with other publ. dates for recently excavated samples (see Broecker and Kulp, 1957; Vogel and Waterbolk, 1963; Leroi-Gourhan and Brezillon, 1966) it was hoped to form general assessment of whether selected samples from old collections can sometimes be used as reliable source of material for dating Upper Palaeolithic industries. Samples were demineralized before burning.

	$11,750 \pm 300$
BM-302.	9800 в.с.

Harpoon fragments (protein fraction).

BM-303.

$11,110 \pm 160$ 9160 в.с.

 $12,070 \pm 180$ 10.120 в.с.

Antler waste (protein fraction). Comparable material has been dated at 2 other sites: Grotte de la Vache, Ariege, France, Level 2. L-336c, $11,600 \pm 200$; GrN-2025, 12,540 ± 105 . Level 4. GrN-2026, 12,850 ± 60 . Pincevent, Montereau, Seine et Marne, France, Lv-291, 10,920 ± 540; Ly-292, 11.610 + 400; Ly-293, 11,310 \pm 330, see refs, above. Comment: dates are fully within expected age range for Magdalenian VI.

BM-304. Montastruc, Bruniquel

Antler waste (protein fraction) from Montastruc rock shelter, Bruniquel, Tarn-et-Garonne, France (44° 03' N Lat, 1° 40' E Long). Coll. ca. 1868 by Peccadeau de Lisle; subm. by G. de G. Sieveking, British Mus. Sample is from level containing Magdalenian IV-V industry, in which fine ivory animal carvings now in British Mus. collection were found. Date fits well with BM-302 and BM-303, above. Sample was demineralized before burning .

D. Great Britain

1610 ± 150

BM-267. Minepit Wood

Charcoal (ref.D5(9), Sample 1) from early iron-working site at Minepit Wood (Orznash), Withyham, Sussex, England (51° 06' N Lat, 0° 10' E Long). Coll. 1964 and subm. by J. H. Money, Sussex Archaeol. Soc.

А.D. 340

Sample from lowest part of early iron-working furnace which was securely sealed by undisturbed soil, and dates last act of smelting in furnace. (P)

Brown Edge series

Three samples of charcoal from Brown Edge, Totley, Derbyshire, England (53° 18' N Lat, 1° 34' W Long), from burial pits 12 to 14 in. deep within low earthwork circle with small central cairn of stones. Coll. 1963 by J. Radley; subm. by G. D. Lewis for Sheffield City Mus., Sheffield, Yorkshire (Radley, 1966). (P)

BM-177. Brown Edge, Burial 1 3000 ± 150 1050 в.с.

Cremation burial contained in "Pennine" urn. Comment (G.D.L.): date is late for collared urn of primary series (Longworth, 1961). See NPL-10, 3355 ± 155 , NPL-11, 3470 ± 145 (Callow *et al.*, 1963). (P)

		3200 ± 150
BM-211.	Brown Edge, Burial 2	1250 в.с.

Un-urned cremation beneath cairn and adjacent to Burial 1 (BM-177). (P)

		3480 ± 150
BM-212.	Brown Edge, Burial 4	1530 в.с.

Un-urned cremation found adjacent to Burials 1 and 2 but not sealed by cairn. (P)

Harland Edge series

Two charcoal samples from filling of pits containing food vessels from Harland Edge, Beeley Moor, Derbyshire, England (53° 13' N Lat, 1° 34' W Long). Coll. 1961-62 by D. N. Riley; subm. by G. D. Lewis for Sheffield City Mus., Sheffield, Yorkshire (Riley, 1966).

		J II U _ 1JU
BM-178.	Harland Edge, Pit 2 (Sample A) (P)	1490 в.с.

 3700 ± 150

 2440 ± 150

BM-210. Harland Edge, Pit 3 (Sample B) (P) 1750 B.C.

Comment (G.D.L.): 1st available radiocarbon dates with food vessel assoc.

BM-179. Barbrook II

3450 ± 150 1500 в.с.

Barbrook II earth circle, Ramsley Moor, Holmesfield, Derbyshire, England (53° 16' N Lat, 1° 35' W Long). Coll. 1963 and subm. by G. D. Lewis, Sheffield City Mus. Charcoal (ref. D.6a, Burial 1) from cremation burial contained in collared urn sealed in pit beneath low stone cairn in low earth circle apparently part of adjacent group of cairns and stone circle. *Comment* (G.D.L.): date agrees with other similar radiocarbon dates for assoc. secondary series collared urns. See NPL-17, 3345 \pm 160, NPL-18, 3355 \pm 160 (Callow *et al.*, 1963); Grn-1686, 3440 \pm 60 (Vogel and Waterbolk, 1964). (P)

BM-213. Chapel Flat Dyke, canoe

Wood from dug-out canoe from Chapel Flat Dyke, Rotherham, Yorkshire, England (53° 25' N Lat, 1° 23' W Long). Found in river mud in ancient bed of Chapel Flat Dyke, at depth 18 ft below present surface, during tunnelling operations. Coll. 1963 and subm. by H. Raymond Singleton, Dir., Sheffield City Mus. No assoc. finds. (P)

BM-315. Eriswell

 3470 ± 115 1520 в.с.

Eriswell, W Suffolk, England (52° 23' N Lat, 0° 30' E Long). Charcoal from wooden object, perhaps bier, on old ground surface under Middle Bronze age barrow and over primary cremation with collared urn in cylindrical hole. Coll. 1966 and subm. by D. P. Dymond, of Extra-Mural Studies, Univ. of Cambridge.

BM-180. Windmill Hill Long Barrow

Antler, ref. B.54, H 419, (326) from primary chalk silt in ditch of Windmill Hill Long Barrow, Avebury, Wiltshire, England (51° 26' N Lat, 1° 52' W Long). Coll. 1961 by P. Ashbee; subm. by G. de G. Sieveking, British Mus. For comparison see dates for Windmill Hill, BM-73, 4910 \pm 150 (Barker and Mackey, 1961); Hembury, BM-130, 5100 \pm 150 (Barker and Mackey, 1963), BM-136, 5190 \pm 150, BM-138, 5280 \pm 150; Fussells Lodge, BM-134, 5180 \pm 150 (Barker and Mackey, 1968), and Willerby Wold, BM-188, 4900 \pm 150, BM-189, 4960 \pm 150 (this date list, below). (\mathbf{P})

Flint mine chronology

Antler picks measured as part of investigation of chronological sequence and period of use of Neolithic flint mines in S England. Picks were left in mine galleries when these were abandoned and subsequently sealed off by backfilling of shaft as adjacent new shafts were dug. Work will be reported and discussed elsewhere. All samples were demineralized so that dates were obtained on protein fraction. Subm. by G. de G. Sieveking, British Mus.

Cissbury area flint mines

BM-181. Church Hill

3390 в.с.

Sample, ref. 61/1584/A from Church Hill, Findon, Sussex, England (50° 51' 42" N Lat, 0° 25' 4" W Long). Coll. ca. 1950 by J. Pull. (P)

BM-182. Harrow Hill

Sample, ref. 4464, H.H. 21.58 from Harrow Hill, Sussex, England (50° 52′ 45″ N Lat, 0° 27′ 45″ W Long). Coll. 1924-25 by E. and E. C. Curwen. (P)

3450 ± 150 1500 в.с.

5190 ± 150 3240 в.с.

 4930 ± 150 2980 в.с.

5340 ± 150

4720 ± 150 2770 b.c.

Sample, ref. 61/1586/A from Cissbury, Sussex, England (50° 51' 30" N Lat, 0° 23' 06" W Long). Coll. ca. 1952 by J. Pull. (P)

BM-184. Cissbury, Harrison's pit 4650 ± 150 2700 в.с.

Sample, ref. 61/1586/A from Cissbury, Sussex, England (50° 51' 30" N Lat, 0° 23' 06" W Long). Coll. 1877 by Park Harrison. (P)

BM-185. Cissbury, Shaft 6 4730 ± 150 2780 в.с.

Sample, ref. 3970 from Cissbury, Sussex, England (50° 51' 30" N Lat, 0° 23' 06" W Long). Coll. 1878 by Park Harrison. (P)

 BM-190. Easton Down flint mine
 4480 ± 150

 2530 B.C.

From Easton Down, Nr. Salisbury, Wiltshire, England (51° 05' N Lat, 1° 48' W Long). Coll. 1934 by J. F. Stone. (P)

Grimes Graves flint mines

BM-377.

BM-290.

BM-183. Cissbury

Samples from Grimes Graves, Thetford, Norfolk, England (52° 25' 39" N Lat, 0° 38' 41" E Long).

3550 ± 150 1600 в.с.

BM-276. Grimes Graves, Pit XII, 1933

Sample from Pit XII. Coll. ca. 1933 by A. L. Armstrong. Intended as check on previous Grimes Graves dates (Barker and Mackey, 1963) for comparison with Cissbury series and expected to give same date as BM-97, 4290 \pm 150, original sample from Pit XII. However, it seems probable that this sample which came from layer in shaft and not sealed gallery, dates later infilling of shaft and not active use of this mine. See BM-377 below. (P)

Grimes Graves, Pit XII, 37/38 4250 ± 130 2300 в.с.

Sample from Pit XII. Coll. 1937-1938 by A. L. Armstrong. Repeat by liquid scintillation method of gas proportional counter measurement BM-97, 4290 \pm 150 (Barker and Mackey, 1963). See BM-276 above.

BM-291. Grimes Graves, Greenwell pit 3810 ± 130 1860 B.C.

Sample from Greenwell pit, Gallery III. Coll. 1870 by Canon Greenwell. *Comment*: date is within expected age range 1600 to 2400 B.C. based on previous series of Grimes Graves dates (Barker and Mackey, 1963).

5090 ± 130 Blackpatch flint mine3140 B.c.

Antler, ref. 61/1585/A, Shaft 4, from Blackpatch, Worthing, Sussex, England (50° 52' N Lat, 0° 27' 15" W Long). Coll. 1923 by J. Pull; subm. by L. M. Bickerton, Worthing Mus.

Willerby Wold long barrow

Two samples of charcoal (Quercus sp.) from charred timber structure beneath E end of Willerby Wold long barrow, near Bridlington, East Riding, Yorkshire, England (54° 09' N Lat, 0° 06' W Long). Coll. 1960 and subm. by T. G. Manby (Manby, 1963).

BM-188. Willerby Wold, A	4900 ± 150 2950 в.с.
Sample A, from base of crematorium deposit. (P)	
BM-189. Willerby Wold, B	4960 ± 150 3010 в.с.

Sample B, from center of facade bedding trench. (P) Comment: dates show that elaborate crematorium long barrows of Yorkshire developed early in British Neolithic period. Features of mortuary enclosure of barrow resemble those of enclosure at Fussell's Lodge (Ashbee, 1958, 1964) BM-134, 5180 \pm 150 (Barker and Mackey, 1968). See also date and Comment above for Windmill Hill long barrow, BM-180, this date list.

High Peak

Two charcoal samples from High Peak, Sidmouth, Devon, England (50° 40' N Lat, 3° 06' W Long), Neolithic site sealed beneath Dark Age ramparts. Coll. 1961 and 1964 by Mrs. S. H. Pollard, Devon Archaeol. Explor. Soc; subm. Dept. of Prehist. and Roman Britain, British Mus. Assoc. archaeol. material consisted of pottery and flint artifacts similar to those from Windmill Hill, BM-73, 4910 \pm 150 (Barker and Mackey, 1961) and Hembury, BM-130, 5100 \pm 150 (Barker and Mackey, 1963).

BM-214.	High Peak, Ref. A.3. (P)	4810±150 2860 в.с.
BM-215.	High Peak, Ref. HB.3. (P)	1550 ± 150 а.д. 400

Comment (S.H.P.): in 1964 this level was thought to be part of Neolithic ditch fill. Further excavation in 1965 revealed that it was later tip of ashy material overlying remains of Neolithic ditch and cut into by Dark Age workings (Pollard, 1966). Date agrees with late 5th to early 6th century A.D. Dark Age amphorae sherds found.

Giants Hills

Two red deer antlers from Giants Hills long barrow Skendleby, Lincolnshire, England (53° 11' N Lat, 0° 05' E Long). Coll. 1935 by C. W. Phillips; subm. by G. de G. Sieveking, British Mus.

BM-191.	Giants Hills, Sample 1,	4410 ± 150
	(protein fraction)	2460 в.с.
Ref. 1935,	4-12, 42. (P)	
BM-192.	Giants Hills, Sample 2,	4320 ± 150
	(protein fraction)	2370 в.с.
Ref. 1935,	4-12, 55. (P)	

BM-271. Aston I

Carbonized grain, ref. A. 058, (probably emmer) from lining inside upper part of small pit in old ground surface sealed by primary beaker barrow at Aston on Trent, Derbyshire, England (52° 52' N Lat, 1° 25' W Long). Coll. 1964 and subm. by D. Reaney, Derbyshire Archaeol. Soc. Expected age ca. 4800 B.P. *Comment*: date agrees well with assoc. Grimston ware. (P) 4680 ± 135

BM-281. Langdale

Charcoal, 90 cm below surface of peat assoc. with several stone implements and chippings at site of Neolithic axe factory at Langdale, Westmoreland, England (54° 27' N Lat, 3° 07' W Long). Coll. 1966 by E. P. Johnson, Cumberland and Westmoreland Archaeol. Soc.; subm. by H. Godwin. Date for manufacture of stone axes at Langdale.

Durrington Walls

Two charcoal samples from late Neolithic henge monument at Durrington Walls, Wiltshire, England (51° 12' N Lat, 1° 47' W Long). Coll. 1966 and subm. by G. J. Wainwright, Inspectorate of Ancient Monuments, Ministry of Public Bldg. and Works (Wainwright, 1967, 1968).

3560 ± 120

BM-285. Durrington Walls, Sample 1 1610 B.C.

Ref. D. W. 1966, D.1, Ditch, Hearth 5. Hearth near base of Neolithic ditch. Intended to date henge and Rusticated Beaker. Expected age ca. 1700 B.C.

				3630 ± 110
BM-286.	Durrington	Walls,	Sample 2	1680 в.с.

Ref. D.W. 1966, D.1, Ditch, Hearth 3. Hearth in secondary silt of Neolithic ditch. Intended to date plain Ebbsfleet ware. Expected age ca. 1700 B.C.

Letchworth

Charcoal from Blackhorse Rd., Letchworth, Hertfordshire, England (51° 59' N Lat, 0° 12' W Long). Coll. 1960 and 1966 and subm. by]. Moss-Eccardt, Letchworth Mus.

BM-186. Letchworth, Sample 1	1470 в.с.
Ref. Pit 7, Layer 3, assoc. with Ebbsfleet ware. (P)	3310 ± 150 1360 в.с.
Ref. Pit 7, Layer 4, assoc. with Ebbsfleet ware. (P)	3830 ± 140
BM-283. Letchworth, Sample 3	1880 в.с.

Ref. Pit 7, Layer 4, assoc. with Ebbsfleet ware. Check on BM-187. Comment: age of these samples was expected to agree with BM-113,

4700 ± 150 2750 в.с.

2730 в.с.

Ebbsfleet 4660 \pm 150 (Barker and Mackey, 1963). Pit from which samples were excavated may have been filled in during Bronze age with refuse which included charcoal samples measured and sherds of older Ebbsfleet ware lying on surface at that time. Mixing of material may also account for poor agreement between BM-187 and BM-283.

BM-284. Letchworth, Sample 4

3590 ± 130 1640 в.с.

 4180 ± 190

Ref. Pit 1, Layer 4, assoc. with pottery assemblage consisting of longnecked beaker, rusticated beaker, and Fengate ware. Date agrees well with expected age of ca. 1700 B.C. and date for similar assemblage at Windmill Hill BM-75, 3500 \pm 150 (Barker and Mackey, 1961).

BM-249. Halling skeleton

2230 в.с. Portion of femur of Halling skeleton, ref. BM-48, British Mus. (Nat. Hist.). Coll. 1912 by W. H. Cook from contracted burial in deposits at Halling, Kent, England (51° 21' N Lat, 0° 27' E Long); subm. by K. P. Oakley, British Mus., (Nat. Hist.). Date is of 2nd, carefully pretreated sample (collagen only) from skeleton previously dated 4100 \pm 180, BM-168 (Barker and Mackey, 1968). Date was much younger than expected (Oakley, 1963a, b) but is now confirmed by this measurement (Oakley et al., 1967). Re-examination of flint artifacts showed that they could fit into either late Mesolithic or Neolithic context (G. de G. Sieveking, 1967, written commun.). (P)

BM-374. Paviland skeleton

Collagen from portions of left femur and tibiae of skeleton of socalled "Red Lady of Paviland," an ochre burial of young male found in Goat's Hole (Paviland Cave), Gower Peninsula, Glamorganshire, S Wales (51° 33' N Lat, 4° 15' W Long). Coll. 1823 by Dean William Buckland, and now in Geol. Collection, Univ. Mus., Oxford. Subm. by K. P. Oakley, British Mus. (Nat. Hist.). (See Buckland, 1823; Sollas, 1913; Garrod, 1926; North, 1942; Oakley, 1968).

Pinhole Cave

Charcoal from Pinhole Cave, Creswell Crags, near Worksop, Nottinghamshire, England (53° 15' 44" N Lat, 1° 12' 01" W Long). Coll. during period 1924-1933 by A. L. Armstrong (1928) and subm. by K. P. Oakley, British Mus. (Nat. Hist.). From series of very small quantities of charcoal from levels containing sequence of Palaeolithic industries claimed to be from Mousterian to "Developed Aurignacian" (now called Creswellian) of expected age from 20,000 to ca. 8000 yr B.P. Although from old excavation it was hoped that samples might prove suitable for dating stone industries in Pinhole Cave.

BM-437. Pinhole Cave 1

4120 ± 140 2170 в.с.

Bulked sample (nos. 7-9) representing so-called Mousterian/Upper Aurignacian boundary.

18.460 ± 340 16.510 в.с.

BM-438. Pinhole Cave 2

3910 ± 120 1960 в.с.

Bulked sample (nos. 1-4) representing so-called Developed Aurignacian (Creswellian). Comment: dates are quite impossible for Mousterian which cannot be less than 20,000 B.P. and are inherently unlikely for Creswellian which is probably not later than 10,000 B.P. and may be much older (McBurney, 1959). Long standing doubts about interpretation of stratigraphy of Pinhole Cave are confirmed by these dates since it is clear that samples measured do not relate to industries represented there.

Mildenhall

Two charcoal samples from Phillips II site, Mildenhall, Suffolk, England (52° 25' N Lat, 0° 31' E Long). Coll. 1962 by Col. T. C. Kelly, USAF: subm. by G. de G. Sieveking, British Mus.

BM-225. Phillips II (1)	3460 ± 150
Sample 1. (P)	1510 в.с.
RM.226 Phillins II (2)	3440 ± 150 1490 в.с.

Sample 2. (P) Comment (G. de G.S.): samples were from late glacial open air habitation site having Upper Palaeolithic flint industry. Expected age was ca. 11,000 yr B.P. Dates suggest that Palaeolithic site was disturbed during Bronze age occupation of adjacent site ca. 100 ft away.

E. Jordan

Jericho series

BM-250.

Charcoal from pre-pottery Neolithic A and B levels at Tell-es Sultan, Jericho, Jordan (31° 53' N Lat, 35° 27' E Long). Coll. 1960 and subm. by Miss K. M. Kenyon, British School of Archaeol., Jerusalem (Kenyon, 1960; Barker and Mackey, 1963; Stuckenrath, 1963).

10.300 ± 500 8350 в.с.

Ref. J.P.D. 303/34 (CS.1036) DI, Phase DDDi. Phase succeeding construction of pre-pottery Neolithic A (PPNA) defenses. See BM-105, 10,250 \pm 200 (Barker and Mackey, 1963), P-378, 9775 \pm 110 (Stuckenrath, 1963). (P)

BM-251. Jericho

Jericho

9390 ± 150 7440 в.с.

Ref. J.P.D. 404/14 (CS.1041) DII, Phase Gi. Stage VI (middle stage) of PPNA defenses. See P-379, 9655 \pm 84 (Stuckenrath, 1963). (P)

BM-252. Jericho

9320 ± 150 7370 в.с.

Ref. J.P.D. 201/38 DI, Phase TTii. Phase succeeding Stage VII (late stage) of PPNA defenses. (P)

BM-253. Jericho

Ref. J.P.D. 209/1a (CS.116) EI, II, V, Phase J. Mid stage of prepottery Neolithic B (PPNB) occupation. See BM-115, 9170 \pm 200 (Barker and Mackey, 1963), P-382, 8956 \pm 103 (Stuckenrath, 1963). (P)

F. Malta

Skorba series

Two charcoal samples from Ghar Dhalam phase at temple site of Li Skorba, Zebbiegh, NW Malta (35° 55' N Lat, 14° 23' 30" E Long). Coll. 1962 and subm. by D. H. Trump, Nat. Mus. Valletta, Malta (Trump, 1966; Barker and Mackey, 1968 and refs.).

BM-216. Skorba

5760 ± 200 3810 в.с.

 8710 ± 150

6760 в.с.

Level SK. AF5 at depth 50 to 70 cms in stratified deposit beside temple. Ghar Dhalam phase. (P)

BM-378. Skorba

6140 ± 160 4190 в.с.

Trench SB, Layer 4, 48 to 62 cms below surface, sealed below clay floor assoc. with pure Ghar Dhalam phase material. *Comment*: dates provide chronologic basis for Maltese sequence of which Ghar Dhalam phase is earliest. Both dates are consistent with other dates of Skorba series (Barker and Mackey, 1968).

Ife series

Charcoal from Ita Yemoo and Orun Oba Ado, Ife, W Nigeria (7° 28' N Lat, 4° 32' E Long). Coll. 1962 by Frank Willett, Northwestern Univ., Illinois.

G. Nigeria

BM-259. Ita Yemoo, Ife

790 ± 130 a.d. 1160

Trench XIII, Layer 19B (IY 2000) from upper part of pit fill beneath potsherd pavement No. 4 and Trench XIII, Layer 6B (IY 866) an occupation deposit on natural subsoil.

BM-261. Ita Yemoo, Ife

990 ± 130 a.d. 960

Trench XIV, Layer II (IY 1427) bottom layer of pit beneath potsherd pavement No. 1. *Comment*: BM-259 and BM-261 agree with estimated date of Classical period of Ife sculpture.

BM-262. Ita Yemoo, Ife

890 ± 130 a.d. 1060

Layer 2 (IY 949, IY 961, IY 963) in which terra cotta sculptures were found and which also represents abandonment of site. *Comment* (F.W.): date confirms estimate that Classical period of Ife art was of 12th to 14th centuries A.D. if not earlier, for sculptures in this deposit were already broken before site was abandoned. There can be no doubt that sculptures were made before 1st Europeans reached Benin.

BM-264. Orun Oba Ado, Ife	A.D. 990
Grave, Pit 6.	1200 + 120
BM-265. Orun Oba Ado, Ife	A.D. 560

Grave, Pit 11. Comment: dates are for site to which heads of Kings of Yoruba dynasty in Benin were returned for burial, estimated to be ca. 1000 to 1300 A.D.

H. West Pakistan

Loebanr I

Charred human bone fragments from pre-Buddhist graves at Loebanr, Swat, W Pakistan (34° 45' N Lat, 72° 30' E Long). Coll. 1962 by Dr. Ch. Silvi-Antonini; subm. by Prof. G. Genna, Inst. of Anthropol., Univ. of Rome. Expected age: 1200 to 300 B.C. (Antonini and Tucci, 1963).

BM-195. Loebanr I, Sample 1 Ref. L, IT, 54. (P)	1030 в.с.
BM-196. Loebanr I, Sample 2 Ref. L, IT, 61. (P)	2850 ± 150 900 в.с.

I. Yugoslavia

BM-379. Lepenski Vir

6900 ± 150 4950 в.с.

 960 ± 130

Charcoal (Quercus sp.) from floor of House 37, 1 of series of trapezoidal houses in pre-Neolithic horizon succeeded by Starcevo Neolithic level at Lepenski Vir on Yugoslav bank of Danube, NW of Donje Milanovac, N Serbia (44° 31' N Lat, 22° 02' E Long). Coll. 1967 by D. Srejović; subm. by J. G. Nandris, Univ. of London, Inst. of Archaeol. *Comment*: date agrees well with expected age and is consistent with other dates for 1st (Starcevo) Neolithic stratified above this level (Nandrus, 1968; Srejović, 1968a, b).

References

Date lists:	
Date lists: Arizona IV British Museum I British Museum III British Museum IV British Museum V Groningen IV Groningen V Lamont IV NPL I Pennsylvania VI	Damon, Long, and Sigalove, 1963 Barker and Mackey, 1959 Barker and Mackey, 1961 Barker and Mackey, 1963 Barker and Mackey, 1968 Vogel and Waterbolk, 1963 Vogel and Waterbolk, 1964 Broecker and Kulp, 1957 Callow, Baker, and Pritchard, 1963 Stuckenrath, 1963 Bagger Fergusson and Libby 1965
UCLA IV UCLA VI	Berger and Libby, 1967

 Antonini, C. S. and Tucci, G., 1963, East and West: v. 14 (new series), no. 1-2, p. 13-28.
 Armstrong, A. L., 1928-1929, Excavations in Pin Hole Caves, Creswell Crags, Derbyshire: Prehist. Soc. Proc., East Anglia, v. 6, p. 330-334. Ashbee, P., 1958, The Fussell's Lodge Long Barrow: Antiquity, v. 32, p. 106-111.

1964, The radiocarbon dating of the Fussell's Lodge Long Barrow: Antiquity, v. 38, p. 139-140.

Barker, Harold, 1953, Radiocarbon dating: Large-scale preparation of acetylene from organic material: Nature, v. 172, p. 631-632.

Barker, Harold and Mackey, C. J., 1959, British Museum natural radiocarbon measurements I: Am. Jour. Sci. Radiocarbon suppl., v. 1, p. 81-86.

1968, British Museum natural radiocarbon measurements V: Radiocarbon, v. 10, p. 1-7.

Barker, Harold, Burleigh, Richard, and Meeks, Nigel, 1969, New method for the combustion of samples for radiocarbon dating: Nature, v. 221, p. 49-50.

Berger, R., Fergusson, G. J., and Libby, W. F., 1965, UCLA radiocarbon dates IV: Radiocarbon, v. 7, p. 336-371.

Berger, R. and Libby, W. F., 1967, UCLA radiocarbon dates VI: Radiocarbon, v. 9, p. 477-504.

Broecker, W. S. and Kulp, J. L., 1957, Lamont natural radiocarbon measurements IV: Science, v. 126, p. 1324-1334.

Buckland, William, 1823, Reliquiae Diluvianae: London, p. 87.

Callow, W. J., Baker, M. J., and Pritchard, D. H., 1963, National Physical Laboratory radiocarbon measurements I: Radiocarbon, v. 5, p. 34-38.

- Damon, P. E., Long, Austin, and Sigalove, J. J., 1963, Arizona radiocarbon dates IV: Radiocarbon, v. 5, p. 283-301.
- Emery, W. B., 1938, The tomb of Hemaka: Excavations at Sakkara: 1938, Government Press, Cairo.

– 1939, Hor-Aha: Excavations at Sakkara, 1937-38: 1939, Government Press, Cairo.

1958, Great tombs of the first Dynasty III: Egypt Explor. Soc., London.

_____ 1961, Archaic Egypt: Harmondsworth, London, p. 1-269.

- 1965, Prelim. Rept. on the Excavations at N. Saqqara, 1964-5: Jour. Egyptian Archaeol., v. 51, p. 7.
- Evans, J. D., 1964a, Excavations in the Neolithic Mound of Knossos, 1958-60: London, Inst. Archaeol. Bull., no. 4, p. 34-60.

______ 1964b, Excavations in the Neolithic Settlement of Knossos, 1957-60: Athens, Ann. of British School, v. 59, p. 132-240.

- Garrod, D. A. E., 1926, The Upper Palaeolithic Age in Britain: Clarendon Press, Oxford, p. 49-64.
- Kenyon, K. M., 1960, Excavations at Jericho 1957-58: Palestine Explor. Quart., July-Dec., 1960.

Leroi-Gourhan, A. and Brezillon, M., 1966, Gallia Prehistoire, v. 9, pt. 1, p. 382.

- Longworth, I. H., 1961, Origins and development of the Collared Urn tradition in England and Wales: Prehist. Soc. Proc., v. 27, p. 263-306.
- Manby, T. G., 1963, The excavations of the Willerby Wold Long Barrow, East Riding of Yorkshire: Prehist. Soc. Proc., v. 29, p. 173-205.
- McBurney, C. B. M., 1959, Report on the first seasons fieldwork on British Upper Palaeolithic cave deposits: Prehist. Soc. Proc., v. 25, p. 260-269.

Nandris, J. G., 1968, Lepenski Vir: Science Jour., Jan. 1968, p. 64-70.

- Noakes, J. E., Kim, S. M., and Stipp, J. J., 1965, Chemical and counting advances in liquid scintillation age dating: Proc. Sixth Internatl. Conf. on Radiocarbon and Tritium Dating, Pullman, Washington. June 7-11, 1965, p. 68-92.
- North, F. J., 1942, Paviland Cave, the "Red Lady", the Deluge, and William Buckland: Annals Sci., v. 91, p. 128.

Oakley, K. P., 1963a, Note on the Antiquity of Halling Man, App. II, *in*: Kerney, M.P., 1963, Late-glacial deposits on the Chalk of south-east England: Royal Soc. (London) Philos. Trans. B., v. 246, no. 730, p. 250-251.

_____ 1963b, Mesolithic men in Britain: Prehist. Soc. Proc., v. 29, p. 426-427.

------- 1968, The date of the "Red Lady" of Paviland: Antiquity, v. 42, p. 306-307.

Oakley, K. P., Barker, H., and Sieveking, G. de G., 1967, The skeleton of Halling Man: Archaeologia Cantiana, v. 82, p. 218-220. Petrie, W. M. Flinders, 1914, Tarkhan II: British School of Archaeol. in Egypt, (London) p. 6.

Pollard, S. H., 1966, Neolithic and Dark Age settlements on High Peak, Sidmouth, Devon: Devon Archaeol. Explor. Soc. Proc., no. 23, p. 35-59.

Radley, J., 1966, A Bronze Age ring-work on Totley Moor and other Bronze Age ring-works in the Pennines: Archaeol. Jour., v. 123, p. 1-26.

Riley, D. N., 1966, An Early Bronze Age cairn on Harland Edge, Beeley Moor, Derbyshire: Derbs. Archaeol. Jour., v. 86, p. 31-53.

Smith, H. S., 1964, Egypt and C-14 dating: Antiquity, v. 38, p. 32-37.

Sollas, W. J., 1913, Paviland Cave: An Aurignacian station in Wales: Jour. Royal Anthropol. Inst., v. 43, p. 337-364.

Srejovic, Dragoslav, 1968a, Lepinski Vir 7,000 years ago: Illus. London News, Jan. 20, 1968.

______ 1968b, Lepenski Vir pottery and sculptures: Illus. London News, Feb. 3, 1968. Stuckenrath, Robert, Jr., 1963, Univ. of Pennsylvania radiocarbon dates VI: Radiocarbon, v. 5, p. 82-103.

Stuiver, Minze and Suess, H. E., 1966, On the relationship between radiocarbon dates and true sample ages: Radiocarbon, v. 8, p. 534-540.

Suess, H. E., 1967, Bristlecone pine calibration of the radiocarbon time scale from 4100 B.C. to 1500 B.C. (SM-87/52): Monaco symposium, Monaco, 2-10 March, 1967.

Trump, D. H., 1966, Skorba and the prehistory of Malta: Soc. of Antiquaries Research Rept. 22, p. 10, 48, 54.

Vogel, J. C. and Waterbolk, H. T., 1963, Groningen radiocarbon dates IV: Radiocarbon, v. 5, p. 163-202.

______ 1964, Groningen radiocarbon dates V: Radiocarbon, v. 6, 349-369.

Wainwright, G. J., 1967, The excavation of the henge monument at Durrington Walls, Wiltshire, 1966: Antiquaries Jour., v. 47, p. 166-184.

_____ 1968, Durrington Walls a ceremonial enclosure of the 2nd millennium B.C.: Antiquity, v. 42, no. 165, p. 20-26.