NICOTINIC ACETYLCHOLINE RECEPTOR $\alpha 4$ SUBUNIT GENE VARIATION ASSOCIATED WITH CHINESE HAN PATIENTS WITH ATTENTION DEFICIT HYPERACTIVITY DISORDER

J. Weidong¹, H. Xuezh², G. Tianyou³, Y. Chuang⁴, L. Qianqian⁵, X. Yong⁶, G. Lanting⁷

¹Shanghai Changning Mental Health Center/Bio-X Center Hospital of Shanghai Jiaotong University, Shanghai, ²Psychiatry, School of Medicine, Tsinghua University, Beijing, ³Department of Psychology, Normal College, Shenzhen University, Shenzhen, ⁴Psychiatry, Medical College of Wenzhou, Wenzhou, ⁵Psychiatry, West China Hospital of Sichuan University, Chengdu, ⁶School of Medicine of Tsinghua University, Beijing, ⁷West China Hospital of Sichuan University, Chengdu, China

Previous pharmacological, human genetical, and animal models have implicated the nicotinic acetylcholine receptor $\alpha 4$ subunit (CHRNA4) gene in the pathogenesis of ADHD. The objective of this study is to examine genetic association between single nucleotide polymorphisms (SNPs) in the CHRNA4 gene (rs2273502, rs1044396, rs1044397 and rs3827020 loci) and ADHD. Both case-control and family-based design were used in this study. Children aged 6 to 16 years were interviewed and assessed with the CBCL and CPRS-R to identify probands. No significant differences in frequency distribution of genotypes or alleles between the case and control groups were found. However, further haplotype analyses showed CCGG haplotype on risk for ADHD in 164 case-control sample and TDT analysis suggested that the allele C of rs2273502 over-transferred in 98 ADHD parent-offspring trios. Our findings suggest that CHRNA4 gene may play a role in the pathogenesis of ADHD, and further work is necessary to replicate and confirm what role the CHRNA4 gene may play in the etiology and pathogenesis of ADHD in large independent samples.