THE GRAVITY-DARKENING OF THE MAIN-SEQUENCE COMPONENTS OF SPECTRAL TYPES A, F AND G IN DETACHED CLOSE BINARY SYSTEMS

Y. NAKAMURA

Department of Science Education

Fukushima University, Fukushima 960-12, Japan

M. KITAMURA

National Astronomical Observatory

Mitaka, Tokyo 181, Japan

(Not reviewed)

Abstract. From analysis of the photometric ellipticity effect in seven well-understood detached close binary systems, empirical values of the exponent α of gravity-darkening have been practically deduced for eleven main-sequence components of spectral types A, F and G which should cover the range of structural change (from radiative to convective) in stellar atmospheres. The result indicate that values of the exponent diminish gradually with decreasing effective temperatures from $\alpha \sim 1.0$ for radiative atmospheres with $T > 8500$ K to $\alpha = 0.2 \sim 0.3$ for convective atmospheres with $T < 6500$ K, in spite of some uncertainty in the reflection correction process.

Fig.1. Empirical values of the exponent α of gravity-darkening deduced for main-sequence components of detached close binary systems. 1, 2 in the parentheses represent the primary and the secondary component, respectively. o expresses α -values deduced with albedo $E = 1.0$ assumed for reflection correction while • with $E = 0.5$.