

A 2.5-5 μm spectroscopic study of hard X-ray selected AGNs with AKARI

A. Castro¹, T. Miyaji^{1,2}, M. Shirahata³, S. Oyabu⁴, D. Clark¹,
K. Ichikawa⁵, M. Imanishi⁶, T. Nakagawa³ and Y. Ueda⁵

¹Instituto de Astronomía - UNAM, Ensenada, Baja California, México,
(mailing address: PO Box 439027, San Diego, CA 92143-9027, USA)
email: acastro@astroesen.unam.mx

²University of California, San Diego, Center for Astrophysics and Space Sciences, USA

³Japan Aerospace Exploration Agency (JAXA), Sagami-hara, Japan

⁴Graduate School of Science, Nagoya University, Nagoya, Japan

⁵Department of Astronomy, Faculty of Science, Kyoto University, Kyoto, Japan

⁶Subaru Telescope, National Astronomical Observatory of Japan, Hilo, USA

Abstract. We explore the relationships between the 3.3 μm polycyclic aromatic hydrocarbon (PAH) feature and active galactic nucleus (AGN) properties of a sample of 54 hard X-ray selected bright AGNs, including both Seyfert 1 and Seyfert 2 type objects, using the InfraRed Camera (IRC) on board the infrared astronomical satellite *AKARI*. The sample is selected from the 9-month *Swift*/BAT survey in the 14-195 keV band and all of them have measured X-ray spectra at $E \lesssim 10$ keV. These X-ray spectra provide measurements of the neutral hydrogen column density (N_{H}) towards the AGNs. We use the 3.3 μm PAH luminosity ($L_{3.3\mu\text{m}}$) as a proxy for star formation activity and hard X-ray luminosity ($L_{14-195\text{keV}}$) as an indicator of the AGN activity. We searched for possible difference of star-formation activity between type 1 (un-absorbed) and type 2 (absorbed) AGNs. Our regression analysis of $\log L_{14-195\text{keV}}$ versus $\log L_{3.3\mu\text{m}}$ shows a positive correlation and the slope seems steeper for type 1/unobscured AGNs than that of type 2/obscured AGNs. The same trend has been found for the $\log(L_{14-195\text{keV}}/M_{\text{BH}})$ versus $\log(L_{3.3\mu\text{m}}/M_{\text{BH}})$ correlation. Our analysis show that the circum-nuclear star-formation is more enhanced in type 2/absorbed AGNs than type 1/un-absorbed AGNs for low X-ray luminosity/low Eddington ratio AGNs.

Keywords. galaxies: active — X-rays: galaxies

1. Description of the research

Infrared spectroscopy is a powerful tool for the diagnostic buried AGNs, because the effect of dust extinction is relatively small (Imanishi *et al.* 2010). It allows to study sources in which both an AGN and a starburst (SB) are present. Very hard X-ray surveys such as available with *Swift* Burst Alert Telescope (BAT), provide efficient way of constructing a fair sample of AGNs including heavily obscured ones. The infrared 2.5-5 μm spectra available with the Infrared Camera (IRC) grism mode on board *AKARI* provide us with rich pieces of information with emission/absorption lines such as the 3.3 μm polycyclic and the broad 3.1 μm H₂O ice, 3.4 μm bare carbonaceous dust, 4.26 μm CO₂ and 4.67 μm CO absorptions. In this work, we use the 3.3 μm PAH feature as a proxy for star formation and explore the relation between AGN activity, type and circum-nuclear star-formation.




Figure 1. (a) $\log(L_{14-195\text{keV}})$ versus $\log(L_{3.3\mu\text{m}})$ and (b) $\log(L_{14-195\text{keV}}/M_{\text{BH}})$ versus $\log(L_{3.3\mu\text{m}}/M_{\text{BH}})$ relations. Open red circles are used for Optical type 1 objects and filled black circles for Optical type 2 objects. Arrows are for upper-limits. The best-fit lines from linear regressions with upper limits are also shown. (c) Distribution of $\log(L_{3.3\mu\text{m}})$.

2. Results and Conclusions

We use a linear regression analysis using the E-M method available within ASURV package (Isobe *et al.* 1986) among observed quantities. We have studied the dependencies between the luminosity of the PAH at $\lambda_{rest} = 3.3 \mu\text{m}$ ($L_{3.3\mu\text{m}}$) emission line to the X-ray luminosity in the 14-195 keV band ($L_{14-195\text{keV}}$) for the entire sample:

$$\log(L_{3.3\mu\text{m}}) = (0.4 \pm 0.1)\{\log(L_{14-195\text{keV}}) - 43.6\} + (40.2 \pm 0.1) \quad (2.1)$$

$$\log(L_{3.3\mu\text{m}}/M_{\text{BH}}) = (0.7 \pm 0.2)\{\log(L_{14-195\text{keV}}/M_{\text{BH}}) - 35.4\} + (31.9 \pm 0.1) \quad (2.2)$$

We have made separate regressions for samples divided in optical classification (type 1 vs. type 2 AGNs) as well as X-ray absorption divided samples ($\log N_{\text{H}}[\text{cm}^{-2}] < 22$ [unabsorbed] or > 22 [absorbed]). Our analysis show that significant correlations have been found between $3.3 \mu\text{m}$ PAH luminosity in the circumnuclear and the 14-195 keV luminosity for type 1/unabsorbed AGNs, while the correlation is weaker for the type 2/absorbed AGNs.

We divided the sample into high and low $L_{14-195\text{keV}}$ (or $L_{14-195\text{keV}}/M_{\text{BH}}$) and compared the mean $\log L_{3.3\mu\text{m}}$ (or $L_{3.3\mu\text{m}}/M_{\text{BH}}$) values of the type 1/unabsorbed and type 2 absorbed AGNs. Our analysis show that the circum-nuclear star-formation is more enhanced in type 2/absorbed AGNs than type 1/un-absorbed AGNs for low X-ray luminosity/low Eddington ratio AGNs, while there is no significant dependence of star-formation activities on the AGN type in the high X-ray luminosities/Eddington ratios. We note that the underlying assumption of applying the EM-algorithm is that the objects are normally distributed around the regression line.

This work is supported by UNAM PAPIIT-IN 104113 and CONACyT Grant 179662.

References

- Imanishi, M., Nakagawa, T., Shirahata, M., & Ohya, Y., Onaka 2010, *ApJ*, 721, 1233
 Isobe, T., Feigelson, E. D., & Nelson, P. I. 1986, *ApJ*, 306, 490