



## Diet and well-being of 1st year Medical Interns during hospital placement

N.M. Simm<sup>1,2,3</sup>, G.M. Williams<sup>2,3</sup>, A. Thorpe<sup>4</sup>, E.C. Hoedt<sup>1,2,3</sup>, S.J. Caban<sup>1,2,3</sup>, K.L. Tooley<sup>5</sup>, G.W. Tyson<sup>6</sup>, S.D. Brown<sup>4</sup>, A. Eidels<sup>4</sup>, S. Keely<sup>1,2,3</sup>, N.J. Talley<sup>2,3,7</sup> and K. Duncanson<sup>2,3,7</sup>

<sup>1</sup>School of Biomedical Sciences & Pharmacy, College of Health, Medicine and Wellbeing, University of Newcastle, NSW, Australia.

<sup>2</sup>NHMRC Centre for Research Excellence in Digestive Health, University of Newcastle, Newcastle, NSW, Australia

<sup>3</sup>Hunter Medical Research Institute, New Lambton Heights, NSW, Australia

<sup>4</sup>School of Psychology, University of Newcastle, Callaghan, NSW, Australia

<sup>5</sup>Human Systems Performance, DST Group, Department of Defence, Edinburgh, Adelaide, SA, Australia

<sup>6</sup>Centre for Microbiome Research, School of Biomedical Sciences, Queensland University of Technology (QUT), Translational Research Institute, Brisbane Australia

<sup>7</sup>School of Medicine & Public Health, College of Health, Medicine and Wellbeing, University of Newcastle, NSW, Australia

Hospital placement is essential training for medical interns, involving shift work and high-pressure environments. This can increase physiological and psychological stress, which may be mediated by metabolites of microbial digestion<sup>(1)</sup>. Nutrients of interest include those accessible to microbial digestion and associated with altered signalling within the microbiota-gut-brain axis (MGBA)<sup>(1)</sup>. Fibre is fermented by gut microbes to produce short-chain fatty acids<sup>(2)</sup> and is associated with improved psychological outcomes<sup>(3)</sup>. Tryptophan, a precursor to gut-derived serotonin<sup>(2)</sup>, has been negatively associated with anxiety<sup>(4)</sup>. Processed foods contain food additives, excess sugars, and saturated fats that may disrupt gut homeostasis<sup>(1)</sup> and impact psychological well-being<sup>(4)</sup>. Lastly, total energy intake may determine the level of substrate available for microbial fermentation<sup>(2)</sup>. Therefore, this research explores how microbiota-accessible food components interact with physical and psychological well-being in a cohort of medical interns undertaking their first-year of hospital placement. Participants were healthy medical interns, during first-year hospital placement (n = 21) from the Hunter New England Local Health District, NSW, Australia. Participants completed diet and wellbeing surveys at baseline and every 2 months over a 10-month period. 24-hour diet diaries were self-recorded from participants using a mobile application (Easy Diet Diary) and analysed using AusNut and the NOVA classification system of ultra-processed foods (ULP). Wellbeing surveys include depression, anxiety, stress scale (DASS), and PROMIS survey for mental (M), physical (P), and sleep well-being. Current data represents an 'in-progress' of the longitudinal data collection. This study utilised Spearman correlation and Tukey's post hoc test for mixed methods analysis. From baseline to timepoint 3 (T3, 4 months) daily energy intake was consistent with cohort estimated energy requirements (EER). However, consumption ranged from 37% to 167% of EER, indicating a large variation of intakes. Energy consumed from ULP ranged from 30% to 34% (p = 0.6875). Baseline tryptophan intake ( $\bar{x}$  = 1139mg) was within the suggested target, whilst fibre intake ( $\bar{x}$  = 23g) was below the recommended intake. Neither saw significant changes from baseline to T3. Fibre intake was positively correlated with mental and physical well-being at baseline ( $\bar{x}$  = 23.1g, M: r = 0.474, p = 0.04, P: r = 0.608, p = 0.007), and timepoint 2 ( $\bar{x}$  = 31.5g, M: r = 0.647, p = 0.026, P: r = 0.780, p = 0.004) but not at T3. In addition, baseline consumption of sugar ( $\bar{x}$  = 18g) and poly-unsaturated fats ( $\bar{x}$  = 15g) were both negatively correlated with mental and physical well-being. Overall, no significant dietary changes were evident from baseline to mid-year collection in a first-year medical intern cohort during hospital placements. Fibre was significantly associated with mental and physical well-being, building on current understanding of fibre's role in the MGBA. Planned metabolite analysis will explore the mechanisms of proposed microbiome-accessible nutrients alongside diet, well-being, and microbiota data. Findings from this study will identify how diet-microbiome interactions change under stress, with wider positive implications on intense workplace environments with the aim to preserve individual wellbeing.

**Keywords:** gut microbiota; mental well-being; microbiome; fibre

### Ethics Declaration

Yes

### Financial Support

This work was supported by the Defence Science and Technology (DST) Group: Next Generation Technology Funding (NGTF) (grant number: G2100173).

### References

1. Foster JA, Rinaman L & Cryan JF (2017) *Neurobiol Stress* 7, 124–136.
2. Debnatha N, Kumarb R *et al.* (2021) *Biotechnol Genet Eng Rev* 37(2), 105–153.
3. Saghafian S, Hajishafiee M *et al.* (2023) *Nutr Neurosci* 26(2), 108–126.