

Abstract prepared by Arthur F. Ramos and Ruy J. G. B. de Queiroz.

E-mail: afr@cin.ufpe.br

MARCUS VINÍCIUS MIDENA RAMOS, *Formalization of Context-Free Language Theory*, Universidade Federal de Pernambuco, Brazil, 2016. Supervised by Ruy J. G. B. de Queiroz. MSC: 03B35, 03D05, 03F65, 68Q42, 68Q45, 68T15. Keywords: proof assistants, Coq, formalization, language theory, context-free languages, context-free grammars.

Abstract

Proof assistants are software-based tools that are used in the mechanization of proof construction and validation in mathematics and computer science, and also in certified program development. Different such tools are being increasingly used in order to accelerate and simplify proof checking, and the Coq proof assistant is one of the most well known and used in large-scale projects. Language and automata theory is a well-established area of mathematics, relevant to computer science foundations and information technology. In particular, context-free language theory is of fundamental importance in the analysis, design, and implementation of computer programming languages. This work describes a formalization effort, using the Coq proof assistant, of fundamental results of the classical theory of context-free grammars and languages. These include closure properties (union, concatenation, and Kleene star), grammar simplification (elimination of useless symbols, inaccessible symbols, empty rules, and unit rules), the existence of a Chomsky Normal Form for context-free grammars and the Pumping Lemma for context-free languages. The result is an important set of libraries covering the main results of context-free language theory, with more than 500 lemmas and theorems fully proved and checked. As it turns out, this is a comprehensive formalization of the classical context-free language theory in the Coq proof assistant and includes the formalization of the Pumping Lemma for context-free languages. The perspectives for the further development of this work are diverse and can be grouped in three different areas: inclusion of new devices and results, code extraction, and general enhancements of its libraries.

Abstract prepared by Marcus Vinícius Midena Ramos.

E-mail: mvmramos@gmail.com

URL: <http://marcusramos.com.br/univasf/tese.pdf>

DOUGLAS ULRICH, *Some Applications of Set Theory to Model Theory*, University of Maryland, College Park, USA, 2018. Supervised by Michael C. Laskowski. MSC: 03C55. Keywords: Keisler's order, Borel complexity.

Abstract

We investigate set-theoretic dividing lines in model theory. In particular, we are interested in Keisler's order and Borel complexity.

Keisler's order \leq is a pre-order on complete countable theories T , measuring the saturation of ultrapowers of models of T . In Chapter 3, we present a self-contained survey on Keisler's order, cast in terms of full Boolean-valued models and pseudosaturation. The cornerstone of our development is a compactness theorem for full Boolean-valued models. As an application, we show that if T is a complete countable theory and \mathcal{B} is a complete Boolean algebra, then λ^+ -saturated \mathcal{B} -valued models of T exist. Moreover, if \mathcal{U} is an ultrafilter on T and \mathbf{M} is a λ^+ -saturated \mathbf{M} -valued model of T , then whether or not \mathbf{M}/\mathcal{U} is λ^+ -saturated just depends on \mathcal{U} and T ; we say that \mathcal{U} λ^+ -saturates T in this case. We show that Keisler's order can be formulated as follows: $T_0 \leq T_1$ if and only if for every cardinal λ , for every complete Boolean algebra \mathcal{B} with the λ^+ -c.c., and for every ultrafilter \mathcal{U} on \mathcal{B} , if \mathcal{U} λ^+ -saturates T_1 , then \mathcal{U} λ^+ -saturates T_0 . We also prove that if \mathcal{B} is a complete Boolean algebra with an antichain of size λ , then there is a λ^+ -good ultrafilter on \mathcal{B} ; conversely, if \mathcal{B} is a complete Boolean algebra with the λ -c.c., then no nonprincipal ultrafilter on \mathcal{B} λ^+ -saturates any unsimple theory, and no \aleph_1 -incomplete ultrafilter on \mathcal{B} λ^+ -saturates any nonlow theory. Finally, we