of functioning at the end of an episode, 9. less symptoms between episodes.

Conclusion: The acute transient psychoses as defined in ICD-10 are good prognosis psychotic disorders differing significantly from schizophrenia. A syndrome-shift to schizophrenia during the long term course is extremely rare. The above findings support the assumption of independence of ATP from schizophrenia.

FC36-6

THE COCHRANE SCHIZOPHRENIA GROUP

L. Roberts¹*, C. Adams¹. ¹Cochrane Schizophrenia Group, Oxford OX2 7LG, UK

Registered in June 1994, the Cochrane Schizophrenia Group (CSG) was the first mental health group within the Cochrane Collaboration, and is a global network with collaborators on every continent. The CSG is concerned with the evaluation of care relevant to those with non-affective, functional, psychotic illnesses.

Aims of the Cochrane Schizophrenia Group:

□ To identify every randomized controlled trial relevant to the care of those with schizophrenia and other related disorders by systematically searching journals and electronic databases

- □ To regularly update the CSG Register of Trials (currently containing c.3,500 trials)
- □ To produce and maintain methodologically sound, clearly written reviews of RCTs relevant to the scope of the Group
- □ To make these reviews accessible to clinicians, researchers, managers, policy makers and recipients of care by ensuring that they are presented in an unintimidating and easily understandable format
- □ To continue to disseminate this information via publication on the Cochrane Library.

We currently (Dec 1997) have 20 reviews and 12 protocols published on the Cochrane Library. These have already been effective in influencing policy and care of those with schizophrenia.

The Cochrane Library will be demonstrated and the CSG's future agenda outlined.

FC36-7

PATIENT AND RELATIVE EDUCATION IN COMMUNITY PSYCHIATRY; A RANDOMIZED TRIAL REGARDING IT'S USEFULNESS

L.B. Merinder¹*, A.G. Viuff, H.D. Laugesen, K. Clemmensen, S. Misfelt, B. Espensen. ¹Department of Psychiatric Demography, Psychiatric Hospital, DK-8240 Risskov, Denmark

Psychoeducation meets increasing accept as a part of the treatment program for schizophrenia.

But do the patients accept psychoeducation as readily as the relatives and the professionals? Are the effects of psychoeducation aimed at patients useful measured by changes in satisfaction with services, knowledge, Expressed emotion scores and compliance.

In relation to a randomized study of a short psychoeducational program for patients with schizophrenia and their relatives in two Community Psychiatric Centres in Aarhus and Viborg, an analysis of eligibility of patients for this kind of program was made.

The representativity of the participating group was analysed through comparison of the characteristics of this group with the group of refusers and with The Aarhus population of patients of patients registered in the National Psychiatric Register.

An analysis of the effects of a short 8 session psychoeducational program for patients and relatives on relevant outcome measures is described. It was shown that changes in specific dimensions of satisfaction with services was found both for patients and relatives and that there was a corresponding increase of knowledge. The selection of the study group made it impossible to detect changes in compliance and insight into psychosis due to ceiling effects. There were no significant effects in the more "hard" outcome measures relapse or psychopathology measured by BPRS.

The results indicate that a short psychoeducational program like this can be seen as an introduction to the rehabilitative work with patients with schizophrenia facilitating other and more pervasive rehabilititative efforts rather than an effective treatment modality in itself.

S37. Research and development in psychiatry in Europe

Chairs: AH Ghodse (UK), P Bech (DK)

S37-1

MENTAL HEALTH RESEARCH & POLICY IN EUROPE

H. Ghodse. Institute of Psychiatry, London, UK

Science and public policy in many countries are as far apart as God and atheism. Yet, both are in need of each other. The challenge of each other's diversity combined with deep-seated suspicions have, in many societies, separated the two to a point of nonreconciliation. This is particularly marked at times of recession and political change and affects all areas of science and policy, particularly in the field of health, and significantly so in mental health sciences. Local and national policies may sometimes conflict with pursuing scientific enquiries, while international conflicts, pressures and compromises also undermine collaboration between scientists. It is equally true that the results of sophisticated laboratory experiments may have far less effect than a non-scientific bilateral/multilateral formal agreement or deal between the political leaders for example. Despite all the diversities of mental health and their related problems in the world, there are certain universal and psycho-biological facts which cross the boundaries of cultures and nations. Psychiatry as well as dealing with its immediate problems, should also be looking ahead. There can be no doubt that over the next century attention will increasingly have to be turned on world-wide scale to the medical implications of human behaviour. Thus, psychiatry can be seen as a meeting place between biological sciences on the one hand, and a variety of social-behavioural sciences on the other and it is impossible to conceive of fruitful research which neglects either of these complementary groups of discipline. The European Professors of Psychiatry Association (Euro PoP) is a forum for academics of Europe in the field of psychiatry to profess the speciality in order to advance research and teaching in mental health and toward the well being of individuals as well as the population. It could become a pathway to bridge the gap between science and policy, not only within individual nations in Europe, but also across cultures by discussion, debate and collaboration.