

Hereditarily frequently hypercyclic operators and disjoint frequent hypercyclicity

FRÉDÉRIC BAYART[†], SOPHIE GRIVAUX[‡], ÉTIENNE MATHERON[§] and
QUENTIN MENET[¶]

[†] *Laboratoire de Mathématiques Blaise Pascal UMR 6620 CNRS,
Université Clermont Auvergne, Campus universitaire des Cézeaux,
3 place Vasarely, 63178 Aubière Cedex, France
(e-mail: frederic.bayart@uca.fr)*

[‡] *Univ. Lille, CNRS, UMR 8524 - Laboratoire Paul Painlevé, F-59000 Lille, France
(e-mail: sophie.grivaux@univ-lille.fr)*

[§] *Univ. Artois, UR 2462 - Laboratoire de Mathématiques de Lens (LML) F-62300 Lens,
France
(e-mail: etienne.matheron@univ-artois.fr)*

[¶] *Service d'Analyse fonctionnelle, Département de Mathématique, Université de Mons,
Place du Parc 20, 7000 Mons, Belgium
(e-mail: quentin.menet@umons.ac.be)*

(Received 17 September 2024 and accepted in revised form 6 March 2025)

Abstract. We introduce and study the notion of hereditary frequent hypercyclicity, which is a reinforcement of the well-known concept of frequent hypercyclicity. This notion is useful for the study of the dynamical properties of direct sums of operators; in particular, a basic observation is that the direct sum of a hereditarily frequently hypercyclic operator with any frequently hypercyclic operator is frequently hypercyclic. Among other results, we show that operators satisfying the frequent hypercyclicity criterion are hereditarily frequently hypercyclic, as well as a large class of operators whose unimodular eigenvectors are spanning with respect to the Lebesgue measure. However, we exhibit two frequently hypercyclic weighted shifts $B_w, B_{w'}$ on $c_0(\mathbb{Z}_+)$ whose direct sum $B_w \oplus B_{w'}$ is not \mathcal{U} -frequently hypercyclic (so that neither of them is hereditarily frequently hypercyclic), and we construct a C -type operator on $\ell_p(\mathbb{Z}_+)$, $1 \leq p < \infty$, which is frequently hypercyclic but not hereditarily frequently hypercyclic. We also solve several problems concerning disjoint frequent hypercyclicity: we show that for every $N \in \mathbb{N}$, any disjoint frequently hypercyclic N -tuple of operators (T_1, \dots, T_N) can be extended to a disjoint frequently hypercyclic $(N + 1)$ -tuple $(T_1, \dots, T_N, T_{N+1})$ as soon as the underlying space supports a hereditarily frequently hypercyclic operator; we construct

a disjoint frequently hypercyclic pair which is not densely disjoint hypercyclic; and we show that the pair (D, τ_a) is disjoint frequently hypercyclic, where D is the derivation operator acting on the space of entire functions and τ_a is the operator of translation by $a \in \mathbb{C} \setminus \{0\}$. Part of our results are in fact obtained in the general setting of Furstenberg families.

Key words: frequent hypercyclicity, Furstenberg families, countable Lebesgue spectrum, disjointness

2020 Mathematics Subject Classification: 47A16, 37A30 (Primary); 37B20 (Secondary)

1. Introduction

This paper is devoted to two different topics, both pertaining to the study of the dynamics of linear operators. First, motivated by some questions regarding the behaviour of direct sums of operators, we introduce a new dynamical property of continuous linear operators on Banach or Fréchet spaces, which appears to be a very natural strengthening of the classical notion of frequent hypercyclicity; we call it *hereditary frequent hypercyclicity*. We believe that this is an interesting notion and we study it in some detail. Second, we address some questions concerning *disjoint frequent hypercyclicity*—also called *diagonal* frequent hypercyclicity. One notable connection between these two topics is that hereditarily frequently hypercyclic operators can be used to extend diagonally frequently hypercyclic tuples (see below).

In what follows, the letter X denotes an infinite-dimensional Polish topological vector space and $\mathfrak{L}(X)$ is the space of continuous linear operators on X . Recall that an operator $T \in \mathfrak{L}(X)$ is said to be hypercyclic if it has a dense orbit, that is, there exists $x \in X$ such that $\{T^n x : n \geq 0\}$ is dense in X ; equivalently, for each non-empty open set $V \subset X$, the ‘visit set’ $\mathcal{N}_T(x, V) := \{n \in \mathbb{N} : T^n x \in V\}$ is infinite. A much stronger property, introduced in [5], is *frequent hypercyclicity*: the operator T is frequently hypercyclic if there exists $x \in X$ such that for each non-empty open set $V \subset X$, the set $\mathcal{N}_T(x, V)$ has positive lower density. We refer the reader to [7, 33] for an in-depth presentation of various aspects of linear dynamics.

More recently, quantitative notions of hypercyclicity have begun to be studied in a very general framework [11, 17, 24]. Let \mathcal{F} be a *Furstenberg family*, that is, a family of non-empty subsets of \mathbb{N} which is hereditary upwards (if $A' \supset A \in \mathcal{F}$, then $A' \in \mathcal{F}$). Following [11], we say that an operator $T \in \mathfrak{L}(X)$ is \mathcal{F} -*hypercyclic* if there exists $x \in X$, called an \mathcal{F} -hypercyclic vector for T , such that for each non-empty open set $V \subset X$, the set $\mathcal{N}_T(x, V)$ belongs to \mathcal{F} . Thus, hypercyclicity corresponds to the family of all infinite subsets of \mathbb{N} , and frequent hypercyclicity corresponds to the family of sets with positive lower density. The set of \mathcal{F} -hypercyclic vectors for T will be denoted by $\mathcal{F}\text{-HC}(T)$. However, in accordance with a well-established notation, we write $\text{HC}(T)$ in the hypercyclic case and $\text{FHC}(T)$ in the frequently hypercyclic case. Also, when \mathcal{F} is the family of all subsets of \mathbb{N} with positive upper density, we say that T is \mathcal{U} -*frequently hypercyclic* and we write $\text{UFHC}(T)$.

The starting point of the paper is the following question.

Question 1.1. Let $T_1 \in \mathcal{L}(X_1)$ and $T_2 \in \mathcal{L}(X_2)$ be two frequently hypercyclic operators; is it true that $T_1 \oplus T_2$ is frequently hypercyclic?

This question seems to have been considered for the first time in [31, §8] and appears as a natural variant of the following well-known open problem in linear dynamics [5]: if T is a frequently hypercyclic operator, is it true that $T \oplus T$ is frequently hypercyclic? Question 1.1 makes sense for \mathcal{F} -hypercyclicity as well; and in the especially interesting case $T_1 = T_2$, the answer is known for the family of all infinite subsets of \mathbb{N} and for the family of sets with positive *upper* density. Indeed, a famous example from [23] shows that hypercyclicity of T does not imply that of $T \oplus T$, whereas it is proved in [24] that \mathcal{U} -frequent hypercyclicity of T does imply that of $T \oplus T$.

Given $T_1 \in \mathcal{L}(X_1)$ and $T_2 \in \mathcal{L}(X_2)$, two frequently hypercyclic operators, a natural way to show that $T_1 \oplus T_2$ is frequently hypercyclic would be the following. Let $(V_i)_{i \in \mathbb{N}}$ be a countable basis of open sets for $X_1 \times X_2$ and assume that each V_i has the form $V_i = V_{i,1} \times V_{i,2}$, where $V_{i,1}$ is open in X_1 and $V_{i,2}$ is open in X_2 . Pick a frequently hypercyclic vector $x_1 \in X_1$ for T_1 . Then, for any $i \in \mathbb{N}$, there exists a set $A_i \subset \mathbb{N}$ with positive lower density such that $T_1^n x_1 \in V_{i,1}$ for all $n \in A_i$. We would be done if we were able to find a vector $x_2 \in X_2$ with the following property: for every $i \in \mathbb{N}$, there exists a set B_i with positive lower density *and contained in* A_i (this is the important point, which cannot be guaranteed if x_2 is simply assumed to be frequently hypercyclic for T_2) such that $T_2^n x_2 \in V_{i,2}$ for all $n \in B_i$. This leads to the following definition.

Definition 1.2. Let $\mathcal{F} \subset 2^{\mathbb{N}}$ be a Furstenberg family. We say that an operator $T \in \mathcal{L}(X)$ is *hereditarily \mathcal{F} -hypercyclic* if, for any countable family $(V_i)_{i \in I}$ of non-empty open subsets of X and any family $(A_i)_{i \in I} \subset \mathcal{F}$ indexed by the same countable set I , there exists a vector $x \in X$ such that $\mathcal{N}_T(x, V_i) \cap A_i \in \mathcal{F}$ for every $i \in I$; in other words, for each $i \in I$, there is a set $B_i \in \mathcal{F}$ such that $B_i \subset A_i$ and $T^n x \in V_i$ for all $n \in B_i$.

When \mathcal{F} is the family of sets with positive lower density, we say (of course) that the operator T is *hereditarily frequently hypercyclic*; and likewise for \mathcal{U} -frequent hypercyclicity. By the above discussion, we get the following observation.

Observation 1.3. Let $\mathcal{F} \subset 2^{\mathbb{N}}$ be a Furstenberg family. If T_1, T_2 are two \mathcal{F} -hypercyclic operators and at least one of them is hereditarily \mathcal{F} -hypercyclic, then $T_1 \oplus T_2$ is \mathcal{F} -hypercyclic.

Note that when \mathcal{F} is the family of all infinite subsets of \mathbb{N} , hereditary \mathcal{F} -hypercyclicity is equivalent to *topological mixing*; see §9.1 for the (easy) proof. So, Observation 1.3 implies in particular that the direct sum of a hypercyclic operator with a topologically mixing operator is hypercyclic; which is of course well known.

We also point out that—perhaps surprisingly—hereditary \mathcal{F} -hypercyclicity automatically implies *dense* hereditary \mathcal{F} -hypercyclicity: given (A_i) and (V_i) as in Definition 1.2 above, there is a dense set of vectors $x \in X$ satisfying the required property; see Proposition 6.4 below.

Having introduced a definition, we are immediately faced with some obvious questions.

• *Are there any hereditarily frequently hypercyclic operators?* We answer this in the affirmative by two different methods. Indeed, there are two ‘standard’ ways of proving that an operator is frequently hypercyclic: either by showing that it satisfies the so-called frequent hypercyclicity criterion (see [7] or [33]) or by exhibiting a large supply of eigenvectors associated with unimodular eigenvalues (see for example [5] or [8]). It turns out that in both cases, one gets in fact hereditary frequent hypercyclicity (Theorems 2.1, 3.1) or hereditary \mathcal{U} -frequent hypercyclicity (Theorem 3.24).

• *Is hereditary frequent hypercyclicity a new notion?* In other words, are there any frequently hypercyclic operators which are not hereditarily frequently hypercyclic? The answer is ‘Yes’, and we prove this in two ways. On the one hand, we construct two frequently hypercyclic weighted shifts B_w and $B_{w'}$ on $c_0(\mathbb{Z}_+)$ such that $B_w \oplus B_{w'}$ is not \mathcal{U} -frequently hypercyclic (Theorem 4.2), so that neither of them can be hereditarily frequently hypercyclic by Observation 1.3. This also gives a strong negative answer to the $T_1 \oplus T_2$ frequent hypercyclicity problem of Question 1.1. On the other hand, with the terminology of [31], we construct a C-type operator on $\ell_p(\mathbb{Z}_+)$, $1 \leq p < \infty$, which is frequently hypercyclic but not hereditarily frequently hypercyclic (Theorem 5.2).

• *What are hereditarily frequently hypercyclic operators good for?* We will use them in the context of ‘disjoint hypercyclicity’. The notion of disjointness in linear dynamics was introduced independently in [10, 14]. Let $N \geq 1$ and let $T_1, \dots, T_N \in \mathcal{L}(X)$. Following [14], we say that T_1, \dots, T_N are *disjoint* or that the tuple (T_1, \dots, T_N) is *diagonally hypercyclic* if there exists $x \in X$ such that the set $\{(T_1^n x, \dots, T_N^n x) : n \geq 0\}$ is dense in X^N ; in other words, the ‘diagonal’ vector $x \oplus \dots \oplus x$ is hypercyclic for $T_1 \oplus \dots \oplus T_N$. Such a vector x is said to be *d-hypercyclic* for the tuple (T_1, \dots, T_N) and the set of *d-hypercyclic* vectors for (T_1, \dots, T_N) will be denoted by $d - \text{HC}(T_1, \dots, T_N)$. Similarly, (T_1, \dots, T_N) is said to be *d-frequently hypercyclic* if there exists $x \in X$ such that $x \oplus \dots \oplus x$ is a frequently hypercyclic vector for $T_1 \oplus \dots \oplus T_N$, and we denote by $d - \text{FHC}(T_1, \dots, T_N)$ the set of *d-frequently hypercyclic* vectors for (T_1, \dots, T_N) .

A natural problem regarding *d-hypercyclicity* is that of the *extension* of *d-hypercyclic* tuples. It was shown in [39] that given any $N \geq 1$, any Banach space X and any $T_1, \dots, T_N \in \mathcal{L}(X)$ such that (T_1, \dots, T_N) is *d-hypercyclic*, there exists $T_{N+1} \in \mathcal{L}(X)$ such that (T_1, \dots, T_{N+1}) is also *d-hypercyclic*. As for *d-frequent hypercyclicity*, the situation is trickier since there exist Banach spaces which do not support any frequently hypercyclic operator [53]. The best one could hope for is that as soon as X supports a frequently hypercyclic operator, then one can extend *d-frequently hypercyclic* tuples. We are unable to prove this, but we show that one can indeed extend *d-frequently hypercyclic* tuples as soon as X supports a *hereditarily frequently hypercyclic* operator (Theorem 6.1).

The preceding discussion has outlined the content of §§2–6 of the paper and hopefully it is clear that §6 makes a transition between our two topics—hereditary frequent hypercyclicity and *d-frequent hypercyclicity*. The next two sections are exclusively devoted to *d-frequent hypercyclicity*. In §7, we show (in the spirit of [51]) that there exists a *d-frequently hypercyclic* pair (T_1, T_2) on some Banach space X which is not densely *d-hypercyclic*, that is, the set $d - \text{HC}(T_1, T_2)$ is not dense in X (Theorem 7.2). In §8, we give a sufficient condition for *d-frequent hypercyclicity* of a tuple (T_1, \dots, T_N) in terms of eigenvectors of the operators T_i (Theorem 8.2); and this allows us for example to show

that the pair (D, τ_a) is d -frequently hypercyclic, where D is the derivation operator on the space of entire functions $H(\mathbb{C})$ and τ_a is the operator of translation by $a \in \mathbb{C} \setminus \{0\}$.

Finally, §9 contains a few additional remarks and a number of open questions originating in a rather natural way from our work.

2. The frequent hypercyclicity criterion

The frequent hypercyclicity criterion (FHCC) is a very efficient tool for showing that a given operator is frequently hypercyclic. Since [5], there have been several versions of it in the literature; we choose here the most widely used [16]: an operator $T \in \mathcal{L}(X)$ satisfies the FHCC provided there exist a dense set $\mathcal{D} \subset X$ and a map $S : \mathcal{D} \rightarrow \mathcal{D}$ such that:

- $TS = I$ on \mathcal{D} ;
- for any $x \in \mathcal{D}$, the series $\sum T^n x$ and $\sum S^n x$ are unconditionally convergent.

In this section, we show that the FHCC implies, in fact, hereditary frequent hypercyclicity.

THEOREM 2.1. *If $T \in \mathcal{L}(X)$ satisfies the FHCC, then T is hereditarily frequently hypercyclic.*

The proof of this theorem will closely mimic the classical proof that an operator satisfying the FHCC is frequently hypercyclic. Recall that the latter depends on the construction of subsets of \mathbb{N} with positive lower density which are ‘well separated’. To obtain hereditary frequent hypercyclicity, we need to control more precisely these subsets and, in particular, we have to be sure that one can find them inside some prescribed subsets of \mathbb{N} (of positive lower density, of course). This is the content of the next lemma, which is useful in other situations as well (see [36] and the very recent [18]). This lemma is actually contained in [36, Lemma 2.2], but we give a proof for completeness (and convenience of the reader).

LEMMA 2.2. *Let $(A_i)_{i \in I}$ be a countable family of subsets of \mathbb{N} with positive lower density and let $(N_i)_{i \in I}$ be a family of positive integers indexed by the same countable set I . There exists a family $(B_i)_{i \in I}$ of pairwise disjoint subsets of \mathbb{N} with positive lower density such that:*

- $B_i \subset A_i$ and $\min(B_i) \geq N_i$ for all $i \in I$;
- for any $i, j \in I$ and any $(n, m) \in B_i \times B_j$ with $n \neq m$, $|n - m| \geq N_i + N_j$.

Proof. We may assume that $I = \mathbb{N}$. For each $i \in \mathbb{N}$, let $M_i := 2 \max_{j \leq i} N_j$.

Enumerate each set A_i as an increasing sequence $(n_i(k))_{k \in \mathbb{N}}$ and for $s \geq 1$, define

$$\begin{aligned} A_{i,s} &:= \{n_i(sk) : k \in \mathbb{N}\}, \\ \tilde{A}_{i,s} &:= (A_{i,s} + [-M_i, M_i]) \cap \mathbb{N}. \end{aligned}$$

Then (by subadditivity of upper density),

$$\overline{\text{dens}}(\tilde{A}_{i,s}) \leq \frac{2M_i + 1}{s}.$$

Since all sets $A_{i,s}$ have positive lower density, it follows that one can construct, by induction, a sequence of positive integers $(s(i))_{i \in \mathbb{N}}$ such that for all $i \in \mathbb{N}$, $s(i) \geq M_i$ and

$$\overline{\text{dens}}(\tilde{A}_{i,s(i)}) \leq \min_{j < i} \frac{1}{4^{i-j}} \underline{\text{dens}}(A_{j,s(j)}).$$

We then set

$$B_i := A_{i,s(i)} \setminus \bigcup_{j > i} \tilde{A}_{j,s(j)},$$

so that

$$\underline{\text{dens}}(B_i) \geq \underline{\text{dens}}(A_{i,s(i)}) \left(1 - \sum_{j > i} \frac{1}{4^{j-i}} \right) > 0.$$

Moreover, B_i is clearly contained in A_i , $\min(B_i) \geq s(i) \geq M_i$, the sets B_i are pairwise disjoint, and if $(n, m) \in B_i \times B_j$ with $n \neq m$ and $j \geq i$, then:

- either $j = i$, in which case $|n - m| \geq s(i) \geq M_i \geq 2N_i = N_i + N_j$;
- or $j > i$, in which case $|n - m| \geq M_j \geq N_i + N_j$ since $n \notin B_j + [-M_j, M_j]$.

□

We can now give the proof of Theorem 2.1.

Proof of Theorem 2.1. Let $(A_p)_{p \in \mathbb{N}}$ be a sequence of subsets of \mathbb{N} with positive lower density and let $(V_p)_{p \in \mathbb{N}}$ be a sequence of non-empty open subsets of X . We have to find a vector $x \in X$ such that $\mathcal{N}_T(x, V_p) \cap A_p$ has positive lower density for all $p \in \mathbb{N}$; and for that, we follow the proof of [7, Theorem 6.18]. Let us fix an F -norm $\|\cdot\|$ defining the topology of X .

Let \mathcal{D} be the dense set given by the FHCC. For each $p \geq 1$, choose a vector $x_p \in \mathcal{D} \cap V_p$ and let $\alpha_p > 0$ be such that $B(x_p, 3\alpha_p) \subset V_p$. Let also $(\varepsilon_p)_{p \geq 1}$ be a summable sequence of positive real numbers such that for all $p \geq 1$,

$$p\varepsilon_p + \sum_{q > p+1} \varepsilon_q < \alpha_p.$$

By unconditional convergence of the series involved in the FHCC, for each $p \geq 1$, one can find a positive integer N_p such that, for any set $F \subset \mathbb{N} \cap [N_p, \infty)$,

$$\left\| \sum_{n \in F} T^n x_i \right\| + \left\| \sum_{n \in F} S^n x_i \right\| \leq \varepsilon_p \quad \text{for all } i \leq p.$$

Now, let $(B_p)_{p \in \mathbb{N}}$ be the sequence of subsets of \mathbb{N} with positive lower density associated with (A_p) and (N_p) by Lemma 2.2. The vector x we are looking for is defined by

$$x := \sum_{p=1}^{\infty} \sum_{n \in B_p} S^n x_p.$$

First, we note that x is well defined. Indeed, each series $\sum_{n \in B_p} S^n x_p$ is convergent and since $B_p \subset [N_p, \infty)$ for all p , we have

$$\sum_{p \geq 1} \left\| \sum_{n \in B_p} S^n x_p \right\| \leq \sum_{p \geq 1} \varepsilon_p < \infty.$$

Let us fix $p \geq 1$ and $n \in B_p$: we show that $T^n x \in V_p$. By definition of x , we have

$$\|T^n x - x_p\| \leq \sum_{q=1}^{\infty} \left\| \sum_{\substack{m \in B_q \\ m > n}} S^{m-n} x_q \right\| + \sum_{q=1}^{\infty} \left\| \sum_{\substack{m \in B_q \\ m < n}} T^{n-m} x_q \right\|.$$

To estimate the first sum, we decompose it as

$$\sum_{q=1}^p \left\| \sum_{\substack{m \in B_q \\ m > n}} S^{m-n} x_q \right\| + \sum_{q=p+1}^{\infty} \left\| \sum_{\substack{m \in B_q \\ m > n}} S^{m-n} x_q \right\|.$$

Since $n \in B_p$, we know that $m - n > \max(N_p, N_q)$ whenever $m \in B_q$ and $m > n$. By the choice of the sequence (N_p) , it follows that

$$\sum_{q=1}^{\infty} \left\| \sum_{\substack{m \in B_q \\ m > n}} S^{m-n} x_q \right\| \leq p\varepsilon_p + \sum_{q=p+1}^{\infty} \varepsilon_q < \alpha_p.$$

Estimating the second sum in the same way, we conclude that

$$\|T^n x - x_p\| < 3\alpha_p,$$

so that $T^n(x) \in V_p$. □

As a direct consequence of Theorem 2.1 and the fact that every frequently hypercyclic weighted backward shift on $\ell_p(\mathbb{Z}_+)$ satisfies the FHCC [9], we obtain the following corollary.

COROLLARY 2.3. *A weighted backward shift on $\ell_p(\mathbb{Z}_+)$, $1 \leq p < \infty$ is frequently hypercyclic if and only if it is hereditarily frequently hypercyclic.*

Remark 2.4. It should be clear from the proof of Theorem 2.1 that the FHCC implies hereditary \mathcal{F} -hypercyclicity for any Furstenberg family \mathcal{F} satisfying Lemma 2.2. For example, this holds true for the family of sets with positive upper density and the family of sets with positive Banach upper density, see [36, Lemma 2.2]; so any operator satisfying the FHCC is hereditarily \mathcal{U} -frequently hypercyclic. It is not clear to us that this could be deduced from Theorem 2.1: indeed, if \mathcal{F} and \mathcal{F}' are two Furstenberg families, the inclusion $\mathcal{F} \subset \mathcal{F}'$ does not formally imply that hereditary \mathcal{F} -hypercyclicity is a stronger property than hereditary \mathcal{F}' -hypercyclicity.

3. Ergodic theory

3.1. Results and general strategy. It is well known that if $T \in \mathfrak{L}(X)$ and if one can find a T -invariant Borel probability measure μ on X with full support with respect to which T is an ergodic transformation, then T is frequently hypercyclic. Let us recall the argument. Let

$(V_p)_{p \in \mathbb{N}}$ be a countable basis of open sets for X . Applying Birkhoff's pointwise ergodic theorem to the characteristic functions $\mathbf{1}_{V_p}$, we obtain a sequence (Ω_p) of subsets of X with $\mu(\Omega_p) = 1$ such that

$$\frac{1}{N} \#(N_T(x, V_p) \cap [0, N-1]) = \frac{1}{N} \sum_{n=0}^{N-1} \mathbf{1}_{V_p}(T^n x) \xrightarrow{N \rightarrow \infty} \mu(V_p) > 0 \quad \text{for every } x \in \Omega_p.$$

Hence, any $x \in \bigcap_{p \in \mathbb{N}} \Omega_p$ is a frequently hypercyclic vector for T .

It is also well known (see for example [5], [7, Ch. 5], [8]) that if X is a complex Banach space and if an operator $T \in \mathcal{L}(X)$ admits 'sufficiently many' \mathbb{T} -eigenvectors (that is, eigenvectors whose associated eigenvalues have modulus 1), then it is indeed possible to find an ergodic measure with full support for T . So, it may seem reasonable to expect that operators with sufficiently many \mathbb{T} -eigenvectors are hereditarily frequently hypercyclic.

Now, if one wants to repeat the above argument to show that a given operator is hereditarily frequently hypercyclic, Birkhoff's ergodic theorem is not enough: what is needed is a pointwise convergence result for averages of quantities of the form $\mathbf{1}_V(T^n x)$ not only along the whole sequence of integers, but in fact along any sequence (n_k) with positive lower density. Specifically, we will use a theorem of Conze [20], which we state in the next section (Theorem 3.3). The assumptions of Conze's theorem are much stronger than merely asking that T is an ergodic transformation; so we will need to impose a rather strong condition on the \mathbb{T} -eigenvectors to be able to conclude that our operator T is indeed hereditarily frequently hypercyclic.

Let us recall a few definitions. Assume that X is a complex Fréchet space. If $T \in \mathcal{L}(X)$, a \mathbb{T} -eigenvector field for T is any bounded map $E : \mathbb{T} \rightarrow X$ such that $TE(z) = zE(z)$ for every $z \in \mathbb{T}$. Given a positive Borel measure σ on \mathbb{T} , we say that a family of \mathbb{T} -eigenvector fields $(E_i)_{i \in I}$ is σ -spanning if $\overline{\text{span}}(E_i(z) : z \in \mathbb{T} \setminus N, i \in I) = X$ for every Borel set $N \subset \mathbb{T}$ such that $\sigma(N) = 0$. Similarly, we say that the \mathbb{T} -eigenvectors of T are σ -spanning if for every Borel set $N \subset \mathbb{T}$ such that $\sigma(N) = 0$, the eigenvectors of T with eigenvalues in $\mathbb{T} \setminus N$ span a dense subspace of X . It follows from [7, Lemma 5.29] that the \mathbb{T} -eigenvectors of T are σ -spanning if and only if there exists a σ -spanning countable family of Borel \mathbb{T} -eigenvector fields for T . One of our main aims in this part of the paper will be to prove the following theorem. (We refer to any book on Banach space theory, for example [2], for the definition of type. Let us just recall that an L_p -space has type 2 if $2 \leq p < \infty$ and type p if $1 \leq p < 2$.)

THEOREM 3.1. *Let X be a separable complex Banach space and let $T \in \mathcal{L}(X)$. Assume that one can find a λ -spanning, finite or countably infinite family $(E_i)_{i \in I}$ of \mathbb{T} -eigenvector fields for T , where λ is the Lebesgue measure on \mathbb{T} . Moreover, assume that one of the following holds true:*

- (a) X has type 2;
- (b) X has type $p \in [1, 2)$ and each eigenvector field E_i is α_i -Hölderian for some $\alpha_i > 1/p - 1/2$.

Then T is hereditarily frequently hypercyclic.

From this, we immediately get the following corollary.

COROLLARY 3.2. *Let X be a separable complex Banach space and let $T \in \mathcal{L}(X)$. If the \mathbb{T} -eigenvectors of T are spanning with respect to the Lebesgue measure and if, moreover, the Banach space X has type 2, then T is hereditarily frequently hypercyclic.*

In view of the above discussion, our strategy for proving Theorem 3.1 should be clear: we will show that under the above assumptions, one can find a T -invariant (Borel probability) measure μ on X for which one can apply Conze's Theorem 3.3 below to get hereditary frequent hypercyclicity. This will yield a more precise result, Theorem 3.9.

The proof of Theorem 3.1 will occupy us from §§3.2 to 3.6. After that, in §3.7, we will come back to the FHCC; in particular, we explain how the ergodic-theoretic point of view allows to give a completely different proof of Theorem 2.1, and we compare the assumptions in Theorems 2.1 and 3.1. Finally, in §3.8, we show that an assumption on the \mathbb{T} -eigenvectors similar to that of Corollary 3.2 but much weaker—namely, the *perfect spanning* property—is sufficient to imply hereditary \mathcal{U} -frequent hypercyclicity (without any additional assumption on the space X).

3.2. Conze's theorem. Before stating Conze's theorem, let us introduce some terminology.

In what follows, by a *measure-preserving dynamical system* (X, \mathcal{B}, μ, T) , we mean a pair consisting of a probability space (X, \mathcal{B}, μ) and a measurable map $T : X \rightarrow X$ such that $\mu \circ T^{-1} = \mu$. Note that here, we are departing from our standing notation: X is an abstract space, not necessarily a topological vector space, and hence, T is not necessarily a linear operator. This ambiguity is in fact intentional and it should cause no confusion.

Given a measure-preserving dynamical system (X, \mathcal{B}, μ, T) , we denote by $U_T : L^2(X, \mathcal{B}, \mu) \rightarrow L^2(X, \mathcal{B}, \mu)$ the associated Koopman operator, which is defined by

$$U_T f := f \circ T.$$

This is an isometry of $L^2(X, \mathcal{B}, \mu)$, and a unitary operator if T is bijective and bimeasurable, in which case we say that T is an *automorphism* of (X, \mathcal{B}, μ) or that the measure-preserving dynamical system (X, \mathcal{B}, μ, T) is *invertible*.

We stress a technical point: all probability spaces (X, \mathcal{B}, μ) under consideration will be assumed to be *standard Borel*, that is, the underlying measurable space (X, \mathcal{B}) is isomorphic to $(Z, \mathcal{B}(Z))$ for some Polish space Z , where $\mathcal{B}(Z)$ is the Borel σ -algebra of Z . Additionally, when X is already a Polish space, we assume that \mathcal{B} is the Borel σ -algebra of X .

A unitary operator $U : H \rightarrow H$ acting on a Hilbert space H is said to have *Lebesgue spectrum* if there exists a family of vectors $(f_i)_{i \in I}$ in H such that $\{U^n f_i : n \in \mathbb{Z}, i \in I\}$ is an orthonormal basis of H . Observe that if H is separable, the family (f_i) has to be finite or countably infinite. When there exists a countably infinite such family (f_i) , we say that U has *countable Lebesgue spectrum*. Finally, we say that an invertible measure-preserving dynamical system (X, \mathcal{B}, μ, T) has (countable) Lebesgue spectrum if the restriction of the Koopman operator U_T to $L_0^2(X, \mathcal{B}, \mu) := \{f \in L^2(X, \mathcal{B}, \mu) : \int_X f \, d\mu = 0\}$ has (countable) Lebesgue spectrum. We may also say that T itself has (countable) Lebesgue spectrum.

Conze's theorem from [20] now reads as follows.

THEOREM 3.3. *Let (X, \mathcal{B}, μ, T) be an invertible measure-preserving dynamical system and assume that T has Lebesgue spectrum. If $(n_k)_{k \geq 0}$ is an increasing sequence of integers with positive lower density then, for any $f \in L^1(\mu)$,*

$$\frac{1}{N} \sum_{k=0}^{N-1} f(T^{n_k} x) \xrightarrow{N \rightarrow \infty} \int_X f d\mu \quad \text{for } \mu\text{-almost every } x \in X.$$

Getting back to the case where X is a separable Banach space, it is easy to check that if $T \in \mathcal{L}(X)$ and if one can find a T -invariant measure with full support μ such that (X, \mathcal{B}, μ, T) satisfies the assumptions of Conze's theorem, then T is hereditarily frequently hypercyclic. However, Conze's theorem is only stated for automorphisms and, in our context, it would be rather restrictive to confine ourselves to the case of automorphisms. We will get round this problem thanks to the notion of *factor*. Recall that a measure-preserving dynamical system (X, \mathcal{B}, μ, T) is a factor of a measure-preserving dynamical system (Y, \mathcal{C}, ν, S) or that (Y, \mathcal{C}, ν, S) is an *extension* of (X, \mathcal{B}, μ, T) , if (possibly after deleting two sets of measure 0 in X and Y) there exists a measurable map $\pi : Y \rightarrow X$ such that $\pi \circ S = T \circ \pi$ and $\mu = \nu \circ \pi^{-1}$. Using suitable extensions, we will be able to apply Conze's theorem to general operators T via the following corollary.

COROLLARY 3.4. *Let $T \in \mathcal{L}(X)$ and assume that there exists a T -invariant probability measure μ on X with full support such that (X, \mathcal{B}, μ, T) is a factor of an invertible measure-preserving dynamical system (Y, \mathcal{C}, ν, S) with Lebesgue spectrum. Then, T is hereditarily frequently hypercyclic. More precisely, given a countable family $(A_i)_{i \in I}$ of subsets of \mathbb{N} with positive lower density and a family $(V_i)_{i \in I}$ of non-empty open sets, μ -almost every $x \in X$ is such that $\mathcal{N}_T(x, V_i) \cap A_i$ has positive lower density for every $i \in I$.*

Proof. It is enough to prove the result for a single pair (A, V) , where $A \subset \mathbb{N}$ has positive lower density and V is a non-empty open set.

Let $(n_k)_{k \geq 1}$ be the increasing enumeration of A . Let also $W := \pi^{-1}(V)$, where $\pi : (Y, \mathcal{C}, \nu, S) \rightarrow (X, \mathcal{B}, \mu, T)$ is a factor map, so that $\nu(W) = \mu(V) > 0$, and let

$$\Omega := \left\{ y \in Y : \frac{1}{N} \sum_{k=1}^N \mathbf{1}_W(S^{n_k} y) \xrightarrow{N \rightarrow \infty} \mu(V) \right\}.$$

By Conze's theorem, we know that $\nu(\Omega) = 1$. Since we are working with standard Borel spaces, it follows that the set $\pi(\Omega)$ is μ -measurable (being an analytic set, it is universally measurable) and that $\mu(\pi(\Omega)) = 1$. So it is enough to show that every $x \in \pi(\Omega)$ is such that $\underline{\text{dens}}(\mathcal{N}_T(x, V) \cap A) > 0$.

Let $x \in \pi(\Omega)$ and write x as $x = \pi(y)$ for some $y \in \Omega$. By definition of Ω , we know that the set $\{k \geq 1 : S^{n_k} y \in W\}$ has positive lower density; enumerate it as an increasing sequence $(m_k)_{k \geq 1}$. Then, $B := \{n_{m_k} : k \geq 1\}$ has positive lower density, it is contained in A and $S^{n_k} y \in \pi^{-1}(V)$ for all $n \in B$. This means that $T^n x \in V$ for all $n \in B$, which concludes the proof. \square

3.3. *Natural extensions.* To apply Corollary 3.4, we need a simple way of extending a given measure-preserving dynamical system (X, \mathcal{B}, μ, T) to a measure-preserving dynamical system $(\tilde{X}, \tilde{\mathcal{B}}, \tilde{\mu}, \tilde{T})$, where \tilde{T} is an automorphism of $(\tilde{X}, \tilde{\mathcal{B}}, \tilde{\mu})$. Fortunately, there is a canonical procedure doing precisely that. Consider the set

$$\tilde{X} := \{(x_k)_{k \geq 0} \in X^{\mathbb{Z}_+} : T(x_{k+1}) = x_k \text{ for all } k \geq 0\},$$

and for $k \geq 0$, let $\pi_k : \tilde{X} \rightarrow X$ denote the projection onto the k th coordinate of \tilde{X} . Endow \tilde{X} with the smallest σ -algebra $\tilde{\mathcal{B}}$ which makes every projection π_k measurable. The Rokhlin's natural extension of T is the measurable transformation $\tilde{T} : \tilde{X} \rightarrow \tilde{X}$ defined by

$$\tilde{T}(x_0, x_1, \dots) := (T(x_0), x_0, x_1, \dots).$$

One can prove that there exists a unique probability measure $\tilde{\mu}$ on $(\tilde{X}, \tilde{\mathcal{B}})$ such that $\tilde{\mu} \circ \pi_k^{-1} = \mu$ for all $k \geq 0$. (Here, the fact (X, \mathcal{B}) is a standard Borel space is needed.) Then, it is a simple exercise to check that $\tilde{T} : \tilde{X} \rightarrow \tilde{X}$ is an automorphism of $(\tilde{X}, \tilde{\mathcal{B}}, \tilde{\mu})$ such that $\pi_k \circ \tilde{T} = T \circ \pi_k$ for all $k \geq 0$. In particular, (X, \mathcal{B}, μ, T) is a factor of $(\tilde{X}, \tilde{\mathcal{B}}, \tilde{\mu}, \tilde{T})$ as witnessed by $\pi_0 : \tilde{X} \rightarrow X$. Note also that if X is a Polish space, then \tilde{X} is a Borel subset of the Polish space $X^{\mathbb{Z}_+}$ endowed with the product topology, and $\tilde{\mathcal{B}}$ is the Borel sigma-algebra of \tilde{X} . For more details, see for example [56, §8.4] or [46].

We will need the following lemma. Recall that if X is a complex Fréchet space, a Borel probability measure μ on X is said to be *Gaussian* if every continuous linear functional on X has a complex symmetric Gaussian distribution when considered as a random variable on (X, \mathcal{B}, μ) .

LEMMA 3.5. *Assume that X is a complex Fréchet space, that $T \in \mathfrak{L}(X)$ and that the measure μ is Gaussian. Then, \tilde{X} is a Fréchet space when endowed with the induced product topology of $X^{\mathbb{Z}_+}$, \tilde{T} is an invertible continuous linear operator on \tilde{X} and the measure $\tilde{\mu}$ is Gaussian.*

Proof. It is clear that \tilde{X} is a closed linear subspace of $X^{\mathbb{Z}_+}$ (and hence a Fréchet space) and that \tilde{T} is an invertible continuous linear operator on \tilde{X} .

Let ϕ be a continuous linear functional on \tilde{X} . By the Hahn–Banach theorem, one can extend ϕ to a continuous linear functional Φ on $X^{\mathbb{Z}_+}$. Since $X^{\mathbb{Z}_+}$ is endowed with the product topology, this linear functional Φ has the form

$$\Phi = \sum_{k=0}^N x_k^* \circ \pi_k,$$

where $x_0^*, \dots, x_N^* \in X^*$. Moreover, by definition of \tilde{X} , we have $\pi_k = T^{N-k} \circ \pi_N$ on \tilde{X} for $k = 0, \dots, N$; so we may write

$$\phi = x^* \circ \pi_N \quad \text{where } x^* = \sum_{k=0}^N x_k^* \circ T^{N-k} \in X^*.$$

Hence, $\tilde{\mu} \circ \phi^{-1} = (\tilde{\mu} \circ \pi_N^{-1}) \circ (x^*)^{-1} = \mu \circ (x^*)^{-1}$. Since μ is a Gaussian measure, it follows that $\tilde{\mu}$ is Gaussian as well. \square

3.4. Gaussian measures and countable Lebesgue spectrum. In this section, we state a general result about Gaussian linear measure-preserving dynamical systems. This result looks very much like [21, Theorem 14.3.2], but it is not clear to us that it can be formally deduced from it; so we will give a more or less complete proof. However, this proof is not necessary for understanding the proof of Theorem 3.1, so it is postponed to §3.6 for the sake of better readability. This means that a reader who is ready to take Proposition 3.6 on faith can safely ignore that section.

Recall that if (X, \mathcal{B}, μ, T) is a measure-preserving dynamical system then, for any $f \in L^2(\mu)$, there is a unique positive Borel measure σ_f on \mathbb{T} with non-negative Fourier coefficients

$$\widehat{\sigma_f}(n) = \langle U_T^n f, f \rangle_{L^2(\mu)}, \quad n \geq 0.$$

This follows from Herglotz's theorem and from the fact that the Koopman operator $U_T : L^2(\mu) \rightarrow L^2(\mu)$ is a contraction operator (so that the map $\mathbb{Z}_+ \ni n \mapsto \langle U_T^n f, f \rangle_{L^2(\mu)}$ can be extended to a positive-definite function on \mathbb{Z} , see for example [55, p. 28]). The measure σ_f is the *spectral measure* of f with respect to U_T .

PROPOSITION 3.6. *Let (X, \mathcal{B}, μ, T) be a measure-preserving dynamical system where X is a separable Fréchet space, T is an invertible linear operator and μ is a Gaussian measure on X . Assume that for any $x^* \in X^* \subset L^2(\mu)$, the spectral measure σ_{x^*} with respect to U_T is absolutely continuous with respect to the Lebesgue measure. Then, (X, \mathcal{B}, μ, T) has countable Lebesgue spectrum.*

Proposition 3.6 has the following consequence for not necessarily invertible Gaussian linear measure-preserving dynamical systems.

COROLLARY 3.7. *Let (X, \mathcal{B}, μ, T) be a measure-preserving dynamical system, where X is a complex separable Fréchet space, T is a continuous linear operator and μ is a Gaussian measure on X . Assume that for any $x^* \in X^* \subset L^2(\mu)$, the spectral measure σ_{x^*} with respect to U_T is absolutely continuous with respect to the Lebesgue measure. Then, the natural extension of (X, \mathcal{B}, μ, T) has countable Lebesgue spectrum.*

Proof. Let $(\tilde{X}, \tilde{\mathcal{B}}, \tilde{\mu}, \tilde{T})$ be the natural extension. By Lemma 3.5, we know that \tilde{T} is an invertible linear operator and $\tilde{\mu}$ is a Gaussian measure. Hence, by Proposition 3.6, it is enough to show that for any continuous linear functional ϕ on \tilde{X} , the spectral measure σ_ϕ (with respect to $U_{\tilde{T}}$) is absolutely continuous with respect to the Lebesgue measure.

By the proof of Lemma 3.5, one can find $N \in \mathbb{Z}_+$ and $x^* \in X^*$ such that $\phi = x^* \circ \pi_N$ on \tilde{X} , where $\pi_N : X^{\mathbb{Z}_+} \rightarrow X$ is the projection onto the N th coordinate. Now, we apply the following purely formal fact, which holds true for any measure-preserving dynamical system (X, \mathcal{B}, μ, T) .

FACT 3.8. *Let $f \in L^2(X, \mathcal{B}, \mu)$ and $F := f \circ \pi_N \in L^2(\tilde{X}, \tilde{\mathcal{B}}, \tilde{\mu})$. Then, the spectral measure σ_F with respect to $U_{\tilde{T}}$ is equal to σ_f , the spectral measure of f with respect to U_T .*

Proof of Fact 3.8. For all $n \geq 0$,

$$\begin{aligned}\langle U_T^n F, F \rangle_{L^2(\tilde{\mu})} &= \langle F \circ \tilde{T}^n, F \rangle_{L^2(\tilde{\mu})} = \langle f \circ \pi_N \circ \tilde{T}^n, f \circ \pi_N \rangle_{L^2(\tilde{\mu})} \\ &= \langle f \circ T^n \circ \pi_N, f \circ \pi_N \rangle_{L^2(\tilde{\mu})} \\ &= \langle f \circ T^n, f \rangle_{L^2(\mu)} \\ &= \langle U_T^n f, f \rangle_{L^2(\mu)}.\end{aligned}\quad \square$$

By Fact 3.8, $\sigma_\phi = \sigma_{x^*}$ is absolutely continuous with respect to the Lebesgue measure, which concludes the proof of Corollary 3.7. \square

3.5. Proof of Theorem 3.1. We can now state and prove very easily the following more precise version of Theorem 3.1.

THEOREM 3.9. *Let X be a separable complex Banach space and let $T \in \mathfrak{L}(X)$. Assume that one can find a λ -spanning, finite or countably infinite family $(E_i)_{i \in I}$ of \mathbb{T} -eigenvector fields for T , where λ is the Lebesgue measure on \mathbb{T} . Moreover, assume that one of the following holds true:*

- (a) X has type 2;
- (b) X has type $p \in [1, 2)$, and each E_i is α_i -Hölderian for some $\alpha_i > 1/p - 1/2$.

Then, there exists a T -invariant Gaussian measure μ on X with full support such that (X, \mathcal{B}, μ, T) is a factor of a measure-preserving system which has countable Lebesgue spectrum. In particular, T is hereditarily frequently hypercyclic; and more precisely: given a countable family $(A_i)_{i \in I}$ of subsets of \mathbb{N} with positive lower density and a family $(V_i)_{i \in I}$ of non-empty open sets, μ -almost every $x \in X$ is such that $\mathcal{N}_T(x, V_i) \cap A_i$ has positive lower density for every $i \in I$.

Proof. Under the assumptions above, there exists a T -invariant Gaussian measure μ on X with full support such that, for all $x^* \in X^*$, the measure σ_{x^*} is absolutely continuous with respect to the Lebesgue measure; this is contained for instance in [7, Lemma 5.35, (4)]. So, the result follows immediately from Corollaries 3.7 and 3.4. \square

3.6. Proof of Proposition 3.6. In this section, we give the promised proof of Proposition 3.6.

3.6.1. Two facts concerning unitary operators. We will need several results on unitary operators. These results are certainly well known, but since they are rather difficult to locate in the literature, we provide some details.

Let U be a unitary operator acting on a complex separable Hilbert space H . By Herglotz's theorem, for any $f \in H$, there exists a unique positive and finite Borel measure σ_f on \mathbb{T} , called the *spectral measure* of f with respect to U , such that

$$\text{for all } n \in \mathbb{Z} : \langle U^n f, f \rangle = \widehat{\sigma_f}(n).$$

Note in particular that σ_f is equal to the Lebesgue measure λ if and only if the sequence $(U^n f)_{n \in \mathbb{Z}}$ is orthonormal. This explains the terminology 'Lebesgue spectrum'.

Denote by $C(f)$ the *cyclic subspace* generated by f , that is, $C(f) = \overline{\text{span}}(U^n f : n \in \mathbb{Z})$. With this notation, one form of the spectral theorem reads as follows (see for example [19, Ch. IX], [47, Appendix, §2] or [49]): there exists a finite or infinite sequence of vectors $f_i \in H$, $0 \leq i < m$, where $m \in \mathbb{N} \cup \{\infty\}$, such that

$$H = \bigoplus_{0 \leq i < m} C(f_i) \quad \text{and} \quad \sigma_{f_0} \gg \sigma_{f_1} \gg \cdots \gg \sigma_{f_i} \gg \cdots.$$

Moreover, these measures are essentially unique in the following sense: for any other sequence $(g_i)_{0 \leq i < m'}$ satisfying $H = \bigoplus_{0 \leq i < m'} C(g_i)$ and $\sigma_{g_0} \gg \sigma_{g_1} \gg \cdots \gg \sigma_{g_i} \gg \cdots$, we have $m' = m$ and $\sigma_{f_i} \sim \sigma_{g_i}$ for all i . The *maximal spectral type* of T is then defined as (the equivalence class of) the measure σ_{f_0} .

Observe that for any $f \in H$, the restriction of U to $C(f)$ is unitary equivalent to the multiplication operator $M_{\sigma_f} : L^2(\mathbb{T}, \sigma_f) \rightarrow L^2(\mathbb{T}, \sigma_f)$ defined by $M_{\sigma_f} u(z) := zu(z)$, $z \in \mathbb{T}$.

Suppose now that U has countable Lebesgue spectrum and let $(f_i)_{i \in \mathbb{N}}$ be a sequence in H such that $\{U^n f_i : n \in \mathbb{Z}, i \in \mathbb{N}\}$ is an orthonormal basis of H . Then, $H = \bigoplus_{i \in \mathbb{N}} C(f_i)$ and $\sigma_{f_i} = \lambda$ for all $i \in \mathbb{N}$. Hence, U is unitarily equivalent to $M_\lambda^{(\infty)} := \bigoplus_{i=1}^\infty M_\lambda$ acting on $\bigoplus_{i=1}^\infty L^2(\mathbb{T}, \lambda)$. Conversely, it is clear that if $U \cong M_\lambda^{(\infty)}$, then U has countable Lebesgue spectrum.

By the uniqueness part of the spectral theorem, it follows in particular that the maximal spectral type of a unitary operator with countable Lebesgue spectrum is the Lebesgue measure. The next lemma is a kind of converse.

LEMMA 3.10. *Let U be a unitary operator on a complex separable Hilbert space H satisfying the following two conditions.*

- (a) *There exists a closed subspace $K \subset H$ such that $U(K) = K$ and $U|_K : K \rightarrow K$ has countable Lebesgue spectrum.*
- (b) *The maximal spectral type of U is the Lebesgue measure.*

Then, $U : H \rightarrow H$ has countable Lebesgue spectrum.

Proof. We will use the following notation: if σ is a positive finite Borel measure on \mathbb{T} and $N \in \mathbb{N} \cup \{\infty\}$, we denote by $M_\sigma^{(N)}$ the operator $\bigoplus_{0 \leq j < N} M_\sigma$ acting on $\bigoplus_{0 \leq j < N} L^2(\mathbb{T}, \sigma)$. Recall also that λ is the Lebesgue measure on \mathbb{T} .

By condition (a), we know that

$$U|_K \cong M_\lambda^{(\infty)}.$$

Consider now the operator $U|_{K^\perp} : K^\perp \rightarrow K^\perp$. By condition (b), its maximal spectral type is absolutely continuous with respect to the Lebesgue measure λ . By the ‘second formulation’ of the spectral theorem (see for example [49] or [19, Ch. IX]), there exist pairwise disjoint Borel sets $\Delta_\infty, \Delta_1, \Delta_2, \dots$ and positive finite measures $\mu_\infty, \mu_1, \mu_2, \dots$ on \mathbb{T} supported on $\Delta_\infty, \Delta_1, \Delta_2, \dots$ and absolutely continuous with respect to λ , such that $U|_{K^\perp}$ is unitarily equivalent to $M_{\mu_\infty}^{(\infty)} \oplus M_{\mu_1}^{(1)} \oplus M_{\mu_2}^{(2)} \oplus \cdots$. Note that some of the measures μ_n may be 0. For $n \in \{\infty\} \cup \mathbb{N}$, we may write $\mu_n = f_n \lambda$ for some non-negative function $f_n \in L^1(\mathbb{T})$ and we may assume that $\Delta_n = \{f_n > 0\}$. Hence, if

we set $\nu_n := \mathbf{1}_{\Delta_n} \lambda$, we get $M_{\mu_n} \cong M_{\nu_n}$ since the measures μ_n and ν_n are equivalent. Therefore,

$$U|_{K^\perp} \cong M_{\nu_\infty}^{(\infty)} \oplus M_{\nu_1}^{(1)} \oplus M_{\nu_2}^{(2)} \oplus \dots$$

Now, let $E := \mathbb{T} \setminus (\Delta_\infty \cup \bigcup_{n \geq 1} \Delta_n)$ and $\nu := \mathbf{1}_E \lambda$. Then, $\lambda = \nu + \nu_\infty + \sum_{n \in \mathbb{N}} \nu_n$ so that

$$M_\lambda \cong M_\nu \oplus M_{\nu_\infty} \oplus M_{\nu_1} \oplus \dots$$

Consequently, we obtain

$$\begin{aligned} U &\cong U|_K \oplus U|_{K^\perp} \\ &\cong M_\nu^{(\infty)} \oplus M_{\nu_\infty}^{(\infty)} \oplus M_{\nu_1}^{(\infty)} \oplus M_{\nu_2}^{(\infty)} \oplus \dots \\ &\quad \oplus M_{\nu_\infty}^{(\infty)} \oplus M_{\nu_1}^{(1)} \oplus M_{\nu_2}^{(2)} \oplus \dots \\ &\cong M_\nu^{(\infty)} \oplus M_{\nu_\infty}^{(\infty)} \oplus M_{\nu_1}^{(\infty)} \oplus M_{\nu_2}^{(\infty)} \dots \\ &\cong M_\lambda^{(\infty)}, \end{aligned}$$

which means that U has countable Lebesgue spectrum. \square

The following observation is useful to check assumption (a) in Lemma 3.10.

LEMMA 3.11. *Let $(f_i)_{i \in I}$ be a countably infinite family of vectors in H . Assume that the cyclic subspaces $C(f_i)$ are pairwise orthogonal and let $K := \bigoplus_{i \in I} C(f_i)$. If σ_{f_i} is equivalent to the Lebesgue measure for all $i \in I$, then $U|_K$ has countable Lebesgue spectrum.*

Proof. For each $i \in I$, there exists $g_i \in C(f_i)$ such that σ_{g_i} is exactly equal to the Lebesgue measure and for any such g_i , we have $C(g_i) = C(f_i)$; see for example [47, pp. 93–94]. Hence, $\{U^n g_i : i \in I, n \in \mathbb{Z}\}$ is an orthonormal basis of K . \square

We will also need the following fact.

LEMMA 3.12. *Let U be a unitary operator acting on a separable Hilbert space H and let σ be a finite, positive Borel measure on \mathbb{T} . If there exists a dense set $\mathcal{D} \subset H$ such that $\sigma_f \ll \sigma$ for all $f \in \mathcal{D}$, then $\sigma_f \ll \sigma$ for all $f \in H$.*

Proof. Denote by $M(\mathbb{T})$ the space of all complex Borel measures on \mathbb{T} . By [49, Corollary 2.2], the map $f \mapsto \sigma_f$ is continuous from H into $M(\mathbb{T})$ endowed with the norm topology. The result follows immediately. \square

3.6.2. Proof of Proposition 3.6. Before really starting the proof, we need to recall some basic facts concerning the L^2 -space of a Gaussian measure. In what follows, μ is a Gaussian measure on the (separable) complex Fréchet space X .

Let $\mathcal{G} := \overline{\text{span}}(\langle x^*, \cdot \rangle : x^* \in X^*)$, where the closure is taken in $L^2(X, \mathcal{B}, \mu)$. This is a Gaussian subspace of $L^2(X, \mathcal{B}, \mu)$, in the sense that any function in \mathcal{G} has symmetric complex Gaussian distribution. Moreover, \mathcal{G} is clearly U_T -invariant.

For $k \geq 0$, let us denote by \mathcal{G}^k the space of homogeneous polynomials of degree k in the elements of \mathcal{G} , with $\mathcal{G}^0 = \mathbb{C}$. The subspaces $\mathcal{G}^k, k \geq 0$, are linearly independent (which is not obvious) and one can orthonormalize them thanks to the so-called *Wick transform* $f \mapsto :f:$ which is defined on $\bigcup_{k \geq 0} \mathcal{G}^k$ as follows:

$$:f: = \begin{cases} f & \text{if } f \text{ is constant,} \\ f - P_k f & \text{if } f \in \mathcal{G}^k, k \geq 1, \end{cases}$$

where we denote by P_k the orthogonal projection onto $\overline{\text{span}}(\mathcal{G}^i : 0 \leq i \leq k-1)$.

With this notation, we have the orthogonal decomposition

$$L^2(X, \mathcal{B}, \mu) = \bigoplus_{k=0}^{\infty} \overline{: \mathcal{G}^k :}.$$

Moreover, for any $f_1, \dots, f_k, g_1, \dots, g_k \in \mathcal{G}$, the scalar product $\langle :f_1 \cdots f_k:, :g_1 \cdots g_k: \rangle$ is given by

$$\langle :f_1 \cdots f_k:, :g_1 \cdots g_k: \rangle = \sum_{s \in \mathfrak{S}_k} \langle f_{s(1)}, g_1 \rangle \cdots \langle f_{s(k)}, g_k \rangle, \quad (3.1)$$

where \mathfrak{S}_k is the permutation group of $\{1, \dots, k\}$.

For details on these general facts, we refer to [48, Ch. 8] or [34, Chs. 2 and 3]. We will also need the following lemma (see [5, Lemma 3.28]).

LEMMA 3.13. *Let $T \in \mathcal{L}(X)$ and let μ be a T -invariant Gaussian measure on X . For any $f_1, \dots, f_k \in \mathcal{G}$, we have*

$$U_T(:f_1 \cdots f_k:) = :(U_T f_1) \cdots (U_T f_k):.$$

In particular, each subspace $: \mathcal{G}^k :$ is U_T -invariant.

We can now finally give the proof of Proposition 3.6.

Proof of Proposition 3.6. Let (X, \mathcal{B}, μ, T) be a measure-preserving dynamical system, where $T \in (X)$ is invertible and the measure μ is Gaussian, such that for any $x^* \in X^* \subset L^2(\mu)$, the spectral measure σ_{x^*} with respect to U_T is absolutely continuous with respect to the Lebesgue measure. We want to show that (X, \mathcal{B}, μ, T) has countable Lebesgue spectrum.

Let us first recall that for any $f, g \in L^2(\mu)$, there is a unique complex Borel measure $\sigma_{f,g}$ on \mathbb{T} with Fourier coefficients

$$\widehat{\sigma_{f,g}}(n) = \langle U_T^n f, g \rangle, \quad n \in \mathbb{Z}.$$

If $f = g$, then $\sigma_{f,f}$ is the spectral measure σ_f ; and the existence of $\sigma_{f,g}$ for arbitrary functions f and g follows from a polarization argument. Indeed, we have

$$\sigma_{f,g} = \frac{1}{4} \sum_{k=0}^3 \mathbf{i}^k \sigma_{f+\mathbf{i}^k g}.$$

This formula and the assumption of the proposition show that σ_{x^*, y^*} is absolutely continuous with respect to the Lebesgue measure for any $x^*, y^* \in X^*$. Note also that the map $(f, g) \mapsto \sigma_{f, g}$ is obviously \mathbb{R} -bilinear.

In what follows, we denote by \star the convolution product for measures on \mathbb{T} and we write the elements of X^* as f, g, \dots rather than x^*, y^*, \dots to avoid the proliferation of stars.

FACT 3.14. *If $f_1, \dots, f_k \in X^*$, then $\sigma_{f_1 \dots f_k} = \sum_{s \in \mathfrak{S}_k} \sigma_{f_{s(1)}, f_1} \star \dots \star \sigma_{f_{s(k)}, f_k}$.*

Proof of Fact 3.14. By Lemma 3.13 and equation (3.1), we have for all $n \in \mathbb{Z}$:

$$\begin{aligned} \widehat{\sigma_{f_1 \dots f_k}}(n) &= \langle (U_T^n f_1) \cdots (U_T^n f_k) : , : f_1 \cdots f_k : \rangle \\ &= \sum_{s \in \mathfrak{S}_k} \langle U_T^n f_{s(1)}, f_1 \rangle \cdots \langle U_T^n f_{s(k)}, f_k \rangle \\ &= \sum_{s \in \mathfrak{S}_k} \widehat{\sigma_{f_{s(1)}, f_1}}(n) \cdots \widehat{\sigma_{f_{s(k)}, f_k}}(n). \end{aligned} \quad \square$$

Now, let us denote by \mathcal{D} the set of all functions $f \in L^2(\mu)$ of the form

$$f = \sum_{k=1}^N : \sum_{j=1}^{m_k} f_{j,1} \cdots f_{j,k} : \quad \text{with } m_k \geq 1 \text{ and } f_{j,i} \in X^*.$$

This is a dense linear subspace of $L_0^2(\mu)$.

FACT 3.15. *If $f \in \mathcal{D}$, then σ_f is absolutely continuous with respect to the Lebesgue measure.*

Proof of Fact 3.15. Let $f \in \mathcal{D}$, so that $f = \sum_{k=1}^N f_k$, where

$$f_k = \sum_{j=1}^{m_k} : f_{j,1} \cdots f_{j,k} : \quad \text{with } f_{j,i} \in X^*, \quad \text{so that } f_k \in : \mathcal{G}^k :.$$

By orthogonality and U_T -invariance of the subspaces $: \mathcal{G}^k :$, we have for all $n \in \mathbb{Z}$,

$$\widehat{\sigma_f}(n) = \langle U_T^n f, f \rangle = \sum_{k=1}^N \langle U_T^n f_k, f_k \rangle = \sum_{k=1}^N \widehat{\sigma_{f_k}}(n),$$

so that $\sigma_f = \sigma_{f_1} + \dots + \sigma_{f_N}$. Hence, it is enough to check that each measure σ_{f_k} is absolutely continuous with respect to the Lebesgue measure. However, this is clear by Fact 3.14 and bilinearity of the map $(f, g) \mapsto \sigma_{f, g}$: indeed, we have

$$\sigma_{f_k} = \sum_{j, j'=1}^{m_k} \sigma_{f_{j,1} \dots f_{j,k} : , : f_{j',1} \dots f_{j',k} :} = \sum_{j, j'=1}^{m_k} \sum_{s \in \mathfrak{S}_k} \sigma_{f_{j, s(1)}, f_{j',1}} \star \dots \star \sigma_{f_{j, s(k)}, f_{j',k}}$$

and the result follows since all measures $\sigma_{f_{j, s(i)}, f_{j',i}}$ are absolutely continuous with respect to the Lebesgue measure. \square

We can now finish the proof of Proposition 3.6. By Fact 3.15 and Lemma 3.12, the maximal spectral type of U_T is absolutely continuous with respect to the Lebesgue

measure. Now, take any $f \in X^* \setminus \{0\}$. Then, σ_f is a non-zero measure which is absolutely continuous with respect to the Lebesgue measure. It follows that there exists $k_0 \geq 1$ such that, for all $k \geq k_0$, the measure $\sigma_{f^k} = k! \sigma_f \star \cdots \star \sigma_f$ (k times) is *equivalent* to the Lebesgue measure (see for example [45, Lemma 3.6], where the symmetry assumption on the measure is in fact not necessary). Let us set

$$K := \overline{\text{span}} \{U_T^n(\cdot f^k) : n \in \mathbb{Z}, k \geq k_0\}.$$

By orthogonality and U_T -invariance of the spaces \mathcal{G}^k , and applying Lemma 3.11, we see that $(U_T)|_K$ has countable Lebesgue spectrum. Hence, by Lemma 3.10, U_T has countable Lebesgue spectrum. \square

3.7. The frequent hypercyclicity criterion again. The tools introduced in the previous sections allow us to give another proof of Theorem 2.1. Arguably, this proof is much less elementary. Let us say that an operator $T \in \mathcal{L}(X)$ has *countable Lebesgue spectrum after extension* if there exists a T -invariant Borel probability measure μ on X with full support such that (X, \mathcal{B}, μ, T) is a factor of a measure-preserving dynamical system which has countable Lebesgue spectrum. By Corollary 3.4, any such operator T is hereditarily frequently hypercyclic.

PROPOSITION 3.16. *Let $T \in \mathcal{L}(X)$ be an operator satisfying the FHCC. Then, T has countable Lebesgue spectrum after extension.*

Proof. It is shown in [44] that there exists a T -invariant measure μ on X with full support such that (X, \mathcal{B}, T, μ) is a factor of a Bernoulli shift; and it is well known that Bernoulli shifts have countable Lebesgue spectrum (see for example [57, Theorems 4.30 and 4.33]). \square

One can also prove the following ‘probabilistic’ version of Proposition 3.16. Let us say that a sequence $(x_n)_{n \in \mathbb{Z}}$ is a *bilateral backward orbit* for an operator T if $Tx_n = x_{n-1}$ for all $n \in \mathbb{Z}$.

PROPOSITION 3.17. *Let X be a complex Fréchet space and let $T \in \mathcal{L}(X)$. Assume that there exists a bilateral backward orbit $(x_n)_{n \in \mathbb{Z}}$ for T such that $\overline{\text{span}}(x_n : n \in \mathbb{Z}) = X$ and the series $\sum g_n x_n$ is almost surely convergent, where (g_n) is a sequence of independent complex standard Gaussian variables. Then, T has countable Lebesgue spectrum after extension. More precisely, there exists a T -invariant Gaussian measure μ on X with full support such that (X, \mathcal{B}, μ, T) is a factor of a measure-preserving dynamical system with countable Lebesgue spectrum.*

The (almost sure) convergence of the bilateral series $\sum g_n x_n$ means that both series

$$\sum_{n \geq 0} g_n x_n \quad \text{and} \quad \sum_{n < 0} g_n x_n$$

are (almost surely) convergent.

Proof. Let μ be the distribution of the random variable $\xi := \sum_{n \in \mathbb{Z}} g_n x_n$. This is a Gaussian measure, which has full support since $\overline{\text{span}}(x_n : n \in \mathbb{Z}) = X$ and which is T -invariant because (x_n) is a bilateral backward orbit for T . By Corollary 3.7, it is enough to show that for any $x^* \in X^* \subset L^2(\mu)$, the spectral measure σ_{x^*} of x^* with respect to U_T is absolutely continuous with respect to the Lebesgue measure.

By orthogonality of the Gaussian variables g_k , we have for all $n \geq 0$,

$$\begin{aligned} \widehat{\sigma_{x^*}}(n) &= \langle U_T^n x^*, x^* \rangle = \sum_{k \in \mathbb{Z}} \langle x^*, T^n x_k \rangle \overline{\langle x^*, x_k \rangle} \\ &= \sum_{k \in \mathbb{Z}} \langle x^*, x_{k-n} \rangle \overline{\langle x^*, x_k \rangle}. \end{aligned}$$

The series is absolutely convergent since the almost sure convergence of the scalar Gaussian series $\sum \langle x^*, x_k \rangle g_k$ implies that $\sum_{k \in \mathbb{Z}} |\langle x^*, x_k \rangle|^2 < \infty$.

Now, let $\varphi \in L^2(\mathbb{T})$ be the function with Fourier coefficients $\widehat{\varphi}(k) := \langle x^*, x_{-k} \rangle$, $k \in \mathbb{Z}$, that is,

$$\varphi(z) \sim \sum_{k \in \mathbb{Z}} \langle x^*, x_{-k} \rangle z^k;$$

and let $g := |\varphi|^2 \in L^1(\mathbb{T})$. By definition, we have for all $n \geq 0$,

$$\widehat{g}(n) = \sum_{k \in \mathbb{Z}} \widehat{\varphi}(k) \widehat{\varphi}(n-k) = \sum_{k \in \mathbb{Z}} \overline{\langle x^*, x_k \rangle} \langle x^*, x_{k-n} \rangle.$$

Since two positive measures on \mathbb{T} with the same non-negative Fourier coefficients must be equal, it follows that $\sigma_{x^*} = g(\lambda) d\lambda$, which concludes the proof. \square

Remark 3.18. The above proof shows in particular that if $(x_n)_{n \in \mathbb{Z}}$ is a bilateral backward orbit for T such that the series $\sum g_n x_n$ is almost surely convergent, then the distribution of the random variable $\xi := \sum_{n \in \mathbb{Z}} g_n x_n$ is a strongly mixing measure for T . This is not specific to Gaussian variables: as shown in [1], the same result holds true if (g_n) is replaced by any sequence of independent, identically distributed random variables.

We mentioned above that Proposition 3.17 is a probabilistic version of Proposition 3.16; let us be a little bit more explicit. The following fact (which was observed independently by A. López-Martínez) can be extracted from the proof of [1, Theorem 4.9].

Fact 3.19. Let X be a Fréchet space. If $T \in \mathfrak{L}(X)$ satisfies the FHCC, then there exists a bilateral backward orbit (x_n) for T such that $\overline{\text{span}}(x_n : n \in \mathbb{Z}) = X$ and the series $\sum x_n$ is unconditionally convergent.

In view of that, the next result is an improvement of Proposition 3.16 when X is a Banach space with non-trivial cotype.

COROLLARY 3.20. *Let X be a Banach space with non-trivial cotype and let $T \in \mathfrak{L}(X)$. Assume that there exists a bilateral backward orbit $(x_n)_{n \in \mathbb{Z}}$ for T such that $\overline{\text{span}}(x_n : n \in \mathbb{Z}) = X$ and the series $\sum \pm x_n$ is convergent for almost every choice of signs \pm . Then, T has countable Lebesgue spectrum after extension.*

Proof. By assumption on X , almost sure convergence of the Rademacher series $\sum \pm x_n$ is equivalent to almost sure convergence of the Gaussian series $\sum g_n x_n$ (this follows from [40, Corollaire 1.3, p. 67]; see also for example [35, Proposition 9.14]). So, the result follows immediately from Proposition 3.17. \square

Remark 3.21. In the same spirit, Proposition 3.17 can be used to show that if an operator T satisfies the ‘probabilistic frequent hypercyclicity criterion’ of [28, Theorem 1.2], then T is hereditarily frequently hypercyclic. Indeed, by [28, Lemma 3.2], there is a bilateral backward orbit $(x_n)_{n \in \mathbb{Z}}$ for T satisfying the assumption of Proposition 3.17.

We now show that, at least for operators on complex Hilbert spaces, Theorem 3.1 is ‘strictly stronger’ than Theorem 2.1. This answers a very natural question asked by the referee.

PROPOSITION 3.22. *Let X be a complex Fréchet space.*

- (1) *If $T \in \mathcal{L}(X)$ satisfies the FHCC, then there exists a family $(E_i)_{i \in I}$ of continuous \mathbb{T} -eigenvector fields for T such that $\overline{\text{span}} \bigcup_{i \in I} E_i(\mathbb{T}) = X$. In particular, this family is spanning with respect to the Lebesgue measure λ , so that if X is a Banach space with type 2, one can apply Theorem 3.1 to conclude that T is hereditarily frequently hypercyclic.*
- (2) *If X is a Hilbert space, there exists $T \in \mathcal{L}(X)$ which admits a λ -spanning \mathbb{T} -eigenvector field (so that T is hereditarily frequently hypercyclic by Theorem 3.1) but does not satisfy the FHCC. In fact, one can even require that T has no periodic vector except 0.*

Proof. (1) This is well known, see for example [8, §8.1].

(2) Let Λ be any compact subset of \mathbb{T} with empty interior, but such that all its non-empty (relative) open subsets have positive Lebesgue measure (to obtain such a set, take a compact set Λ_0 with empty interior and positive Lebesgue measure, and remove all open subsets of Λ_0 with Lebesgue measure 0). Let $T_\Lambda : H_\Lambda \rightarrow H_\Lambda$ be the Kalish operator associated with Λ (see for example [7, Ch. 5] for the definition). The operator T_Λ has the following properties: its point spectrum is equal to Λ and it admits a continuous \mathbb{T} -eigenvector field $E : \Lambda \rightarrow H_\Lambda$ such that $\overline{\text{span}} E(\Lambda) = H_\Lambda$. By continuity and since $\Lambda \setminus N$ is dense in Λ for every set $N \subset \mathbb{T}$ with Lebesgue measure 0, this eigenvector field is λ -spanning. However, T does not satisfy the FHCC. Indeed, assume otherwise. The continuous \mathbb{T} -eigenvector fields $E_i : \mathbb{T} \rightarrow H_\Lambda$ given by item (1) cannot vanish identically outside Λ (unless they are identically 0 on \mathbb{T}) since $\mathbb{T} \setminus \Lambda$ is dense in \mathbb{T} ; so T_Λ has eigenvalues outside Λ , which is a contradiction since $\sigma_p(T_\Lambda) = \Lambda$. Moreover, if Λ contains no roots of unity, then T_Λ has no periodic vector except 0 since no root of unity can be an eigenvalue of T_Λ . Recall that any operator (on a real or complex space) satisfying the FHCC is chaotic, see for example [7, Theorem 6.10]. \square

Finally, the ergodic-theoretic point of view allows us to give an ‘extreme’ example of a hereditarily frequently hypercyclic operator T which does not satisfy the FHCC.

Example 3.23. There exists a separable complex Banach space X supporting a hereditarily frequently hypercyclic operator with no eigenvalues.

Proof. Let T be the Kalish operator acting on the space $X := C_0([0, 2\pi])$ of continuous functions on $[0, 2\pi]$ vanishing at the point 0 (see [5, Example 4.2] or [7, §5.5.4]). The operator T has no eigenvalues. However, T admits a Gaussian invariant measure μ with full support such that all the spectral measures σ_x^* , $x \in X^*$ relative to the Koopman operator $U_T : L^2(\mu) \rightarrow L^2(\mu)$ are absolutely continuous with respect to the Lebesgue measure; so T has countable Lebesgue spectrum after extension by Corollary 3.7 and hence it is hereditarily frequently hypercyclic. \square

3.8. Perfect spanning and hereditary UFHC. The links between properties of unimodular eigenvectors of an operator T and frequent hypercyclicity of T have been very much studied since [5]. The strongest available result may be the following (see [8, 29]).

Let X be a separable complex Fréchet space and let $T \in \mathfrak{L}(X)$. If the \mathbb{T} -eigenvectors of T are perfectly spanning, then T is frequently hypercyclic and, in fact, there exists a Gaussian T -invariant measure μ with full support such that T is weakly mixing with respect to μ .

The perfect spanning assumption means that for any countable set $N \subset \mathbb{T}$, the eigenvectors of T with eigenvalues in $\mathbb{T} \setminus N$ span a dense linear subspace of X ; equivalently (see [30, Proposition 6.1]), the \mathbb{T} -eigenvectors of T are σ -spanning for some continuous probability measure σ on \mathbb{T} . It is plausible that under this assumption, the operator T is, in fact, hereditarily frequently hypercyclic; but we are very far from being able to prove that. We would be already happy enough if we could weaken the assumptions of Theorem 3.1 and prove that for any complex Banach space X , an operator $T \in \mathfrak{L}(X)$ is hereditarily frequently hypercyclic as soon as the \mathbb{T} -eigenvectors of T are spanning with respect to the Lebesgue measure—but again, this seems out of reach for the moment. However, we do have the following result.

THEOREM 3.24. *Let X be a complex Fréchet space and let $T \in \mathfrak{L}(X)$. If the \mathbb{T} -eigenvectors of T are perfectly spanning, then T is hereditarily \mathcal{U} -frequently hypercyclic.*

For the proof, we will need the following variant of [20, Lemme 5].

LEMMA 3.25. *Let (X, \mathcal{B}, μ, T) be a measure-preserving dynamical system and assume that T is weakly mixing with respect to μ . Let also $(n_k)_{k \geq 1}$ and $(k_i)_{i \geq 0}$ be two increasing sequence of integers. Assume that $n_{k_i} = O(k_i)$ as $i \rightarrow \infty$. Then, for any measurable set $V \subset X$,*

$$\frac{1}{k_i} \sum_{k=1}^{k_i} \mathbf{1}_V \circ T^{n_k} \xrightarrow{L^2} \mu(V) \quad \text{as } i \rightarrow \infty.$$

Proof. The proof is similar to that of the classical Blum–Hanson theorem [15]. We have

$$\begin{aligned} \left\| \frac{1}{k_i} \sum_{k=1}^{k_i} \mathbf{1}_V \circ T^{n_k} - \mu(V) \right\|_2^2 &= \frac{1}{k_i^2} \sum_{r,s=1}^{k_i} (\mu(T^{-n_r}(V) \cap T^{-n_s}(V)) - \mu(V)^2) \\ &= \frac{2}{k_i^2} \sum_{1 \leq r < s \leq k_i} (\mu(V \cap T^{-(n_s - n_r)}(V)) - \mu(V)^2) + O\left(\frac{1}{k_i}\right). \end{aligned}$$

So it is enough to check that

$$\sum_{1 \leq r < s \leq k_i} |\mu(V \cap T^{-(n_s - n_r)}(V)) - \mu(V)^2| = o(k_i^2). \quad (3.2)$$

In what follows, we put

$$\gamma_{s,r} := |\mu(V \cap T^{-(n_s - n_r)}(V)) - \mu(V)^2|.$$

Since T is weakly mixing with respect to μ , there is a set $D \subseteq \mathbb{N}$ with $\text{dens}(D) = 1$ such that

$$\mu(V \cap T^{-d}(V)) - \mu(V)^2 \rightarrow 0 \quad \text{as } d \rightarrow \infty, d \in D. \quad (3.3)$$

Write

$$\sum_{1 \leq r < s \leq k_i} \gamma_{s,r} = \sum_{r=1}^{k_i} \sum_{\substack{r < s \leq k_i \\ n_s - n_r \in D}} \gamma_{s,r} + \sum_{r=1}^{k_i} \sum_{\substack{r < s \leq k_i \\ n_s - n_r \notin D}} \gamma_{s,r} =: \alpha_i + \beta_i.$$

Using equation (3.3) and since $\gamma_{r,s} \leq 1$ for all r, s , it is not hard to check that

$$\sum_{\substack{r < s \leq k_i \\ n_s - n_r \in D}} \gamma_{s,r} = o(k_i) \quad \text{as } i \rightarrow \infty, \text{ uniformly in } r;$$

and it follows that $\alpha_i = o(k_i^2)$. Moreover, since $\text{dens}(D) = 1$, we see that

$$\beta_i \leq \sum_{r=1}^{k_i} \#\{s \in (r, k_i] : n_s - n_r \notin D\} \leq k_i \times \#((0, n_{k_i}] \setminus D) = k_i \times o(n_{k_i});$$

so $\beta_i = o(k_i^2)$ since we are assuming that $n_{k_i} = O(k_i)$. This proves equation (3.2). \square

COROLLARY 3.26. *Let (X, \mathcal{B}, μ, T) be a measure-preserving dynamical system and assume that T is weakly mixing with respect to μ . Let also $A \subset \mathbb{N}$ with $\overline{\text{dens}}(A) > 0$. If $V \subset X$ is a measurable set such that $\mu(V) > 0$, then $\overline{\text{dens}}(A \cap \mathcal{N}_T(x, V)) > 0$ for μ -almost every $x \in X$.*

Proof. Let $(n_k)_{k \geq 1}$ be the increasing enumeration of A . Since $\overline{\text{dens}}(A) > 0$, one can find an increasing sequence of integers $(k_i)_{i \geq 0}$ such that $n_{k_i} = O(k_i)$. By Lemma 3.25, one can find a subsequence (k'_i) of (k_i) such that $(1/k'_i) \sum_{k=1}^{k'_i} \mathbf{1}_V \circ T^{n_k} \rightarrow \mu(V)$ μ -almost everywhere. In other words: for μ -almost every $x \in X$,

$$\frac{1}{k'_i} \#\{k \in [1, k'_i] : n_k \in \mathcal{N}_T(x, V)\} \rightarrow \mu(V).$$

Since $\#\{k \in [1, k'_i] : n_k \in \mathcal{N}_T(x, V)\} = \#([1, n_{k'_i}] \cap A \cap \mathcal{N}_T(x, V))$ and $n_{k'_i} = O(k'_i)$, it follows that $\overline{\text{dens}}(A \cap \mathcal{N}_T(x, V)) > 0$ for μ -almost every $x \in X$. \square

Proof of Theorem 3.24. Assume that the \mathbb{T} -eigenvectors of T are perfectly spanning. By [8], there exists a T -invariant Gaussian measure μ on X with full support such that T is weakly mixing with respect to μ . Let $(A_i)_{i \in I}$ be a countable family of subsets of \mathbb{N} with

positive upper density and let $(V_i)_{i \in I}$ be a family of non-empty open subsets of X . It follows immediately from Corollary 3.26 that one can find $x \in X$ (in fact, μ -almost every $x \in X$ will do) such that $\overline{\text{dens}}(A_i \cap \mathcal{N}_T(x, V_i)) > 0$ for all $i \in I$. \square

Let us point out one consequence of Theorem 3.24.

COROLLARY 3.27. *If X is a complex Banach space admitting an unconditional Schauder decomposition, then X supports a hereditarily \mathcal{U} -frequently hypercyclic operator.*

Proof. It is shown in [22] that such a space X supports an operator with a perfectly spanning set of \mathbb{T} -eigenvectors. \square

Remark 3.28. The proof of Theorem 3.24 shows the following: if $T \in \mathfrak{L}(X)$ and if there exists a T -invariant Borel measure with full support μ on X such that T is weakly mixing with respect to μ , then T is hereditarily \mathcal{U} -frequently hypercyclic. It would be interesting to know if the weak mixing assumption can be replaced by ergodicity. Incidentally, we do not know any example of an operator T admitting an ergodic measure with full support but no weakly mixing measure with full support.

4. The $T_1 \oplus T_2$ frequent hypercyclicity problem

One of the most intriguing open problems regarding frequent hypercyclicity is to decide whether $T \oplus T$ is frequently hypercyclic whenever T is frequently hypercyclic [5]. Note that, by [24], the corresponding question for \mathcal{U} -frequent hypercyclicity has a positive answer. A related problem is Question 1.1, which asks whether $T_1 \oplus T_2$ is frequently hypercyclic for every frequently hypercyclic operator T_1 and T_2 . This question is also open if we replace frequent hypercyclicity by \mathcal{U} -frequent hypercyclicity. As observed in [31], $T_1 \oplus T_2$ is hypercyclic as soon as T_1 and T_2 are \mathcal{U} -frequently hypercyclic. In the opposite direction, it seems that the best known result is [31, Theorem 7.33] which deals with infinite sums: there exists a sequence $(T_n)_{n \geq 1}$ of frequently hypercyclic operators on $\ell_p(\mathbb{N})$, $p > 1$, such that the operator $T = \bigoplus_{n \geq 1} T_n$ acting on the ℓ_p -sum $X = \bigoplus_{n \geq 1} \ell_p(\mathbb{N})$ is not \mathcal{U} -frequently hypercyclic.

As mentioned in §1, things are much simpler if we consider hereditarily frequently hypercyclic operators. We now give the detailed proof for the convenience of the reader.

PROPOSITION 4.1. *Let $\mathcal{F} \subset 2^{\mathbb{N}}$ be a Furstenberg family and let X_1, X_2 be two Polish topological vector spaces. Let also $T_1 \in \mathfrak{L}(X_1)$ and $T_2 \in \mathfrak{L}(X_2)$. If T_1 is \mathcal{F} -hypercyclic and T_2 is hereditarily \mathcal{F} -hypercyclic, then $T_1 \oplus T_2$ is \mathcal{F} -hypercyclic. If both T_1 and T_2 are hereditarily \mathcal{F} -hypercyclic, then $T_1 \oplus T_2$ is hereditarily \mathcal{F} -hypercyclic.*

Proof. Assume that T_1 is \mathcal{F} -hypercyclic and T_2 is hereditarily \mathcal{F} -hypercyclic. Let $(V_i)_{i \in I}$ be a countable basis of open sets for $X_1 \times X_2$. Without loss of generality, we may assume that each V_i has the form $V_i = V_{i,1} \times V_{i,2}$, where $V_{i,1}, V_{i,2}$ are open in X_1, X_2 . Let $x_1 \in X_1$ be any \mathcal{F} -hypercyclic vector for T_1 . Then, for each $i \in I$, the set $A_i := \mathcal{N}_{T_1}(x_1, V_{i,1})$ belongs to \mathcal{F} . Since T_2 is hereditarily \mathcal{F} -hypercyclic, it follows that one can find a vector $x_2 \in X_2$ such that $B_i := A_i \cap \mathcal{N}_{T_2}(x_2, V_{i,2}) \in \mathcal{F}$ for all $i \in I$. Then, $x := (x_1, x_2)$ is a frequently hypercyclic vector for $T := T_1 \oplus T_2$ since $\mathcal{N}_T(x, V_i) \supset B_i$ for all $i \in I$. \square

The proof of the second part of the proposition is essentially the same. \square

Using weighted backward shifts on $c_0(\mathbb{Z}_+)$, we now find a counterexample to the $T_1 \oplus T_2$ frequent hypercyclicity problem, and thus answer Question 1.1 in the negative. This counterexample also solves the $T_1 \oplus T_2\mathcal{U}$ -frequent hypercyclicity problem.

THEOREM 4.2. *There exist two frequently hypercyclic weighted shifts $B_w, B_{w'}$ on $c_0(\mathbb{Z}_+)$ such that $B_w \oplus B_{w'}$ is not \mathcal{U} -frequently hypercyclic.*

From this theorem and Proposition 4.1, we immediately deduce the following result, which is of course to be compared with Corollary 2.3.

COROLLARY 4.3. *There exist weighted shifts on $c_0(\mathbb{Z}_+)$ which are frequently hypercyclic but not hereditarily frequently hypercyclic.*

Let us also point out another consequence of Theorem 4.2 and Remark 3.28.

COROLLARY 4.4. *There exist frequently hypercyclic weighted shifts on $c_0(\mathbb{Z}_+)$ which admit no weakly mixing invariant measure with full support, and hence no ergodic invariant Gaussian measure with full support.*

This is not really a new result: the Gaussian part has been known since [6] (with arguably a more complicated example than the one we are about to present here) and it was proved in [30] that there exist frequently hypercyclic *bilateral* weighted shifts on $c_0(\mathbb{Z})$ which admit no ergodic invariant measure with full support.

In the proof of Theorem 4.2, we shall use the following lemma, which gives a simple characterization of frequent hypercyclicity for weighted shifts on $c_0(\mathbb{Z}_+)$ whose weight sequence is bounded below (see [9] or [17, Corollary 34]).

LEMMA 4.5. *Let $w = (w_n)_{n \geq 1}$ be a bounded sequence of positive real numbers and assume that $\inf_{n \geq 1} w_n > 0$. Then, the associated weighted shift B_w is frequently hypercyclic on $c_0(\mathbb{Z}_+)$ if and only if there exist a sequence $(M(p))_{p \geq 1}$ of positive real numbers tending to infinity and a sequence $(E_p)_{p \geq 1}$ of disjoint subsets of \mathbb{N} with positive lower density such that:*

- (a) $\lim_{n \rightarrow \infty, n \in E_p} w_1 \cdots w_n = \infty$ for all $p \geq 1$;
- (b) for all $p, q \geq 1$, for all $m \in E_p$ and $n \in E_q$ with $m > n$,

$$w_1 \cdots w_{m-n} \geq \max(M(p), M(q)).$$

We will also need the following elementary lemma, which is almost the same as [9, Lemma 6.1]. For $\varepsilon > 0$, $a > 1$ and $u \in \mathbb{N}$, we let

$$I_u^{a,\varepsilon} := [(1 - \varepsilon)a^u, (1 + \varepsilon)a^u].$$

LEMMA 4.6. *There exist $\varepsilon > 0$ and $a > 1$ such that, for any integers $u > v \geq 1$,*

$$I_u^{a,4\varepsilon} \cap I_v^{a,4\varepsilon} = \emptyset, \quad I_u^{a,2\varepsilon} - I_v^{a,2\varepsilon} \subset I_u^{a,4\varepsilon} \quad \text{and} \quad I_v^{a,\varepsilon} + [-v, v] \subset I_v^{a,2\varepsilon}.$$

Proof. Provided that $\varepsilon \in (0, 1/4)$, the first condition is equivalent to saying that

$$(1 + 4\varepsilon)a^u < (1 - 4\varepsilon)a^{u+1} \quad \text{for all } u \geq 1,$$

that is,

$$\frac{1 + 4\varepsilon}{(1 - 4\varepsilon)a} < 1.$$

The second one is satisfied as soon as

$$(1 - 2\varepsilon)a^u - (1 + 2\varepsilon)a^{u-1} \geq (1 - 4\varepsilon)a^u \quad \text{for all } u \geq 2,$$

which is equivalent to

$$\frac{2\varepsilon a}{1 + 2\varepsilon} \geq 1.$$

The last condition is satisfied if $(1 - \varepsilon)a^v - v \geq (1 - 2\varepsilon)a^v$ for all $v \geq 1$, in other words,

$$\varepsilon a^v \geq v \quad \text{for all } v \geq 1.$$

Therefore, one can choose, for example, $\varepsilon := 1/8$ and then take a sufficiently large. \square

Proof of Theorem 4.2. Our construction is inspired by that of [9, §6]. In what follows, we fix once and for all $\varepsilon > 0$ and $a > 1$ satisfying the conclusions of Lemma 4.6.

For $k \geq 1$, let

$$A_k := 2^{k-1}\mathbb{N} \setminus 2^k\mathbb{N}.$$

Note that each A_k is a syndetic set, that is, it has bounded gaps, and the sets A_k are pairwise disjoint. Moreover, since $2^{k-1} \geq k$, we have $I_v^{a,\varepsilon} + [-k, k] \subset I_v^{a,2\varepsilon}$ for each $k \geq 1$ and all $v \in A_k$.

We also fix an increasing sequence of positive integers $(b_p)_{p \geq 1}$ such that

$$\lim_{p \rightarrow \infty} \overline{\text{dens}} \left[\bigcup_{q \geq p} (b_q\mathbb{N} + [-q, q]) \right] = 0.$$

Finally, we set for $p \geq 1$,

$$E_p := \bigcup_{u \in A_{2p}} (I_u^{a,\varepsilon} \cap b_p\mathbb{N}),$$

$$F_p := \bigcup_{u \in A_{2p+1}} (I_u^{a,\varepsilon} \cap b_p\mathbb{N}).$$

We note that since A_{2p} and A_{2p+1} are syndetic, we have

$$\underline{\text{dens}}(E_p) > 0 \quad \text{and} \quad \underline{\text{dens}}(F_p) > 0 \quad \text{for all } p \geq 1.$$

Indeed, for all $p \geq 1$, there exists $\delta_p > 0$ such that, for all u sufficiently large,

$$\#(I_u^{a,\varepsilon} \cap b_p\mathbb{N}) \geq \delta_p a^u.$$

Let R_p be such that if u and v are two consecutive elements of A_{2p} , then $v - u \leq R_p$. If now n is very large and if we consider u and v as two consecutive elements of A_{2p} such that

$$(1 + \varepsilon)a^u < n \leq (1 + \varepsilon)a^v,$$

then we see that

$$\frac{\#(E_p \cap [1, n])}{n} \geq \frac{\#(I_u^{a,\varepsilon} \cap b_p \mathbb{N})}{(1 + \varepsilon)a^v} \geq \frac{\delta_p}{(1 + \varepsilon)a^{R_p}}.$$

Hence, $\underline{\text{dens}}(E_p) > 0$. A similar argument shows that $\underline{\text{dens}}(F_p) > 0$.

We now construct our weight sequences w and w' .

For $p \geq 1$, we first define a sequence $(w_n^p)_{n \geq 1} \subset \{1/2, 1, 2\}$ such that, for all $n \geq 1$,

$$w_1^p \cdots w_n^p = \begin{cases} 1 & \text{if } n \notin I_u^{a,2\varepsilon}, u \in A_{2p}, \\ 2^u & \text{if } n \in I_u^{a,\varepsilon}, u \in A_{2p}. \end{cases}$$

This is possible since $I_u^{a,\varepsilon} + [-u, u] \subset I_u^{a,2\varepsilon}$. These sequences will be used for handling condition (a) in Lemma 4.5.

For $p \geq 1$, we also define a sequence $(\omega_n^p)_{n \geq 1} \subset \{1/2, 1, 2\}$ such that, for all $n \geq 1$,

$$\omega_1^p \cdots \omega_n^p = \begin{cases} 1 & \text{if } n \notin b_p \mathbb{N} + [-p, p], \\ 2^p & \text{if } n \in b_p \mathbb{N}. \end{cases}$$

These sequences will help us to verify condition (b) in Lemma 4.5 for $p = q$.

Finally, for $u > v \geq 1$ with $u \in A_{2p}$ and $v \in A_{2q}$ for some $p, q \geq 1$, we define a sequence $(w_n^{u,v})_{n \geq 1} \subset \{1/2, 1, 2\}$ such that, for all $n \geq 1$,

$$w_1^{u,v} \cdots w_n^{u,v} = \begin{cases} 1 & \text{if } n \notin I_u^{a,4\varepsilon}, \\ \max(2^p, 2^q) & \text{if } n \in I_u^{a,\varepsilon} - I_v^{a,\varepsilon}. \end{cases}$$

This is possible since

$$\begin{aligned} I_u^{a,\varepsilon} - I_v^{a,\varepsilon} + [-\max(p, q), \max(p, q)] &\subset (I_u^{a,\varepsilon} + [-p, p]) - (I_v^{a,\varepsilon} + [-q, q]) \\ &\subset I_u^{a,2\varepsilon} - I_v^{a,2\varepsilon} \\ &\subset I_u^{a,4\varepsilon}. \end{aligned}$$

These sequences will be needed to check condition (b) in Lemma 4.5 for $p \neq q$.

We finally define the weight sequence $w = (w_n)_{n \geq 1}$ as follows: for all $n \geq 1$, we require that

$$w_1 \cdots w_n = \max_{p, u > v} (w_1^p \cdots w_n^p, \omega_1^p \cdots \omega_n^p, w_1^{u,v} \cdots w_n^{u,v}).$$

It is not difficult to check that $w_n \in [1/2, 2]$ for all $n \geq 1$. Indeed, assume for instance that $w_1 \cdots w_n = w_1^p \cdots w_n^p$. Then, $w_1 \cdots w_{n-1} \geq w_1^p \cdots w_{n-1}^p$ and

$$w_n \leq w_n^p \leq 2.$$

The same argument works for the other cases; for the lower bound, assume for instance that $w_1 \cdots w_{n-1} = w_1^p \cdots w_{n-1}^p$. Then, $w_1 \cdots w_n \geq w_1^p \cdots w_n^p$ so that

$$w_n \geq w_n^p \geq \frac{1}{2}.$$

We define in a similar way a weight sequence $w' = (w'_n)_{n \geq 1} \subset [1/2, 2]$, replacing everywhere A_{2p} by A_{2p+1} and A_{2q} by A_{2q+1} .

Let us first show that w and w' satisfy the conditions of Lemma 4.5, so that B_w and $B_{w'}$ are frequently hypercyclic. It is clearly enough to do that for w .

- (a) If $n \in E_p$, there is a unique $u = u(n) \in A_{2p}$ such that $n \in I_u^{a,\varepsilon}$. Then, $w_1 \cdots w_n \geq w_1^p \cdots w_n^p \geq 2^{u(n)}$, which shows that $w_1 \cdots w_n \rightarrow \infty$ as $n \rightarrow \infty$, $n \in E_p$.
- (b) Let $p, q \geq 1$, and let us fix $m \in E_p$ and $n \in E_q$ with $m > n$. If $p = q$, then $m - n \in b_p \mathbb{N}$, so that $w_1 \cdots w_{m-n} \geq w_1^p \cdots w_{m-n}^p \geq 2^p$. If $p \neq q$, there exist $u > v \geq 1$ such that $m \in I_u^{a,\varepsilon}$ and $n \in I_v^{a,\varepsilon}$, and then $w_1 \cdots w_{m-n} \geq w_1^{u,v} \cdots w_{m-n}^{u,v} \geq \max(2^p, 2^q)$.

Now, let us show that $B_w \oplus B_{w'}$ is not \mathcal{U} -frequently hypercyclic. Denote by $(e_j)_{j \geq 0}$ the canonical basis of $c_0(\mathbb{Z}_+)$. We show that for any vector $x \in c_0(\mathbb{Z}_+) \oplus c_0(\mathbb{Z}_+)$, the set $E_x := \{n \in \mathbb{N} : \|(B_w \oplus B_{w'})^n x - (e_0, e_0)\| < 1/2\}$ has upper density equal to 0.

Towards a contradiction, assume that $\overline{\text{dens}}(E_x) > 0$ for some vector x . It is easy to check that

$$\lim_{n \rightarrow \infty, n \in E_x} w_1 \cdots w_n = \lim_{n \rightarrow \infty, n \in E_x} w'_1 \cdots w'_n = \infty.$$

It follows that if we set

$$G_p := \{n \in \mathbb{N} : w_1 \cdots w_n \geq 2^p \text{ and } w'_1 \cdots w'_n \geq 2^p\},$$

then $E_x \setminus G_p$ is finite and hence $\overline{\text{dens}}(G_p) \geq \overline{\text{dens}}(E_x) > 0$ for all $p \geq 1$.

Now, the construction of the weight sequence w yields that if $n \in G_p$, then either $n \in b_q \mathbb{N} + [-q, q]$ for some $q \geq p$, or $n \in I_u^{a,4\varepsilon}$ for some $u \in \bigcup_{q \geq 1} A_{2q}$. Similarly, by construction of the sequence w' , we also know that if $n \in G_p$, then either $n \in b_q \mathbb{N} + [-q, q]$ for some $q \geq p$, or $n \in I_v^{a,4\varepsilon}$ for some $v \in \bigcup_{q \geq 1} A_{2q+1}$. By disjointness of the sets $I_u^{a,4\varepsilon}$ and $I_v^{a,4\varepsilon}$ for $u \neq v$, it follows that

$$G_p \subset \bigcup_{q \geq p} (b_q \mathbb{N} + [-q, q]).$$

By our choice of the sequence (b_p) , we get a contradiction with $\overline{\text{dens}}(G_p) \geq \overline{\text{dens}}(E_x) > 0$. \square

Remark 4.7. The weighted shift B_w cannot serve as a counterexample to the $T \oplus T$ frequent hypercyclicity problem. Indeed, it can be shown (see [32, Theorem 18]) that any weighted shift on $c_0(\mathbb{Z}_+)$ satisfying the assumptions of Lemma 4.5 is such that any finite direct sum $B_w \oplus \cdots \oplus B_w$ is itself frequently hypercyclic.

5. FHC operators on $\ell_p(\mathbb{Z}_+)$ which are not hereditarily FHC

5.1. *The result.* In this section, we use the machinery developed in [31], following the construction in [41] of chaotic operators which are not frequently hypercyclic, to produce an operator on $\ell_p(\mathbb{Z}_+)$, $1 \leq p < \infty$, which is frequently hypercyclic but not hereditarily frequently hypercyclic. We will, in fact, obtain a formally stronger result.

Definition 5.1. Let $A \subset \mathbb{N}$ be a set with $\underline{\text{dens}}(A) > 0$. We say that an operator $T \in \mathcal{L}(X)$ is *frequently hypercyclic along A* if the sequence $(T^n)_{n \in A}$ is frequently hypercyclic: there exists $x \in X$ such that $\underline{\text{dens}}(A \cap \mathcal{N}_T(x, V)) > 0$ for every non-empty open set $V \subset X$.

Obviously, if an operator is hereditarily frequently hypercyclic, then it is frequently hypercyclic along any set $A \subset \mathbb{N}$ with positive lower density. Our aim is to prove the following theorem.

THEOREM 5.2. *Let $1 \leq p < \infty$. There exist an operator T on $\ell_p(\mathbb{Z}_+)$ and a set $A \subset \mathbb{N}$ with $\underline{\text{dens}}(A) > 0$ such that T is frequently hypercyclic and chaotic, but not frequently hypercyclic along A (and thus not hereditarily frequently hypercyclic).*

With the terminology of [31], the operator we are looking for will be a $C_{+,1}$ -type operator. So we will need to recall the definition of $C_{+,1}$ -type operators, and more generally of C -type and C_+ -type operators. However, before that, we will prove a general result allowing to check in a simple way that an operator is not frequently hypercyclic along some set with positive lower density.

5.2. *How not to be hereditarily FHC.* The next theorem gives simple conditions ensuring that an operator is not hereditarily frequently hypercyclic.

THEOREM 5.3. *Let X be a Banach space admitting a Schauder basis $(e_k)_{k \geq 0}$ and let $T \in \mathcal{L}(X)$. Denoting by π_K , $K \geq 1$ the canonical projection onto $\text{span}(e_k : 0 \leq k \leq K - 1)$, assume that there exist increasing sequences of integers $(K_n)_{n \geq 1}$ and $(J_n)_{n \geq 1}$ such that for every n :*

- (a) $T^{J_n} \pi_{K_n} = \pi_{K_n}$;
- (b) $\|\pi_{K_n} T^j (I - \pi_{K_n})x\| \leq \|(I - \pi_{K_n})x\|$ for all $x \in X$ and $0 \leq j \leq (n+1)^{2^{J_n}} J_n$.

Then, there exists a set $A \subset \mathbb{N}$ with positive lower density such that T is not hereditarily frequently hypercyclic along A .

Proof. Extracting subsequences of (K_n) and (J_n) if necessary, we may assume without loss of generality that $J_{n+1} \geq 2(n+1)^{2^{J_n}} J_n$ for all n .

Let us denote by $\mathcal{F}_n \subset 2^{\mathbb{N}}$ the family of all finite sets $F \subset [0, J_n)$ such that $\#F \geq J_n/2$. Let $C_n := \#\mathcal{F}_n$ and let $(F_{n,j})_{0 \leq j < C_n}$ be an enumeration of \mathcal{F}_n . We set $M_n := (n+1)^{C_n} J_n$ and we remark that $M_n \leq (n+1)^{2^{J_n}} J_n \leq J_{n+1}/2$.

We now construct the set A by induction. To start, let

$$A_1 := [0, J_1) \cup \bigcup_{0 \leq j < C_1} \bigcup_{0 \leq l < 2^{J_1} - 2^j} ((2^j + l)J_1 + F_{1,j}).$$

Given $k \geq 2$, if A_1, \dots, A_{k-1} have been defined already, we define A_k by setting

$$A_k := [M_{k-1}, J_k) \cup \bigcup_{0 \leq j < C_k} \bigcup_{0 \leq l < (k+1)^{j+1} - (k+1)^j} ((k+1)^j + l)J_k + F_{k,j}.$$

Observe that the sets $(k+1)^j + l)J_k + F_{k,j}$ involved in this definition are pairwise disjoint, since they are contained in successive intervals. More precisely,

$$A_k \cap [sJ_k, (s+1)J_k) = sJ_k + F_{k,j}$$

for every $0 \leq j < C_k$ and every $(k+1)^j \leq s < (k+1)^{j+1}$, and $\max(A_k) < (k+1)^{C_k} J_k = M_k$. Hence, $A_k \subseteq [M_{k-1}, M_k)$. Finally, we let

$$A := \bigcup_{k \geq 1} A_k.$$

CLAIM 5.4. *We have $\#(A \cap [0, N])/(N+1) \geq 1/4$ for all $N \geq 0$. In particular, $\underline{\text{dens}}(A) > 0$.*

Proof of Claim 5.4. We will check by induction on $k \geq 1$ that

$$\frac{\#(A \cap [0, N])}{(N+1)} \geq \frac{1}{4} \quad \text{for all } N < M_k. \quad (5.1)$$

If $N < M_1$, there exists $0 \leq s < 2^{C_1}$ such that $sJ_1 \leq N < (s+1)J_1$. Then, $\#(A \cap [0, N])/(N+1) \geq 1$ if $s = 0$ and

$$\frac{\#(A \cap [0, N])}{N+1} \geq \frac{\#(A_1 \cap [0, sJ_1])}{(s+1)J_1} \geq \frac{s}{2(s+1)} \geq \frac{1}{4} \quad \text{if } s \geq 1,$$

because the sets involved in the definition of A_1 are pairwise disjoint and $\#F_{1,j} \geq J_1/2$ for every $j < C_1$.

Assume that the inequality (5.1) has been proved for $k-1$. To get the result for k , it is enough to check that

$$\frac{\#(A \cap [0, N])}{N+1} \geq \frac{1}{4} \quad \text{for every } M_{k-1} \leq N < M_k.$$

If $M_{k-1} \leq N < J_k$ then, since $[M_{k-1}, N] \subset [M_{k-1}, J_k) \subset A_k$, we get by the induction assumption that

$$\frac{\#(A \cap [0, N])}{N+1} \geq \frac{\#(A \cap [0, M_{k-1}])}{M_{k-1}} \geq \frac{1}{4}.$$

Moreover, we even have $\#(A \cap [0, J_k))/J_k \geq \frac{1}{2}$ since $[M_{k-1}, J_k) \subset A_k$ and $M_{k-1} \leq J_k/2$. However, if $J_k \leq N < M_k$, there exists $1 \leq s < (k+1)^{C_k}$ such that $sJ_k \leq N < (s+1)J_k$ and we obtain in this case that

$$\frac{\#(A \cap [0, N])}{N+1} \geq \frac{\#(A \cap [0, J_k))}{(s+1)J_k} + \frac{\#(A_k \cap [J_k, sJ_k])}{(s+1)J_k} \geq \frac{1}{2(s+1)} + \frac{s-1}{2(s+1)} \geq \frac{1}{4}.$$

This proves Claim 5.4. \square

Let us now get back to the proof of Theorem 5.3. Our aim is to show that under assumptions (a) and (b), T is not hereditarily frequently hypercyclic along the set A that we just constructed. To this end, we consider for any $c > 0$, the open sets

$$U_c := \{y \in X : |\langle e_0^*, y \rangle| < c\} \quad \text{and} \quad V_c = \{y \in X : |\langle e_0^*, y \rangle| > c\},$$

and we show that for every $x \in X$, either $\mathcal{N}_T(x, U_{1/2}) \cap A$ or $\mathcal{N}_T(x, V_{3/2}) \cap A$ has a lower density equal to 0. Let $x \in X$. Since $|\langle e_0^*, u \rangle| \leq C \|\pi_{K_n} u\|$ for some absolute constant $C > 0$, it follows from assumption (b) that for n sufficiently large, we have

$$\begin{aligned} \mathcal{N}_T(x, U_{1/2}) \cap [0, (n+1)^{2^{J_n}} J_n] &\subset \mathcal{N}_T(\pi_{K_n} x, U_{3/4}) \cap [0, (n+1)^{2^{J_n}} J_n] \\ &\subset \mathcal{N}_T(x, U_1) \cap [0, (n+1)^{2^{J_n}} J_n] \end{aligned}$$

and

$$\begin{aligned} \mathcal{N}_T(x, V_{3/2}) \cap [0, (n+1)^{2^{J_n}} J_n] &\subset \mathcal{N}_T(\pi_{K_n} x, V_{4/3}) \cap [0, (n+1)^{2^{J_n}} J_n] \\ &\subset \mathcal{N}_T(x, V_1) \cap [0, (n+1)^{2^{J_n}} J_n]. \end{aligned}$$

Moreover, since $\mathcal{N}_T(x, U_1) \cap \mathcal{N}_T(x, V_1) = \emptyset$, we have, for any $n \geq 1$, that

$$\text{either } \#(\mathcal{N}_T(x, U_1) \cap [0, J_n]) \leq J_n/2 \quad \text{or} \quad \#(\mathcal{N}_T(x, V_1) \cap [0, J_n]) \leq J_n/2.$$

Hence, either $\#(\mathcal{N}_T(x, U_1) \cap [0, J_n]) \leq J_n/2$ for infinitely many n values or $\#(\mathcal{N}_T(x, V_1) \cap [0, J_n]) \leq J_n/2$ for infinitely many n values. Without loss of generality, we assume that $\#(\mathcal{N}_T(x, U_1) \cap [0, J_n]) \leq J_n/2$ for infinitely many n values (the other case being similar). Hence, there exists an increasing sequence $(n_k)_{k \geq 1}$ of integers and a sequence $(j_k)_{k \geq 1}$ of integers such that

$$\mathcal{N}_T(x, U_1) \cap [0, J_{n_k}) \cap F_{n_k, j_k} = \emptyset \quad \text{for every } k \geq 1.$$

Now, since $T^{J_{n_k}} \pi_{K_{n_k}} = \pi_{K_{n_k}}$ by assumption (a), we have for every $k \geq 1$,

$$\begin{aligned} \mathcal{N}_T(x, U_{1/2}) \cap [0, (n_k+1)^{2^{J_{n_k}}} J_{n_k}) &\subset \mathcal{N}_T(\pi_{K_{n_k}} x, U_{3/4}) \cap [0, (n_k+1)^{2^{J_{n_k}}} J_{n_k}) \\ &= \bigcup_{0 \leq l < (n_k+1)^{2^{J_{n_k}}}} (l J_{n_k} + \mathcal{N}_T(\pi_{K_{n_k}} x, U_{3/4}) \cap [0, J_{n_k})) \\ &\subset \bigcup_{0 \leq l < (n_k+1)^{2^{J_{n_k}}}} (l J_{n_k} + \mathcal{N}_T(x, U_1) \cap [0, J_{n_k})). \end{aligned}$$

Intersecting with A and observing that $(n_k+1)^{j_k+1} \leq (n_k+1)^{2^{J_{n_k}}}$, we get that

$$\begin{aligned} (\mathcal{N}_T(x, U_{1/2}) \cap A) \cap [0, (n_k+1)^{j_k+1} J_{n_k}) &\subset A \cap \bigcup_{0 \leq s < (n_k+1)^{j_k+1}} (s J_{n_k} + \mathcal{N}_T(x, U_1) \cap [0, J_{n_k})). \end{aligned}$$

Now, by definition of A , we have

$$A \cap \bigcup_{(n_k+1)^{j_k} \leq s < (n_k+1)^{j_k+1}} [sJ_{n_k}, (s+1)J_{n_k}) = \bigcup_{(n_k+1)^{j_k} \leq s < (n_k+1)^{j_k+1}} (sJ_{n_k} + F_{n_k, j_k}).$$

Since $\mathcal{N}_T(x, U_1) \cap [0, J_{n_k}) \cap F_{n_k, j_k} = \emptyset$, it follows that

$$(\mathcal{N}_T(x, U_{1/2}) \cap A) \cap [(n_k+1)^{j_k} J_{n_k}, (n_k+1)^{j_k+1} J_{n_k}) = \emptyset,$$

so that

$$(\mathcal{N}_T(x, U_{1/2}) \cap A) \cap [0, (n_k+1)^{j_k+1} J_{n_k}) \subset [0, (n_k+1)^{j_k} J_{n_k}).$$

So we see that

$$\frac{\#((\mathcal{N}_T(x, U_{1/2}) \cap A) \cap [0, (n_k+1)^{j_k+1} J_{n_k}))}{(n_k+1)^{j_k+1} J_{n_k}} \leq \frac{(n_k+1)^{j_k} J_{n_k}}{(n_k+1)^{j_k+1} J_{n_k}}.$$

The right-hand side of this inequality tends to 0 as n tends to infinity, and this shows that $\text{dens}(\mathcal{N}_T(x, U_{1/2}) \cap A) = 0$. \square

Remark 5.5. Assumption (b) in Theorem 5.3 can be weakened: it is enough to assume that there exists a non-zero linear functional $x^* \in X^*$ such that $|\langle x^*, T^j(I - \pi_{K_n})x \rangle| \leq \|(I - \pi_{K_n})x\|$ for all $x \in X$ and $j \leq (n+1)^{2^{j_n}} J_n$. This is apparent from the above proof.

5.3. C -type operators. We recall here very succinctly some basic facts concerning C -type operators, and we refer the reader to [31, §§6 and 7] for more on this class of operators. In what follows, we denote by $(e_k)_{k \geq 0}$ the canonical basis of $\ell_p(\mathbb{Z}_+)$, $1 \leq p < \infty$.

Let us consider four ‘parameters’ v, w, φ and b , where:

- $v = (v_n)_{n \geq 1}$ is a sequence of non-zero complex numbers such that $\sum_{n \geq 1} |v_n| < \infty$;
- $w = (w_j)_{j \geq 1}$ is a sequence of complex numbers such that $0 < \inf_{k \geq 1} |w_k| \leq \sup_{k \geq 1} |w_k| < \infty$;
- φ is a map from \mathbb{Z}_+ into itself, such that $\varphi(0) = 0$, $\varphi(n) < n$ for every $n \geq 1$, and the set $\varphi^{-1}(l) = \{n \geq 0; \varphi(n) = l\}$ is infinite for every $l \geq 0$;
- $b = (b_n)_{n \geq 0}$ is a strictly increasing sequence of positive integers such that $b_0 = 0$ and $b_{n+1} - b_n$ is a multiple of $2(b_{\varphi(n)+1} - b_{\varphi(n)})$ for every $n \geq 1$.

If w and b are such that

$$\inf_{n \geq 0} \prod_{b_n < j < b_{n+1}} |w_j| > 0,$$

then, by [31, Lemma 6.2], there is a unique bounded operator $T_{v, w, \varphi, b}$ on $\ell_p(\mathbb{Z}_+)$ such that

$$T_{v, w, \varphi, b} e_k = \begin{cases} w_{k+1} e_{k+1} & \text{if } k \in [b_n, b_{n+1} - 1), n \geq 0, \\ v_n e_{b_{\varphi(n)}} - \left(\prod_{j=b_n+1}^{b_{n+1}-1} w_j \right)^{-1} e_{b_n} & \text{if } k = b_{n+1} - 1, n \geq 1, \\ - \left(\prod_{j=b_0+1}^{b_1-1} w_j \right)^{-1} e_0 & \text{if } k = b_1 - 1. \end{cases}$$

Any such operator $T_{v, w, \varphi, b}$ is called a *C-type operator*. A notable fact to be pointed out immediately is that C-type operators have lots of periodic points; indeed, we have the following fact, which is [31, Lemma 6.4].

Fact 5.6. If $T = T_{v, w, \varphi, b}$ is a C-type operator, then

$$T^{2(b_{n+1}-b_n)} e_k = e_k \quad \text{if } k \in [b_n, b_{n+1}), n \geq 0.$$

It follows that every finitely supported vector is periodic for $T_{v, w, \varphi, b}$; in particular, a C-type operator is chaotic as soon as it is hypercyclic.

A *C₊-type operator* is a C-type operator for which the parameters satisfy the following additional conditions: for every $k \geq 1$:

- φ is increasing on each interval $[2^{k-1}, 2^k)$ with $\varphi([2^{k-1}, 2^k)) = [0, 2^{k-1})$, that is,

$$\varphi(n) = n - 2^{k-1} \quad \text{for every } n \in [2^{k-1}, 2^k);$$

- the blocks $[b_n, b_{n+1}), n \in [2^{k-1}, 2^k)$ all have the same size, which we denote by $\Delta^{(k)}$:

$$b_{n+1} - b_n = \Delta^{(k)} \quad \text{for every } n \in [2^{k-1}, 2^k);$$

- the sequence v is constant on the interval $[2^{k-1}, 2^k)$: there exists $v^{(k)}$ such that

$$v_n = v^{(k)} \quad \text{for every } n \in [2^{k-1}, 2^k);$$

- the sequences of weights $(w_{b_n+i})_{1 \leq i < \Delta^{(k)}}$ are independent of $n \in [2^{k-1}, 2^k)$: there exists a sequence $(w_i^{(k)})_{1 \leq i < \Delta^{(k)}}$ such that

$$w_{b_n+i} = w_i^{(k)} \quad \text{for every } n \in [2^{k-1}, 2^k) \text{ and } 1 \leq i < \Delta^{(k)}.$$

Finally, a *C_{+,1}-type operator* is a C₊-type operator whose parameters are such that for all $k \geq 1$,

$$v^{(k)} = 2^{-\tau^{(k)}} \quad \text{and} \quad w_i^{(k)} = \begin{cases} 2 & \text{if } 1 \leq i \leq \delta^{(k)}, \\ 1 & \text{if } \delta^{(k)} < i < \Delta^{(k)}, \end{cases}$$

where $(\tau^{(k)})_{k \geq 1}$ and $(\delta^{(k)})_{k \geq 1}$ are two increasing sequences of integers with $\delta^{(k)} < \Delta^{(k)}$ for each $k \geq 1$.

These operators have been studied in detail in [31, §7]. In particular, we have the following crucial fact [31, Theorem 7.1].

Fact 5.7. A $C_{+,1}$ -type operator $T_{v,w,\varphi,b}$ is frequently hypercyclic as soon as

$$\limsup_{k \rightarrow \infty} \frac{\delta^{(k)} - \tau^{(k)}}{\Delta^{(k)}} > 0. \quad (5.2)$$

5.4. Proof of Theorem 5.2. Let $T = T_{v,w,\varphi,b}$ be an operator of $C_{+,1}$ -type on $\ell_p(\mathbb{Z}_+)$, so that v and w are given by

$$v^{(k)} = 2^{-\tau^{(k)}} \quad \text{and} \quad w_i^{(k)} = \begin{cases} 2 & \text{if } 1 \leq i \leq \delta^{(k)}, \\ 1 & \text{if } \delta^{(k)} < i < \Delta^{(k)}. \end{cases}$$

We assume that $\Delta^{(k)} \in 8\mathbb{N}$ for all $k \geq 1$, and we choose

$$\delta^{(k)} := \frac{1}{4}\Delta^{(k)} \quad \text{and} \quad \tau^{(k)} := \frac{1}{8}\Delta^{(k)}.$$

So the only ‘free’ parameter is now the sequence $(\Delta^{(k)})_{k \geq 1}$.

By Fact 5.7, the operator T is frequently hypercyclic (and hence chaotic since it is a C -type operator). So we just have to show that if the sequence $(\Delta^{(k)})$ is suitably chosen, then T satisfies the assumptions of Theorem 5.3. We will in fact show that this holds as soon as the sequence $(\Delta^{(k)})$ grows sufficiently rapidly. Let us set

$$K_n := b_{2^n} \quad \text{and} \quad J_n := 2\Delta^{(n)} \quad \text{for every } n \geq 1.$$

With this choice of the sequences (K_n) and (J_n) , condition (a) in Theorem 5.3 is satisfied by Fact 5.6. So the only thing to check is condition (b).

Let $\gamma_k := 2^{\delta^{(k-1)} - \tau^{(k)}} (\Delta^{(k)})^{1-1/p}$. If $(\Delta^{(k)})$ grows sufficiently rapidly, then the sequence (γ_k) is decreasing and

$$2^n \sum_{k \geq n+1} 2^{k-1} \gamma_k \leq 1 \quad \text{for all } n \geq 0.$$

Let us also define a sequence $(\beta_l)_{l \geq 1}$ as follows:

$$\beta_l := 4\gamma_k \quad \text{if } l \in [2^{k-1}, 2^k).$$

Finally, for any $l \geq 1$, let P_l be the projection of $\ell_p(\mathbb{Z}_+)$ defined by

$$P_l x = \sum_{k=b_l}^{b_{l+1}-1} x_k e_k \quad \text{for every } x \in \ell_p(\mathbb{Z}_+).$$

As in the proof of [31, Theorem 7.2], one can show that the following estimate holds for every $k \geq 0$, every $l \in [2^{k-1}, 2^k[$, every $0 \leq m < l$ and every $0 \leq j \leq \Delta^{(k)} - \delta^{(k)} = \frac{3}{4}\Delta^{(k)}$:

$$\|P_m T^j P_l x\| \leq \frac{\beta_l}{4} \left(\prod_{i=\Delta^{(k)}-j+1}^{\Delta^{(k)}-1} |w_i^{(k)}| \right) \|P_l x\| \leq \frac{\beta_l}{4} \|P_l x\|.$$

Hence, we have for all n and $j \leq \frac{3}{4}\Delta^{(n+1)}$,

$$\begin{aligned}\|\pi_{K_n} T^j (I - \pi_{K_n})x\| &\leq \sum_{m < 2^n} \sum_{l \geq 2^n} \|P_m T^j P_l x\| \\ &\leq \sum_{m < 2^n} \sum_{l \geq 2^n} \frac{\beta_l}{4} \|P_l x\| \\ &\leq 2^n \left(\sum_{l \geq 2^n} \frac{\beta_l}{4} \right) \|(I - \pi_{K_n})x\| \\ &\leq 2^n \left(\sum_{k \geq n+1} 2^{k-1} \gamma_k \right) \|(I - \pi_{K_n})x\| \leq \|(I - \pi_{K_n})x\|.\end{aligned}$$

So, if we take care to ensure that $\frac{3}{4}\Delta^{(n+1)} \geq (n+1)^{2^{J_n}} J_n = (n+1)^{2^{2\Delta^{(n)}}} 2\Delta^{(n)}$ for all $n \geq 1$, then condition (b) in Theorem 5.3 is satisfied. This concludes the proof of Theorem 5.2.

6. Extending frequently d -hypercyclic tuples

Let us recall the definition of d - \mathcal{F} -hypercyclicity, for a given Furstenberg family $\mathcal{F} \subset 2^{\mathbb{N}}$: a tuple of operators (T_1, \dots, T_N) is d - \mathcal{F} -hypercyclic if there exists $x \in X$ such that $x \oplus \dots \oplus x$ is \mathcal{F} -hypercyclic for $T_1 \oplus \dots \oplus T_N$.

In this section, our aim is to prove the following result, which is a natural analogue of [39, Theorem 2.1] for d - \mathcal{F} -hypercyclicity. Let us denote by SOT the *strong operator topology* on $\mathcal{L}(X)$, that is, the topology of pointwise convergence.

THEOREM 6.1. *Let $\mathcal{F} \subset 2^{\mathbb{N}}$ be a Furstenberg family and let X be a Banach space supporting a hereditarily \mathcal{F} -hypercyclic operator. Let $T_1, \dots, T_N \in \mathcal{L}(X)$ and assume that (T_1, \dots, T_N) is d - \mathcal{F} -hypercyclic. Then, for any countable and linearly independent set $Z \subset d$ - \mathcal{F} -HC(T_1, \dots, T_N), the set*

$$\{T \in \mathcal{L}(X) : Z \subset d\text{-}\mathcal{F}\text{-HC}(T_1, \dots, T_N, T)\}$$

is SOT-dense in $\mathcal{L}(X)$.

Applying this result to $Z = \{x\}$ with $x \in d$ - \mathcal{F} -HC(T_1, \dots, T_N), we get the following corollary.

COROLLARY 6.2. *Let X be a Banach space supporting a hereditarily frequently hypercyclic operator. Let $T_1, \dots, T_N \in \mathcal{L}(X)$ and assume that (T_1, \dots, T_N) is d -frequently hypercyclic. Then, there exists $T_{N+1} \in \mathcal{L}(X)$ such that (T_1, \dots, T_{N+1}) is d -frequently hypercyclic.*

If (T_1, \dots, T_N) is *densely* d - \mathcal{F} -hypercyclic then, applying Theorem 6.1 with any dense linearly independent set $Z \subset X$ contained in d - \mathcal{F} -HC(T_1, \dots, T_N), we obtain the following corollary.

COROLLARY 6.3. *Let $\mathcal{F} \subset 2^{\mathbb{N}}$ be a Furstenberg family and let X be a Banach space supporting a hereditarily \mathcal{F} -hypercyclic operator. Let $T_1, \dots, T_N \in \mathcal{L}(X)$ and assume that (T_1, \dots, T_N) is *densely* d - \mathcal{F} -hypercyclic. Then, the set*

$$\{T \in \mathfrak{L}(X) : (T_1, \dots, T_N, T) \text{ is densely } d\text{-}\mathcal{F}\text{-hypercyclic}\}$$

is SOT-dense in $\mathfrak{L}(X)$.

In the proof of Theorem 6.1, we will need the following fact (already mentioned in §1).

PROPOSITION 6.4. *If $T \in \mathfrak{L}(X)$ is hereditarily \mathcal{F} -hypercyclic, then it is in fact densely hereditarily \mathcal{F} -hypercyclic: given a countable family $(A_i)_{i \in I} \subset \mathcal{F}$ and a family $(V_i)_{i \in I}$ of non-empty open sets in X , there is a dense set of $x \in X$ such that $\mathcal{N}_T(x, V_i) \cap A_i \in \mathcal{F}$ for all $i \in I$.*

Proof. Let U be a non-empty open set in X : we want to find $x \in U$ such that $\mathcal{N}_T(x, V_i) \cap A_i \in \mathcal{F}$ for all $i \in I$. For $(i, N) \in I \times \mathbb{N}$, define $V_{i,N} := T^{-N}(V_i)$ and $A_{i,N} := A_i$. Since T is hereditarily \mathcal{F} -hypercyclic, one can find $x_0 \in X$ and sets $B_{i,N} \in \mathcal{F}$ such that $B_{i,N} \subset A_{i,N} = A_i$ and $T^n x_0 \in V_{i,N}$ for all $n \in B_{i,N}$. Since we may assume that the family (V_i) is a basis of open sets for X , the vector x_0 is in particular a hypercyclic vector for T . So one can find an integer N_U such that $x := T^{N_U} x_0 \in U$. Then, for all $n \in B_{i,N_U}$, we see that $T^n x = T^{N_U} (T^n x_0) \in V_i$. \square

Proof of Theorem 6.1. Let us denote by $\text{GL}(X)$ the set of all invertible operators on X . The core of the proof is contained in the following fact.

FACT 6.5. *Let $T_1, \dots, T_N \in \mathfrak{L}(X)$ and assume that (T_1, \dots, T_N) is $d\text{-}\mathcal{F}$ -hypercyclic. Let $T \in \mathfrak{L}(X)$ be a hereditarily \mathcal{F} -hypercyclic operator. For any countable and linearly independent set $Z \subset d\text{-}\mathcal{F}\text{-HC}(T_1, \dots, T_N)$, for any $S \in \text{GL}(X)$ and any $\varepsilon > 0$, one can find $L \in \text{GL}(X)$ such that $\|L - S\| < \varepsilon$ and $Z \subset d\text{-}\mathcal{F}\text{-HC}(T_1, \dots, T_N, L^{-1}TL)$.*

Proof of Fact 6.5. Without loss of generality, we assume that Z is infinite; we enumerate Z as a sequence $(z_l)_{l \in \mathbb{N}}$, without repetition. Let us fix $S \in \text{GL}(X)$ and $\varepsilon > 0$. Since $\text{GL}(X)$ is $\|\cdot\|$ -open in $\mathfrak{L}(X)$, we may assume that any operator $L \in \mathfrak{L}(X)$ such that $\|L - S\| < \varepsilon$ is invertible.

Let $(W_p)_{p \in \mathbb{N}}$ be a countable basis of open sets for $X^{N+1} = X^N \times X$, and assume that each set W_p has the form $W_p = U_p \times V_p$ where U_p is open in X^N and V_p is open in X . For each $(l, p) \in \mathbb{N} \times \mathbb{N}$, we may fix a set $A_{l,p} \in \mathcal{F}$ such that

$$(T_1^n z_l, \dots, T_N^n z_l) \in U_p \quad \text{for all } n \in A_{l,p}.$$

We construct, by induction, a sequence $(L_l)_{l \geq 0}$ in $\mathfrak{L}(X)$ with $L_0 = S$, a sequence $(x_l)_{l \geq 1}$ of vectors of X and a family $(B_{l,p})_{l,p \geq 1}$ of sets in \mathcal{F} , such that the following holds true for every $l, p \geq 1$:

- (i) $B_{l,p} \subset A_{l,p}$ and $T^n x_l \in V_p$ for all $n \in B_{l,p}$;
- (ii) $L_l(z_s) = x_s$ for all $s \leq l$;
- (iii) $\|L_l - L_{l-1}\| < 4^{-l} \varepsilon$.

The inductive step is as follows. Choose a linear functional $v_l^* \in X^*$ such that $v_l^*(z_s) = 0$ for all $s < l$ and $v_l^*(z_l) = 1$, which is possible by linear independence of Z .

Next, since T is *densely* hereditarily \mathcal{F} -hypercyclic by Proposition 6.4, we can find a vector $x_l \in X$ and sets $B_{l,p} \subset A_{l,p}$ with $B_{l,p} \in \mathcal{F}$ for each $p \geq 1$, such that

$$\|x_l - L_{l-1}(z_l)\| < \frac{\varepsilon}{4^l \|v_l^*\|} \quad \text{and} \quad T^n x_l \in V_p \text{ for all } n \in B_{l,p}.$$

Then, define $L_l := L_{l-1} + v_l^* \otimes (x_l - L_{l-1}(z_l)) \in \mathfrak{L}(X)$, that is,

$$L_l(x) = L_{l-1}(x) + v_l^*(x) (x_l - L_{l-1}(z_l)).$$

Clearly, $L_l(z_s) = L_{l-1}(z_s)$ for all $s < l$, so that $L_l(z_s) = x_s$ by the induction hypothesis, $L_l(z_l) = x_l$ and $\|L_l - L_{l-1}\| < 4^{-l}\varepsilon$.

By properties (ii) and (iii), the sequence (L_l) converges to some $L \in \mathfrak{L}(X)$, which satisfies $L(z_l) = x_l$ for all $l \in \mathbb{N}$ and $\|L - S\| < \varepsilon$; in particular, L is invertible. Moreover, $T^n x_l \in V_p$ for all $l, p \geq 1$ and $n \in B_{l,p}$. Since $B_{l,p} \subset A_{l,p}$, it follows that

$$(T_1^n z_l, \dots, T_N^n z_l, (L^{-1}TL)^n z_l) \in U_p \times L^{-1}(V_p) =: \widetilde{W}_p \quad \text{for all } n \in B_{p,l}.$$

Now, $(\widetilde{W}_p)_{p \in \mathbb{N}}$ is a basis of the topology of $X^{N+1} = X^N \times X$ because $I \oplus L^{-1}$ is a homeomorphism of X^{N+1} ; so we see that $z_l \in d\text{-}\mathcal{F}\text{-HC}(T_1, \dots, T_N, L^{-1}TL)$ for each $l \geq 1$. \square

To conclude the proof of Theorem 6.1, we observe that since the operator T is hypercyclic, the similarity orbit of T , that is, the set $\{S^{-1}TS : S \in \text{GL}(X)\}$, is SOT-dense in $\mathfrak{L}(X)$; see for example [7, Proposition 2.20]. By Fact 6.5, it follows that the set

$$\{L^{-1}TL : Z \subset d\text{-}\mathcal{F}\text{-HC}(T_1, \dots, T_N, L^{-1}TL), L \in \text{GL}(X)\}$$

is SOT-dense in $\mathfrak{L}(X)$. \square

Remark 6.6. Our proof of Theorem 6.1 differs from that of [39, Theorem 2.1] regarding d -hypercyclicity, where a Baire category argument was used; and it must be so at least for d -frequent hypercyclicity, since $\text{FHC}(T)$ is always meagre in X , for any operator $T \in \mathfrak{L}(X)$ (see [43] or [9]). However, there may be a Baire category proof of Theorem 6.1 when \mathcal{F} is the family of sets with positive upper density (or, more generally, an ‘upper’ Furstenberg family in the sense of [17]).

To apply Theorem 6.1, in the frequently hypercyclic case, it would be nice to exhibit a class of Banach spaces as large as possible supporting hereditarily frequently hypercyclic operators. It is easy to see that any Banach space with a symmetric Schauder basis has this property: it suffices to take $T := 2B$, where B is the backward shift associated with the basis. In view of Corollary 3.27, a natural (and much broader) class would be that of complex Banach spaces admitting an unconditional Schauder decomposition, but we are not able to prove that every such space has the required property. In any event, we can use the method of [54] to prove the existence of d -frequently hypercyclic tuples of arbitrary length for this class of spaces.

PROPOSITION 6.7. *Let X be a complex separable infinite-dimensional Banach space with an unconditional Schauder decomposition. For any $N \geq 1$, there exist $T_1, \dots, T_N \in \mathfrak{L}(X)$ such that (T_1, \dots, T_N) is d -frequently hypercyclic.*

Proof. By [22], X supports an operator T with a perfectly spanning set of \mathbb{T} -eigenvectors. Then, $T \oplus \cdots \oplus T$ has the same property and, in particular, it is frequently hypercyclic; let $x_1 \oplus \cdots \oplus x_N$ be a frequently hypercyclic vector for $T \oplus \cdots \oplus T$. Now, let $y \in X \setminus \{0\}$ be arbitrary. Since $\text{GL}(X)$ acts transitively on X , we may choose $S_1, \dots, S_N \in \text{GL}(X)$ such that $S_i(x_i) = y$ for $i = 1, \dots, N$. Then, setting $T_i := S_i T S_i^{-1}$, we see that $y \in d\text{-FHC}(T_1, \dots, T_N)$. \square

7. Frequent d -hypercyclicity versus dense frequent d -hypercyclicity

Despite the similarity of the definitions, there are strong differences between hypercyclicity and d -hypercyclicity. For instance, if $T \in \mathcal{L}(X)$ is hypercyclic, then $\text{HC}(T)$ is always dense in X , and $\text{HC}(T) \cup \{0\}$ contains a dense linear subspace of X . In contrast, for two operators T_1 and T_2 , the set $d\text{-HC}(T_1, T_2) \cup \{0\}$ may be equal to some finite-dimensional subspace (see [51, Theorem 3.4]). In particular, d -hypercyclic tuples are not necessarily ‘densely d -hypercyclic’.

However, since frequent d -hypercyclicity is a strong form of d -hypercyclicity, it is natural to ask whether some properties that are not true for d -hypercyclic tuples might be true for d -frequently hypercyclic tuples. In this spirit, the following question was asked in [36–38].

Question 7.1. Let (T_1, T_2) be a d -frequently hypercyclic pair of operators on a Banach space X . Is (T_1, T_2) necessarily densely d -hypercyclic?

The next theorem provides a solution to this problem.

THEOREM 7.2. *There exist a Banach space X and $T_1, T_2 \in \mathcal{L}(X)$ such that (T_1, T_2) is d -frequently hypercyclic but not densely d -hypercyclic.*

Our proof is inspired by [51], where the authors construct a d -hypercyclic pair which is not densely d -hypercyclic. A key role will be played by the similarity orbit of some well-chosen operator T . The next two lemmas point out the relationship between (frequently) hypercyclic vectors of T and $T \oplus T$, and (frequently) d -hypercyclic vectors of (T_1, T_2) when T_1 and T_2 belong to the similarity orbit of T .

LEMMA 7.3. *Let $T \in \mathcal{L}(X)$ and $L_1, L_2 \in \text{GL}(X)$, and set $T_m := L_m^{-1} T L_m$, $m = 1, 2$. Let also $x \in X$. If $x \in d\text{-HC}(T_1, T_2)$, then $L_2 x - L_1 x \in \text{HC}(T)$.*

Proof. This is [51, Lemma 3.1]. \square

LEMMA 7.4. *Let $T \in \mathcal{L}(X)$ and $L_1, L_2 \in \text{GL}(X)$, and set $T_m := L_m^{-1} T L_m$. Let also $\mathcal{F} \subset 2^{\mathbb{N}}$ be a Furstenberg family and let $x \in X$. Then, $x \in d\text{-}\mathcal{F}\text{-HC}(T_1, T_2)$ if and only if $(L_1 x, L_2 x) \in \mathcal{F}\text{-HC}(T \oplus T)$.*

Proof. It is identical to the proof of [51, Lemma 2.1]. Just observe that for any positive integer n and any pair of non-empty open subsets (U, V) in X ,

$$(T^n L_1 x, T^n L_2 x) \in U \times V \iff (T_1^n x, T_2^n x) \in L_1^{-1}(U) \times L_2^{-1}(V),$$

and apply the definition of \mathcal{F} -hypercyclicity. \square

The operators T_1 and T_2 that we are going to construct will be such that $T_2 = (cI + R)^{-1}T_1(cI + R)$ for some $R \in \mathfrak{L}(X)$ and $c > 0$. By Lemma 7.3 (with $T = T_1$ and $L_1 = cI$), any $x \in d\text{-HC}(T_1, T_2)$ is such that $Rx \in \text{HC}(T_1)$. We shall construct T_1 and R in such a way that this condition prevents $d\text{-HC}(T_1, T_2)$ from being dense in X . The following result will be useful to prove that the pair (T_1, T_2) is d -frequently hypercyclic.

LEMMA 7.5. *Let $T_1, R \in \mathfrak{L}(X)$ and $c > 0$ be such that $L_2 = cI + R$ is invertible, and let $T_2 := L_2^{-1}T_1L_2$. Let also $\mathcal{F} \subset 2^{\mathbb{N}}$ be a Furstenberg family. If $x \in X$ is such that $(x, Rx) \in \mathcal{F}\text{-HC}(T_1 \oplus T_1)$, then $x \in d\text{-}\mathcal{F}\text{-HC}(T_1, T_2)$.*

Proof. Let us set $L_1 := cI$. By Lemma 7.4, it suffices to show that the condition $(x, Rx) \in \mathcal{F}\text{-HC}(T_1 \oplus T_1)$ implies $(L_1x, L_2x) \in \mathcal{F}\text{-HC}(T_1 \oplus T_1)$. Now it is clear that $(cx, Rx) \in \mathcal{F}\text{-HC}(T_1 \oplus T_1)$. Let U, V be two non-empty open subsets of X , and let $U' \subset U$ and W be non-empty open sets such that $U' + W \subset V$. There exists a set $A \in \mathcal{F}$ such that $T_1^n(cx) \in U'$ and $T_1^n(Rx) \in W$ for all $n \in A$. Then, for all $n \in A$, we have

$$T_1^n L_1 x = T_1^n(cx) \in U' \quad \text{and} \quad T_1^n L_2 x = T_1^n(cx + Rx) \in U' + W \subset V. \quad \square$$

We now go into the details of the construction. First, we define the Banach space X as

$$X := \left(\bigoplus_{l \geq 1} X(l) \right)_{c_0} \quad \text{where } X(l) = \ell_1(\mathbb{Z}_+) \text{ for every } l \geq 1.$$

(Following a standard notation, the subscript ' c_0 ' indicates that the direct sum is a c_0 -sum.)

Next, we introduce the following operator $T \in \mathfrak{L}(X)$: denoting by B the canonical backward shift on $\ell_1(\mathbb{Z}_+)$, let

$$T := \bigoplus_{l \geq 1} T(l), \quad \text{where } T(l) = I + 2^{-l}B \in \mathfrak{L}(X(l)) \text{ for every } l \geq 1.$$

LEMMA 7.6. *The operator $T \oplus T$ is frequently hypercyclic.*

Proof. It is enough to prove that T has a perfectly spanning set of \mathbb{T} -eigenvectors. Indeed, $T \oplus T$ will have the same property and hence will be frequently hypercyclic. \square

We now define our first operator T_1 :

$$T_1 := \bigoplus_{l \geq 1} (I + B) \in \mathfrak{L}(X).$$

Note that the same proof as that of Lemma 7.6 shows that $T_1 \oplus T_1$ is frequently hypercyclic. However, we will use the above operator T to produce a frequently hypercyclic vector for $T_1 \oplus T_1$ with specific properties.

In what follows, we denote by $(e_k(l))_{k \geq 0}$ the canonical basis of the l th component $X(l) = \ell_1(\mathbb{Z}_+)$ of X and by $(e_k^*(l))_{k \geq 0}$ the associated sequence of coordinate functionals, which we will consider as linear functionals on X . A vector $x \in X$ will be written as $x = (x(l))_{l \geq 1}$ and we will use the notation $x_k(l) = \langle e_k^*(l), x \rangle$ for every $k \geq 0$.

LEMMA 7.7. *There exists $(u, v) \in \text{FHC}(T_1 \oplus T_1)$ such that $u_0(1) \neq 0$ and $|v_k(l)| \leq 2^{-lk}$ for all $k \geq 0$ and all $l \geq 1$.*

Proof. For $l \geq 1$, Let us consider the diagonal operator $D(l)$ on $X(l)$ defined by

$$D(l)(x(l)) := \sum_{k \geq 0} 2^{-lk} x_k(l) e_k(l)$$

and set

$$D := \bigoplus_{l \geq 1} D(l) \in \mathfrak{L}(X).$$

It is easy to check that $(I + B)D(l) = D(l)(I + 2^{-l}B)$ for each $l \geq 1$, so that $T_1 D = D T$. Moreover, the operator D has dense range. So T_1 is a quasi-factor of T with quasi-factoring map D and, hence, $T_1 \oplus T_1$ is a quasi-factor of $T \oplus T$ with quasi-factoring map $D \oplus D$. Let $(x, y) \in \text{FHC}(T \oplus T)$ with $\|y\| \leq 1$ and $x_0(1) \neq 0$, and let us set $(u, v) := (Dx, Dy)$. Then, $(u, v) \in \text{FHC}(T_1 \oplus T_1)$. Moreover, $u_0(1) = x_0(1) \neq 0$ and, for $k \geq 0$ and $l \geq 1$,

$$|v_k(l)| = |2^{-lk} y_k(l)| \leq 2^{-lk}. \quad \square$$

We now give a result which provides a condition preventing a vector from being hypercyclic for T_1 .

LEMMA 7.8. *Let $x \in X$. Assume that there exist $l \geq 1$ and $\lambda \neq 0$ such that, for all $k \geq 1$ sufficiently large, $\Re \langle e_k^*(l), x/\lambda \rangle \geq 0$. Then, $x \notin \text{HC}(T_1)$.*

Proof. Since $x \in \text{HC}(T_1)$ if and only if $x/\lambda \in \text{HC}(T_1)$, we may assume $\lambda = 1$. Now, if $\Re \langle x_k(l), x \rangle = \Re \langle e_k^*(l), x \rangle \geq 0$ for all sufficiently large k , then the arguments of [51] (see the proof of Claim 1, p. 845) show that either $x(l)$ is finitely supported or

$$\Re \left\langle e_0^*(l), (I + B)^n(x(l)) \right\rangle \geq 0 \quad \text{for all sufficiently large } n.$$

Therefore, $x(l)$ cannot be a hypercyclic vector for $I + B$ and, hence, $x \notin \text{HC}(T_1)$. \square

Let us fix a sequence of positive real numbers $(\varepsilon_l)_{l \geq 1}$ going to zero. Let also $V : \ell_1(\mathbb{Z}_+) \rightarrow \ell_1(\mathbb{Z}_+)$ be the (bounded) operator defined by

$$V y := \sum_{k \geq 1} \left(\sum_{j \geq 1} 2^{-jk} y_j \right) e_k \quad \text{for every } y \in \ell_1(\mathbb{Z}_+)$$

and let $R_0 : X \rightarrow X$ be the (bounded) operator on X defined by

$$R_0(x) := (\varepsilon_1 V(x(1)), \dots, \varepsilon_l V(x(1)), \dots).$$

The operator R_0 satisfies the following crucial estimates.

LEMMA 7.9. *Let $x \in X$ and $m \geq 1$ be such that $\langle e_m^*(1), x \rangle \neq 0$ and $\langle e_j^*(1), x \rangle = 0$ for $1 \leq j < m$. Then, there exists $\delta : \mathbb{Z}_+ \rightarrow \mathbb{R}_+$ such that $\delta(k) \rightarrow 0$ as $k \rightarrow \infty$ and such that for all $l \geq 1$ and $k \geq 0$, we can write*

$$\langle e_k^*(l), R_0(x) \rangle = \varepsilon_l \langle e_m^*(1), x \rangle 2^{-mk} (1 + \delta(k)).$$

Proof. We just write

$$\begin{aligned}\langle e_k^*(l), R_0(x) \rangle &= \varepsilon_l \sum_{j=1}^{\infty} 2^{-jk} \langle e_j^*(1), x \rangle \\ &= \varepsilon_l \sum_{j=m}^{\infty} 2^{-jk} \langle e_j^*(1), x \rangle \\ &= \varepsilon_l 2^{-mk} \langle e_m^*(1), x \rangle \left(1 + \sum_{j=1}^{\infty} 2^{-jk} \frac{\langle e_{j+m}^*(1), x \rangle}{\langle e_m^*(1), x \rangle} \right);\end{aligned}$$

and we conclude because

$$\left| \sum_{j=1}^{\infty} 2^{-jk} \frac{\langle e_{j+m}^*(1), x \rangle}{\langle e_m^*(1), x \rangle} \right| \leq \frac{2^{-k}}{\langle e_m^*(1), x \rangle} \|x(1)\|_1 \xrightarrow{k \rightarrow \infty} 0. \quad \square$$

We consider (u, v) given by Lemma 7.7. We set, for $x \in X$,

$$R(x) := R_0(x) + \frac{x_0(1)}{u_0(1)}(v - R_0(u)).$$

This defines a bounded operator such that $R(u) = v \in \text{HC}(T_1)$. It turns out that there are not so many vectors $x \in X$ such that $R(x) \in \text{HC}(T_1)$.

LEMMA 7.10. *Let $x \in X$ be such that $R(x) \in \text{HC}(T_1)$. Then, $x(1)$ is a scalar multiple of $u(1)$.*

Proof. Let us set $z := x - (x_0(1)/u_0(1))u$, so that

$$R(x) = R_0(z) + \frac{x_0(1)}{u_0(1)}v.$$

Assume first that $z(1) \notin \text{span}(e_0(1))$. Then, there exists $m \geq 1$ such that $\langle e_m^*(1), z \rangle \neq 0$, whereas $\langle e_j^*(1), z \rangle = 0$ for $1 \leq j < m$. Let $l > m$. By Lemma 7.9, it follows that

$$\langle e_k^*(l), R_0(z) \rangle = \varepsilon_l \langle e_m^*(1), z \rangle 2^{-mk} (1 + \delta(k)),$$

so that

$$\begin{aligned}\langle e_k^*(l), R(x) \rangle &= \varepsilon_l \langle e_m^*(1), z \rangle 2^{-mk} (1 + \delta(k)) + \frac{x_0(1)}{u_0(1)} v_k(l) \\ &= \varepsilon_l \langle e_m^*(1)(z) 2^{-mk} (1 + \delta_l(k)),\end{aligned}$$

where for all $l > m$, $\delta_l(k) \rightarrow 0$ as $k \rightarrow \infty$, since $|v_k(l)| \leq 2^{-lk} = o(2^{-mk})$. By Lemma 7.8, $R(x) \notin \text{HC}(T_1)$, which is a contradiction.

Hence, there exists a complex number $\alpha(1)$ such that

$$x(1) - \frac{x_0(1)}{u_0(1)}u(1) = \alpha(1)e_0(1).$$

Applying the functional $e_0^*(1)$ to this equation, we easily get $\alpha(1) = 0$, which implies that $x(1)$ belongs to $\text{span}(u(1))$. \square

We can now give the proof.

Proof of Theorem 7.2. Let us fix $c > \|R\|$. We set $L_2 := cI + R$ (which is invertible) and $T_2 := L_2^{-1}TL_2$. We show that the pair (T_1, T_2) is d -frequently hypercyclic but not densely d -hypercyclic.

By construction, $(u, Ru) = (u, v) \in \text{FHC}(T_1 \oplus T_1)$. Hence, $u \in d\text{-FHC}(T_1, T_2)$ by Lemma 7.5. Moreover, setting $T = T_1$ and $L_1 = cI$, Lemma 7.3 implies that if $x \in d\text{-HC}(T_1, T_2)$, then $R(x) \in \text{HC}(T_1)$. By Lemma 7.10, it follows that $x(1) \in \text{span}(u(1))$ for every $x \in d\text{-HC}(T_1, T_2)$. In particular, $d\text{-HC}(T_1, T_2)$ cannot be dense in X . \square

8. Eigenvectors and d -frequent hypercyclicity

In this section, we give a criterion relying on properties of the eigenvectors for showing that a tuple of operators is (densely) d -frequently hypercyclic. The initial motivation was the following question asked by K. Grosse-Erdmann: let D be the derivation operator acting on the space of entire functions $H(\mathbb{C})$, and for every $a \in \mathbb{C} \setminus \{0\}$, denote by τ_a the operator of translation by a on $H(\mathbb{C})$, defined by $\tau_a f(z) := f(z + a)$. It is well known (see [7] or [33]) that both D and τ_a are frequently hypercyclic. Now one can ask the following question.

Question 8.1. Do the operators D and τ_a have common frequently hypercyclic vectors?

It will follow from the next theorem that the answer to Question 8.1 is positive.

THEOREM 8.2. *Let $N \geq 2$, let X be a complex Fréchet space and let $T_1, \dots, T_N \in \mathfrak{L}(X)$. Assume that there exist a holomorphic vector field $E : O \rightarrow X$, defined on some connected open set $O \subset \mathbb{C}$, and non-constant holomorphic functions ϕ_1, \dots, ϕ_N , defined on some connected open set containing O , such that:*

- $\overline{\text{span}} E(O) = X$;
- $T_i E(z) = \phi_i(z)E_i(z)$ for every $i = 1, \dots, N$ and $z \in O$;
- $O \cap \phi_i^{-1}(\mathbb{T}) \cap \bigcap_{j \neq i} \phi_j^{-1}(\mathbb{D}) \neq \emptyset$ for every $i = 1, \dots, N$.

Then, the N -tuple (T_1, \dots, T_N) is densely d -frequently hypercyclic.

Before proving this result, let us state two consequences and give some examples.

COROLLARY 8.3. *Let D be the derivation operator on $X := H(\mathbb{C})$. If ϕ_1 and ϕ_2 are two entire functions of exponential type such that $\phi_1^{-1}(\mathbb{T}) \cap \phi_2^{-1}(\mathbb{D}) \neq \emptyset$ and $\phi_2^{-1}(\mathbb{T}) \cap \phi_1^{-1}(\mathbb{D}) \neq \emptyset$, then the pair $(\phi_1(D), \phi_2(D))$ is densely d -frequently hypercyclic.*

Proof. Let $E : \mathbb{C} \rightarrow X$ be the holomorphic vector field defined by $E(z) := e^z \cdot$. We have $DE(z) = zE(z)$ for all $z \in \mathbb{C}$ and $\overline{\text{span}} E(\mathbb{C}) = X$; so we may apply Theorem 8.2 to the operators $T_i := \phi_i(D)$. \square

Since $\tau_a = \Phi_a(D)$, where $\phi_a(z) := e^{az}$, Corollary 8.3 applies to pairs of operators involving D and τ_a :

Example 8.4. Taking $\phi_1(z) := z$ and $\phi_2(z) := e^{az}$, we see that for any $a \neq 0$, the pair (D, τ_a) is densely d -frequently hypercyclic (so that, in particular, $\text{FHC}(D) \cap \text{FHC}(\tau_a) \neq \emptyset$). Indeed, any complex number z such that $|z| = 1$ and $\Re e(az) < 0$ belongs to

$\phi_1^{-1}(\mathbb{T}) \cap \phi_2^{-1}(\mathbb{D})$, while any $z \in i\bar{a}\mathbb{R}$ such that $|z| < 1$ belongs to $\phi_2^{-1}(\mathbb{T}) \cap \phi_1^{-1}(\mathbb{D})$. Similarly, if $a, b \neq 0$ and $a/b \notin \mathbb{R}$, then (τ_a, τ_b) is densely d -frequently hypercyclic.

COROLLARY 8.5. *Let B be the canonical backward shift acting on $X = \ell_p(\mathbb{Z}_+)$ or $c_0(\mathbb{Z}_+)$. If ϕ_1 and ϕ_2 are two holomorphic functions defined in a neighbourhood of the closed unit disk $\overline{\mathbb{D}}$ such that $\mathbb{D} \cap \phi_1^{-1}(\mathbb{T}) \cap \phi_2^{-1}(\mathbb{D}) \neq \emptyset$ and $\mathbb{D} \cap \phi_2^{-1}(\mathbb{T}) \cap \phi_1^{-1}(\mathbb{D}) \neq \emptyset$, then the pair $(\phi_1(B), \phi_2(B))$ is densely d -frequently hypercyclic.*

Proof. The operators $T_i := \phi_i(B)$ are well defined since $\sigma(B) = \overline{\mathbb{D}}$. Let $E : \mathbb{D} \rightarrow X$ be the holomorphic vector field defined by $E(z) := \sum_{n=0}^{\infty} z^n e_n$. We have $BE(z) = zE(z)$ for all $z \in \mathbb{D}$ and $\overline{\text{span}} E(\mathbb{C}) = X$; so Theorem 8.2 applies. \square

Example 8.6. If $|\lambda| > 1$ and $0 < |\alpha| < 2|\lambda|$, the pair $(\lambda B, I + \alpha B)$ is densely d -frequently hypercyclic. If $|\lambda| > 1$ and $a \neq 0$, the pair $(\lambda B, e^{aB})$ is densely d -frequently hypercyclic. However, Theorem 8.2 is completely inefficient to show, for example, that the pair (aB, bB^2) is d -frequently hypercyclic if $1 < a < b$, which is nevertheless true by [36].

In the proof of Theorem 8.2, we will make use of the following straightforward observation.

Fact 8.7. Let $T_1, \dots, T_N \in \mathcal{L}(X)$ and let $x_1, \dots, x_N \in X$. Assume that $(x_1, \dots, x_N) \in \text{FHC}(T_1 \oplus \dots \oplus T_N)$ and that $T_j^m x_i \rightarrow 0$ as $m \rightarrow \infty$ whenever $i \neq j$. Then, $x := x_1 + \dots + x_N$ is a d -frequently hypercyclic vector for (T_1, \dots, T_N) .

Proof of Theorem 8.2. We first note that for $i = 1, \dots, N$, there is a non-empty open set $V_i \subset O$ and $r_i \in (0, 1)$ such that $(\phi_i)|_{V_i}$ is a diffeomorphism (onto its range), $\phi_i(V_i) \cap \mathbb{T} \neq \emptyset$ and $\phi_j(V_i) \subset D(0, r_i)$ for $j \neq i$. Indeed, let $a \in O$ be such that $\phi_i(a) \in \mathbb{T}$ and $\phi_j(a) \in \mathbb{D}$ for $j \neq i$. Choose an open neighbourhood W of a and $r_i \in (0, 1)$ such that $\phi_j(W) \subset D(0, r_i)$ for all $j \neq i$. By the open mapping theorem, $\phi_i(W)$ is an open set intersecting \mathbb{T} , so $\phi_i(W) \cap \mathbb{T}$ is uncountable. Hence, one can find $b \in W$ such that $\phi_i(b) \in \mathbb{T}$ and $\phi'_i(b) \neq 0$; and the claim follows from the inverse function theorem.

Taking the open set V_i smaller if necessary, we may assume that $\Lambda_i := \phi_i(V_i) \cap \mathbb{T}$ is a proper open arc of \mathbb{T} . We choose a ‘cut-off’ function $\chi_i \in C^\infty(\mathbb{T})$ such that $\chi_i(\lambda) = 0$ outside Λ_i and $\chi_i(\lambda) > 0$ on Λ_i , and (with the obvious abuse of notation) we define $F_i : \mathbb{T} \rightarrow X$ by setting

$$F_i(\lambda) := \chi_i(\lambda) E(\phi_i^{-1}(\lambda)) \quad \text{for every } \lambda \in \mathbb{T}.$$

Thus, F_i is a C^∞ -smooth \mathbb{T} -eigenvector field for T_i , that is, $T_i F_i(\lambda) = \lambda F_i(\lambda)$ for every $\lambda \in \mathbb{T}$. Let us denote by $\widehat{F}_i(n)$ the Fourier coefficients of F_i :

$$\widehat{F}_i(n) = \int_{\mathbb{T}} F_i(\lambda) \lambda^{-n} d\lambda, \quad n \in \mathbb{Z}.$$

Since F_i is a \mathbb{T} -eigenvector field for T_i , we have $T_i \widehat{F}_i(n) = \widehat{F}_i(n-1)$ for all $n \in \mathbb{Z}$, that is, the sequence $(\widehat{F}_i(n))_{n \in \mathbb{Z}}$ is a bilateral backward orbit for T_i . Moreover, since $\overline{\text{span}} E(O) = X$, it follows from the Hahn–Banach theorem, together with the identity principle for analytic functions, that $\overline{\text{span}} (\widehat{F}_i(n) : n \in \mathbb{Z}) = X$.

In the remainder of the proof, we fix a family $(g_{i,n})_{1 \leq i \leq N, n \in \mathbb{Z}}$ of independent, standard complex Gaussian variables defined on some probability space $(\Omega, \mathcal{A}, \mathbb{P})$.

CLAIM 8.8. *For every $i = 1, \dots, N$, the series*

$$\sum_{n \in \mathbb{Z}} g_{i,n} \widehat{F}_i(n)$$

is almost surely convergent and defines an X -valued random variable ξ_i on $(\Omega, \mathcal{A}, \mathbb{P})$, which is such that

$$\text{for every } i, j = 1, \dots, N \text{ with } j \neq i, \quad T_j^m \xi_i \xrightarrow{m \rightarrow \infty} 0 \text{ almost surely.}$$

Proof of Claim 8.8. Since F_i is \mathcal{C}^2 -smooth, two integrations by parts show that for any continuous semi-norm \mathbf{q} on X , we have $\mathbf{q}(\widehat{F}_i(n)) = O(1/n^2)$ as $|n| \rightarrow \infty$, so that

$$\sum_{n=-\infty}^{\infty} \mathbb{E}(\mathbf{q}(g_{i,n} \widehat{F}_i(n))) < \infty.$$

This implies that the series $\sum_{n \in \mathbb{Z}} g_{i,n} \widehat{F}_i(n)$ is almost surely convergent.

Let us fix $j \neq i$. By the definition of F_i , we have $T_j F_i(\lambda) = \psi_{i,j}(\lambda) F_i(\lambda)$ for every $\lambda \in \Lambda_i$, where $\psi_{i,j}(\lambda) := \phi_j(\phi_i^{-1}(\lambda))$; and $T_j F_i(\lambda) = 0$ if $\lambda \notin \Lambda_i$. Hence, for almost every $\omega \in \Omega$ and every $m \in \mathbb{N}$, we have

$$T_j^m(\xi_i(\omega)) = \sum_{n \in \mathbb{Z}} g_n(\omega) \int_{\Lambda_i} \psi_{i,j}(\lambda)^m F_i(\lambda) \lambda^{-n} d\lambda.$$

Let \mathbf{q} be a continuous semi-norm on X . Since $|\psi_{i,j}(\lambda)| < r_i$ for every $\lambda \in \Lambda_i$ by definition of ψ , two integrations by parts show that there is a constant $C_{\mathbf{q}}$ such that

$$\mathbf{q}\left(\int_{\Lambda} \psi_{i,j}(\lambda)^m F(\lambda) \lambda^{-n} d\lambda\right) \leq C_{\mathbf{q}} \times \frac{m^2 r_i^m}{1+n^2} \quad \text{for every } m \geq 0 \text{ and every } n \in \mathbb{Z}.$$

Moreover, it follows from the Borel–Cantelli lemma that for almost every $\omega \in \Omega$, there exists an integer $N(\omega)$ such that

$$\text{for all } |n| > N(\omega) : |g_n(\omega)| \leq \sqrt{n}.$$

Hence, given a continuous semi-norm \mathbf{q} on X , one can find for almost every $\omega \in \Omega$, some constant $M_{\mathbf{q},\omega}$ such that

$$\text{for all } m \in \mathbb{N} : \mathbf{q}(T_j^m \xi_i(\omega)) \leq M_{\mathbf{q},\omega} m^2 r_i^m.$$

Hence, $\mathbf{q}(T_j^m \xi_i(\omega)) \rightarrow 0$ almost surely as $m \rightarrow \infty$ for any given continuous semi-norm \mathbf{q} , that is, $T_j^m \xi_i \rightarrow 0$ almost surely. \square

We can now conclude the proof of Theorem 8.2. For $i = 1, \dots, N$, let us denote by μ_i the distribution of the random variable $\xi_i : \Omega \rightarrow X$. By definition, μ_i is a T_i -invariant Gaussian measure with full support; and by [1], T_i is *strongly mixing* with respect to μ_i . Hence, the measure $\mu_1 \otimes \dots \otimes \mu_N$ is a $(T_1 \oplus \dots \oplus T_N)$ -invariant measure on X^N with full support and $T_1 \oplus \dots \oplus T_N$ is mixing with respect to $\mu_1 \otimes \dots \otimes \mu_N$. Since $\mu_1 \otimes \dots \otimes \mu_N$ is the distribution of the random vector $\xi := (\xi_1, \dots, \xi_N)$ by

independence of ξ_1, \dots, ξ_N , it follows that the vector $\xi(\omega)$ is almost surely frequently hypercyclic for $T_1 \oplus \dots \oplus T_N$. Moreover, by Claim 8.8, we see that $T_j^m \xi_i(\omega) \rightarrow 0$ almost surely as $m \rightarrow \infty$, whenever $i \neq j$. By Fact 8.7, it follows that the vector $\xi_1(\omega) + \dots + \xi_N(\omega)$ is almost surely d -frequently hypercyclic for (T_1, \dots, T_N) . In other words, $(\mu_1 \star \dots \star \mu_N)$ -almost every $x \in X$ is d -frequently hypercyclic for (T_1, \dots, T_N) . Since the measure $\mu_1 \star \dots \star \mu_N$ has full support, this terminates the proof of Theorem 8.2. \square

9. Remarks and questions

9.1. Hereditary frequent hypercyclicity in a weak sense. Another natural definition for hereditary \mathcal{F} -hypercyclicity could be the following: an operator $T \in \mathcal{L}(X)$ is *hereditarily \mathcal{F} -hypercyclic in the weak sense* if, for every $A \in \mathcal{F}$, the sequence $(T^n)_{n \in A}$ is \mathcal{F} -hypercyclic, that is, there exists $x \in X$ such that $\mathcal{N}_T(x, V) \cap A \in \mathcal{F}$ for all non-empty open sets $V \subset X$. Equivalently, T is \mathcal{F}_A -hypercyclic for every $A \in \mathcal{F}$, where $\mathcal{F}_A := \{B \subset \mathbb{N} : B \cap A \in \mathcal{F}\}$. Of course, hereditary \mathcal{F} -hypercyclicity implies hereditary \mathcal{F} -hypercyclicity in the weak sense. Note also that Theorem 5.2 says precisely that there exist frequently hypercyclic operators which are not hereditarily frequently hypercyclic in the weak sense.

When \mathcal{F} is the family of all infinite subsets of \mathbb{N} , an operator T is hereditarily \mathcal{F} -hypercyclic in the weak sense if and only if it is ‘hereditarily hypercyclic with respect to the whole sequence of integers’ in the sense of [13]; and this means exactly that T is *topologically mixing* (see for example [27, Lemma 2.2]). The next result shows that this is also equivalent to hereditary \mathcal{F} -hypercyclicity.

PROPOSITION 9.1. *Let \mathcal{F} be a Furstenberg family with the following property: for any operator T and any $A \subset \mathbb{N}$, the set $\mathcal{F}_A\text{-HC}(T)$ is either empty or comeagre in the underlying space. Then, hereditary \mathcal{F} -hypercyclicity and hereditary \mathcal{F} -hypercyclicity in the weak sense are equivalent. In particular, when \mathcal{F} is the family of all infinite subsets of \mathbb{N} , an operator T is hereditarily \mathcal{F} -hypercyclic if and only if it is topologically mixing.*

Proof. Assume that T is hereditarily \mathcal{F} -hypercyclic in the weak sense. Let $(A_i)_{i \in I}$ be a countable family of sets in \mathcal{F} and let $(V_i)_{i \in I}$ be a family of non-empty open subsets of X . By assumption on \mathcal{F} , for each $i \in I$, the set G_i of \mathcal{F} -hypercyclic vectors for the sequence $(T^n)_{n \in A_i}$ is comeagre in X ; so $G := \bigcap_{i \in I} G_i$ is non-empty. Then, any $x \in G$ satisfies the required property: for every $i \in I$, there is a set $B_i \in \mathcal{F}$ such that $B_i \subset A_i$ and $T^n x \in V_i$ for all $n \in B_i$. \square

We can now ask the following question.

Question 9.2. For which Furstenberg families \mathcal{F} do hereditary \mathcal{F} -hypercyclicity and hereditary \mathcal{F} -hypercyclicity in the weak sense coincide?

In view of the results of [17], *upper Furstenberg families* may be good candidates. However, we are unable to handle even the case of sets with positive upper density. Proposition 9.1 leads naturally to the following question.

Question 9.3. Let us denote by $\overline{\mathcal{D}}$ the family of all sets $A \subset \mathbb{N}$ with positive upper density. Is it true that if $A \in \overline{\mathcal{D}}$ and $T \in \mathcal{L}(X)$ is $\overline{\mathcal{D}}_A$ -hypercyclic, then $\overline{\mathcal{D}}_A$ -HC(T) is comeagre in X ?

However, it might seem more than plausible that the two notions are not equivalent in the case of frequent hypercyclicity, that is, when \mathcal{F} is the family of sets with positive lower density. However again, we do not know how to prove this.

One may also think of ‘local’ versions of hereditary frequent hypercyclicity. For example, one could say that an operator $T \in \mathcal{L}(X)$ is:

- hereditarily \mathcal{F} -hypercyclic with respect to some sequence $(\Lambda_i)_{i \in \mathbb{N}} \subset \mathcal{F}$ if, for any sequence $(A_i) \subset \mathcal{F}$ with $A_i \subset \Lambda_i$ and for any sequence of non-empty open sets (V_i) in X , one can find a vector $x \in X$ such that $\mathcal{N}_T(x, V_i) \cap A_i \in \mathcal{F}$ for all $i \in \mathbb{N}$;
- hereditarily \mathcal{F} -hypercyclic with respect to some set $\Lambda \in \mathcal{F}$ if it is hereditarily \mathcal{F} -HC with respect to the constant sequence $\Lambda_i = \Lambda$;
- hereditarily \mathcal{F} -hypercyclic in the weak sense with respect to some set $\Lambda \in \mathcal{F}$ if it is \mathcal{F}_A -hypercyclic for any $A \in \mathcal{F} \cap 2^\Lambda$.

When \mathcal{F} is the family of all infinite subsets of \mathbb{N} , hereditary \mathcal{F} -hypercyclicity in the weak sense with respect to some set $\Lambda = \{n_k : k \geq 0\}$ is the same as hereditary hypercyclicity with respect to the sequence (n_k) in the sense of [13]; and hence, by [13, Theorem 2.3], an operator T is hereditarily \mathcal{F} -hypercyclic in the weak sense with respect to some set Λ if and only if it is topologically weakly mixing, that is, $T \oplus T$ is hypercyclic. Also, the proof of Proposition 4.1 makes it clear that if T is hereditarily \mathcal{F} -hypercyclic with respect to some sequence (Λ_i) , then $T \oplus T$ is \mathcal{F} -hypercyclic. This leads to the following question.

Question 9.4. If $T \in \mathcal{L}(X)$ is hereditarily \mathcal{F} -hypercyclic in the weak sense with respect to some set $\Lambda \in \mathcal{F}$, does it follow that $T \oplus T$ is \mathcal{F} -hypercyclic? And conversely?

In the same spirit and with [24] in mind, one may ask another question.

Question 9.5. Does \mathcal{U} -frequent hypercyclicity imply some weak form of hereditary \mathcal{U} -frequent hypercyclicity, yet strong enough to ‘explain’ why $T \oplus T$ is \mathcal{U} -frequently hypercyclic as soon as T is?

9.2. \mathcal{F} -transitivity and hereditary \mathcal{F} -transitivity. In topological dynamics, there is a natural notion of ‘transitivity’ associated with a given Furstenberg family \mathcal{F} (see for example [12, 26] in the linear setting): if X is a topological space, a continuous map $T : X \rightarrow X$ is said to be \mathcal{F} -transitive if $\mathcal{N}_T(U, V) \in \mathcal{F}$ for every pair (U, V) of non-empty open sets in X , where

$$\mathcal{N}_T(U, V) := \{n \in \mathbb{N} : T^n(U) \cap V \neq \emptyset\}.$$

Following [12], one can consider a ‘hereditary’ version of \mathcal{F} -transitivity: let us say that an operator $T \in \mathcal{L}(X)$ is *hereditarily \mathcal{F} -transitive* if $\mathcal{N}(U, V) \cap A \in \mathcal{F}$ for every $A \in \mathcal{F}$ and all non-empty open sets U, V . There is an obvious link with hereditary \mathcal{F} -hypercyclicity.

Remark 9.6. Hereditarily \mathcal{F} -hypercyclic operators are hereditarily \mathcal{F} -transitive.

Proof. By Proposition 6.4, we know that if T is hereditarily \mathcal{F} -hypercyclic, then T is densely hereditarily \mathcal{F} -hypercyclic. Let U, V be non-empty open sets in X and let $A \in \mathcal{F}$. By dense hereditary \mathcal{F} -hypercyclicity, there exists $x \in U$ such that $N(x, V) \cap A \in \mathcal{F}$. In particular, $N(U, V) \cap A \in \mathcal{F}$, so T is hereditarily \mathcal{F} -transitive. \square

The converse is definitely not true in general, for the following reason: there exist topologically mixing operators that are not frequently hypercyclic. In particular, any such operator is hereditarily \underline{D} -transitive, where \underline{D} is the family of sets with positive lower density, but not frequently hypercyclic (that is, not \underline{D} -hypercyclic). This leads to the following questions (the third one was suggested by the referee).

Question 9.7. Are there operators which are frequently hypercyclic and topologically mixing, but not hereditarily frequently hypercyclic?

Question 9.8. Are there at least operators which are hereditarily \underline{D} -transitive and frequently hypercyclic, but not hereditarily frequently hypercyclic?

Question 9.9. Assume that $T \in \mathcal{L}(X)$ admits an invariant measure with full support with respect to which it is a strongly mixing transformation. Does it follow that T is hereditarily frequently hypercyclic? What if, additionally, the measure is Gaussian?

Given a Furstenberg family \mathcal{F} , one can define the *dual family* \mathcal{F}^* as the collection of all subsets A of \mathbb{N} such that $A \cap B \neq \emptyset$ for every $B \in \mathcal{F}$. It is clear by definition that every hereditarily \mathcal{F} -transitive operator is \mathcal{F}^* -transitive; and it is also clear that $(\underline{D})^* = \overline{D}_1$, the family of sets with upper density equal to 1. Hence, every hereditarily frequently hypercyclic operator is \overline{D}_1 -transitive. It is natural to wonder if every frequently hypercyclic operator is \overline{D}_1 -transitive too. The next proposition shows that this is not the case. This is an improvement of [12, Proposition 5.1], where it is shown that reiterative hypercyclicity does not imply \overline{D}_1 -transitivity. Moreover, the example we give is any of the weighted shifts introduced in the proof of Theorem 4.2; so this provides another proof that these shifts are not hereditarily frequently hypercyclic.

PROPOSITION 9.10. *There exists a frequently hypercyclic weighted shift B_w on $c_0(\mathbb{Z}_+)$ which is not \overline{D}_1 -transitive.*

Proof. Let B_w be one of the weighted shifts introduced in the proof of Theorem 4.2. By [12, Proposition 3.3], to show that B_w is not \overline{D}_1 -transitive, it is enough to find $M > 0$ such that

$$C_M := \{n \in \mathbb{N} : |w_1 \cdots w_n| > M\} \notin \overline{D}_1.$$

With the notation of the proof of Theorem 4.2, we know that for every $p \geq 1$,

$$C_{2^p} \subset \bigcup_{q \geq p} (b_q \mathbb{N} + [-q, q]) \cup \bigcup_{q \geq 1} \bigcup_{u \in A_{2q}} I_u^{a, 4\epsilon}.$$

Moreover, by assumption on (b_q) , we have

$$\lim_{p \rightarrow \infty} \overline{\text{dens}} \left(\bigcup_{q \geq p} (b_q \mathbb{N} + [-q, q]) \right) = 0;$$

and we also have

$$\bigcup_{q \geq 1} \bigcup_{u \in A_{2q}} I_u^{a, 4\varepsilon} \subset \bigcup_{u \geq 1} I_u^{a, 4\varepsilon}.$$

Since

$$\overline{\text{dens}} \left(\bigcup_{u \geq 1} I_u^{a, 4\varepsilon} \right) \leq \lim_{u \rightarrow \infty} \frac{\sum_{k=1}^u 8\varepsilon a^k}{(1 + 4\varepsilon)a^u} = \frac{8\varepsilon}{1 + 4\varepsilon} \sum_{k=0}^{\infty} a^{-k} < 1 \quad \text{if } a \text{ is sufficiently big,}$$

it follows that if p is sufficiently big, then

$$\overline{\text{dens}} C_{2^p} < 1.$$

This concludes the proof of Proposition 9.10. \square

9.3. About disjointness. The original definition of disjointness in topological dynamics goes back to Furstenberg's seminal paper [25]. The setting is that of compact dynamical systems (X, T) , that is, X is a compact metric space and $T : X \rightarrow X$ is a continuous map. Two compact dynamical systems (X_1, T_1) and (X_2, T_2) are said to be disjoint if the only closed, $(T_1 \times T_2)$ -invariant set $\Gamma \subset X_1 \times X_2$ such that $\pi_{X_1}(\Gamma) = X_1$ and $\pi_{X_2}(\Gamma) = X_2$ is $\Gamma = X_1 \times X_2$. Note that since the spaces are compact, one could replace $\pi_{X_i}(\Gamma)$ by $\overline{\pi_{X_i}(\Gamma)}$ in the definition. For dynamical systems (X, T) whose underlying space is not necessarily compact, both definitions make sense and lead to *a priori* different notions of disjointness (the one 'with closure' being stronger than the one 'without closure'). In particular, one could consider these notions in the linear setting. However, there are no disjoint pairs of linear dynamical systems in this sense, even 'without closures'. Indeed, if $T_1 \in \mathfrak{L}(X_1)$ and $X_2 \in \mathfrak{L}(X_2)$, then $\Gamma := (\{0\} \times X_2) \cup (X_1 \times \{0\})$ shows that disjointness cannot be met. One can get round this difficulty by changing the definitions a little bit as follows: instead of $\pi_{X_i}(\Gamma) = X_i$, require that $\pi_{X_i}(\Gamma \cap (X_1 \setminus \{0\}) \times (X_2 \setminus \{0\})) = X_i \setminus \{0\}$; and likewise for the definition 'with closures'.

Even though these definitions of disjointness are likely to be artificial, one can try to play with them a little. For example, copying out the relevant parts of [25]—namely, the proofs of Theorems II.1 and II.2—one gets the following results. Let us say that a linear dynamical system (X, T) is *minimal apart from 0* if every non-zero vector $x \in X$ is hypercyclic for T ; equivalently, if the only closed T -invariant subsets of X are $\{0\}$ and X . Famous examples of Read [50] show that this can indeed happen.

PROPOSITION 9.11. *Let (X_1, T_1) and (X_2, T_2) be two linear dynamical systems. If (X_1, T_1) and (X_2, T_2) are disjoint 'without closures', then at least one of them is minimal apart from 0.*

Proof. Assume that (X_1, T_1) and (X_2, T_2) are not minimal apart from 0. Then, for $i = 1, 2$, one can find a closed T_i -invariant set $C_i \subset X_i$ such that $C_i \neq X_i$ and $C_i \cap (X_i \setminus \{0\}) \neq \emptyset$; and $\Gamma := (C_1 \times X_2) \cup (X_1 \times C_2)$ shows that (X_1, T_1) and (X_2, T_2) are not disjoint ‘without closures’. \square

PROPOSITION 9.12. *Let (X_1, T_1) and (X_2, T_2) be two linear dynamical systems. Assume that the periodic points of T_1 are dense in X_1 and that (X_2, T_2) is minimal apart from 0. Then, (X_1, T_1) and (X_2, T_2) are disjoint ‘without closures’.*

Proof. Let $\Gamma \subset X_1 \times X_2$ be a closed, $(T_1 \times T_2)$ -invariant set such that $\pi_{X_i}(\Gamma \cap (X_1 \setminus \{0\}) \times (X_2 \setminus \{0\})) = X_i \setminus \{0\}$ for $i = 1, 2$. We have to show that $\Gamma = X_1 \times X_2$; and since the periodic points of T_1 are dense in X_1 , it is enough to show that $(\text{Per}(T_1) \setminus \{0\}) \times X_2 \subset \Gamma$.

Let $u \in X_1$ be any non-zero periodic point of T_1 and choose $d \in \mathbb{N}$ such that $T_1^d u = u$. By assumption on Γ , one can find $v \in X_2 \setminus \{0\}$ such that $(u, v) \in \Gamma$. Then, $(u, T_2^{dn} v) \in \Gamma$ for all $n \in \mathbb{N}$ by $(T_1 \times T_2)$ -invariance of Γ . Moreover, $v \in \text{HC}(T_2)$ by assumption on (X_2, T_2) . Hence, by Ansari’s theorem [3], v is also a hypercyclic vector for T_2^d ; and since Γ is closed in $X_1 \times X_2$, it follows that $\{u\} \times X_2 \subset \Gamma$. \square

Proposition 9.12 implies in particular that linear dynamical systems which are disjoint ‘without closure’ do exist. We do not know if this is also true ‘with closure’; so one could think of possible weakenings of the definition of disjointness ‘with closures’. For hypercyclic operators, one possible such weakening could be the following: one could say that two hypercyclic operators $T_1 \in \mathcal{L}(X_1)$ and $T_2 \in \mathcal{L}(X_2)$ are *pseudo-disjoint* (just to give a name) if, whenever x_1 is a hypercyclic vector for T_1 and x_2 is a hypercyclic vector for T_2 , it follows that (x_1, x_2) is hypercyclic for $T_1 \times T_2$. This is indeed weaker than the definition of disjointness ‘with closures’ (consider $\Gamma := \overline{\text{Orb}((x_1, x_2), T_1 \times T_2)}$), yet formally much stronger than the disjointness notion introduced in [10, 14], that is, diagonal hypercyclicity. We are not much further ahead since we do not know if there are any pseudo-disjoint pairs of linear operators (whereas there are lots of interesting examples for diagonal hypercyclicity). So we ask the following question.

Question 9.13. Are there any pseudo-disjoint pairs of operators, that is, pairs of hypercyclic operators (T_1, T_2) such that $\text{HC}(T_1) \times \text{HC}(T_2) \subset \text{HC}(T_1 \times T_2)$?

Regarding this question, one may observe that two linear operators T_1 and T_2 are trivially pseudo-disjoint if it happens that every vector $x \in X_1 \times X_2$ with non-zero coordinates is hypercyclic for $T_1 \times T_2$. This leads to the following ‘strong’ form of Question 9.13.

Question 9.14. Are there pairs of operators (T_1, T_2) such that every $x \in (X_1 \setminus \{0\}) \times (X_2 \setminus \{0\})$ is hypercyclic for $T_1 \times T_2$?

Let us point out the following amusing fact: if such a pair (T_1, T_2) can be found, then the operator $T = T_1 \times T_2$ acting on $X = X_1 \times X_2$ is a hypercyclic operator such that $\text{HC}(T)$

is an open set but $HC(T) \neq X \setminus \{0\}$. We do not know of any example of operators with that property.

Finally, we note that if one extends the definition of pseudo-disjointness to possibly nonlinear systems in the obvious way, it follows from the main result of [52] that any irrational rotation of the circle is pseudo-disjoint from any hypercyclic operator. In view of that, one may consider the following variant of Question 9.13.

Question 9.15. Are there natural classes of hypercyclic operators $\mathfrak{C}_1, \mathfrak{C}_2$ such that any $T_1 \in \mathfrak{C}_1$ is pseudo-disjoint from any $T_2 \in \mathfrak{C}_2$?

9.4. *Other questions.* We conclude the paper by adding some other possibly interesting questions motivated by the results obtained in the paper.

The first question asks for a converse to Observation 1.3.

Question 9.16. Let \mathcal{F} be a Furstenberg family and let $T \in \mathfrak{L}(X)$. Assume that $S \oplus T$ is \mathcal{F} -hypercyclic for every \mathcal{F} -hypercyclic operator S . Does it follow that T is hereditarily \mathcal{F} -hypercyclic?

The next two questions are related to an already mentioned ‘trap’ into which it is easy to fall: if \mathcal{F} and \mathcal{F}' are two Furstenberg families, the fact that $\mathcal{F} \subset \mathcal{F}'$ does not formally imply that hereditary \mathcal{F} -hypercyclicity is a stronger property than hereditary \mathcal{F}' -hypercyclicity.

Question 9.17. Are there hereditarily frequently hypercyclic operators which are not topologically mixing?

Question 9.18. Does hereditary frequent hypercyclicity imply hereditary \mathcal{U} -frequent hypercyclicity?

In the theory of frequently hypercyclic operators, there are non-trivial counterexamples to some tempting ‘conjectures’. It is natural to ask if these examples are in fact hereditarily frequently hypercyclic or if it is possible to modify them to get hereditarily frequently hypercyclic examples. In particular, with [4, 42] in mind, this leads to the following questions (the second one is a strengthening of Question 9.17).

Question 9.19. Are there invertible hereditarily frequently hypercyclic operators whose inverse is not frequently hypercyclic?

Question 9.20. Are there operators which are both hereditarily frequently hypercyclic and chaotic but not topologically mixing?

The next two questions are related to the sufficient conditions we found for hereditary frequent hypercyclicity. Observe first that in addition to the FHCC and the unimodular eigenvectors machinery, there are other criteria to prove frequent hypercyclicity (see [11] or [31, Theorem 5.35]). They do not imply hereditarily frequent hypercyclicity. Indeed, the

criterion of [11] is equivalent to frequent hypercyclicity for weighted shifts on c_0 , whereas the C -type operator of Theorem 5.2 satisfies [31, Theorem 5.35].

Question 9.21. If $T \in \mathcal{L}(X)$ is such that the \mathbb{T} -eigenvectors of T are spanning with respect to the Lebesgue measure, does it follow that one can find a T -invariant measure μ with full support such that (X, \mathcal{B}, μ, T) is a factor of a dynamical system with countable Lebesgue spectrum? Does it follow at least that T is hereditarily frequently hypercyclic?

Question 9.22. Let $T \in \mathcal{L}(X)$. Is it true that if the \mathbb{T} -eigenvectors of T are perfectly spanning, then T is hereditarily frequently hypercyclic?

Concerning the invariant measure business, the next two questions seem natural. The first one is motivated by Theorem 3.24.

Question 9.23. Does there exist an operator T which is not hereditarily \mathcal{U} -frequently hypercyclic but admits an ergodic measure with full support?

Question 9.24. If X is a reflexive Banach space, then any frequently hypercyclic operator T on X admits a continuous invariant probability measure with full support (see [30]). Is it possible to improve this result if T is assumed to be hereditarily frequently hypercyclic?

The next question is, of course, strongly reminiscent of the Bès–Peris theorem [13], according to which the hypercyclicity criterion characterizes topological weak mixing.

Question 9.25. Given a Furstenberg family \mathcal{F} , is there some ‘ \mathcal{F} -hypercyclicity criterion’ characterizing the operators T such that $T \oplus T$ is \mathcal{F} -hypercyclic?

Finally, our last three questions concern the links between (hereditary) frequent hypercyclicity and the geometry of the underlying space X .

Question 9.26. On which spaces X is it possible to find hereditarily frequently hypercyclic operators? Is it possible at least on any complex Banach space admitting an unconditional Schauder decomposition?

Question 9.27. Are there spaces X which support frequently hypercyclic operators, but no hereditarily frequently hypercyclic operator?

Question 9.28. On which Banach spaces X is it possible to construct d -frequently hypercyclic pairs (T_1, T_2) which are not densely d -frequently hypercyclic?

Acknowledgements. We gratefully acknowledge the support of the Mathematisches Forschungsinstitut Oberwolfach, where part of this work was carried out. The second author was also supported in part by the Labex CEMPI (ANR-11-LABX-0007-01). The fourth author is a Research Associate of the Fonds de la Recherche Scientifique - FNRS. Finally, we would like to thank the anonymous referee for suggesting two questions and helping us to improve the presentation in §3.

REFERENCES

- [1] K. Agneessens. Frequently hypercyclic random vectors. *Proc. Amer. Math. Soc.* **151** (2023), 1103–1117.
- [2] F. Albiac and N. J. Kalton. *Topics in Banach Space Theory*. Springer, New York, 2006.
- [3] S. I. Ansari. Hypercyclic and cyclic vectors. *J. Funct. Anal.* **128** (1995), 374–383.
- [4] C. Badea and S. Grivaux. Unimodular eigenvalues, uniformly distributed sequences and linear dynamics. *Adv. Math.* **211** (2007), 766–793.
- [5] F. Bayart and S. Grivaux. Frequently hypercyclic operators. *Trans. Amer. Math. Soc.* **358** (2006), 5083–5117.
- [6] F. Bayart and S. Grivaux. Invariant Gaussian measures for operators on Banach spaces and linear dynamics. *Proc. Lond. Math. Soc. (3)* **94** (2007), 181–210.
- [7] F. Bayart and É. Matheron. *Dynamics of Linear Operators (Cambridge Tracts in Mathematics, 179)*. Cambridge University Press, Cambridge, 2009.
- [8] F. Bayart and É. Matheron. Mixing operators and small subsets of the circle. *J. Reine Angew. Math.* **715** (2016), 75–123.
- [9] F. Bayart and I. Ruzsa. Difference sets and frequently hypercyclic weighted shifts. *Ergod. Th. & Dynam. Sys.* **35** (2015), 691–709.
- [10] L. Bernal-González. Disjoint hypercyclic operators. *Studia Math.* **182**(2) (2007), 113–131.
- [11] J. P. Bès, Q. Menet, A. Peris and Y. Puig. Recurrence properties of hypercyclic operators. *Math. Ann.* **366** (2016), 545–572.
- [12] J. P. Bès, Q. Menet, A. Peris and Y. Puig. Strong transitivity properties for operators. *J. Differential Equations* **266** (2019), 1313–1337.
- [13] J. P. Bès and A. Peris. Hereditarily hypercyclic operators. *J. Funct. Anal.* **167** (1999), 94–112.
- [14] J. P. Bès and A. Peris. Disjointness in hypercyclicity. *J. Math. Anal. Appl.* **336** (2007), 297–315.
- [15] J. R. Blum and D. L. Hanson. On the mean ergodic theorem for subsequences. *Bull. Amer. Math. Soc.* **66** (1960), 308–311.
- [16] A. Bonilla and K.-G. Grosse-Erdmann. Frequently hypercyclic operators and vectors. *Ergod. Th. & Dynam. Sys.* **27** (2007), 383–404.
- [17] A. Bonilla and K.-G. Grosse-Erdmann. Upper frequent hypercyclicity and related notions. *Rev. Mat. Complut.* **31** (2018), 673–711.
- [18] R. Cardeccia and S. Muro. Frequently recurrent backward shifts. *Preprint*, 2024, [arXiv:2407.11799](https://arxiv.org/abs/2407.11799).
- [19] J. B. Conway. *A Course in Functional Analysis (Graduate Texts in Mathematics, 96)*. Springer, New York, 1985.
- [20] J.-P. Conze. Convergence des moyennes ergodiques pour des sous-suites. *Mém. Soc. Math. Fr. (N.S.)* **35** (1973), 7–15.
- [21] I. P. Cornfeld, S. V. Fomin and Ya. G. Sinai. *Ergodic Theory (Grundlehren der mathematischen Wissenschaften, 245)*. Springer, New York, 1982.
- [22] M. De la Rosa, L. Frerick, S. Grivaux and A. Peris. Frequent hypercyclicity, chaos, and unconditional Schauder decompositions. *Israel J. Math.* **190** (2012), 389–399.
- [23] M. De la Rosa and C. Read. A hypercyclic operator whose direct sum $T \oplus T$ is not hypercyclic. *J. Operator Theory* **61** (2009), 369–380.
- [24] R. Ernst, C. Esser and Q. Menet. \mathcal{U} -frequent hypercyclicity notions and related weighted densities. *Israel J. Math.* **241** (2021), 817–848.
- [25] H. Furstenberg. Disjointness in ergodic theory, minimal sets, and a problem in Diophantine approximation. *Math. Systems Theory* **1** (1967), 1–49.
- [26] E. Glasner. Classifying dynamical systems by their recurrence properties. *Topol. Methods Nonlinear Anal.* **24** (2004), 21–40.
- [27] S. Grivaux. Hypercyclic operators, mixing operators and the bounded steps problem. *J. Operator Theory* **54** (2005), 147–168.
- [28] S. Grivaux. A probabilistic version of the frequent hypercyclicity criterion. *Studia Math.* **176** (2006), 279–290.
- [29] S. Grivaux. A new class of frequently hypercyclic operators. *Indiana Univ. Math. Math. J.* **60** (2011), 1177–1201.
- [30] S. Grivaux and É. Matheron. Invariant measures for frequently hypercyclic operators. *Adv. Math.* **265** (2014), 371–427.
- [31] S. Grivaux, É. Matheron and Q. Menet. Linear dynamical systems on Hilbert spaces: typical properties and explicit examples. *Mem. Amer. Math. Soc.* **269** (2021), 147 pp.
- [32] K. Grosse-Erdmann. Frequently hypercyclic bilateral weighted shifts. *Glasg. Math. J.* **61** (2019), 271–286.
- [33] K. Grosse-Erdmann and A. Peris, *Linear Chaos (Universitext)*. Springer, London, 2011.
- [34] S. Janson, *Gaussian Hilbert Spaces (Cambridge Tracts in Mathematics, 129)*. Cambridge University Press, Cambridge, 1997.

- [35] M. Ledoux and M. Talagrand. *Probability in Banach Spaces (Ergebnisse der Mathematik und ihrer Grenzgebiete, 23)*. Springer, Berlin, 1991.
- [36] Ö. Martin, Q. Menet and Y. Puig. Disjoint frequently hypercyclic pseudo-shifts. *J. Funct. Anal.* **283** (2022), 109474.
- [37] Ö. Martin and Y. Puig. Existence of disjoint frequently hypercyclic operators which fail to be disjoint weakly mixing. *J. Math. Anal. Appl.* **500** (2021), 125106.
- [38] Ö. Martin and R. Sanders. Disjoint supercyclic weighted shifts. *Integral Equ. Oper. Theory* **85** (2016), 191–220.
- [39] Ö. Martin and R. Sanders. Extending families of disjoint hypercyclic operators. *J. Math. Anal. Appl.* **539** (2024), 128539.
- [40] B. Maurey and G. Pisier. Séries de variables aléatoires vectorielles indépendantes et propriétés géométriques des espaces de Banach. *Studia Math.* **58** (1976), 45–90.
- [41] Q. Menet. Linear chaos and frequent hypercyclicity. *Trans. Amer. Math. Soc.* **369** (2017), 4977–4994.
- [42] Q. Menet. Inverse of frequently hypercyclic operators. *J. Inst. Math. Jussieu* **21** (2022), 1867–1886.
- [43] T. K. S. Moothathu. Two remarks on frequent hypercyclicity. *J. Math. Anal. Appl.* **408** (2013), 843–845.
- [44] M. Murillo-Arcila and A. Peris. Strong mixing measures for linear operators and frequent hypercyclicity. *J. Math. Anal. Appl.* **398** (2013), 462–465.
- [45] D. Newton and W. Parry. On a factor automorphism of a normal dynamical system. *Ann. Math. Statist.* **37** (1966), 1528–1533.
- [46] M. Nicol and K. Petersen. Ergodic theory: basic examples and constructions. *Mathematics of Complexity and Dynamical Systems*. Ed. R. A. Meyers. Springer, New York, 2012, pp. 264–288.
- [47] W. Parry. *Topics in Ergodic Theory*. Cambridge University Press, Cambridge, 1981.
- [48] V. Peller. *Hankel Operators and Their Applications (Springer Monographs in Mathematics)*. Springer, New York, 2003.
- [49] M. Queffélec. *Substitution Dynamical Systems – Spectral Analysis (Lecture Notes in Mathematics, 1294)*. 2nd edn. Springer, 2010.
- [50] C. J. Read. The invariant subspace problem for a class of Banach spaces. II. Hypercyclic operators. *Israel J. Math.* **63** (1988), 1–40.
- [51] R. Sanders and S. Shkarin. Existence of disjoint weakly mixing operators that fail to satisfy the Disjoint Hypercyclicity Criterion. *J. Math. Anal. Appl.* **417** (2014), 834–855.
- [52] S. Shkarin. Universal elements for nonlinear operators and their applications. *J. Math. Anal. Appl.* **348** (2008), 193–210.
- [53] S. Shkarin. On the spectrum of frequently hypercyclic operators. *Proc. Amer. Math. Soc.* **137** (2009), 123–134.
- [54] S. Shkarin. A short proof of existence of disjoint hypercyclic operators. *J. Math. Anal. Appl.* **367** (2010), 713–715.
- [55] B. Szőkefalvi-Nagy, C. Foias, H. Bercovici and L. Kérchy. *Harmonic Analysis of Operators on Hilbert Space*. Springer, New York, 2010.
- [56] M. Urbański, M. Roy and S. Munday. *Non-invertible Dynamical Systems*. Vol. 1. De Gruyter, Berlin, 2022.
- [57] P. Walters. *An Introduction to Ergodic Theory (Graduate Texts in Mathematics, 79)*. Springer, Berlin, 1982.