

Conclusion: The evidence regarding the effectiveness of interventions to reduce C-spine imaging in adult ED patients with stable neck trauma is inconclusive. Given the national and international focus on improving appropriateness and reducing unnecessary imaging through campaigns such as Choosing Wisely®, additional interventional research in this field is warranted.

Keywords: diagnostic imaging, cervical spine, intervention

LO84

Computed tomography use for headache presentations to emergency departments in Alberta: regional, site and physician level variation

L. Krebs, MPP, MSc, C. Alexiu, BSc, C. Villa-Roel, MD, MSc, PhD, S.W. Kirkland, MSc, L. Gaudet, BSc, MSc, B.R. Holroyd, MD, MBA, M. Ospina, PhD, C. Pryce, BScN, MN, J. Bakal, PhD, S.E. Jelinski, PhD, DVM, B.H. Rowe, MD, MSc, University of Alberta, Edmonton, AB

Introduction: Headaches are a common emergency department (ED) presentation. Evidence demonstrates that computed tomography (CT) imaging varies significantly within and across sites. This study explored CT ordering and variation among headache presentations across Alberta EDs. **Methods:** Administrative health data for Alberta were obtained from the National Ambulatory Care Reporting System (NACRS) for all adult (>17 years) headache (ICD-10-CA: G44, G43, R51) ED visits from 2011-2015. Patients with a primary or secondary diagnosis code of headache were included. Exclusions were: sites without CT scanners, Canadian Triage and Acuity Scale score of 1, patients with trauma or external mechanism of injury (e.g., ICD-10-CA S,T,V,W,X,Y), or enhanced/contrast CTs. NACRS data were linked with Alberta Health Services' (AHS) diagnostic imaging data. Preliminary analysis on imaging variation at the zone, ED site, and physician level was completed using SAS (v.9.4). Physicians who saw less than an average of 10 headache patients per year were excluded. **Results:** Overall, 98,804 headache presentations were recorded (~20,000/year; 8.5% average annual increase) in 30 EDs. The average proportion of visits receiving CT was 25.1% with an average 6.2% increase per year. CT ordering varied across AHS zones (Variation [V]:23%; range:9.6-32.7%). Site ordering variation was more dramatic (V:45%; range:1.4-46.5%). The greatest variation was observed among physicians (V:84 %; range: 0.0-83.7%) with mean ordering proportion of 28.7%. **Conclusion:** From 2011-2015, headache presentations and CT imaging for these patients in the ED increased. Substantial variation in CT ordering exists at multiple levels in Alberta. Further exploration of CT appropriateness is urgently needed.

Keywords: diagnostic imaging, headache, emergency department

LO85

Substantial variation in CTPE ordering patterns and diagnostic yield in a large group of specialty-trained emergency physicians

E. Lang, MD, J. Andruchow, MD, MSc, D. Grigat, MA, G. Innes, MD, A. McRae, MD, University of Calgary, Calgary, AB

Introduction: Computerized tomography for pulmonary embolism (CTPE) has come under increased scrutiny with recommendations for evidence-based use found on Choosing Wisely lists in both Canada and the US. However practice variation in ordering patterns and diagnostic yield have not been well-reported for the Canadian context. Our objective was to investigate practice variation in CTPE ordering and rule-in rates within a large group of specialty-trained emergency physicians. **Methods:** We undertook an analysis of a computerized physician order entry database from four tertiary care EDs covering a 12-month period from August 1, 2016 to July 30, 2016 with 31 419 visits for potential

pulmonary embolism (PE) as determined by a previously validated algorithm based on presenting complaints. CTPE utilization and diagnostic yield were determined for 149 physicians who ordered at least 10 studies over that time period. Outcomes of interest included CT utilization as determined by electronic order entry and a confirmed diagnosis of PE based on ICD-10 coding of the emergency visit. Descriptive statistics using medians, IQR and 95% confidence intervals are reported. This study is approved through REB14-0650 and is a component of a larger cluster RCT to improve CTPE utilization. **Results:** During the study period 2670 CTPE studies were ordered for potential PE patients representing 8.5% of the total with relevant complaints. We observed a 10-fold variation in CTPE ordering among physicians with rates as low as 2.7% and as high as 25%. The median rate of CTPE ordering for potential PE was 8.8% with an IQR of 6.0% to 11.7%. A total of 4146 CTPE studies were ordered during the study period with physicians ordering an average of 28 CTPE studies each; range 10-90. In terms of diagnostic yield, 591/4146 studies, or 14.3% (95% CI 13.2-15.3%) were associated with a diagnosis of PE. Diagnostic yield per physician ranged from 0 to 50%, with a median of 13.5% and an IQR of 7.6% to 21.4%. **Conclusion:** In this large, robust administrative dataset from four Canadian urban EDs, threshold for CTPE ordering varies widely among physicians as does diagnostic yield. Efforts to improve appropriate utilization are justified with an eye to reducing unnecessary radiation, costs and incidental findings.

Keywords: computerized tomography, Choosing Wisely, pulmonary embolism

LO86

Overutilization of computed tomography as a first-line investigation for patients presenting with suspected recurrent nephrolithiasis in the emergency department: a retrospective cohort study

J. Himelfarb, BSc, J.S. Lee, MD, MSc, D. Shelton, MD, University of Toronto, Toronto, ON

Introduction: Computed tomography (CT) has increasingly been used as a standard initial investigation for patients presenting to the Emergency Department (ED) with suspected nephrolithiasis. Compared to ultrasound, CT has increased system-level costs, ionizing radiation exposure and frequently does not alter management. For these reasons, Choosing Wisely (CW) recommends avoiding CT imaging of otherwise healthy patients younger than age 50 years presenting with symptoms of uncomplicated renal colic that have a known history of nephrolithiasis or ureterolithiasis. We aimed to evaluate the degree of utilization of CT imaging for this subgroup of patients in a tertiary care centre ED. **Methods:** A retrospective chart review was performed for all patients younger than 50 years who visited Sunnybrook Health Sciences Centre ED for six months between December 2015 and May 2016 with renal colic symptoms and a history of nephrolithiasis. Demographic data, relevant past medical history, clinical presentation, lab values, urology consultation, ED treatments administered, diagnostic imaging orders and dispositions were recorded for each eligible patient. **Results:** Out of 130 reviewed patient charts, 73 patients were identified with a previous history of nephrolithiasis and a presentation consistent with uncomplicated renal colic. 54 patients received ultrasound, KUB x-ray, or no imaging. The other 19 (26.0%) of these patients received an abdominal/pelvic CT with an indication of looking for renal or ureteral stones. Of the patients that received CT, none demonstrated significant findings warranting hospital admission or leading to identifiable changes in ED management. Five (26.3%) of these 19 patients had received a total of three to four CTs for renal colic during past Sunnybrook ED visits, while one had previously received 13 CTs. **Conclusion:** CT scans are often used as an initial diagnostic modality for suspected renal colic