

## The effect of dietary supplements of $\omega$ 3 polyunsaturated fatty acids on the fatty acid composition of platelets and plasma choline phosphoglycerides

BY T. A. B. SANDERS AND KATHERINE M. YOUNGER

*Department of Nutrition, Queen Elizabeth College, Campden Hill Road, London W8 7AH*

(Received 7 October 1980 – Accepted 25 November 1980)

1. The effects of dietary supplements of  $\omega$ 3 polyunsaturated fatty acids on the fatty acid composition of platelets and plasma choline phosphoglycerides were studied in vegans and in omnivores.
2. A supplement of 18:3 $\omega$ 3 led to an increase in 20:5 $\omega$ 3 but was less effective than one of 20:5 $\omega$ 3+22:6 $\omega$ 3.

Although it is not known if linolenic acid (18:3 $\omega$ 3) is essential its derivatives are important (Tinoco *et al.* 1979). Eicosapentaenoic acid (20:5 $\omega$ 3) is the precursor of the triene prostaglandins (Gryglewski *et al.* 1979) and when incorporated into platelet lipids may influence bleeding time (Sanders *et al.* 1980). Docosahexaenoic acid (22:6 $\omega$ 3) is a major component of human brain and retinal lipids and is found in its highest concentrations in the phosphoglycerides of synaptic membranes and rod outer segments implying that it has a role in neural transmission.

Radiotracer experiments *in vitro* suggest that man can convert 18:3 $\omega$ 3 to 20:5 $\omega$ 3 and 22:6 $\omega$ 3 (De Gomez Dumm & Brenner, 1975; Aerberhard *et al.* 1978) but this capacity may be limited (Dyerberg *et al.* 1980). Consequently dietary 20:5 $\omega$ 3 and 22:6 $\omega$ 3 may be important (Crawford & Sinclair, 1972), the only significant sources being fish, fish oil and offal (Shepherd *et al.* 1978; Southgate & Paul, 1978). Indeed, vegans whose diets are devoid of 20:5 $\omega$ 3 and 22:6 $\omega$ 3 have very much lower proportions of these fatty acids in their plasma choline phosphoglycerides than do omnivores (Sanders *et al.* 1977). It was suggested that the high value for the ratio, 18:2 $\omega$ 6:18:3 $\omega$ 3 in the vegan diets suppressed the transformation of 18:3 $\omega$ 3 to 20:5 $\omega$ 3 and 22:6 $\omega$ 3 and instead favoured the conversion of 18:2 $\omega$ 6 to 20:4 $\omega$ 6 and 22:4 $\omega$ 6. This being so then a dietary supplement of 18:3 $\omega$ 3 should reduce the value for 18:2 $\omega$ 6:18:3 $\omega$ 3 and lead to an increase in the proportions of 20:5 $\omega$ 3 and 22:6 $\omega$ 3 in this lipid fraction. In order to test this hypothesis, vegan and omnivore subjects were given a supplement of linseed oil, rich in 18:3 $\omega$ 3 for 2 weeks and changes in their blood lipids were monitored. So that a comparison could be made between dietary linolenate and its long-chain derivatives, the effect of a fish oil supplement, rich in 20:5 $\omega$ 3 and 22:6 $\omega$ 3, was also studied in the omnivores.

### EXPERIMENTAL

#### *Subjects, materials and methods*

Vegan subjects (aged 26–37 years) were contacted through the Vegan Society (47 Highlands Road, Leatherhead, Surrey) and omnivores (aged 23–47 years) were recruited from the staff and students of the Nutrition Department, Queen Elizabeth College. All the subjects considered themselves to be healthy. Consent was obtained from individuals and from the College Ethical Committee for all procedures.

In the first experiment, four vegan (three male, one female) and five omnivore subjects (three male, two female) were instructed to take 5 ml refined and deodorized linseed oil four times daily with meals for 2 weeks. In the second experiment, two male and two female omnivore subjects, who had 7 weeks previously taken the linseed oil supplement, took a

Table 1. *Fatty acid composition (wt%) of the two oils fed to the subjects*

Fatty acid	Maxepa	Linseed oil
14:0	7.01	0.03
14:1	0.36	—
15:0	0.63	—
15:1 + 15:0 Br	0.14	—
16:0	17.03	5.28
16:1	8.68	—
17:0	2.24	0.12
17:1 + 17:0 Br	1.11	—
18:0	4.79	3.53
18:1	12.75	19.79
18:2 $\omega$ 6	1.79	18.19
18:3 $\omega$ 3	0.46	53.93
20:0	2.19	—
20:1	1.57	—
20:5 $\omega$ 3	17.43	—
22:1	0.73	—
22:5 $\omega$ 3	3.20	—
22:6 $\omega$ 3	16.84	—

fish oil supplement (Maxepa) for 2 weeks (5 ml four times daily with meals). The oils used were provided by British Cod-liver Oils, Hull and were analysed for fatty acids by gas-liquid chromatography before use (Table 1). Fresh bottles of oil were issued each week and oil consumption was assessed by weighing the bottles. All subjects completed a 7-d-weighted dietary intake record. Nutrient intakes were calculated by computer using the tables of Southgate & Paul (1978). For foods not included in these tables additional values were obtained from other published sources (Platt, 1962; Watt & Merrill, 1963; Miller & Mumford, 1972; US Department of Health & Welfare & FAO, 1972; Shepherd *et al.* 1978) or from information provided by the manufacturers of proprietary products.

Venous blood samples were drawn from subjects after an overnight fast with minimum venostasis before and after taking the supplements. The blood samples were anticoagulated with EDTA, chilled to 4° and processed immediately. Platelet and plasma choline phosphoglycerides (Sanders *et al.* 1978) were prepared and transmethylated with sodium methoxide after elution from the thin-layer chromatography absorbent (Christie, 1973). The resulting methyl esters were analysed by gas-liquid chromatography on a Pye model 204 chromatograph equipped with flame ionization detectors: a 1.8 m  $\times$  4 mm internal diameter glass column packed with 100 g Silar 10C/kg Gaschrom Q (100–120 mesh) was used; oven temperature 200°, injection temperature 210° and detector temperature 260°; nitrogen carrier gas flow-rate 40 ml/min. Chromatographic peaks were integrated using a DP 88 integrator (W. G. Pye, Cambridge). Methyl esters were identified by comparison with mixtures of known composition. Analyses were confirmed on a less polar column packed with 100 g Silar 5C/kg Chromosorb W HP (80–100 mesh) operated at 210°. Statistical analyses of the results were performed using a one-tailed paired sample *t* test, with each subject acting as his or her own control. A two-sample two-tailed *t* test was used for between group comparisons.

#### RESULTS AND DISCUSSION

The vegans had followed their diet for an average of 10 years (range 5–14). Their diets comprised unrefined cereals, nuts, fruit, pulses and vegetables and were supplemented with

vitamin B<sub>12</sub>. Such diets do not result in detectable deficiency (Sanders, 1978). Their mean energy intake was 10 MJ (range 6–14 MJ) compared with 8 MJ (range 6–10 MJ) in the omnivores; all the diets provided at least one-tenth of the dietary energy as protein. Although their average fat intake was similar to that of the omnivores ( $84 \pm 5$  v.  $87 \pm 5$  g/d), its composition was very different: their intake of saturated fatty acids was much lower ( $16 \pm 1$  v.  $38 \pm 3$  g/d in the omnivores) and that of linoleic acid (18:2 $\omega$ 6) was almost four times greater ( $28 \pm 2$  v.  $7 \pm 1$  g/d); their average intake of linolenic acid (18:3 $\omega$ 3) was similar ( $2.0 \pm 0.6$  v.  $1.3 \pm 0.2$  g/d). Therefore, the ratio, 18:2 $\omega$ 6:18:3 $\omega$ 3 was much greater in the vegans than in the omnivores ( $16 \pm 3.5$  v.  $6 \pm 1.0$ ). This high value would be expected to suppress the conversion of 18:3 $\omega$ 3 to its long-chain derivatives. Indeed, the differences in the fatty acid composition of the plasma choline phosphoglycerides between vegans and omnivores were similar to those previously reported (Sanders *et al.* 1978): the proportion of 20:5 $\omega$ 3 and especially 22:6 $\omega$ 3 were much lower (Table 2) and those of 20:4 $\omega$ 6 ( $13.9 \pm 0.31$  v.  $11.3 \pm 0.65$  wt%;  $P < 0.05$ ) and 18:2 $\omega$ 6 ( $29.7 \pm 1.5$  v.  $24.5 \pm 1.39$  wt%;  $P < 0.05$ ) were greater. Comparable differences were noted in the platelet phosphoglycerides.

Table 2. Proportions (wt%) of eicosapentaenoic (20:5 $\omega$ 3) and docosahexaenoic (22:6 $\omega$ 3) acids in plasma choline phosphoglycerides (CPG) and platelet phosphoglycerides (PG) of vegans and omnivores before and after 2 weeks of taking an oil supplement

(Results are expressed as mean values with their standard errors)

Group	No. of subjects	Supplement	20:5 $\omega$ 3				22:6 $\omega$ 3			
			Before		After		Before		After	
			Mean	SE	Mean	SE	Mean	SE	Mean	SE
Plasma CPG										
Omnivores	5	Linseed oil	1.3	0.37	2.7*	0.50	4.3	0.33	3.7	1.03
Vegans	4	Linseed oil	0.3†	0.06	1.0*†	0.24	1.3†	0.30	1.3†	0.46
Omnivores	4	Maxepa†	1.5	0.27	6.7*†	0.96	3.9	0.45	4.5*	0.34
Platelet PG										
Omnivores	5	Linseed oil	0.6	0.10	1.2*	0.18	3.0	0.33	2.9	0.09
Vegans	4	Linseed oil	0.3†	0.04	0.3†	0.04	0.9†	0.17	1.2†	0.26
Omnivores	3	Maxepa†	1.2	0.17	4.1*†	0.10	2.7	0.26	3.9*†	0.27

Statistical significance of difference between mean values; \*  $P < 0.05$  compared with before the supplement; †  $P < 0.05$  compared with the corresponding values obtained for the omnivores who took the linseed oil supplement.

† British Cod-liver Oils, Hull.

The linseed oil supplement, which provided an average of 6.5 g 18:3 $\omega$ 3/d, reduced the ratio, 18:2 $\omega$ 6:18:3 $\omega$ 3 from six to one in the omnivores. This led to a doubling of the proportion of 20:5 $\omega$ 3 in both plasma choline phosphoglycerides and in the platelets. The same supplement reduced the value for the ratio from sixteen to three in the vegans and led to a threefold increase in the proportion of 20:5 $\omega$ 3 in the plasma choline phosphoglycerides but not in the platelets. These observations provide evidence that man can convert 18:3 $\omega$ 3 to 20:5 $\omega$ 3. The higher proportion of linoleic acid in the adipose tissue of vegans compared with omnivores (Sanders *et al.* 1978) would tend to buffer any short-term attempt to reduce the ratio 18:2 $\omega$ 6:18:3 $\omega$ 3 in the body. Consequently a more marked increase in the proportion of 20:5 $\omega$ 3 might have occurred in the vegans had the supplement been continued for longer.

The failure of the linseed oil supplement to increase the proportion of 22:6 $\omega$ 3 in these lipid fractions in both vegans and omnivores is consistent with the hypothesis that the rate of  $\Delta$ 4 desaturation is slow in man (Sanders & Naismith, 1979).

The fish oil supplement (Maxepa) provided 2.5 g/d each of 20:5 $\omega$ 3 and 22:6 $\omega$ 3. This led to a fourfold increase in the proportion of 20:5 $\omega$ 3 and a less marked increase in 22:6 $\omega$ 3 in both the plasma choline phosphoglycerides and the platelets in the omnivores; these changes occurred mainly at the expense of 18:2 $\omega$ 6 in the plasma choline phosphoglycerides and 20:4 $\omega$ 6 and 22:4 $\omega$ 6 in the platelets. Similar changes have been reported in subjects given a cod-liver oil supplement (Sanders *et al.* 1980). The relatively small amounts of 20:5 $\omega$ 3 and 22:6 $\omega$ 3 provided by the Maxepa supplement had a far greater effect than the larger amount of 18:3 $\omega$ 3 provided by the linseed oil. Thus the effect of administering different  $\omega$ 3 polyunsaturated fatty acids is not equivalent and dietary 20:5 $\omega$ 3 and 22:6 $\omega$ 3 are a more effective means of increasing the proportions of these acids in tissue lipids than a reduction in the ratio 18:2 $\omega$ 6:18:3 $\omega$ 3 in the diet. We were unable to make a trustworthy estimate of the 20:5 $\omega$ 3 and 22:6 $\omega$ 3 in the diets of the omnivore subjects owing to the lack of reliable food composition information. However, our results imply that the major source of 22:6 $\omega$ 3 in membrane phospholipids is obtained preformed rather than indirectly from dietary linolenic acid. This would explain the higher proportion of 22:6 $\omega$ 3 in the lipids of the omnivores compared with the vegans. Further research is necessary to determine the optimum levels of C<sub>20-22</sub>  $\omega$ 3 polyunsaturated fatty acids in tissues.

The authors are grateful to British Cod-liver Oils.

#### REFERENCES

- Aerberhard, E. E., Corbo, L. & Menkes, J. H. (1978). *Lipids* **13**, 758.  
 Christie, W. W. (1973). *Lipid Analysis*. Oxford: Pergamon.  
 Crawford, M. A. & Sinclair, A. J. (1972). In *Lipids, Malnutrition and the Developing Brain*, p. 267 [K. Elliot and J. Knight, editors]. Amsterdam: Elsevier.  
 De Gomez Dumm, I. N. T. & Brenner, R. R. (1975). *Lipids* **10**, 315.  
 Dyerberg, J., Bang, H. O. & Aagard, O. (1980). *Lancet* **i**, 199.  
 Gryglewski, R. J., Salmon, J. A., Ubatuba, F. B., Weatherley, B. C., Moncada, S. & Vane, J. R. (1979). *Prostaglandins* **18**, 453.  
 Miller, D. S. & Mumford, P. (1972). *Pl. Fd hum. Nutr.* **2**, 201.  
 Platt, B. S. (1962). *Med. Res. Counc. Spec. Rep. Ser.* no. 302.  
 Sanders, T. A. B. (1978). *Pl. Fds Man* **2**, 181.  
 Sanders, T. A. B., Ellis, F. R. & Dickerson, J. W. T. (1977). *Lancet* **i**, 751.  
 Sanders, T. A. B., Ellis, F. R. & Dickerson, J. W. T. (1978). *Am. J. clin. Nutr.* **31**, 805.  
 Sanders, T. A. B., Haines, A. P. & Vickers, M. (1980). *Lancet* **i**, 1189.  
 Sanders, T. A. B. & Naismith, D. J. (1979). *Br. J. Nutr.* **41**, 619.  
 Shepherd, A. J., Iverson, J. & Weihrauch, J. L. (1978). In *Handbook of Lipid Research*, vol. 1, p. 341 [A. Kuksis, editor]. London: Plenum Press.  
 Southgate, D. A. T. & Paul, A. A. (1978). *McCance & Widdowson's The Composition of Foods*. London HM Stationery Office.  
 Tinoco, J., Babcock, R., Hincenbergs, I., Medwadowski, B. & Miljanich, P. (1979). *Lipids* **14**, 166.  
 US Department of Health & Welfare & FAO. (1972). *Food Composition Tables for Use in S.E. Asia*. Rome: FAO.  
 Watt, B. K. & Merrill, A. G. (1963). *Composition of foods. Handbook* no. 8. Washington: US Department of Agriculture.