INSTRUCTION TO CONTRIBUTORS

The Journal of Helminthology publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology, London School of Hygiene and Tropical Medicine, Winches Farm Field Station, 395 Hatfield Road, St Albans, Herts AL4 0XQ, England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0×20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0×20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecology* and *Physiology of parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains of Schistosoma haematobium. Journal of Helminthology, 47, 49–59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Journal of Helminthology volume 57 • no. 4 • dec 1983

S. GHATAK

Pages

Contents

Some properties of the surface of nematode larvae. (Research Note). MICHELE

JUNGERY, C. D. MACKENZIE and BRIDGET M. OGILVIE 291-295 Studies on the cryopreservation of eggs of Angiostrongylus cantonensis. SHOJI UGA, KAZUYOSHI ARAKI, TAKEO MATSUMURA and NOBORU IWAMURA 297-303 The use of lyophilized Schistosoma mansoni eggs as antigenic particles in a radioimmunoassay. J. HAMBURGER, H. MOSCOVICI and S. LUSTIGMAN 305-312 Monoamine oxidase in adult Ascaridia galli. S. K. MISHRA, RAMANUJ SEN and 313-318 Cryopreservation of Dirofilaria immitis microfilariae and third-stage larvae. (Research Note). J. B. LOK, M. MIKA-GRIEVE and R. B. GRIEVE 319-324 Oesophagostomum columbianum: gamma-irradiated third-stage larvae for immunization in lambs. R. L. SHARMA and D. N. DHAR 325-330 Description of musculature differences in spargana of Spirometra (Cestoda; Pseudophyllidea) and tetrathyridia of Mesocestoides (Cestoda; Cyclophyllidea) and their value in identification. (Research Note). KARIN I. ANDERSEN 331-334 Uptake of adenosine 3',5'-cyclic monophosphate and isoproterenol by filarial parasites. KHALID MASOOD, V. M. L. SRIVASTAVA, D. P. SINGH, R. K. CHATTERJEE and A. B. SEN 335-338 Surface morphology of Gastrodiscoides hominis Lewis & McConnell, 1876) Leiper, 1913 (Trematoda: Digenea) as revealed by scanning electron microscopy. VEENA TANDON and S. C. MAITRA 220 242

TANDON and S. C. MAITRA	339-342
Nematospiroides dubius: factors affecting the primary response to SRBC in infected mice. NAJAH M. H. ALI and JERZY M. BEHNKE	343-353
Onchocerca raillieti: adult location and skin distribution of the microfilaria in Sudanese donkeys. SAAD E. EL SAMMANI and HUSSEIN S. HUSSEIN	355-360
Subject Index	361
Author Index	364
Contents of Volume	i-III

and published by the London School of Hygiene and Tropical Medicine