

## EPV0111

**Diagnostic difficulties in bipolar disorder type II**

L. Núñez\*, C. A. del Val, A. F. Merlos, E. X. G. Vivero, C. C. Carrascosa and M. H. Naval

Psychiatry, Instituto de Psiquiatria y Salud Mental, Hospital Clínico San Carlos, Madrid, Spain

\*Corresponding author.

doi: 10.1192/j.eurpsy.2023.1468

**Introduction:** Bipolar disorder is one of the top 10 medical causes of disability according to the WHO and despite this, its diagnosis can be delayed up to 10 years after the appearance of the first symptoms of the disease. A major reason for the difficult diagnosis is the challenge of differentiating bipolar disorder type II from unipolar depression and borderline personality disorder, especially in those patients with no clear history of hypomania.

**Objectives:** To present a case report of a bipolar disorder undiagnosed for years to remark the importance of recognizing premorbid symptoms of the disease in order to implement an early intervention that potentially improves the prognosis of patients.

**Methods:** We compiled the patient's complete medical history and we carried out a non-systematic review of literature containing the key-words "bipolar disorder type II" and "diagnosis".

**Results:** We present the case of a 48-year-old woman going through a depressive episode, multiple suicide attempts and more than 10 admissions in the Acute Inpatient Psychiatric Unit. For 3 years, the evolution was torpid with a significant multidomain cognitive impairment in a previously functional patient. Different antidepressant treatments were tested, however they were not tolerated due to adverse effects such as anxiety, insomnia and nervousness. After considering multiple differential diagnoses, bipolar disorder type II was finally diagnosed. A hypomanic episode that took place after 3 sessions of electroconvulsive therapy during an admission for depression, allowed to guide the diagnosis and after the introduction of Lithium and Quetiapine as treatment, the patient experienced a complete remission of the symptoms.

**Conclusions:**

1. It is important to consider the differential diagnosis of bipolar disorder type II due to its impact on the patient's life.
2. An early diagnosis improves the course and prognosis of the disease.
3. Patients resistant or intolerant to antidepressant treatment could have undiagnosed bipolar disorder.

**Disclosure of Interest:** None Declared

## EPV0112

**Syndrome of Irreversible Lithium-Effectuated NeuroToxicity: SILENT, but not innocent**

M. S. Bicho\*, J. M. Coelho and H. J. Fontes

Psychiatry, Hospital do Divino Espírito Santo de Ponta Delgada, EPER, Ponta Delgada, Portugal

\*Corresponding author.

doi: 10.1192/j.eurpsy.2023.1469

**Introduction:** Lithium is one of the main drugs used in Bipolar Affective Disorder. However, it has a narrow therapeutic window,

which requires close monitoring and progressive dose adjustment, according to serum levels, clinical response and the appearance of side effects. The term 'SILENT' explains descriptively persistent neurological sequelae related to lithium salt intoxication when symptoms persist for more than 2 months after stopping treatment. SILENT Syndrome is more common in females, at ages ranging from 21 to 77 years and is characterized mainly by avermian-type cerebellar disorder, persistent extrapyramidal syndrome, brainstem dysfunction and dementia of varying severity. It can also result in apraxia of the body, changes in the coordination and balance, dysarthria, as well as intentional and kinetic cerebellar tremor, involuntary movements of orofacial dyskinesias or resting tremor.

**Objectives:** The authors intend to review the relevant and current literature in order to extend the knowledge about this condition and find the best conducts for clinical practice.

**Methods:** Non-systematic literature review.

**Results:** Complications from the use of lithium known in the medical literature include mainly nephrotoxicity, endocrine alterations and neurotoxicity.

The neurotoxic effects of lithium usually occur at high serum concentrations. However, they can also occur with lithium in the therapeutic range, and memory, attention and ataxia impairment may be some of the permanent sequelae.

The etiopathogenesis is unclear, but demyelination has been detected in multiple brain regions, mainly in the cerebellum. The mechanism of lithium-induced cerebellar injury is believed to be mediated by the entry of calcium into the cells of this organ.

The main factors that predispose to greater side effects and risk of toxicity are patients with decreased renal function, advanced age, use of diuretics, dementia, pregnancy, low sodium intake and physical illness with vomiting and/or diarrhea.

**Conclusions:** Lithium is a drug used mostly in affective disorders and given the narrow therapeutic window, it requires close monitoring in order to avoid side effects that can be permanent. In this way, it is important to review the factors that increase the lithium toxicity and make recommendations about it.

**Disclosure of Interest:** None Declared

## EPV0113

**PEAKS AND VALLEYS: BIPOLAR DISORDER, RAPID CYCLERS AND ENERGY DRINKS CONSUMPTION**

M. Calvo Valcárcel\*, M. A. Andreo Vidal, P. Martinez Gimeno, P. Pando Fernández, B. Rodriguez Rodriguez, N. Navarro Barriga, M. Fernández Lozano, M. J. Mateos Sexmero, M. D. C. Vallecillo Adame, T. Jimenez Aparicio, C. de Andres Lobo, M. Queipo de Llano de la Viuda, A. A. Gonzaga Ramirez and G. Guerra Valera

PSYCHIATRY, Hospital Clínico Universitario de Valladolid, VALLADOLID, Spain

\*Corresponding author.

doi: 10.1192/j.eurpsy.2023.1470

**Introduction:** Bipolar Disorder (BD) is considered a serious mental disorder characterized by a changing mood that fluctuates between two completely opposite poles. It causes pathological and recurrent mood swings, alternating periods of exaltation and grandiosity with periods of depression. We talk about rapid cyclers when four or more manic, hypomanic or depressive episodes have occurred