Current perspectives on global sugar consumption: definitions, recommendations, population intakes, challenges and future direction

Janette Walton¹ , Haley Bell⁴, Roberta Re⁴ and Anne P Nugent^{2,3}

¹Department of Biological Sciences, Munster Technological University, Cork, Republic of Ireland

²Institute for Global Food Security, School of Biological Sciences, Queens University Belfast, Belfast, Northern Ireland

³Institute of Food and Health, School of Agriculture and Food Sciences, University College Dublin, Belfield, Dublin 4, Ireland

⁴Cambridge Food Science, Cambridge, UK

Abstract

Currently, there is considerable emphasis on the relationship between dietary sugar consumption and various health outcomes, with some countries and regions implementing national sugar reduction campaigns. This has resulted in significant efforts to quantify dietary sugar intakes, to agree on terms to describe dietary sugars and to establish associated recommendations. However, this information is infrequently collated on a global basis and in a regularised manner. The present review provides context regarding sugar definitions and recommendations. It provides a global review of the available data regarding dietary sugar intake, considering forms such as total, free and added sugars. A comprehensive breakdown of intakes is provided by age group, country and sugar form. This analysis shows that free sugar intakes as a percentage of total energy (%E) are the highest for children and adolescents (12–14%E) and the lowest for older adults (8%E). This trend across lifecycle stages has also been observed for added sugars. The available data also suggest that, while some reductions in sugar intake are observed in a few individual studies, overall intakes of free/added sugars remain above recommendations. However, any wider conclusions are hampered by a lack of detailed high-quality data on sugar intake, especially in developing countries. Furthermore, there is a need for harmonisation of terms describing sugars (ideally driven by public health objectives) and for collaborative efforts to ensure that the most up-to-date food composition data are used to underpin recommendations and any estimates of intake or modelling scenarios.

Keywords: Free sugar: Added sugar: Dietary surveys: Dietary sugars: Sugar intake/consumption

(Received 26 July 2019; revised 6 July 2021; accepted 29 July 2021)

Introduction

Sugar has been described as a 'ubiquitous term', difficult to both define and measure, which may be in part due to its dual function as both a nutrient and as an ingredient⁽¹⁾. A form of carbohydrate, chemically, the term 'sugars' usually describes mono- and disaccharides, with examples of monosaccharides being glucose, galactose and fructose and examples of disaccharides being sucrose, lactose and maltose. Examples of some natural contributing food sources include fruits (fructose, glucose, sucrose), honey (fructose, glucose, sucrose), milk and milk products (galactose, lactose), and table sugar (sucrose). Such '-ose' sugars can also form the building blocks of other carbohydrate types, such as oligosaccharides and polysaccharides. In addition, sugars in many of the above forms may be added as an ingredient to foods during processing or preparation to preserve food and/or to confer functional properties, such as viscosity, texture, body and colour, or at the table by consumers to add sweetness^(1,2). As a result, there is a broad range of foods containing sugars with varying quantities depending on the desired function, individual

preference and food form. As a carbohydrate form, sugars, regardless of type, contribute 3.75 kcal/g, which is equivalent to the contribution from starch.

Whilst the relationship between the frequency of consumption of foods containing sugars (particularly sucrose) and dental caries is well established^(3–5), there is debate concerning the relationship between dietary sugar intake and other health outcomes such as obesity, type 2 diabetes and cardiovascular disease. The evidence has been critiqued and systematically reviewed both in scientific publications^(6,7) and by advisory bodies and competent authorities^(3–5,8,9) with no firm conclusion that dietary sugars are directly related to diet-related diseases other than as a source of dietary energy particularly implicated as a contributor towards positive energy balance. As an individual dietary sugar, fructose has been identified of note, particularly with respect to metabolic health and hyperuricemia; however, as yet, no agreed conclusion has been reached regarding fructose^(10,11). The strongest available evidence of any link between dietary sugars and health is for body weight, and following the consumption of sugar-sweetened beverages rather than sugars or a single sugar

* Corresponding author: Janette Walton, email janette.walton@mtu.ie

Table 1. Definitions of sugars as used in dietary recommendations

Term	Definition	Issuing body
Total sugars	Glycaemic carbohydrates composed of one to two monomers found in food. Includes monosaccharides (glucose, fructose, galactose) and disaccharides (sucrose, lactose, maltose, trehalose)	FAO/WHO, 1998 ⁽²²⁾ EFSA, 2010 ⁽³⁾
Added sugars	Sugars (mono- and disaccharides) used as ingredients in processed and prepared foods and sugars eaten separately or added to foods at the table	USDA/HHS, 2020 ⁽²³⁾ IoM, 2005 ⁽²⁴⁾ EFSA 2010 ⁽³⁾ Nordic Council of Ministers, 2012 ⁽²⁵⁾
Free sugars	Includes monosaccharides (glucose, fructose and galactose) and disaccharides (sucrose, lactose, maltose and trehalose) added to foods by the manufacturer, cook, or consumer plus sugars naturally present in honey, syrups, fruit juices and fruit juice concentrates	WHO 2003, 2015 ^(4,26) SACN 2015 ⁽⁵⁾
Non-milk extrinsic sugars	Sugars not located within the cellular structure of a food, such as those found in fruit juice, honey and syrups, and those added to processed foods, excluding lactose in milk. Definition originated in UK and in 2015 use was superseded by use of 'free sugars'	UK DoH 1991 ⁽²⁷⁾

FAO/WHO (Food and Agriculture Organization and the World Health Organization), 1998. Carbohydrates in human nutrition. Report of a Joint FAO/WHO Expert Consultation, Rome, Italy, 14–18 April 1997. EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2010. Scientific Opinion on Dietary Reference Values for carbohydrates and dietary fibre.

EFSA Journal 2010;8(3): 1462, 77 pp. <https://doi.org/10.2903/j.efsa.2010.1462>.

IoM (Institute of Medicine), 2002. Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein, and amino acids.

per se^(5,12,13). At the request of a number of countries in the wider Nordic region, the European Food Safety Authority (EFSA) is currently (2020) undertaking a renewed review of the evidence to provide scientific advice on dietary sugars and to derive a tolerable upper intake level (UL) for total/added/free dietary sugars⁽¹⁴⁾. A draft for public consultation is expected in mid-2021.

In light of this continuing focus on sugars and their role in health, it is important to review current estimates of intake globally and to place these in context against recommendations. This allows generation of a benchmark for ongoing surveillance. Previously, there have been three reviews covering dietary sugars intakes, one with a global perspective⁽¹⁵⁾ a second focused on European countries⁽¹⁶⁾ and a third focusing on Latin American countries⁽¹⁷⁾. However, there exists a need to capture more recent data on a global basis and in a consistent balanced manner. The current review provides an up-to date critique of sugars in terms of their classification and dietary recommendations. A secondary analysis of the available data on dietary sugar intakes worldwide is described, followed by a perspective on the role of sugars in the diet globally.

Classification and definitions of dietary sugars

The term 'sugars' is conventionally used to define the mono- and disaccharides in foods (also known as total sugars). This includes naturally occurring sugars as well as those added to a food or drink (such as sucrose, fructose, maltose, lactose, honey, syrup, corn syrup, high-fructose corn syrup, molasses and fruit juice concentrate) but does not include polyols. Other definitions of dietary sugars are typically either linked to function in terms of establishing dietary recommendations, or they inform regulations and food labelling. With respect to definitions used to underpin dietary recommendations, the key terms are 'total sugars' (as defined above), 'free sugars', 'added sugars' and, historically, 'non-milk extrinsic sugars' (UK only). Table 1 provides a brief description of these definitions and issuing guidance body.

Of note, the term 'total sugars' provides the most encompassing measure of dietary sugars intake, used widely in food labelling and for informing consumers about the total sugars content of foods and drinks. The use of the term 'total sugars' in food labelling is mainly due to there currently being no accepted method for empirically measuring free sugars or non-milk extrinsic sugars in a food/drink. However, it is less frequently used in dietary recommendations⁽¹⁸⁾. Terms that are used more frequently in dietary recommendations include 'free' or 'added' sugars⁽¹⁹⁾. Both EFSA (Europe) and the Institute of Medicine (United States) define added sugars as all sugars that are added to foods and beverages by the manufacturer, cook and consumer, but do not include natural sources of sugars such as those included in unsweetened fruit juices^(3,20). On the other hand, free sugars, as defined by WHO and EFSA, include all added sugars but also those sugars naturally present in honey, syrups and fruit juices^(4,21). Such divergence in definitions for sugars is reflected in varying descriptions when reporting on dietary intake, in varying use of the terms in food regulations and even in defining the composition of formulae and foods for infants and young children⁽¹⁾.

Dietary guidelines for sugars consumption

Universally, most food-based dietary guidelines make the sensible and pragmatic suggestion for populations to limit the consumption of foods of low nutrient density, such as foods of low dietary quality that are high in calories, sugar, salt and/or fat, especially saturated fat. In recent years, many government and health organisations worldwide have revisited their guidance for dietary sugar intake, particularly in relation to maximum recommended intakes for added or free sugars. Some of these goals are average population maximum targets, and some are recommendations for intakes at the individual level. Examples of current recommendations relating to quantitative guidelines for dietary sugars are included in Table 2 and are summarised briefly below.

For free sugars, the World Health Organisation (WHO) recommends that all individuals, both adults and children, reduce their intake of free sugars to <10% energy (E) intake and suggest a further conditional recommendation of reduction to <5% E for further health benefits⁽⁴⁾. This <5%E value is also proposed by both the UK Scientific Advisory Committee on Nutrition (SACN) (as an average population maximum target) and by the European Society for Paediatric Gastroenterology Hepatology and Nutrition (ESPGHAN) (as individual guidance). Of note, there is a differing body of evidence underpinning these recommendations. Whilst SACN focus on intervention and prospective cohort studies relating to the intake of sugars to energy intake and to an increased risk of dental caries and type 2 diabetes mellitus⁽⁵⁾, the ESPGHAN recommendation⁽¹⁾ is informed by the conclusions of both WHO and SACN^(4,5) rather than by independent analyses. Of note also, ESPGHAN suggest children under 2 years of age should have even lower intakes than the <5%E⁽¹⁾.

For added sugars, both the Nordic countries and the United States recommend limiting the intake of added sugars to <10% E^(23,25). In both cases, these recommendations were in the context of healthy dietary patterns rather than any link with disease risk, a major difference to recommendations for free sugars. Whilst the American Heart Association (AHA) in the United States have listed a UL for children⁽²⁹⁾, this was not reflected in US national guidelines, which were unable to establish this guidance value owing to lack of evidence. In the absence of such scientific evidence, the IoM suggested that intakes above 25%E from added sugars may be associated with displacement of foods that are major sources of essential micronutrients, but they were unable to set a UL at that time (2002) and have not set one since⁽⁹⁾. Of note, the French Agency for Food, Environmental and Occupational Health & Safety (ANSES) have set an upper limit for total sugars of 100 g/d for adults, but this excludes lactose and galactose naturally present in milk and dairy products⁽⁸⁾. As mentioned previously, the EFSA is currently mandated to provide scientific advice on dietary sugars and to establish a UL if the available data allow it, having previously being unable to set such a benchmark in 2010⁽³⁾. It is unknown yet how an establishment of such a UL may influence national dietary guidelines within Europe: currently, whilst some European countries have quantitative limits established, such as the Nordic countries, the UK and France, others such as Germany have adopted the WHO recommendations⁽²⁸⁾. What is currently known regarding the EFSA scientific evaluation is that the exposure assessment will focus on intake of free sugars and that the disease endpoints of interest will include incidence/severity of dental caries and incidence of metabolic diseases. Other less direct endpoints will include measures of body fatness, ectopic fat distribution, glucose homeostasis and blood lipids among others⁽¹⁴⁾. As we await the findings of this EFSA scientific opinion, recent reviews suggest that, for numerical dietary sugar recommendations, more explicit quantification, justifications and communication of uncertainty factors, together with greater international harmonisation for both exposures and outcomes, would help to minimise any variation in recommendations^(17,19,30). An update of these reviews is included below.

Sugar intake data

Selection of data

Recent reviews of dietary surveys (published between 2014 and 2018) were used to identify a list of countries that have undertaken dietary surveys and the dates that these surveys were performed^(31–34). These were then supplemented with searches of databases such as Science Direct, PubMed and Google Scholar using terms such as ‘national dietary survey’ and ‘sugar(s) consumption’. As nutrient and food intakes derived from national food consumption survey data are typically published in reports, rather than peer-reviewed papers, the data presented in the present review were primarily collected from various reports and publications of national dietary surveys. Surveys and reports known to the authors of the present study were also included. Peer review publications were used to obtain additional data, for countries including New Zealand, Mexico, Spain, Brazil, the United States and Ireland. Studies were included if the data were (i) nationally representative, (ii) available in the public domain, (iii) summarised (no raw data were analysed) and (iv) collected at the individual level, and if (v) sugar intake was reported as a nutrient rather than a food group. There was no limit to the study size, and data from all available direct dietary assessment methodology were included, including food diaries, diet recalls or food frequency questionnaires.

Presentation of data

In an attempt to assess the available evidence which quantifies dietary sugar intake, a secondary analysis of published intake data for all sugar forms has been completed, and the findings (as reported in their original form) are reported in Tables 3–7 and summarised below. Energy values have been included to allow others to make crude estimates of sugars intakes as a percentage of energy where these data were not in the original reports.

Values are reported for estimated intakes of energy, total sugars, added sugars, free sugars or sucrose as available. Where energy was provided in kcal, it was converted to MJ using a factor of 238.85. Data are presented as provided in the reports and publications, and where possible, data have been divided by age range and not subdivided by sex. No statistical analysis has been performed on these data sets, and the results are purely descriptive. Under-reporting of food consumption can influence the accuracy of dietary intake data, resulting in an underestimate of energy and nutrients. The data presented in the tables do not account for under-reporting as under-reporters are not typically excluded from data presented in reports of food and nutrient consumption. Additionally, the data are usually presented as a percentage of energy intake, which partially accounts for this but will still result in an underestimate of sugar intake if foods/drinks that are high in sugar are selectively under-reported. It should also be noted that there are significant limitations in comparing sugar intakes across countries (from different time points) since the composition of individual foods and beverages may have changed drastically over that time and may not be reflective of current consumption.

Table 2. Examples of dietary reference values for sugars globally

Organisation	Population group	Year	Recommendation	Evidence base	Reference
German Nutrition Society	Children and adults	2019	Maximum free sugar intake of no more than 10 % of total energy intake	Adopted from WHO	Ernst <i>et al.</i> 2019 ⁽²⁸⁾
European Society for Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN) Committee on Nutrition	Children	2017	Intakes of free sugars should be reduced in children and adolescents (2–18 years) with a desirable upper limit of less than 5 % energy intake and should be even lower in those younger than 2 years	Built on the conclusions of other authoritative bodies (WHO, 2015, SACN, 2015, Vos, 2016)	Fidler <i>et al.</i> 2017 ⁽¹⁾
American Heart Association (AHA)	Children	2017	Children should consume ≤ 25 g (100 kcal or ≈ 6 tea-spoons) of added sugars per day, and added sugars should be avoided for children < 2 years of age		Vos <i>et al.</i> 2017 ⁽²⁹⁾
French Agency for Food, Environmental and Occupational Health & Safety (ANSES)	Adult population	2016	Upper intake limit for total sugars of 100 g/d (excluding lactose and galactose naturally present in milk and dairy products)	Based on calculation from minimum daily intake of fructose for which a significant increase in blood triacylglycerol were observed (in intervention studies)	ANSES, 2016 ⁽⁸⁾
World Health Organisation (WHO)	Children and adults	2015	Intake of free sugars should not exceed 10 % of total energy intake (strong recommendation) and should be less than 5 % of total energy intake (conditional recommendation)	Based on risk of non-communicable diseases with a particular focus on weight gain and dental caries	WHO, 2015 ⁽⁴⁾
U.S. Department of Health and Human Services and U.S. Department of Agriculture (2015)	General Population	2020	Consume less than 10 % of calories per day from added sugars	Target based on food pattern modelling and national data on intakes of calories from added sugars that demonstrate the public health need to limit calories from added sugars to meet food group and nutrient needs within calorie limits	USDA, 2020 ⁽²³⁾
Scientific Advisory Committee on Nutrition (SACN) UK	2 years and over	2015	Recommended that the average population intake of free sugars should not exceed 5 % of total energy intake for age groups from 2 years upwards	Based on evidence of increased energy from associations with an increased risk of dental caries. Additional evidence related to that of sugar-sweetened beverages and weight gain in children and adolescents and an increased risk of type 2 diabetes in adults	SACN, 2015 ⁽⁵⁾
Nordic Nutrition Recommendations (NNR)	General Nordic population	2012	Limit the intake of added sugars (sucrose, fructose, and starch hydrolysates) to < 10 % of the total energy intake	Based on ensuring adequate intakes of dietary fibre and micronutrients	NNR, 2012 ⁽²⁵⁾
European Food Safety Authority (EFSA)	General population	2010	Insufficient data to set an upper limit for (added) sugars intake.		EFSA, 2010 ⁽³⁾
US Institute Of Medicine (IoM)	General population	2005	Maximum intake level of ≤ 25 % total energy	Based on prevention of displacement of foods that are a major source of micronutrients	IoM, 2005 ⁽²⁰⁾

Table 3. Energy and sugar intake of infants and young children (<4 years)

Country	Assessment year	Age range (years)	n	Method	Energy		Total sugars				Added sugars				Sucrose				Free sugars			
					MJ	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD
Australia ^(35,36)	2011–2012	2–3 (M)		24-h recall	6.0		94.3		24.9		32.5		8.4						42.2		11.0	
		2–3 (F)		24-h recall	5.9		88.7		24.1		32.0		8.1						39.1		10.0	
Belgium ⁽³⁷⁾	2014–2015	3–5 (M)	230	24-h recall	6.0		102.0		28.0													
		3–5 (F)	224	24-h recall	5.6		98.0		27.0													
Chile ⁽³⁸⁾	2010	2–5	257	24-h recall			74.4															
Denmark ^{*(39)}	2006–2007	6–7 months (M)	157	7-d record	3.5	0.8					4.4	5.0	2.1	2.2								
		8–9 months (M)	153	7-d record	3.8	0.9					5.9	6.3	2.6	2.4								
		10–11 months (M)	114	7-d record	4.2	0.9					10.0	11.0	4.1	4.0								
		12–23 months (M)	221	7-d record	4.7	1.0					18.0	12.0	6.5	4.0								
		24–36 months (M)	228	7-d record	5.3	1.0					28.0	15.0	9.0	4.2								
		6–7 months (F)	159	7-d record	2.9	0.8					2.7	3.4	1.6	1.9								
		8–9 months (F)	136	7-d record	3.5	0.9					5.8	5.4	2.7	2.2								
		10–11 months (F)	107	7-d record	3.9	0.9					7.6	7.0	3.2	2.7								
		12–23 months (F)	253	7-d record	4.5	1.0					16.0	11.0	5.9	3.4								
		24–36 months (F)	215	7-d record	5.0	1.0					26.0	11.0	8.8	3.7								
		2–5 (M)	277	24-h recall	6.0										48.0							
		2–5 (F)	302	24-h recall	5.6										46.0							
Finland ⁽⁴¹⁾	2003–2005	1 BF (M)	55	3-d record	2.8	0.8	43.0	16.0							10.5	5.9						
		1 FF (M)	257	3-d record	3.9	0.7	69.0	18.0							13.0	9.4						
		2 (M)	112	3-d record	4.6	1.0	78.0	23.0							34.3	16.9						
		3 (M)	236	3-d record	5.4	1.0	90.0	27.0							41.7	18.3						
		1 BF (F)	57	3-d record	2.6	0.8	38.0	15.0							9.0	6.0						
		1 FF (F)	198	3-d record	3.6	0.6	64.0	16.0							11.7	8.0						
		2 (F)	118	3-d record	4.5	0.9	77.0	21.0							31.5	14.7						
		3 (F)	235	3-d record	5.1	1.0	85.0	23.0							40.3	17.0						
France ⁽⁴²⁾	2014–2015	0–11 months (M)	80	24-h recall	3.1	1.3	65.4	25.6														
		0–11 months (F)		24-h recall	2.8	0.5	65.8	13.5														
		1–3 (M)	229	24-h recall	5.2	0.8	92.4	17.0														
		1–3 (F)		24-h recall	4.9	1.1	85.6	18.2														
Iceland ^(43,44)	2005–2007	9 months (M)	93	3-d record	3.2						3.4	4.2										
		12 months (M)	82	3-d record	3.6						7.6	5.4										
		9 months (F)	102	3-d record	3.0						3.6	5.2										
		12 months (F)	88	3-d record	3.3						9.7	7.2										
Ireland ⁽⁴⁵⁾	2007	3	225	3-d record	5.8	1.1					26.0	13.0										
		1	126	4-d record	4.2	0.9	70.7	5.3	26.2	1.3									25.1	3.8	9.0	1.2
		2	124	4-d record	4.7	1.2	74.3	5.7	24.8	1.4									33.4	4.9	11.0	1.5
		3	126	4-d record	4.8	0.9	75.9	5.9	24.9	1.4									41.4	5.7	13.4	1.7
Israel ⁽⁴⁶⁾	2015–2016	2–5 (M)		24-h recall	6.0	2.2	68.0	38.0	19.6	8.7												
		2–5 (F)		24-h recall	5.5	1.9	61.0	32.0	18.8	7.7												
Italy ⁽⁴⁷⁾	2005–2006	0–3	52	3-d record	4.7	1.8	71.4	36.0	24.9	8.8												
Mexico ⁽⁴⁸⁾	2012	1–4 (M)	2113	24-h recall	5.8									12.1								
		1–4 (F)		24-h recall	5.5									12.1								
Netherlands ⁽⁴⁹⁾	2012–2016	1–3 (M)	332	24-h recall	5.5		102.0		30.7		45.8		12.7						54.9		15.6	
		1–3 (F)	340	24-h recall	5.2		98.0		31.4		47.2		14.1						57.6		17.4	

Current perspectives on global sugars consumption

Table 3. (Continued)

Country	Assessment year	Age range (years)	n	Method	MJ	Energy		Total sugars			Added sugars			Sucrose			Free sugars		
						SD	g	SD	g	SD	g	SD	g	SD	g	SD	g		
Norway ^(50,51)	2006–2007	7 months to 1 year	881	24-h recall	5.1	1.6					12.2	14.0	4.0	4.1					
		14 months to 2 years	1674	24-h recall	5.7	1.5					22.5	16.0	6.7	4.2					
JK ⁽⁵²⁾	2014/15–2015/16	1.5–3	250	4-d record	4.5	1.0													
United States ^{†(53–55)}	2017–2018	2–5 (M)	260	24-h recall	6.9	0.2	106.0	5.8									32.6	17.8	
		2–5 (F)	280	24-h recall	6.0	0.2	95.0	4.8										11.3	
	2015–2016	6–11 months (M)									3.8		1.6					5.4	
		6–11 months (F)									3.8		1.5						
		12–23 months (M)									24.8		7.7						
		12–23 months (F)									24.4		7.4						

M: male; F: female.
BF: breastfed; FF: formula fed.
* Energy values per 10 MJ.
† United States reports SEM.

Children (<4 years). For infants and very young children (<4 years), sugar intake data are available for 16 countries: Australia, Belgium, Chile, Denmark, Estonia, Finland, France, Iceland, Ireland, Israel, Italy, Mexico, the Netherlands, Norway, the UK and the United States (Table 3). These include intakes of total sugars (nine countries), added sugars (seven countries), free sugars (four countries) and sucrose (two countries).

Total sugar intake expressed as a percentage of total energy (%E) ranged from 18.8 % for 2–5-year-old girls in Israel to 31.4 % for 1–3-year-old girls in the Netherlands. Intake of added sugars (%E) ranged from 1.5 % in 6–11-month-old girls in the United States to 14.1 % in 1–3-year-old girls in the Netherlands. Intake of free sugars (%E) ranged from 9 % in 1-year-olds in Ireland to 17.4 % in 1–3-year-old girls in the Netherlands.

Children (4–12 years). For children aged 4–12 years, sugar intake data are available for 25 countries: Australia, Austria, Belgium, Brazil, Canada, Chile, China, Denmark, Estonia, Finland, France, Germany, Iceland, India, Ireland, Israel, Italy, Mexico, the Netherlands, New Zealand, Norway, Portugal, Spain, the UK and the United States (Table 4). These include intakes of total sugars (eighteen countries), added sugars (eleven countries), free sugars (six countries), sucrose (four countries) and intrinsic sugars (Spain only).

Total sugar intake (%E) ranged from 8 % in 4–13-year-olds in China to 28.6 % in 4–8-year-old girls in the Netherlands. Intakes of added sugars (%E) ranged from 3 % in 4–13-year-old children in China to 17.0 % in 4–8-year-old girls in the Netherlands. Intakes of free sugars (%E) ranged from 9.1 % in 3-month-old-to-10-year-old boys in Portugal to 19.5 % for 4–8-year-old girls in the Netherlands.

Adolescents (13–18 years). For older children (13–18 years), sugar intake data were available for 24 countries: Andorra, Australia, Austria, Belgium, Brazil, Canada, Chile, Denmark, Estonia, France, Germany, Iceland, India, Ireland, Israel, Italy, Mexico, the Netherlands, New Zealand, Portugal, Slovenia, Spain, the UK and the United States (Table 5). These include intakes of total sugars (sixteen countries), added sugars (twelve countries), free sugars (eight countries), sucrose (four countries) and intrinsic sugars (Spain only).

Total sugar intake (%E) ranged from 15.4 % in 10–18-year-old boys in Italy to 30 % in 15–16-year-old girls in Slovenia. Intakes of added sugars (%E) ranged from 9.1 % in 10–17-year-old girls in Portugal to 16.2 % in 9–18-year-old children in the United States. Intakes of free sugars (%E) ranged from 9.1 % in 10–17-year-old boys in Portugal to 17.9 % for 9–13-year-old boys in the Netherlands.

Adults (19–64 years). For adults, sugar intake data are available for 29 countries: Andorra, Australia, Austria, Belgium, Brazil, Canada, Chile, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, India, Ireland, Israel, Italy, Lithuania, Mexico, the Netherlands, New Zealand, Norway, Portugal, Spain, Switzerland, the UK and the United States (Table 6). These include intakes of total sugars (eighteen countries), added sugars (fifteen countries), free sugars (eleven countries), sucrose (four countries) and intrinsic sugars (Spain only).

Table 4. Energy and sugar intake of children (4–12 years)

Country	Assessment year	Age range (years)	n	Method	Energy		Total sugars				Added sugars				Sucrose				Free sugars			
					MJ	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD
Australia ^(35,36)	2011–2012	4–8 (M)		24-h recall	7.6		110.7		23.3		52.2		10.8						60.4		12.5	
		9–13 (M)			9.2		125.4		21.7		66.9		11.4						75.3		12.9	
		4–8 (F)			6.4		94.8		23.8		43.5		10.5						50.7		12.4	
		9–13 (F)			8.0		115.3		23.5		61.0		12.3						68.2		13.8	
Austria ⁽⁵⁶⁾	2012	7–9 (M)	67	3-d record	8.0												12.0					
		10–12 (M)	83		8.1												10.0					
		7–9 (F)	57		8.0												11.0					
		10–12 (F)	81		7.2												11.0					
Belgium ⁽³⁷⁾	2014–2015	6–9 (M)	279	24-h recall	7.4		113.0		26.0													
		10–13 (M)	210		8.4		118.0		23.0													
		6–9 (F)	259		6.5		97.0		25.0													
		10–13 (F)	239		6.9		95.0		24.0													
Brazil ⁽⁵⁷⁾	2008–2009	10–13 (M)	1515	24-h recall	8.1	2.8					70.0	41.0	14.0	6.0								
		10–13 (F)	1566		7.8	2.5					73.0	41.0	15.0	7.0								
Canada ⁽⁵⁸⁾	2015	2–8		24-h recall			101.0		25.9													
Chile ^{*(38)}	2010	6–13	673	24-h recall			100.2															
		6–13 (M)			8.2																	
		6–13 (F)			6.8																	
China ^{†(59)}	2011	4–13	1460	24-h recall	5.8	0.1	26.0	0.6	8.0	0.1	9.0	0.3	3.0	0.1								
Denmark ⁽⁶⁰⁾	2011–2013	4–5 (M)	66	7-d record	7.9	1.6					40.0	19.2	9.0	3.7								
		6–9 (M)	150		8.7	1.8					55.0	26.8	11.0	4.6								
		10–13 (M)	134		9.6	2.2					66.0	29.7	12.0	4.6								
		4–5 (F)	64		7.0	1.3					39.0	19.1	9.0	4.0								
		6–9 (F)	141		8.0	1.7					50.0	25.7	10.0	4.4								
		10–13 (F)	135		8.1	1.8					55.0	29.8	11.0	4.8								
Estonia ⁽⁴⁰⁾	2015	6–9 (M)	168	24-h recall	7.7										61.0							
		10–13 (M)	93		8.3										59.0							
		6–9 (F)	179		6.8										56.0							
		10–13 (F)	89		6.7										55.0							
Finland ⁽⁴¹⁾	2003–2005	4 (M)	307	3-d record	5.8	1.1	97.0	26.0							46.9	18.0						
		6 (M)	364		6.7	1.2	108.0	28.0							51.5	19.9						
		4 (F)	247		5.5	1.0	92.0	22.0							43.6	16.0						
		6 (F)	349		6.0	1.1	99.0	27.0							48.7	19.3						
France ⁽⁴²⁾	2014–2015	4–6 (M)	454	24-h recall	6.6	1.7	104.1	31.2														
		4–6 (F)			6.1	1.4	98.0	27.5														
		7–10 (M)	643		8.1	1.8	115.2	32.8														
		7–10 (F)			7.2	1.9	104.5	34.0														
Germany ^{*(61)}	2006	6 (M)	106	3-d record	7.2	1.4	121.3	40.0														
		7–9 (M)	321		7.8	1.6	126.5	41.0														
		10–11 (M)	199		8.0	1.8	123.0	42.0														
		6 (F)	102		6.3	1.3	101.0	29.0														
		7–9 (F)	308		7.0	1.4	114.7	41.0														
		10–11 (F)	198		7.6	1.6	119.6	47.0														
Iceland ^(44,62,63)	2007	5	231	3-d record	6.3	1.2					34.0	16.0										
	2011–2012	6	162	3-d record	6.5	1.4							11.2									
	2003–2004	9	175	24-h recall	8.2	1.8	125.0	42.0			66.0	37.0										
India (rural) ⁽⁶⁴⁾	2011–2012	5–14 (M)	5265	24-h recall							10.5	16.3										
		5–14 (F)	4965								10.1	14.1										

Table 4. (Continued)

Country	Assessment year	Age range (years)	n	Method	Energy		Total sugars				Added sugars				Sucrose				Free sugars			
					MJ	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD
India (urban) ⁽⁶⁴⁾	2015–2016	5–14 (M)	4526	24-h recall							12.3	13.0										
		5–14 (F)	3961								11.5	13.0										
Ireland ^(45,65)	2010–2011	4	124	4-d record	5.3	1.0	83.4	6.2	24.7	1.4									49.0	6.4	14.4	1.8
	2017–2018	5–12	600	4-d record	6.3	1.2	73.7	19.4	18.7	4.0									38.4	14.9	9.5	3.4
Israel ⁽⁶⁶⁾	2015–2016	6–11 (M)		24-h recall	6.7	2.3	67.0	39.0	17.0	8.2												
		6–11 (F)			6.0	2.1	63.0	35.0	17.7	7.8												
Italy ⁽⁴⁷⁾	2005–2006	3–10	193	3-d record	8.0	2.0	86.0	30.0	17.0	5.0												
Mexico ^{†(48)}	2012	5–11 (M)	2753	24-h recall	7.8										11.7							
		5–11 (F)			7.3										12.3							
		4–13 [†]	3980	24-h recall	7.7	0.1	92.0	1.5	20.0	0.3	55.0	1.1	12.0	0.2								
Netherlands ⁽⁴⁹⁾	2012–2016	4–8 (M)	261	24-h recall	7.6		129.0		28.4		81.0		16.6						92.1		19.0	
		4–8 (F)	259		6.8		116.0		28.6		71.0		17.0						80.6		19.5	
New Zealand ⁽⁶⁷⁾	2002	5–6 (M)	383	24-h recall	7.6		110.0		24.2						58.5							
		7–10 (M)	738		9.0		130.0		24.2						70.8							
		5–6 (F)	309		6.8		105.0		25.8						55.6							
		7–10 (F)	687		7.8		115.0		24.5						63.6							
Norway ^(68,69)	2016	4 (M)	204	1-d record	6.1	1.2					30.0	18.0	8.6	4.6								
	2015	9–13 (M)	332		8.6	2.7	64.0	54.0	11.9	7.9												
	2016	4 (F)	195		5.5	1.4					28.0	16.0	8.6	4.3								
	2015	9–13 (F)	355		7.4	2.6	60.0	48.0	12.8	7.1												
Portugal ⁽⁷⁰⁾	2015–2016	3 months to 10 years (M)	1470	2-d record	6.9		98.6		25.3		41.6		8.8						46.2		9.1	
		3 months to 10 years (F)	1438		6.6		89.2		24.3		34.5		8.5						39.2		9.6	
Spain ^{‡(71,72)}	2013	9–12 (M)	126	3-d record	8.4	1.9	93.7	35.3	18.8	5.4									50.5	25.9	9.9	4.1
		9–12 (F)	87		7.9	1.6	88.4	30.1	18.8	5.0									45.9	20.6	9.6	3.8
UK ⁽⁷³⁾	2014/15–2015/16	4–10 (M)	276	4-d record	6.3	1.4													54.5	24.4	13.6	5.4
		4–10 (F)	238		5.8	1.3													49.9	23.2	13.4	4.6
United States ^{†(53,55)}	2017–2018	6–11 (M)	383	24-h recall	8.4	0.2	118.0	3.1														
		6–11 (F)	412		7.8	0.2	112.0	4.2														
	2009–2012	2–8	2871	24-h recall							61.0		14.3									
		4–13	3647	24-h recall	7.8	0.1	124.0	1.5	26.0	0.3	76.0	1.2	16.0	0.2							9.1	

M: male; F: female.

* Chile and Germany report median intakes.

† China, Mexico (4–13) and United States report SEM.

‡ Spain energy intake from ANIBES (2014). Sugar intake reported from Ruiz *et al.* (2017).

Table 5. Energy and sugar intake of adolescents (13–18 years)

Country	Assessment year	Age range (years)	n	Method	Energy		Total sugars				Added sugars				Sucrose				Free sugars			
					MJ	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD
Andorra ^{†(74)}	2004–2005	12–24 (M)	78	24-h recall	10.0	1.7	104.8	37.1	18.2	6.6												
		12–24 (F)	64		7.8	1.2	88.6	20.4	19.4	3.5												
Australia ^(35,36)	2011–2012	14–18 (M)		24-h recall	10.2		134.1		21.4		82.2		13.0						92.1		14.5	
		14–18 (F)			8.1		109.2		21.4		62.8		12.0						70.3		13.5	
Austria ⁽⁵⁶⁾	2012	13–14 (M)	19	3-d record	8.6												10.0					
		13–14 (F)	25		7.5												11.0					
Belgium ⁽³⁷⁾	2014–2015	14–17 (M)	240	24-h recall	9.1		119.0		21.0													
		14–17 (F)	239		7.0		93.0		23.0													
Brazil ⁽⁵⁷⁾	2008–2009	14–18 (M)	1905	24-h recall	9.1	3.2					77.0	47.0	14.0	6.0								
		14–18 (F)	1811		8.0	2.6					78.0	45.0	16.0	7.0								
Canada ^(58,75)	2015	9–18		24-h recall			115.0		22.5													
Chile ⁽³⁸⁾	2010	14–18	403	24-h recall			121.2															
Denmark ⁽⁶⁰⁾	2011–2013	14–17 (M)	117	7-d record	10.3	3.3					69.0	39.0	11.0	5.4								
		14–17 (F)	123		7.4	2.8					51.0	38.8	11.0	5.2								
Estonia ⁽⁴⁰⁾	2015	14–17 (M)	80	24-h recall	9.4														66.0			
		14–17 (F)	117		6.6														51.0			
France ⁽⁴²⁾	2014–2015	11–14 (M)	736	24-h recall	8.7	2.2	112.0	36.9														
		11–14 (F)			8.3	3.3	108.0	46.7														
		15–17 (M)	555		9.2	3.0	110.0	44.3														
		15–17 (F)			6.6	2.2	82.9	35.3														
Germany ^{†(61,76)}	2006	12 (M)	114	FFQ	10.6	3.2	167.0	71.8														
		13–14 (M)	214	FFQ	11.7	3.8	187.6	83.1														
		15–17 (M)	294	FFQ	14.3	5.4	221.3	113.4														
	2005–2006	14–80 (M)	7093	24-h recall	10.9 [†]		124 [†]		19.3													
		12 (F)	103	FFQ	9.3	3.2	154.5	86.0														
		13–14 (F)	230	FFQ	9.5	2.7	160.3	70.9														
		15–17 (F)	317	FFQ	9.9	3.8	175.1	96.2														
	2005–2006	15–19 (M)	506	24-h recall	9.2														10.8			
		15–19 (F)	536	24-h recall	6.5														12.8			
	2005–2006	14–80 (F)	8278	24-h recall	6.8 [†]		113 [†]		23.6													
	2006–2016	3–18 (M)	2163	3-d record	7.2				25.6					12.0							16.1	
		3–18 (F)	1953		6.4				25.5					11.8							16.0	
Iceland ⁽⁶³⁾	2003–2004	15	150	24-h recall	10.3	2.7	167.0	68.0			101.0	64.0										
India (rural) ⁽⁶⁴⁾	2011–2012	15–19 (M)	2344	24-h recall							12.0	16.6										
		15–19 (F)	2447								11.8	15.8										
India (urban) ⁽⁶⁴⁾	2015–2016	15–19 (M)	2328	24-h recall							13.8	14.0										
		15–19 (F)	2188								12.8	13.0										
Ireland ^(77,78)	2005–2006	13–17	441	7-d diary	8.3	2.4	108.5	43.0	20.4	5.0	65.7	32.0	12.4	4.9								
Israel ⁽⁷⁹⁾	2015–2016	12–15 (M)	1020	24-h recall	10.0	4.5	146.0	85.0														
		12–15 (F)	1165		9.6	4.6	149.0	89.0														
		16–18 (M)	767		10.5	4.6	144.0	89.0														
		16–18 (F)	954		9.4	4.4	142.0	85.0														
Italy ⁽⁴⁷⁾	2005–2006	10–18 (M)	108	3-d record	10.8	3.1	107.6	53.7	15.4	4.7												
		10–18 (F)	139		8.7	2.2	88.4	35.6	15.8	5.2												
Mexico ⁽⁴⁸⁾	2012	12–19 (M)	2056	24-h recall	9.8																	
		12–19 (F)			7.8									12.7								
														14.4								
Netherlands ⁽⁴⁹⁾	2012–2016	9–13 (M)	259	24-h recall	9.2		137.0		25.8		88.4		15.9						99.3		17.9	
		14–18 (M)	270		10.1		138.0		24.0		87.8		14.8						97.6		16.6	
		9–13 (F)	260		7.5		114.0		26.0		70.6		15.3						79.3		17.4	
		14–18 (F)	254		7.8		111.0		24.3		66.4		14.0						75.2		15.8	

Table 5. (Continued)

Country	Assessment year	Age range (years)	n	Method	Energy		Total sugars			Added sugars			Sucrose			Free sugars		
					MJ	SD	g	SD	%E	g	SD	%E	g	SD	%E	g	SD	%E
New Zealand [†] (67,80,81)	2008–2009	15–18 (M)	326	24-h recall	11.2		143.0		20.2	73.3		11.2	71.8		9.5	85.6		13.0
		15–18 (F)	373		7.9		118.0		23.9	57.4		12.4	62.7		11.6	68.8		15.2
	2002	11–14 (M)	576		10.5		144.0		22.9				76.6					
		11–14 (F)	582		8.4		124.0		24.7				65.2					
Portugal ⁽⁷⁰⁾	2015–2016	10–17 (M)	661	24-h recall	9.4		109.4		20.1	54.0		9.6				60.2		9.1
		10–17 (F)	670		7.6		88.2		20.2	41.9		9.1				44.9		10.1
Slovenia ⁽⁸²⁾	2012	15–16 (M)	1010	FFQ	12.7	6.9	170.0	58.0	26.0	9.0						103.0	56.0	16.0
		15–16 (F)	1214		9.7	5.1	195.0	65.0	30.0	10.0						110.0	60.0	17.0
Spain ^(71,72)	2013	13–17 (M)	137	3-d record	8.9	2.2	90.8	37.2	16.9	5.6						52.0	27.8	9.6
		13–17 (F)	74		7.6	1.8	86.6	31.0	19.2	5.4						48.7	21.0	10.8
UK ⁽⁷³⁾	2014/15–2015/16	11–18 (M)	270	4-d record	7.8	2.4										71.6	41.4	13.9
		11–18 (F)	272		6.5	1.9										62.4	35.3	14.4
United States [§] (63,55)	2017–2018	12–19 (M)	531	24-h recall	9.5	0.2	126.0	3.4										
		12–19 (F)	514		7.4	0.2	97.0	2.9										
	2009–2012	9–18	3238							78.7		16.2						

M: male; F: female.

[†] Total sugars for Andorra reported as digestible sugars.[‡] Germany and New Zealand report median values.[§] United States reports SEM.

Total sugars intake (%E) ranged from 12.7 % in 35–49-year-old men and women in Lithuania to 26 % in 19–30-year-old women in the Netherlands. Intakes of added sugars (%E) ranged from 6.3 % in 50–59-year-old men in Norway to 14 % in 19–59-year-old women in Brazil. Intakes of free sugars (%E) ranged from 6.9 % in 18–64-year-old men in Portugal to 18.1 % in 51–64-year-old women in Austria.

Adults (>65 years). For older adults, sugar intake data are available for 26 countries: Andorra, Australia, Austria, Brazil, Chile, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, India, Ireland, Israel, Italy, Lithuania, Mexico, the Netherlands, New Zealand, Norway, Portugal, Spain, the UK and the United States (Table 7). These include intakes of total sugars (fifteen countries), added sugars (eleven countries), free sugars (seven countries), sucrose (five countries) and intrinsic sugars (Spain only).

Total sugar intake (%E) ranged from 13.2 % in 65–75-year-old adults in Lithuania to 23.3 % in women ≥71 years old in New Zealand. Intakes of added sugars (%E) ranged from 4 % in men aged 65–84 years in Portugal to 12 % in women ≥60 years old in Brazil. Intakes of free sugars (%E) ranged from 4.4 % in 65–84-year-old men in Portugal to 12.5 % for men aged >75 years old in the UK.

Patterns of dietary sugars intake. Our analysis shows that intakes (expressed as %E) of both free and added sugars are highest in children and adolescents and are lowest for older adults.

Based on data as reported and one data point per age category per country, crude global estimates of mean intakes of free sugars (%E) were 12 % for 1–4-year-olds, 12 % for 4–12-year-olds, 14 % for 13–18-year-olds, 11 % for 19–64-year-olds and 8 % for >65-year-olds. For added sugars, crude global estimates of mean intakes were 8 % for 1–4-year-olds, 11 % for 4–12-year-olds, 13 % for 13–18-year-olds, 10 % for 19–64-year-olds and 8 % for >65-year-olds. These findings indicate that, where quantified, intakes of both added and free sugars are generally higher than recommended dietary guidelines at all life stages.

Such patterns of added sugars intake across the lifecycle are similar to previous findings, with the highest intakes in school-aged children and adolescents and the lower intakes observed in young children and adults (particularly older adults)^(15,97). Findings from individual countries such as Australia, Canada, the UK and Ireland have shown some reduction in intakes of added/free sugars in children and adolescents over the last ~15–20 years^(35,73,75,98). For example, free sugar intake in Ireland has decreased from 16%TE (2003–2004) to 9%TE (2017–2018)⁽⁹⁸⁾. These studies have suggested that this reduction is largely due to reformulation, particularly in the sweetened beverage categories which are commonly consumed by children and adolescents. It is difficult to draw stronger conclusions beyond this due to differing terminology and methodologies used between countries and over time.

In brief, two observations are apparent from this analysis of published dietary sugars intake data globally. The first is the high intake of added/free sugars compared with recommendations at all life stages but particularly in school-aged children and

Table 6. Energy and sugar intake of adults (19–64 years)

Country	Assessment year	Age range (years)	n	Method	Energy		Total sugars				Added sugars				Sucrose				Free sugars			
					MJ	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD
Andorra ^{†(74)}	2004–2005	25–44 (M)	222	24-h recall	8.8	1.2	88.4	15.8	17.5	2.4												
		45–64 (M)	85		8.0	1.2	83.7	11.6	18.2	2.2												
		25–44 (F)	215		6.9	1.4	75.1	18.9	18.4	3.5												
		45–64 (F)	143		6.8	1.3	76.8	18.8	19.1	4.1												
Australia ^(35,36)	2011–2012	19–30 (M)		24-h recall	11.0		131.0		19.1		75.5		10.8						85.4		12.3	
		31–50 (M)			10.2		119.4		18.9		61.9		9.4						69.9		10.7	
		51–70 (M)			9.3		101.0		17.3		46.8		7.8						54.1		9.0	
		19–30 (F)			7.9		99.0		20.5		53.1		10.8						60.4		12.3	
		31–50 (F)			7.5		91.1		19.7		42.5		8.9						48.5		10.2	
		51–70 (F)			7.3		87.0		19.2		36.8		7.7						42.0		8.8	
Austria ⁽⁸³⁾	2017	19–24 (M)	89	24-h recall	11.1	3.4															16.4	8.1
		25–50 (M)	478		10.2	2.8															16.4	7.4
		51–64 (M)	169		10.0	2.7															16.8	7.0
		19–24 (F)	181		7.6	2.3															16.8	7.6
		25–50 (F)	856		7.6	2.2															17.5	7.0
		51–64 (F)	245		7.5	2.2															18.1	6.8
Belgium ⁽³⁷⁾	2014–2015	18–39 (M)	205	24-h recall	9.8		113.0		19.0													
		40–64 (M)	284		9.3		99.0		18.0													
		18–39 (F)	315		7.1		87.0		20.0													
		40–64 (F)	322		6.7		77.0		19.0													
Brazil ⁽⁵⁷⁾	2008–2009	19–59 (M)	10 287	24-h recall	8.9	3.0					67.0	41.0	13.0	6.0								
		19–59 (F)	11 344		7.2	2.3					61.0	36.0	14.0	6.0								
Canada ^{†(58,75)}	2015	19+ (M)		24-h recall			94.0		17.7													
		19+ (F)					75.0		18.8													
Chile ^{§(38)}	2010	19+	11817	24-h recall			86.9	1.0	18.8	0.2	41.7	0.8	8.6	0.1					47.5	0.9	9.9	0.1
		19–29	660	24-h recall			89.1															
		30–49	1076				71.2															
		50–64	945				52.3															
Denmark ⁽⁶⁰⁾	2011–2013	14–64 (M)		7-d record	9.4																	
		14–64 (F)			6.7																	
		18–24 (M)	170		10.7	3.4					61.0	47.4	10.0	6.5								
		25–34 (M)	190		11.6	3.7					68.0	46.8	10.0	5.7								
		35–44 (M)	253		11.4	2.9					63.0	49.2	9.0	5.7								
		45–54 (M)	297		11.4	3.3					53.0	45.9	7.0	4.9								
		55–64 (M)	292		11.0	3.0					46.0	34.6	7.0	4.7								
		18–24 (F)	176		8.1	2.5					52.0	40.0	11.0	6.9								
		25–34 (F)	185		8.8	2.9					50.0	34.2	9.0	4.8								
		35–44 (F)	289		8.8	2.2					47.0	32.0	9.0	5.3								
		45–54 (F)	318		8.5	2.1					42.0	27.6	8.0	5.0								
		55–64 (F)	322		8.3	2.1					33.0	21.4	7.0	3.6								
Estonia ⁽⁴⁰⁾	2015	18–24 (M)	135	24-h recall	9.7										58.0							
		25–29 (M)	77		9.5										55.0							
		30–34 (M)	85		8.6										52.0							
		35–39 (M)	84		9.5										55.0							
		40–44 (M)	69		8.7										45.0							
		45–49 (M)	67		8.7										47.0							
		50–54 (M)	73		8.9										46.0							
		55–59 (M)	75		8.2										44.0

Table 6. (Continued)

Country	Assessment year	Age range (years)	n	Method	Energy		Total sugars				Added sugars				Sucrose				Free sugars			
					MJ	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD
Finland ⁽⁸⁴⁾	2017	18–24 (F)	192	24-h recall	6.8										48.0							
		25–29 (F)	137		7.6										54.0							
		30–34 (F)	158		7.4										49.0							
		35–39 (F)	160		7.2										48.0							
		40–44 (F)	167		6.4										41.0							
		45–49 (F)	168		6.2										38.0							
		50–54 (F)	136		6.3										42.0							
		55–59 (F)	160		6.4										38.0							
		18–24 (M)	47		10.9										37.0		6.0					
		25–44 (M)	221		10.1										47.0		7.8					
France ⁽⁴²⁾	2014–2015	45–64 (M)	308	24-h recall	9.1										43.0		8.1					
		18–24 (F)	52		7.0										38.0		9.3					
		25–44 (F)	259		7.7										39.0		8.5					
		45–64 (F)	317		7.5										41.0		9.3					
		18–44 (M)	1287		10.6	2.9	119.2	44.1														
Germany ^{§(85)}	2005–2006	18–44 (F)	1134	24-h recall	7.8	2.4	89.6	38.7														
		45–64 (M)	1134		10.4	3.2	99.8	44.6														
		45–64 (F)	1134		7.4	2.6	79.9	41.3														
		19–25 (M)	469		9.5												10.7					
		25–35 (M)	614		9.5												10.4					
		35–51 (M)	1946		9.9												9.9					
		51–65 (M)	1460		9.3												8.8					
		19–25 (F)	486		7.0												12.8					
		25–35 (F)	852		7.3												12.3					
		35–51 (F)	2648		7.3												11.5					
Greece ⁽⁸⁶⁾	2005–2007	51–65 (F)	1740	24-h recall	7.0												11.1					
		15–80 (M)					120.0		20.0										78.0		13.0	
		15–80 (F)					107.0		24.0										61.0		14.0	
		35–44 (M)	1311				87.3															
		45–54 (M)					85.4															
Hungary ⁽⁸⁷⁾	2014	55–64 (M)		3-d record			98.2															
		35–44 (F)	1373				85.1															
		45–54 (F)					87.4															
		55–64 (F)					88.5															
		18–34 (M)	103		12.4									9.3								
Iceland ⁽⁸⁸⁾	2010–2011	35–64 (M)	199	24-h recall	11.3									7.3								
		18–34 (F)	112		8.9									11.3								
		35–64 (F)	263		8.5									8.0								
		18–30 (M)	131		11.0	3.7	129.0	54.0			76.0	47.0										
		31–60 (M)	350		10.0	3.1	105.0	57.0			56.0	47.0										
India (rural) ⁽⁶⁴⁾	2011–2012	18–30 (F)	119	24-h recall	7.9	2.1	108.0	47.0			55.0	41.0										
		31–60 (F)	394		7.5	2.2	86.0	42.0			40.0	32.0										
		20–24 (M)	2031								12.9	18.2										
		25–29 (M)	2011								11.4	13.8										
		30–34 (M)	1875								12.5	21.4										
		35–39 (M)	1800								12.1	16.3										
		40–44 (M)	1539								11.9	15.7										
		45–49 (M)	1511								13.3	18.4

Table 6. (Continued)

					Energy		Total sugars				Added sugars				Sucrose				Free sugars																						
Country	Assessment year	Age range (years)	n	Method	MJ	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD																			
India (urban) ⁽⁶⁴⁾	2015–2016	50–59 (M)	2106	24-h recall							13.8	20.5																													
		60–69 (M)	1578								14.4	18.7																													
		20–24 (F)	2367								12.1	15.2																													
		25–29 (F)	2330								12.8	17.4																													
		30–34 (F)	1890								12.0	14.5																													
		35–39 (F)	2015								12.9	16.0																													
		40–44 (F)	1541								13.6	17.7																													
		45–49 (F)	1667								13.5	15.0																													
		50–59 (F)	2125								13.4	15.7																													
		60–69 (F)	1528								13.6	17.4																													
		20–24 (M)	2049								14.0	13.6																													
		25–29 (M)	1830								14.5	13.4																													
		30–34 (M)	1877								14.3	12.8																													
		35–39 (M)	1961								14.4	15.0																													
		40–44 (M)	1935								15.4	14.0																													
		45–49 (M)	1803								15.6	15.0																													
		50–59 (M)	2641								16.6	16.0																													
		60–69 (M)	1366								16.5	15.0																													
		20–24 (F)	2139								14.7	14.0																													
		25–29 (F)	2163								15.6	14.0																													
30–34 (F)	2177	16.8	15.0																																						
35–39 (F)	2163	18.2	18.0																																						
40–44 (F)	2139	17.6	15.0																																						
45–49 (F)	1810	18.3	16.0																																						
50–59 (F)	2284	17.8	16.0																																						
60–69 (F)	1339	17.1	15.0																																						
18–35 (M)	276	4-d record	10.7																			2.9	105.6	41.6	16.0	4.9	53.9	32.0	8.1	4.3	60.7	34.3	9.2	4.7							
36–50 (M)	205		9.7																			2.5	97.1	39.1	15.8	4.8	46.4	28.8	7.5	4.1	50.3	30.2	8.2	4.3							
51–64 (M)	153		9.3																			2.4	96.5	39.1	16.2	4.9	43.1	27.5	7.1	4.0	47.6	29.1	7.9	4.2							
18–35 (F)	255		7.5																			2.3	83.8	35.5	17.8	5.1	45.6	28.2	9.2	4.7	50.3	29.9	10.2	4.9							
36–50 (F)	232		7.1																			1.9	78.8	34.4	17.1	5.0	36.4	24.4	7.5	4.1	40.2	26.0	8.3	4.3							
51–64 (F)	153		7.0																			1.7	84.4	35.8	18.7	5.2	45.4	24.0	7.4	4.1	38.8	25.4	8.1	4.3							
Israel ⁽⁹¹⁾	2014–2016	18–34 (M)																				24-h recall	9.0	4.4	87.0	58.0	16.5	9.0													
		35–44 (M)																				7.9	3.3	79.0	49.0	16.9	8.5														
		45–64 (M)																				7.1	3.8	69.0	56.0	16.6	9.6														
		18–34 (F)																				6.2	3.0	69.0	45.0	19.1	9.7														
		35–44 (F)																				6.2	2.7	62.0	39.0	17.3	8.7														
		45–64 (F)																				5.3	2.5	64.0	45.0	21.0	11.3														
		18–65 (M)	1068																			3-d record	10.0	2.7	86.0	38.0	13.5	4.7													
Italy ⁽⁴⁷⁾	2005–2006	18–65 (F)	1244																				8.1	2.2	79.5	33.0	15.4	5.1													
		19–34	772																			24-h recall	8.1		58.1		12.8														
Lithuania ⁽⁹²⁾	2013–2014	35–49	692																				7.8		55.5		12.7														
		50–64	749																				7.4		54.8		13.2														
		20+ (M)	3174																			24-h recall	9.4					12.8													
Mexico ⁽⁴⁸⁾	2012	20+ (F)																					7.2								12.2										
		19–30 (M)	260	24-h recall	11.1		135.0		25.8	81.4	13.0	90.7	14.6																												
Netherlands ⁽⁴⁹⁾	2012–2016	31–50 (M)	259		11.4		123.0		17.7		67.8		10.6						75.5		12.0																				
		51–70 (M)	264		10.3		109.0		17.4		55.5		8.9					62.8		10.1																					

Current perspectives on global sugars consumption

Table 6. (Continued)

Country	Assessment year	Age range (years)	n	Method	Energy		Total sugars				Added sugars				Sucrose				Free sugars			
					MJ	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD
New Zealand ^{§(80,81)}	2008–2009	19–30 (F)	256	24-h recall	8.0		106.0		26.0		59.8		12.3						67.6		13.9	
		31–50 (F)	264		7.9		98.0		20.1		50.5		10.4						57.0		11.8	
		51–70 (F)	258		7.6		92.0		19.8		42.7		9.1						48.7		10.4	
		19–30 (M)	284		11.9		147.0		20.1		69.7		10.8		73.2		9.4		78.1		12.1	
		31–50 (M)	598		11.5		133.0		19.5		9.5		64.5		8.9		70.3		10.9			
		51–70 (M)	378		9.4		108.0		19.5		8.4		49.0		8.4		54.4		10.1			
		19–30 (F)	434		8.4		123.0		23.8		12.2		65.8		11.1		62.4		14.1			
Norway ⁽⁹³⁾	2010	31–50 (F)	746	24-h recall	7.9		98.0		40.1		9.1		45.3		9.6		47.9		10.8			
		51–70 (F)	517		7.2		95.0		21.7		35.7		8.6		40.4		9.3		43.2		9.7	
		18–29 (M)	138		12.8	4.0					69.0	55.0	9.2	6.9								
		30–39 (M)	136		11.5	3.5					49.0	41.0	7.2	6.0								
		40–49 (M)	179		10.6	3.1					51.0	45.0	7.7	6.0								
		50–59 (M)	192		10.4	3.1					41.0	38.0	6.3	5.1								
		18–29 (F)	143		8.1	2.5					46.0	41.0	9.3	6.5								
		30–39 (F)	169		8.4	2.4					42.0	34.0	8.2	5.6								
		40–49 (F)	256		8.1	2.4					32.0	24.0	6.6	4.4								
		50–59 (F)	193		7.9	2.3					33.0	27.0	6.8	4.9								
Portugal ⁽⁷⁰⁾	2015–2016	18–64 (M)	3056	24-h recall	9.8		92.3		16.2		36.6		6.5						40.0		6.9	
		18–64 (F)	3556		6.9		78.0		19.4		30.0		7.1						32.6		7.9	
Spain ^(71,72)	2013	18–64 (M)	798	3-d record	8.2	2.3	78.4	36.7	16.0	5.8									35.2	27.1	7.0	4.8
		18–64 (F)	857		7.0	1.8	71.7	30.5	17.3	5.7					31.6	22.3			7.3	4.5		
Switzerland ⁽⁹⁴⁾	2014–2015	18–75 (M)	933	24-h recall			114.0	49.0	18.1		61.0	41.0	9.4						73.0	44.0	11.4	
		18–75 (F)	1124				99.0	37.0	20.5		46.0	28.0	9.3						57.0	32.0	11.5	
		18–29	400				113.0	47.0	19.6		63.0	41.0	10.5						75.0	45.0	12.7	
		30–64	1319				107.0	44.0	19.2		53.0	36.0	9.3						65.0	39.0	11.4	
		65–75	338				99.0	38.0	19.5		42.0	25.0	8.1						53.0	28.0	10.2	
UK ⁽⁷³⁾	2014/15–2015/16	19–64 (M)	450	4-d record	8.3	2.4													64.3	46.6	11.1	6.4
		19–64 (F)	632		6.6	2.0													50.0	34.4	11.2	6.1
United States ^{** (53,55)}	2017–2018	20–29 (M)	335	24-h recall	10.1	0.3	105.0	3.6														
		30–39 (M)	329		11.1	0.3	124.0	5.8														
		40–49 (M)	340		10.9	0.5	128.0	9.5														
		50–59 (M)	375		10.7	0.2	128.0	7.1														
		20–29 (F)	359		8.3	0.2	99.0	4.8														
		30–39 (F)	407		7.9	0.2	96.0	4.0														
		40–49 (F)	367		8.0	0.3	106.0	5.8														
		50–59 (F)	419		7.6	0.4	97.0	8.2														
		2009–2012																				
		19 +	10 697		24-h recall						62.1		13.1									

M: male; F: female.

† Total sugars for Andorra reported as digestible sugars.

‡ Canada variance reported as standard error (SE), not SD. Intakes for adults not split by sex, based on the first day recall, not usual intake.

§ Chile, Germany and New Zealand report median values.

|| Greece *n*-values include older adults, but data are for age groups as reported.

¶ Hungary mono- and disaccharides added to foods, foodstuffs during production or preparation, and natural sugars in honey.

** United States reports SEM.

Table 7. Energy and sugar intake of older adults (65+ years)

Country	Assessment year	Age range (years)	n	Method	Energy		Total sugars				Added sugars				Sucrose				Free sugars			
					MJ	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD
Andorra [†] (74)	2004–2005	65–75 (M)	45	24-h recall	7.0	0.9	79.6	14.6	19.1	2.8												
		65–75 (F)	48		6.4	1.0	82.5	19.1	22.2	3.9												
Australia ^(35,36)	2011–2012	71+ (M)		24-h recall	8.2		101.8		19.8		45.3		8.5						52.9		10.0	
		71+ (F)			6.6		84.4		20.3		34.2		8.0						40.7		9.5	
Austria ⁽⁸³⁾	2012	65–80 (M)	76	3-d record	8.0	0.0											8.0					
		65–80 (F)	100		7.0												9.0					
Brazil ⁽⁵⁷⁾	2008–2009	60+ (M)	1993	24-h recall	7.7	2.5					52.0	34.0	11.0	6.0								
		60+ (F)	2328		6.5	2.1					46.0	30.0	12.0	6.0								
Chile ⁺ (38)	2010	65+	906	24-h recall			42.9															
		65+ (M)			7.1																	
		65+ (F)			5.5																	
Denmark ⁽⁶⁰⁾	2011–2013	65–75 (M)	262	7-d record	10.8	3.1					53.0	41.3	8.0	4.9								
		65–75 (F)	262		8.3	2.2					38.0	25.9	8.0	4.1								
Estonia ⁽⁴⁰⁾	2015	60–64 (M)	85	24-h recall	8.1										40.0							
		65–69 (M)	83		7.8										34.0							
		70–74 (M)	74		7.6										36.0							
		60–64 (F)	187		6.2										39.0							
		65–69 (F)	194		6.3										36.0							
		70–74 (F)	147		5.6										32.0							
Finland ⁽⁸⁴⁾	2017	65–74 (M)	204	24-h recall	8.0										38.0		8.2					
		65–74 (F)	247		6.6										34.0		8.8					
France ⁽⁴²⁾	2014–2015	65–79 (M)	724	24-h recall	9.3	3.4	89.4	47.7														
		65–79 (F)			6.9	2.7	77.6	38.5														
Germany ⁽⁹⁵⁾	2005–2006	65–80 (M)	1165	24-h recall	8.9												9.1					
		65–80 (F)	1331		6.8												11.3					
Greece [§] (86)	1995–2000	65–74 (M)		24-h recall			92.1															
		65–74 (F)					86.7															
Hungary ⁽⁸⁷⁾	2014	≥65 (M)	372	3-d record	8.1								6.0									
		≥65 (F)	485		10.0								7.4									
Iceland ⁽⁸⁸⁾	2010–2011	61–80 (M)	151	24-h recall	8.7	2.8	80.0	44.0			35.0	31.0										
		61–80 (F)	167		6.7	2.0	74.0	32.0			51.0	24.0										
India (rural) ⁽⁶⁴⁾	2011–2012	70–79 (M)	560	24-h recall							14.0	14.0										
		80+ (M)	180								12.4	11.0										
		70–79 (F)	571								12.5	16.0										
		80+ (F)	192								11.9	10.0										
India (urban) ⁽⁶⁴⁾	2015–2016	70–79 (M)	486	24-h recall							17.2	16.0										
		80+ (M)	143								15.1	13.0										
		70–79 (F)	561								16.9	16.0										
		80+ (F)	207								16.5	12.0										
Ireland ^(89,90)	2008–2010	65+ (M)	106	4-d record	8.3	2.6	89.5	37.0	16.8	4.9	39.5	25.7	7.2	4.0					43.1	27.1	7.8	4.2
		65+ (F)	120		6.5	1.6	83.4	35.8	19.6	5.4	35.3	24.1	7.7	4.3					39.5	25.8	8.8	4.5
Israel ⁽⁹¹⁾	2014–2015	65–74 (M)		24-h recall	6.7	2.9	67.0	48.0	16.1	8.9												
		75–84 (M)			5.9	2.5	56.0	37.0	15.7	8.8												
		85+ (M)			5.2	2.1	59.0	40.0	18.5	10.6												
		65–74 (F)			5.5	2.4	62.0	38.0	17.8	7.8												
		75–84 (F)			4.8	1.9	50.0	30.0	16.9	7.8												
		85+ (F)			4.6	1.7	55.0	37.0	18.6	9.8												

Current perspectives on global sugars consumption

Table 7. (Continued)

Country	Assessment year	Age range (years)	n	Method	Energy		Total sugars				Added sugars				Sucrose				Free sugars			
					MJ	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD	g	SD	%E	SD
Italy ⁽⁴⁷⁾	2005–2006	65+ (M)	202	3-d record	9.6	2.3	81.6	35.0	13.3	5.0												
		65+ (F)	316		7.7	2.0	78.6	32.0	16.2	5.4												
Lithuania ⁽⁹²⁾	2013–2014	65–75	300	24-h recall	6.7		51.1		13.2													
Mexico ^{†(96)}	2012	65+ (M)	237	SFFQ	7.1		87.8															
		65+ (F)	289		5.5 [‡]		71.5 [‡]															
Netherlands ⁽⁴⁹⁾	2012–2016	71–79 (M)	260	24-h recall	9.1		99.0		19.0		49.1		8.0						55.4		9.0	
		71–79 (F)	257		7.4		88.0		20.6		38.9		8.5						44.2		9.8	
New Zealand ^{‡(81)}	2008–2009	71+ (M)	480	24-h recall	8.1		105.0		21.1		45.9		8.8		45.9		9.1		51.9		10.1	
		71+ (F)	585		6.1		84.0		23.3		32.2		8.4		33.5		9.7		36.8		9.9	
Norway ⁽⁹³⁾	2010	60–70 (M)	217	24-h recall	9.9	2.9					39.0	32.0	6.3	4.4								
		60–70 (F)	164		7.4	2.2					30.0	24.0	6.5	4.2								
Portugal ⁽⁷⁰⁾	2015–2016	65–84 (M)	857	24-h recall	8.3		72.3		15.6		20.9		4.0						22.6		4.4	
		65–84 (F)	790		6.3		71.9		19.9		19.1		5.2						23.3		6.2	
Spain ^(71,72)	2013	65–75 (M)	99	3-d record	7.4	2.0	74.2	37.4	16.7	6.1									21.3	15.2	4.8	3.1
		65–75 (F)	107		6.2	1.5	71.8	30.6	19.8	6.9									20.1	14.7	5.4	3.7
UK ⁽⁷³⁾	2014/15–2015/16	65–74 (M)	71	4-d record	7.7	1.7													62.0	34.3	11.8	5.8
		75+ (M)	70		7.4	2.0													60.5	35.7	12.5	6.4
		65–74 (F)	110		6.1	1.8													42.1	30.0	10.4	7.1
		75+ (F)	84		5.6	1.5													37.7	20.1	10.4	4.9
United States ⁽⁵⁵⁾	2017–2018	60–69 (M)	500	24-h recall	10.3	0.3	116.0	4.3														
		70+ (M)	428		9.0	0.2	111.0	4.0														
		60–69 (F)	479		7.5	0.3	90.0	7.0														
		70+ (F)	404		7.0	0.2	88.0	2.4														

* M: male; F: female.

† Total sugars for Andorra reported as digestible sugars.

‡ Chile, Mexico and New Zealand report median.

§ n-Values combined for 19–59 and 65–74 years for Greece.

|| United States reports SEM.

adolescents. The second outcome is that more data (both in terms of quantity and harmonisation) are required to give a complete picture of sugars intake globally. Caution is needed when interpreting these dietary intake data across sugar types and making inter-country comparisons, due to differences in the dietary assessment methodologies, differences in the time periods in which dietary intake was assessed and differences in national clustering of age groups. Future studies could examine the feasibility of identifying and compensating for selected under-reporting (which may be worse for sugar-sweetened foods and beverages) by using biomarkers in a subset of the population.

Interpretation and application of dietary sugar data

Challenges of data availability and quality

Whilst this review serves to update data on sugar intake from two previous reviews, for some countries there have been no updates since the time of the previous reviews, and hence these earlier data are included here also. New data are, however, available and included for countries such as Andorra, Australia, Chile, China, Estonia, Germany, Greece, India, Lithuania, Mexico, New Zealand, Portugal, Slovenia and Switzerland, and updated data are available for some age groups, including for Belgium, Canada, Denmark, Finland, France, Germany, Ireland, Israel, Netherlands, Norway, the UK and the United States.

Notwithstanding the addition of new data, the findings suggest that relatively few countries globally provide any kind of dietary data on sugar intake. The greatest amount of intake data was available for adults, but even then, data were only available for 29 countries; mostly in the European Union (EU) and greater European region, Australia and New Zealand, the United States and some of the Americas. Of note, there are 194 countries in the world today⁽⁹⁹⁾ and 100 countries which have food-based dietary guidelines⁽²⁶⁾. Significant gaps remain in examining sugars intakes globally, mirroring gaps in dietary surveillance in general⁽¹⁰⁰⁾.

A notable difference between this 2020 review and previous reviews^(15,16) is the inclusion of available data for dietary free sugar intake. This is in part due to newly available methods and definitions to estimate free sugars in databases of food consumption^(101,102). These new data allow comparisons against nutrient goals for dietary sugars, which are typically based on free sugars intake. In particular, the inclusion of these data highlights the challenges of meeting dietary recommendations for free sugars, particularly at 5%E. For example, of the twenty-seven surveys that provided an estimate for free sugars, only one survey reported an estimate of mean intake below that value (Spain, where older male adults reported sugars intakes of 4.8%E). Understanding which food groups are the key contributors to free sugars at each lifecycle stage will help to inform strategies to reduce these intakes⁽¹⁰³⁾.

Nevertheless, there remain considerable challenges in estimating dietary sugar intake data. Standard challenges exist relating to population sampling, selection bias and the assumptions and uncertainties inherent to the dietary collection methods used (24 h recall, food records and food frequency

questionnaires)^(15,100). Further, for sugars, obtaining reliable estimates of added and free sugar consumption continues to be a challenge for researchers. Despite the development of protocols to estimate added/free sugar composition^(101,104), detailed food and composition data at brand level are necessary for these purposes^(15,102). As part of EFSA's ongoing assessment to provide scientific advice on an tolerable upper intake level for sugars, harmonised data on dietary sugars intake will be compared for countries in the European region; however, challenges remain for assuring valid comparisons in other global regions⁽¹⁴⁾.

Application of dietary sugar data

For dietary intake researchers, data managers and risk assessors, the need to survey and accurately quantify dietary sugars intake remains a priority. However, it also remains problematic, given the lack of global harmonisation of definitions, classifications and recommendations and the lack of biomarkers to differentiate between the dietary intake of different sugars types. Future situations that could overcome this lack of harmony, but that could also offer objective biomarkers of intake and quantitative measures of the amounts of the various sugar forms present within foods, would be welcome. However, where available, these are currently only in their infancy and, indeed for biomarkers at least, may be a futile effort^(105–108).

For policy makers, current dietary sugar intake patterns are needed for consistent longitudinal monitoring and may enforce concerns and underline the importance of policy initiatives to reduce population sugar and energy (calorie) intakes. To date, there have been a variety of approaches undertaken by different countries, ranging from education strategies, taxation policies on single food types (e.g. taxes on sugar-sweetened beverages) to more comprehensive initiatives affecting multiple food types (e.g. UK initiatives targeting reformulation, portion size and altering consumer purchasing behaviour⁽¹⁰⁹⁾, or indeed a mix of all three^(110,111)). Although there is some information to suggest that such efforts may reduce purchase and, in some cases, intake, any impact on outcomes other than absolute dietary intakes (e.g. health outcomes such as obesity risk) remain unknown, despite numerous modelling exercises^(112–117). A challenge remains in balancing the need to have regular robust population dietary assessments to underpin and evaluate policy with the costs and logistics that accompany such surveillance programmes⁽¹⁰⁰⁾.

Food compositional data for sugars are also used on food labels as applied by the food industry. The values on such labels can be used to underpin estimates of dietary sugars intake by researchers. For most countries, there is no mandatory labelling of added or free sugars, and only total sugars is required to be declared on the label, mainly due the current lack of an accepted analytical method for manufacturers to assess added/free sugar content in foods/drinks. Exceptions include the United States, where added sugars (g, % daily value) will be included on the food label from 2021⁽¹¹⁸⁾, and Mexico, who have recently (2020) also updated their labelling guidelines to include added sugars⁽¹¹⁹⁾. Notwithstanding this, challenges remain: for example, in the United States, single-ingredient sugars such as syrups are treated differently with respect to labelling requirements and are not labelled as added sugars unless added

as an ingredient in other foods. Given such disparities, continued challenges will remain when sourcing and applying food composition data to underpin estimates or modelling of dietary sugars intake, particularly for foods which are either new to market or have been recently reformulated for sugars. It is promising to see a recently proposed method to quantify free sugars in foods and beverages which, if workable, will be a valuable addition for assessment of free sugars and will be useful in product development and for public health bodies, industry and regulatory agencies⁽¹⁰⁷⁾.

A further challenge relates to consumer awareness and understanding of the meaning of the terms 'added' or 'free' sugars in the context of dietary guidance. As mentioned, globally, food labels mostly only provide the interested consumer with information on quantitative amounts of total sugars present. Recent studies have suggested confusion in recognising dietary sources of added or free sugars, with results reiterating the need to have a readily understood term for sugars relating to label information and to educate and inform consumers of such^(120,121). With disconnection between dietary guidance and food labelling regulations with respect to total/free/added sugars, it will be a challenge for consumers to understand the relevant sugar content of foods from the nutrition label.

Future directions and conclusion

With the role of dietary sugars under scrutiny, this review aimed to provide a current picture of dietary guidance and actual sugar intake worldwide. It shows that, globally, a minority of countries provide any estimate of sugar intake. Further, intakes of the sugar forms upon which the greatest emphasis is placed (free and added sugars) generally exceed existing dietary guidelines across all population groups and are highest in children and adolescents and lowest for older adults. Whilst individual studies suggest some reductions in intakes of added sugars (%E) in children and adolescents, this review suggests that there is a need for more data to understand the complete picture. Furthermore, this review suggests a greater need for harmonisation of terms and for collaborative efforts to ensure that the most up-to-date food composition data are used to underpin any estimates of intake or modelling scenarios. Development and consensus on robust, valid biomarkers which provide accurate measures of dietary sugars intake, and preferably in a manner which is cost-effective, non-invasive and quick, would be welcome. In the interim, continued focus on monitoring of both dietary intakes and the efficacy of any public health initiatives remains paramount.

Acknowledgements

We wish to thank the authors of the dietary survey reports in different countries who assisted us with data related queries.

Financial support

This work was supported by the World Sugar Research Organisation (WSRO). WSRO had no role in the analysis or writing of this review.

J.W. received financial support from the WSRO for the initial draft of the manuscript. No funding was received for later drafts or for final manuscript. R.R. and H.B. are previous employees of WSRO and were employees at the time that the data were collected.

R.R. and J.W. conceived the idea of this review paper. H.B. collated the data. J.W. and A.N. interpreted the data and drafted the manuscript. All authors read and approved the final version of the manuscript.

References

1. Fidler Mis N, Braegger C, Bronsky J, *et al.* (2017) Sugar in infants, children and adolescents: a position paper of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. *J Pediatr Gastroenterol Nutr* **65**, 681–696.
2. Clemens RA, Jones JM, Kern M, *et al.* (2016) Functionality of sugars in foods and health. *Comprehensive Reviews in Food Science and Food Safety*. Blackwell Publishing Inc.; **15**, 433–470.
3. European Food Safety Authority (2010) Scientific opinion on dietary reference values for carbohydrates and dietary fibre. *EFSA J [Internet]* **8**, 1462. Available from: http://www.efsa.europa.eu/sites/default/files/scientific_output/files/main_documents/1462.pdf.
4. World Health Organization (2015) *WHO Guideline: sugars intake for adults and children*. Geneva: WHO.
5. Scientific Advisory Committee on Nutrition (2015) Carbohydrates and Health. http://www.gov.uk/government/uploads/system/uploads/attachment_data/file/445503/SACN_Carbohydrates_and_Health.pdf (accessed January 2020).
6. Prinz P (2019) The role of dietary sugars in health: molecular composition or just calories? *Eur J Clin Nutr* **73**, 1216–1223.
7. Johnson RK, Appel LJ, Brands M, *et al.* (2009) Dietary sugars intake and cardiovascular health: a scientific statement from the American Heart Association. *Circulation* 1011–1120.
8. Tappy L, Morio B, Azzout-Marniche D, *et al.* (2018) French recommendations for sugar intake in adults: a novel approach chosen by ANSES. *Nutrients* **10**, 989.
9. Institute of Medicine (2005) *Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein, and amino acids (macronutrients)*. Washington, DC: The National Academies Press.
10. Jamnik J, Rehman S, Blanco Mejia S, *et al.* (2016) Fructose intake and risk of gout and hyperuricemia: a systematic review and meta-analysis of prospective cohort studies. *BMJ Open* **6**, e013191.
11. Dornas WC, de Lima WG, Pedrosa ML, *et al.* (2015) Health implications of high-fructose intake and current research. *Adv Nutr* **6**, 729–737.
12. Lobstein T (2014) WHO | Reducing consumption of sugar-sweetened beverages to reduce the risk of childhood overweight and obesity [Internet]. World Health Organization [cited 2020 Jun 14]. Available from: https://www.who.int/elena/titles/ssbs_childhood_obesity/en/
13. Te Morenga L, Mallard S & Mann J (2013) Dietary sugars and body weight: systematic review and meta-analyses of randomised controlled trials and cohort studies. *BMJ* **346**, e7492.
14. European Food Safety Authority (2018) Protocol for the scientific opinion on the Tolerable Upper Intake Level of dietary sugars. *EFSA J* **16**, 5393–5440.

15. Newens KJ & Walton J (2016) A review of sugar consumption from nationally representative dietary surveys across the world. *J Hum Nutr Diet* **29**, 225–40.
16. Azaïs-Braesco V, Sluik D, Mailliot M, *et al.* (2017) A review of total & added sugar intakes and dietary sources in Europe. *Nutr J* **16**, 6.
17. Fisberg M, Kovalskys I, Gómez G, *et al.* (2018) Total and added sugar intake: assessment in eight Latin American countries. *Nutrients* **10**, 389.
18. Mela DJ & Elizabeth MW (2018) Perspective: total, added, or free? What kind of sugars should we be talking about? *Adv Nutr* **9**, 63–69.
19. Buyken AE, Mela DJ, Dussort P, *et al.* (2018) Dietary carbohydrates: a review of international recommendations and the methods used to derive them. *Eur J Clin Nutr* **72**, 1625–1643.
20. DRI (2005) *Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein, and amino acids (macronutrients)*. Washington DC: The National Academies Press.
21. European Food Safety Authority (2018) Protocol for the scientific opinion on the Tolerable Upper Intake Level of dietary sugars. *EFSA J* **16**, 5393.
22. WHO/FAO (1998) Carbohydrates in human nutrition. Report of a Joint FAO/WHO Expert Consultation. *FAO Food Nutr. Pap* **66**, 1–140.
23. U.S. Department of Agriculture and U.S. Department of Health and Human Services (2020) *Dietary Guidelines for Americans, 2020–2025*, 9th Edn. [Internet]. Available from: DietaryGuidelines.gov.
24. Institute of Medicine (2005) *Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein, and amino acids*. Washington, DC: National Academies Press, 1–1331.
25. Nordic Council of Ministers (2012) *Nordic nutrition recommendations 2012: integrating nutrition and physical activity*. Copenhagen: Nordic Council of Ministers, Narayna Press.
26. FAO (2003) Food based dietary guidelines in the WHO European Region Nutrition and Food Security Programme. WHO Regional Office for Europe [Internet]. Available from: <http://www.euro.who.int>.
27. Committee on Medical Aspects of Food Policy (COMA) (1991) *Dietary reference values for food energy and nutrients for the United Kingdom*. London: HMSO.
28. Ernst JB, Arens-Azevêdo U, Bitzer B, *et al.* (2019) Quantitative Empfehlung zur Zuckerzufuhr in Deutschland Kurzfassung des Konsensuspapiers der Deutschen Adipositas-Gesellschaft e. V. (DAG), der Deutschen Diabetes Gesellschaft e. V. (DDG) und der Deutschen Gesellschaft für Ernährung e. V. (DGE). Ernährungs Umschau [Internet], [cited 2020 Jun 16] 66, 26–34. Available from: www.dge.de/fileadmin/public/
29. Vos MB, Kaar JL, Welsh JA, *et al.* (2017) Added sugars and cardiovascular disease risk in children: a scientific statement from the American Heart Association. *Circulation* **135**, e1017–e1034.
30. Erickson J, Sadeghirad B, Lytvyn L, *et al.* (2017) The scientific basis of guideline recommendations on sugar intake: a systematic review. *Ann Intern Med* **166**, 257–267.
31. Rippin HL, Hutchinson J, Evans CEL, *et al.* (2018) National nutrition surveys in Europe: a review on the current status in the 53 countries of the WHO European region. *Food Nutr Res* **62**, 10.2919.
32. Newens KJ & Walton J (2016) A review of sugar consumption from nationally representative dietary surveys across the world. *J Hum Nutr Diet* **29**, 225–240.
33. Huybrechts I, Aglago EK, Mullee A, *et al.* (2017) Global comparison of national individual food consumption surveys as a basis for health research and integration in national health surveillance programmes. *Proc Nutr Soc* **76**, 549–567.
34. Song SJ & Song WO (2014) National nutrition surveys in Asian countries: surveillance and monitoring efforts to improve global health. *Asia Pac J Clin Nutr* **23**, 514–523.
35. Australian Bureau of Statistics (2016) Australian Health Survey: Consumption of Added Sugars Australia. Australian Bureau of Statistics [Internet]. Available from: [http://www.ausstats.abs.gov.au/ausstats/subscriber.nsf/0/58042B7FE6A65821CA2581F4000D1020/\\$File/4364.0.55.011 - australian health survey - consumption of added sugars.pdf](http://www.ausstats.abs.gov.au/ausstats/subscriber.nsf/0/58042B7FE6A65821CA2581F4000D1020/$File/4364.0.55.011 - australian health survey - consumption of added sugars.pdf).
36. Australian Bureau of Statistics (2014) Australian health survey: nutrition first results – food and nutrients, 2011–12 [Internet]. 4364.0.55.007, 1–78. Available from: <http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/4364.0.55.0072011-12?OpenDocument>.
37. Ridder K, Bel S, Brocatus L, Cuypers K, Lebacqz T, Moyersoen I, Ost C, & Teppers E Enquête de Consommation Alimentaire 2014–2015 Rapport 4: La Consommation Alimentaire. [Internet]. 2016. Available from: https://fcs.wiv-isp.be/nl/Gedeelde-documenten/FRANS/Rapport4/Resume_rapport_4_finaal_finaal.pdf.
38. Ministry of Health Chile (2021) *Chile National Food Consumption Survey 2010–2011*. Santiago, Chile: Ministry of Health Chile.
39. Trolle E, Gondolf UH, Ege M, Kørup K, Ygil KH & Christensen T (2013) Danskernes kostvaner Spæd- og småbørn 2006–2007 [Internet]. Available from: http://orbit.dtu.dk/files/84054319/Rapport_Danskernes_kostvaner_sp_d_og_sm_b_rn_rev_12_12_13_1_.pdf.
40. Pitsi T, Zilmer M, Vaask S, *et al.* (2015) Eesti toitumis- ja liikumissoovitused [Internet]. Available from: https://intra.tai.ee/images/prints/documents/149019033869_eesti_toitumis- ja_liikumissoovitused.pdf.
41. Kyttälä P, Ovaskainen M, Kronberg-Kippilä C, *et al.* (2008) The diet of Finnish preschoolers (in Finnish, abstract in English) [Internet]. Available from: <http://www.julkari.fi/bitstream/handle/10024/78163/2008b32.pdf>.
42. Agence nationale de sécurité sanitaire de l'alimentation, de travail, de l'environnement et du (2017) Étude Individuelle Nationale des Consommations Alimentaires 3 (INCA3) (2014–2015) Rapport [Internet]. Available from: <https://www.anses.fr/fr/system/files/NUT2014SA0234Ra.pdf>.
43. Þórsdóttir I, Þórisdóttir ÁV & Pálsson GI (2008) The diet of Icelandic infants: results from a research on infants' diet, growth and iron status 2005–2007 [Internet]. Available from: <http://www.landlaeknir.is/servlet/file/store93/item11592/Ungbarnaskysla.pdf>.
44. Gunnarsdóttir I, Eysteinsdóttir T, Þórsdóttir T (2008) Hvað borða íslensk börn á leikskólaaldrí? Könnun _a mataræði 3ja og 5 á ra barna [Internet]. Available from: http://www.landlaeknir.is/servlet/file/store93/item14897/3ja_og_5_ara_skyrsla_181208.pdf.
45. Irish Universities Nutrition Alliance (2012) National Pre-School Nutrition Survey 2010–11 [Internet]. Available from: <https://irp-cdn.multiscreensite.com/46a7ad27/files/uploaded/The National Pre-School Nutrition Survey Summary Report June 2012.pdf>.
46. Israel Ministry of Health (2017) MABAT Kids – The National Survey on the Health and Nutrition Status of Children Ages 2–11 in Israel: 2015–2016, 79. Available from: https://www.health.gov.il/publicationsfiles/mabat_kids2_11_2015-2016.pdf.

47. Sette S, Le Donne C, Piccinelli R, *et al.* (2011) The third Italian National Food Consumption Survey, INRAN-SCAI 2005-06 – Part 1: nutrient intakes in Italy. *Nutr Metab Cardiovasc Dis* **21**, 922–932.
48. Sánchez-Pimienta TG, Batis C, Lutter CK, *et al.* (2016) Sugar-sweetened beverages are the main sources of added sugar intake in the Mexican population. *J Nutr* **146**, 1888S–1896S.
49. National Institute for Public Health and the Environment Ministry of Health Welfare and Sport (2019) Food consumption survey 2012–2016 [Internet]. Available from: <https://www.rivm.nl/bibliotheek/rapporten/2020-0083.pdf>.
50. Helsedirektoratet (2009) Spedkost – 12 måneder 2006–2007 [Internet]. Available from: [https://www.helsedirektoratet.no/rapporter/spedkost-og-smabarnskost-landsomfattende-kostholdsundersokelser/Spedkost 12 måneder – landsomfattende kostholdsundersøkelse 2007-2008.pdf/_/attachment/inline/1544ff4a-db55-4db1-b24b-f3efa19dd26b:20d97036e4aa5f5c92](https://www.helsedirektoratet.no/rapporter/spedkost-og-smabarnskost-landsomfattende-kostholdsundersokelser/Spedkost%2012%20måneder%20landsomfattende%20kostholdsundersokelse%202007-2008.pdf/_/attachment/inline/1544ff4a-db55-4db1-b24b-f3efa19dd26b:20d97036e4aa5f5c92).
51. Helsedirektoratet (2009) Småbarnskost – 2 år Landsomfattende kostholdsundersøkelse blant 2 år gamle barn. Available from: [https://www.helsedirektoratet.no/rapporter/spedkost-og-smabarnskost-landsomfattende-kostholdsundersokelser/Småbarnskost 2007 – landsomfattende kostholdsundersøkelse blant 2 år gamle barn.pdf/_/attachment/inline/54fa23d7-f5cc-4ce8-bc21-41418c21a126:5dee89c](https://www.helsedirektoratet.no/rapporter/spedkost-og-smabarnskost-landsomfattende-kostholdsundersokelser/Småbarnskost%202007%20landsomfattende%20kostholdsundersokelse%20blant%2020år%20gamle%20barn.pdf/_/attachment/inline/54fa23d7-f5cc-4ce8-bc21-41418c21a126:5dee89c).
52. Public Health England, Food Standards Agency (2018) National Diet and Nutrition Survey: results from Years 1–4 (combined) of the Rolling Programme [Internet]. Available from: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/699241/NDNS_results_years_7_and_8.pdf.
53. Bailey RL, Fulgoni VL, Cowan AE, *et al.* (2018) Sources of added sugars in young children, adolescents, and adults with low and high intakes of added sugars. *Nutrients* **10**, 102.
54. Herrick KA, Fryar CD, Hamner HC, *et al.* (2020) Added sugars intake among US infants and toddlers. *J Acad Nutr Diet* **120**, 23–32.
55. US Department of Agriculture (2020) Nutrient intakes from food: mean amounts consumed per individual, by gender and age, what we eat in America, NHANES 2017–2018. Available from: www.ars.usda.gov/nea/bhnrc/fsrg.
56. Elmfada I (2012) Österreichischer Ernährungsbericht 2012 [Internet]. Available from: https://ernaehrungsbericht.univie.ac.at/fileadmin/user_upload/dep_ernaehrung/forschung/ernaehrungsberichte/oesterr_ernaehrungsbericht_2012.pdf.
57. Souza RAG, Yokoo EM, Sichieri R, *et al.* (2015) Energy and macronutrient intakes in Brazil: results of the first nationwide individual dietary survey. *Public Health Nutr* **18**, 3086–3095.
58. Statistics Canada (2018) *Canadian community health survey – nutrition*. Ottawa: Health Canada.
59. Afeiche MC, Koyraty BNS, Wang D, *et al.* (2018) Intakes and sources of total and added sugars among 4 to 13-year-old children in China, Mexico and the United States. *Pediatr Obes* **13**, 204–212.
60. Pedersen AN, Christensen T, Matthiessen J, Knudsen VK, Sørensen MR, Biloft-Jensen AP, Fagt S (2015) Danskernes kostvaner 2011–2013 [Internet]. Available from: http://www.food.dtu.dk/-/media/Institut/Foedevareinstituttet/Publikationer/Pub-2015/Rapport_Danskernes-Kostvaner-2011-2013.ashx?la=da.
61. Mensink GBM, Helmut H, Stahl A, Richer A & Vohmann C (2007) Die aktuelle Nährstoffzufuhr von Kindern und Jugendlichen in Deutschland – Ergebnisse aus EsKiMo, Ernährungs Umschau [Internet]. Available from: https://www.ernaehrungs-umschau.de/fileadmin/Ernaehrungs-Umschau/pdfs/pdf_2007/11_07/EU11_636_646.qxd.pdf.
62. Gunnarsdottir I, Helgadóttir H, Thorisdóttir B, *et al.* (2013) Diet of six-year-old Icelandic children – National dietary survey 2011–2012. *Laeknabladid* Laeknafelag Islands; **99**, 17–23.
63. Unit for Nutrition Research (2006) Iceland dietary survey of 9 and 15 year old children and adolescents 2003–2004 [Internet]. Available from: http://www.landlaeknir.is/servlet/file/store93/item11594/hvad_borda_isl_bom_og_ungl.pdf.
64. International Life Sciences Institute India (2019) Consumption levels of sugar among rural and urban population in India [Internet]. Available from: <http://www.ilsa-india.org/> (Accessed December 2020).
65. Irish Universities Nutrition Alliance (2019) National Children's Food Consumption Survey II (NCFS II) 2017–2018 [Internet]. Available from: [rp-cdn.multiscreensite.com/46a7ad27/files/uploaded/The National Children's Food Survey II Summary Report – September 2019.pdf](http://rp-cdn.multiscreensite.com/46a7ad27/files/uploaded/The%20National%20Children's%20Food%20Survey%20II%20Summary%20Report%20September%202019.pdf).
66. National Center for Disease Control Israel National Survey on the Subject of Health and Nutrition Status in children aged 2–11 in Israel [Internet]. Available from: https://www.health.gov.il/publicationsfiles/mabat_kids2_11_2015-2016.pdf.
67. Ministry of Health (2003) 2002 National Children's Nutrition Survey (CNS02) [Internet]. Wellington. Available from: <https://www.health.govt.nz/system/files/documents/publications/nzfoodnzchildren.pdf>.
68. FHI (2017) UNGKOST 3 Landsomfattende kostholdundersøkelse blant 4-åringer i Norge, 2016 [Internet]. Available from: <https://www.fhi.no/globalassets/dokumenterfiler/rapporter/2016/rapport-ungkost-3-landsomfattende-kostholdsundersokelse-blant-4-aringer-i-norge-2016.pdf>.
69. FHI (2016) UNGKOST 3 Landsomfattende kostholdsundersøkelse blant elever i 4. -og 8. klasse i Norge, 2015 [Internet]. Available from: <https://www.fhi.no/globalassets/dokumenterfiler/rapporter/2016/ungkost-rapport-24.06.16.pdf>.
70. IAN-AF (2017) National Food and Physical Activity Survey (IAN-AF) 2015–2016. ANEXO 2 – INGESTÃO NUTRICIONAL [Internet]. Available from: https://ian-af.up.pt/en/consultade-dados?shs_term_node_tid_depth=All&field_indicador_tid=All&field_estratificacao_tid=All%0A (accessed December 2020).
71. FEN (2014) Ingesta, perfil y fuentes de energía en la población española: Resultados obtenidos del estudio científico ANIBES. Available from: http://www.fen.org.es/anibes/archivos/documentos/ANIBES_numero_3.pdf.
72. Ruiz E, Rodríguez P, Valero T, *et al.* (2017) Dietary intake of individual (free and intrinsic) sugars and food sources in the Spanish population: findings from the ANIBES study. *Nutrients* **9**, 275.
73. Roberts C, Steer T, Maplethorpe N, *et al.* (2018) National diet and nutrition survey: results from Years 7 and 8 (combined) of the Rolling Programme (2014/2015–2015/2016). London: Public Health England.
74. d'Andorra G, de Salut M & i Família BS (2007) Evaluation of the nutritional status of the Andorran population [Internet]. Available from: http://www.salut.ad/images/microsites/AvaluacioNutricional_04-05/index.html (accessed December 2020).
75. Langlois K, Garriguet D, Gonzalez A, *et al.* (2019) Change in total sugars consumption among Canadian children and adults. *Health Rep* **30**, 10–19.
76. Perrar I, Schadow AM, Schmitting S, *et al.* (2020) Time and age trends in free sugar intake from food groups among children and adolescents between 1985 and 2016. *Nutrients* **12**, 20.

77. Irish Universities Nutrition Alliance (IUNA) (2007) National Teens' Food Survey 2005–2006. [Internet]. Available from: [https://irp-cdn.multiscreensite.com/46a7ad27/files/uploaded/The National Teens%27 Food Survey %282005-2006%29.pdf](https://irp-cdn.multiscreensite.com/46a7ad27/files/uploaded/The%20National%20Teens%20Food%20Survey%202005-2006%29.pdf).
78. Joyce T, McCarthy SN & Gibney MJ (2008) Relationship between energy from added sugars and frequency of added sugars intake in Irish children, teenagers and adults. *Br J Nutr* **99**, 1117–1126.
79. Israel Center for Disease Control (2017) Mabat Youth Second National Health and Nutrition Survey of 7th–12th grade students 2015–2016. Publication 373 [Internet]. Available from: https://health.gov.il/publicationsfiles/mabat_kids2_11_2015-2016-eng.pdf%0A%0A.
80. Kibblewhite R, Nettleton A, McLean R, *et al.* (2017) Estimating free and added sugar intakes in New Zealand. *Nutrients* **9**, 1292.
81. University of Otago and Ministry of Health (2011) *A focus on nutrition: key findings of the 2008/09 New Zealand Adult Nutrition Survey*. Wellington: Ministry of Health.
82. Mis NF, Kobe H, Štítec M. (2012) Dietary intake of macro- and micronutrients in Slovenian adolescents: comparison with reference values. *Ann Nutr Metab* **61**, 305–313.
83. Rust P, Hasenegger V & König J (2017) Österreichischer Ernährungsbericht [Internet]. Available from: <https://broschuerenservice.sozialministerium.at/Home/Download?publicationId=528%0A%0A> (accessed December 2020).
84. Valsta L, Kaartinen N, Tapanainen H, Männistö S & Sääksjärvi K (2018) Ravitsemus Suomessa – FinRavinto 2017 -tutkimus. Terveystien ja hyvinvoinnin laitos (THL). Raportti 12/2018, 239 sivua [Internet]. Available from: https://www.julkari.fi/bitstream/handle/10024/137433/URN_ISBN_978-952-343-238-3.pdf?sequence=1&isAllowed=y%0A%0A%0A (accessed December 2020).
85. Max Rubner-Institut (2008) Nationale Verzehrsstudie II – Ergebnisbericht Teil 1; Max Rubner-Institut Karlsruhe: Karlsruhe [Internet]. Available from: https://www.bmel.de/SharedDocs/Downloads/Ernaehrung/NVS_ErgebnisberichtTeil2.pdf?__blob=publicationFile%0A%0A (accessed December 2020).
86. Cust AE, Skilton MR, van Bakel MME, *et al.* (2009) Total dietary carbohydrate, sugar, starch and fibre intakes in the European prospective investigation into cancer and nutrition. *Eur J Clin Nutr* **63**, S37–S60.
87. Eszter SN, Márta B, Éva I, *et al.* (2017) Országos táplálkozás és tápláltsági állapot vizsgálat – OTÁP2014. II. A magyar lakosság energia-és makrotápanyag-bevitele. *Orv Hetil* **158**, 587–597.
88. Directorate of Health (2011) Hvað borða Íslendingar? Könnun á mataræði Íslendinga 2010–2011 Helstu niðurstöður 2010–2011. Helstu niðurstöður [Internet]. Available from: [http://www.landlaeknir.is/servlet/file/store93/item14901/Hvað borða Íslendingar_april 2012.pdf](http://www.landlaeknir.is/servlet/file/store93/item14901/Hvað%20borða%20Íslendingar_april%202012.pdf).
89. Irish Universities Nutrition Alliance (IUNA) (2011) National adult nutrition survey: food and nutrient intakes, physical measurements, physical activity patterns and food choice motives. *Summary Report*. [Internet]. Available from: [https://irp-cdn.multiscreensite.com/46a7ad27/files/uploaded/The%20 National%20Adult%20Nutri](https://irp-cdn.multiscreensite.com/46a7ad27/files/uploaded/The%20National%20Adult%20Nutri).
90. Walton J, Kehoe L, McNulty BA, *et al.* (2017) Intakes and sources of dietary sugars in a representative sample of Irish adults (18–90y). *Proc Nutr Soc* [Internet]. Cambridge University Press; [cited 2021 Jan 4]; 76, E65. Available from: https://www.cambridge.org/core/product/identifier/S0029665117001380/type/journal_article.
91. Israel Center for Disease Control (2019) Rav Mabat Adult Second National Health and Nutrition Survey 2014–2016. Publication 383 [Internet]. Available from: https://health.gov.il/publicationsfiles/mabat_adults_2014_2016_383_en.pdf.
92. Barzda A, Bartkeviciute R, Baltusyte I, Stukas R & Bartkeviciute S (2016) Suaugusių ir pagyvenusių Lietuvos gyventojų, faktin̄ es mitybos ir mitybos i,prōciu, tyrimas. *Visuom. Sveik* **72**, 85–94. [Internet]. Available from: [http://www.hi.lt/uploads/pdf/visuomenes_sveikata/2016.01.72/VS_2016_1\(72\)_ORIG_Mitybos_iprociai.pdf%0A%0A](http://www.hi.lt/uploads/pdf/visuomenes_sveikata/2016.01.72/VS_2016_1(72)_ORIG_Mitybos_iprociai.pdf%0A%0A).
93. Helsedirektoratet (2012) Norkost 3 En landsomfattende kostholdsundersøkelse blant menn og kvinner i Norge I alderen 18–70ar, 2010–11 [Internet]. Available from: <https://helsedirektoratet.no/Lists/Publikasjoner/Attachments/301/Norkost-3-en-landsomfattende-kostholdsundersokelse-blant-menn-og-kvinner-i-norge-i-alderen-18-70-ar-2010-11-IS-2000.pdf>.
94. Chatelan A, Gaillard P, Kruseman M, *et al.* (2019) Total, added, and free sugar consumption and adherence to guidelines in Switzerland: results from the First National Nutrition Survey menuCH. *Nutrients* [Internet]. MDPI AG; [cited 2020 Jun 10]; 11. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/31109151>.
95. Heuer T (2017) Estimation of “free” sugar intake in Germany based on the data of the German National Nutrition Survey II. Available from: https://www.openagrar.de/receive/openagrar_mods_00030594.
96. De la Cruz-Góngora V, Martínez-Tapia B, Cuevas-Nasu L, *et al.* (2017) Dietary intake and adequacy of energy and nutrients in Mexican older adults: Results from two National Health and Nutrition Surveys. *Salud Publica Mex* **59**, 285–298.
97. Azaïs-Braesco V, Sluik D, Maillot M, *et al.* (2017) A review of total & added sugar intakes and dietary sources in Europe. *Nutr J BioMed Central Ltd*. **16**, 6.
98. IUNA (2019) National Children's Food Survey II: summary report [Internet], 1–44. Available from: www.iuna.net.
99. Hayes M (2019) How many countries are there in the world? [Internet]. WorldAtlas, [cited 2020 Jun 15]. Available from: <https://www.worldatlas.com/articles/how-many-countries-are-in-the-world.html>.
100. Micha R, Coates J, Leclercq C, *et al.* (2018) Global dietary surveillance: data gaps and challenges. *Food Nutr Bull* **39**, 175–205.
101. Louie JCY, Moshtaghan H, Boylan S, *et al.* (2015) A systematic methodology to estimate added sugar content of foods. *Eur J Clin Nutr* **69**, 154–161.
102. Swan GE, Powell NA, Knowles BL, *et al.* (2018) A definition of free sugars for the UK. *Public Health Nutr* **21**, 1636–1638.
103. Amoutzopoulos B, Page P, Steer T, *et al.* (2020) Free and added sugar consumption and adherence to guidelines: the UK national diet and nutrition survey (2014/15–2015/16). *Nutrients* MDPI AG, **12**, 393.
104. Amoutzopoulos B, Steer T, Roberts C, *et al.* (2018) A disaggregation methodology to estimate intake of added sugars and free sugars: an illustration from the UK national diet and nutrition survey. *Nutrients* MDPI AG; **10**, 1177.
105. Davy B & Jähren H (2016) New markers of dietary added sugar intake. *Curr Opin Clin Nutr Metab Care* **19**, 282–288.
106. Abreu TC, Hulshof PJM, Boshuizen HC, *et al.* (2020) Validity coefficient of repeated measurements of urinary marker of sugar intake is comparable to urinary nitrogen as marker of protein intake in free-living subjects. *Cancer Epidemiol Biomarkers Prev* **30**, 193–202.
107. Mela DJ (2020) A proposed simple method for objectively quantifying free sugars in foods and beverages. *Eur J Clin Nutr* [Internet] **74**, 1366–1368. doi: 10.1038/s41430-020-0575-x

108. Louie JCY (2020) Objective biomarkers for total added sugar intake – are we on a wild goose chase? *Adv Nutr* **11**, 1429–1436.
109. Capacci S, Allais O, Bonnet C, *et al.* (2019) The impact of the French soda tax on prices and purchases. An ex post evaluation. *PLoS One* **14**, e0223196.
110. World Health Organisation (2017) Reformulation measures for foods and beverages – European Health Information Gateway [Internet], [cited 2020 Jun 15]. Available from: https://gateway.euro.who.int/en/indicators/gnpr_survey_17-reformulation-measures-for-foods-and-beverages/.
111. Public Health England (2018) Sugar reduction and wider reformulation programme: report on progress towards the first 5 % reduction and next steps May 2018 About Public Health England [Internet]. Available from: www.facebook.com/PublicHealthEngland.
112. Wright A, Smith KE & Hellowell M (2017) Policy lessons from health taxes: a systematic review of empirical studies. *BMC Public Health* **17**, 583.
113. Redondo M, Hernández-Aguado I & Lumberras B (2018) The impact of the tax on sweetened beverages: a systematic review. *Am J Clin Nutr* **108**, 548–563.
114. Briggs ADM, Mytton OT, Kehlacher A, *et al.* (2013) Overall and income specific effect on prevalence of overweight and obesity of 20% sugar sweetened drink tax in UK: econometric and comparative risk assessment modelling study. *BMJ* **347**, f6189.
115. Blakely T, Nghiem N, Genc M, *et al.* (2020) Modelling the health impact of food taxes and subsidies with price elasticities: the case for additional scaling of food consumption using the total food expenditure elasticity. *PLoS One* **15**, e0230506.
116. Ma Y, He FJ, Yin Y, *et al.* (2016) Gradual reduction of sugar in soft drinks without substitution as a strategy to reduce overweight, obesity, and type 2 diabetes: a modelling study. *Lancet Diabetes Endocrinol* **4**, 105–114.
117. Amies-Cull B, Briggs ADM & Scarborough P (2019) Estimating the potential impact of the UK government's sugar reduction programme on child and adult health: modelling study. *BMJ* **365**.
118. Center for Food Safety and Applied Nutrition (2018) *Labeling & nutrition – changes to the nutrition facts label*. Washington DC: United States Department of Health & Human Services.
119. U.S. Department of Agriculture & Global Agricultural Information Network (2020) Implementation of the New Mexican Regulation for front-of-package nutrition label.
120. Tierney M, Gallagher AM, Giotis ES, *et al.* (2017) An online survey on consumer knowledge and understanding of added sugars. *Nutrients* **9**, 37.
121. Tang Q, Lin Q, Yang Q, *et al.* (2020) Knowledge, attitude, and practice of adolescent parents on free sugar and influencing factors about recognition. *Int J Environ Res Public Health* **17**, 4003.