

to assess psychological distress, social anxiety and physical activity, respectively. Data were analysed using SPSS v25, with application of multivariate logistic regression to assess association of various factors with psychological distress.

Results. Out of 256 medical students, 73 (28.5%) had severe psychological distress and 105 (41.0%) had mild to moderate distress. A logistic regression model to assess the effect of social anxiety disorder, level of physical activity and gender with the likelihood of having severe psychological distress was statistically significant ($p < 0.001$) with overall accuracy of 73.8%. The risk of developing severe psychological distress was higher among females [OR 2.13 (95% CI 1.17–3.87), $p = 0.013$] and those with social anxiety disorder [OR 4.56 (95% CI 2.27–9.16), $p < 0.001$]. Low physical activity was not a significant risk factor for psychological distress [OR 0.88 (95% CI 0.35–2.23), $p = 0.794$].

Conclusion. This study shows that COVID-19 imposed lockdown has adversely affected the mental health of medical students. Female students and those with social anxiety disorder are at more risk of developing severe psychological distress.

A Feasibility Evaluation of Discovery Group: Determining the Acceptability and Potential Outcomes of a Patient-led Research Group in a Secure Mental Health Inpatient setting

Dr Mudassar Arslan*, Dr Anne Aboaja,
Dr Oluwatosin Atewogboye, Miss Lucia Parry-Newton
and Miss Lindsey Wilson

Tees, Esk and Wear Valleys NHS Foundation Trust, Middlesbrough,
United Kingdom

*Presenting author.

doi: 10.1192/bjo.2022.170

Aims. Patient and public involvement and engagement (PPIE) is recognised as an essential part of health research. It provides an opportunity for patients to shape health research and acquire research skills, in the inpatient mental health setting, PPIE may have additional value in providing meaningful activity and enhancing recovery, as defined using connectedness, hope, identity, meaning and empowerment (CHIME) principles. An eight-session PPIE programme ("Discovery Group") was designed to support patient-led research in a secure mental health hospital. This feasibility study aims to evaluate the acceptability of the programme from the perspective of patients and identify potential outcomes.

Methods. A retrospective single-arm post-programme evaluation of Discovery Group was undertaken. Participants attended an evaluation workshop where they were interviewed individually to complete an acceptability questionnaire designed using the domains of the Theoretical Framework of Acceptability. Participants also completed an outcomes questionnaire, which included CHIME-based recovery items. Quantitative data were analysed descriptively. Direct content analysis was applied to qualitative data.

Results. In our sample, eight participants attended at least one session of the discovery group with one patient attending all sessions. Most of the participants felt positive about taking part in the group and expressed interest in joining another group in future. All participants experienced some burden from the effort required during group sessions, but a low level of opportunity cost in terms of the extent to which they perceived they had to forfeit benefits to participate in the programme. Some described the group as effective in helping them learn about research. Of the five CHIME recovery domains, only connectedness was reported as a benefit of the group. The participants valued the opportunity to use their time.

Conclusion. Discovery Group is a novel intervention that offers high level, non-tokenistic PPIE suitable for use in secure mental health inpatient settings. It produces research of value to patients through a programme of high acceptability and provides them with potential benefits of recovery as well as research knowledge and skills, and an activity that alleviates boredom, enhances autonomy, breaks down some important power and paternalistic barriers that can be experienced by patients detained in secure mental health settings. Finally, a future evaluation study that involves patients during the design, implementation, evaluation and writing stages, aiming to measure the potential outcomes identified in the present study using pre- and post-testing with a control group would reliably demonstrate the effectiveness of the revised Discovery Group and ensure meaningful involvement with patients as co-researchers.

'Sheldon' in the Medical Field: Emotional Intelligence and Its Associated Factors in Medical Students of Pakistan

Dr Unsa Athar*, Dr Muhammad Bilal Rehman,
Mr Ahsan Nawaz, Ms Aimen Siddique, Ms Ambreen Fatima,
Mr Alamgir Aslam Khan and Dr Muhammad Luqman Nagi
Shalamar Medical and Dental College, Lahore, Pakistan

*Presenting author.

doi: 10.1192/bjo.2022.171

Aims. The dearth of emotional intelligence leads to medical students' inability to handle the pressures of medical education, subsequently causing burnout and mental illnesses. Poor emotional intelligence in young doctors also begets increased mistrust from the public, lowering the quality of healthcare delivery. Emotional intelligence of Pakistani students, similar to global context, is impacted by a myriad of psycho-socio-economic factors. It is pertinent to find out the detrimental and/or protective factors, and design interventions to enhance emotional intelligence as a soft skill. With this aim in mind, we explored the relationship of emotional intelligence with adverse childhood experiences and prevalent mental illnesses (depression and anxiety) amongst the medical students of one of the most populous cities of Pakistan; Lahore.

Methods. A cross-sectional study was conducted including currently enrolled MBBS (Bachelor of Medicine and Surgery) students from first year to final year in 2 medical schools of Lahore. An online google form was constructed by combining Modified Adverse Childhood Experiences Score Scale (ACES), Brief Emotional Intelligence Scale (BEIS-10) and Hospital Depression and Anxiety Scale (HADS). Data were exported to SPSS version 25.0 for descriptive and analytical analysis. Pearson's chi-square analysis and logistic regression analysis were used to study the association between the outcome and dependent variables; Odds ratio (OR) with 95% Confidence Intervals (CI) were calculated.

Results. Participants (N = 324) belonging to two different medical colleges in Lahore, namely King Edward Medical College (public) and Shalamar Medical and Dental College (private), took part in the study.

Pearson's chi-square showed significant association of emotional intelligence with early private schooling ($p = 0.029$), nuclear family system (0.044) and the presence of symptoms of depression (0.005). The adjusted logistic regression model showed that people who studied in a private sector school (OR: 2.12, CI: 1.01–4.45) and people who lived in a nuclear family (OR: 2.02, CI: 1.00–4.08) had significantly twice the likelihood of having high emotional intelligence. Also noteworthy is that respondents who