



Tensorially absorbing inclusions of C^* -algebras

Paweł Sarkowicz

Abstract. When \mathcal{D} is strongly self-absorbing, we say an inclusion $B \subseteq A$ of C^* -algebras is \mathcal{D} -stable if it is isomorphic to the inclusion $B \otimes \mathcal{D} \subseteq A \otimes \mathcal{D}$. We give ultrapower characterizations and show that if a unital inclusion is \mathcal{D} -stable, then \mathcal{D} -stability can be exhibited for countably many intermediate C^* -algebras concurrently. We show that such unital embeddings between unital \mathcal{D} -stable C^* -algebras are point-norm dense in the set of all unital embeddings, and that every unital embedding between \mathcal{D} -stable C^* -algebras is approximately unitarily equivalent to a \mathcal{D} -stable embedding. Examples are provided.

1 Introduction

The study of inclusions of C^* -algebras has been of recent interest. There is no short supply of research concerning inclusions relating to noncommutative dynamics [8, 18, 30, 42, 44], as well as inclusions of simple C^* -algebras [51]. There has also been work done regarding the passage of properties from a subalgebra to a larger algebra using tracial approximations [38]. We discuss inclusions from the lens of tensorially absorbing a strongly self-absorbing C^* -algebra \mathcal{D} [66].

When speaking of tensorial absorption with a strongly self-absorbing C^* -algebra, central sequences play a role akin to McDuff’s characterization of when a II_1 von Neumann algebra absorbs the unique hyperfinite II_1 factor \mathcal{R} [40]. Central sequences have been studied since the inception of operator algebras, being used by Murray and von Neumann to exhibit non-isomorphic II_1 factors by showing that $\mathcal{L}(\mathbb{F}_2)$ does not have property Γ [41]. They were also used in Connes’ theorem concerning the uniqueness of \mathcal{R} [13], and the classification of automorphisms on hyperfinite factors [11, 13]. In [2, 3], Bisch considered the central sequence algebra $\mathcal{N}^\omega \cap \mathcal{M}'$ associated with an (irreducible) inclusion of II_1 factors $\mathcal{N} \subseteq \mathcal{M}$ and characterized when there was an isomorphism $\Phi : \mathcal{M} \simeq \mathcal{M} \overline{\otimes} \mathcal{R}$ such that $\Phi(\mathcal{N}) = \mathcal{N} \overline{\otimes} \mathcal{R}$ in terms of the existence of non-commuting sequences in \mathcal{N} which asymptotically commute with the larger von Neumann algebra \mathcal{M} (in the $\|\cdot\|_2$ -norm). As pointed out by Izumi [31], there are similar central characterizations for unital inclusions of separable C^* -algebras which tensorially absorb a strongly self-absorbing C^* -algebra \mathcal{D} (it was at least pointed out for \mathcal{D} being one of M_{n^∞} , \mathcal{O}_2 , \mathcal{O}_∞).

Received by the editors May 28, 2023; revised February 19, 2024; accepted April 1, 2024.

Published online on Cambridge Core April 12, 2024.

AMS subject classification: 46L06, 46L35.

Keywords: C^* -algebras, tensor products, inclusions, strongly self-absorbing C^* -algebras.



regularity conditions, we show that we can witness \mathcal{L} -stability of all such intermediate C^* -algebras concurrently.

Theorem 1.3 (Corollary 5.6) *Let A be a unital, simple, separable, nuclear \mathcal{L} -stable C^* -algebra and $G \curvearrowright^\alpha A$ be an action of a finite group with the weak tracial Rokhlin property. There exists an isomorphism $\Phi : A \rtimes_\alpha G \simeq (A \rtimes_\alpha G) \otimes \mathcal{L}$ such that whenever C is a unital C^* -algebra satisfying either:*

- (1) $A^\alpha \subseteq C \subseteq A$ or
- (2) $A \subseteq C \subseteq A \rtimes_\alpha G$,

we have $\Phi(C) = C \otimes \mathcal{L}$.

This paper is structured as follows. We discuss various local properties in Section 3, and then formalize the notion of a \mathcal{D} -stable embedding in Section 4, examining several properties and consequences. In Section 5, we show how several examples arising from noncommutative dynamical systems fit into the framework of \mathcal{D} -stable inclusions. We finish with several examples in Section 6.

2 Preliminaries

2.1 Notation

We use capital letters A, B, C, D to denote C^* -algebras and usually a calligraphic \mathcal{D} to denote a strongly self-absorbing C^* -algebra. Generally, small letters $a, b, c, d, \dots, x, y, z$ will denote operators in C^* -algebras. A_+ will denote cone of positive elements in a C^* -algebra A . If $\varepsilon > 0$ and a, b are elements in a C^* -algebra, we will write

$$(2.1) \quad a \approx_\varepsilon b$$

to mean that $\|a - b\| < \varepsilon$. This will make some approximations more legible.

The symbol \otimes will denote the minimal tensor product of C^* -algebras, while \odot will mean the algebraic tensor product. We use the minimal tensor product throughout, and it is common for us to deal with nuclear C^* -algebras so there should not be any ambiguity. The symbol $\bar{\otimes}$ will denote the von Neumann tensor product.

We will denote by M_n the C^* -algebra of $n \times n$ matrices, and M_{n^∞} the uniformly hyperfinite (UHF) C^* -algebra associated with the supernatural number n^∞ . We will write \mathcal{Q} for the universal UHF algebra $\mathcal{Q} = \bigotimes_{n \in \mathbb{N}} M_n$.

By $G \curvearrowright^\alpha A$, we will mean that the (discrete) group G acts on A by automorphisms, i.e., $\alpha : G \rightarrow \text{Aut}(A)$ is a homomorphism. $A \rtimes_{r, \alpha} G$ will denote the reduced crossed product, which we will just write as $A \rtimes_\alpha G$ if it is clear from context that the group is amenable and A is nuclear (e.g., if G is finite). We will denote by A^α the fixed point subalgebra of the action (or A^G if the action is clear from context).

For a map $f : X \rightarrow Y$ between sets X and Y , we will write $f : X \hookrightarrow Y$ to mean that f is injective and $f : X \twoheadrightarrow Y$ to mean that f is surjective. This will usually be done in the context of $*$ -homomorphisms.

2.2 Ultrapowers, central sequences, and central sequence algebras

Fix a free ultrafilter $\omega \in \beta\mathbb{N}$. Throughout, we will use ultrapowers to describe asymptotic behavior. Alternatively, one can use sequence algebras, although this comes down to a matter of taste and one can swap between the two if desired, as we will provide local characterizations. This also means that all of what we do will be independent of the specific ultrafilter ω .

For a C^* -algebra A , the ultrapower of A is the C^* -algebra

$$(2.2) \quad A_\omega := \ell^\infty(A)/c_{0,\omega}(A),$$

where $c_{0,\omega} := \{(a_n)_{n \in \mathbb{N}} \in \ell^\infty(A) \mid \lim_{n \rightarrow \omega} \|a_n\| = 0\}$ is the ideal of ω -null sequences. We can embed A into A_ω canonically by means of constant sequences: we identify $a \in A$ with the equivalence class of the constant sequence $(a)_{n \in \mathbb{N}}$.

To ease notation, we will usually write elements of A_ω as sequences $(a_n)_{n \in \mathbb{N}}$, keeping in mind that these are equivalence classes without explicitly stating it every time. We note that the norm on A_ω is given by $\|(a_n)_{n \in \mathbb{N}}\| = \lim_{n \rightarrow \omega} \|a_n\|$.

Kirchberg's ε -test [35, Lemma A.1] is essentially the operator algebraists' Łoś' theorem without having to turn to (continuous) model theory. Heuristically, it says that if certain things can be done approximately in an ultrapower, then certain things can be done exactly in an ultrapower.

Lemma 2.1 (Kirchberg's ε -test) *Let $(X_n)_n$ be a sequence of sets and suppose that for each n , there is a sequence $(f_n^{(k)})_{k \in \mathbb{N}}$ of functions $f_n^{(k)} : X_n \rightarrow [0, \infty)$. For $k \in \mathbb{N}$, let*

$$(2.3) \quad f_\omega^k(s_1, s_2, \dots) := \lim_{n \rightarrow \omega} f_n^{(k)}(s_n).$$

Suppose that for every $m \in \mathbb{N}$ and $\varepsilon > 0$, there is $s \in \prod_n X_n$ with $f_\omega^{(k)}(s) < \varepsilon$ for $k = 1, \dots, m$. Then there exists $t \in \prod_n X_n$ with $f_\omega^{(k)}(t) = 0$ for all $k \in \mathbb{N}$.

The above is useful, although if one so wishes, one can usually construct exact objects from approximate objects by using standard diagonalization arguments (under some separability assumptions). These sorts of arguments work in both the ultrapower setting and the sequence algebra setting.

Finally, if $\alpha \in \text{Aut}(A)$ is an automorphism, there is an induced automorphism on A_ω , which we will denote by α_ω , given by

$$(2.4) \quad \alpha_\omega((a_n)_{n \in \mathbb{N}}) := (\alpha(a_n))_{n \in \mathbb{N}}.$$

2.3 Central sequences and central sequence subalgebras

For a unital C^* -algebra A , the C^* -algebra of ω -central sequences is

$$(2.5) \quad A_\omega \cap A' = \{x \in A_\omega \mid [x, a] = 0 \text{ for all } a \in A\},$$

where we are identifying $A \subseteq A_\omega$ with the constant sequences. If $B \subseteq A$ is a unital C^* -subalgebra and $S \subseteq A_\omega$ is a subset, we can associate the relative commutant of S in B_ω :

$$(2.6) \quad B_\omega \cap S' = \{b \in B_\omega \mid [b, s] = 0 \text{ for all } s \in S\}.$$

Of particular interest will be when $S = A$, and $B \subseteq A$ is a unital inclusion of separable C^* -algebras.

2.4 Strongly self-absorbing C^* -algebras

A unital separable C^* -algebra \mathcal{D} is strongly self-absorbing if $\mathcal{D} \neq \mathbb{C}$ and there is an isomorphism $\phi : \mathcal{D} \rightarrow \mathcal{D} \otimes \mathcal{D}$ which is approximately unitarily equivalent to the first factor embedding $d \mapsto d \otimes 1_{\mathcal{D}}$ (see [66]). All known strongly self-absorbing C^* -algebras are: the Jiang–Su algebra \mathcal{Z} [32], the Cuntz algebras \mathcal{O}_2 and \mathcal{O}_{∞} [15], UHF algebras of infinite type, and \mathcal{O}_{∞} tensor a UHF algebra of infinite type. Strongly self-absorbing C^* -algebras have approximately inner flip and therefore have K -theoretic restrictions (see [20, 61]). They are also nuclear, simple, and have at most one tracial state [66].

Tensorial absorption with strongly self-absorbing C^* -algebras gives rise to many regular properties, for example, in terms of K -theory, traces, and the Cuntz semi-group [32, 46, 47, 50]. Of paramount interest is the Jiang–Su algebra \mathcal{Z} . An accumulation of work has successfully classified all (unital) separable, simple, nuclear, infinite-dimensional, \mathcal{Z} -stable C^* -algebras satisfying the Universal Coefficient Theorem (UCT) of Rosenberg and Schochet [53] by means of K -theory and traces (see [9] and the references therein). We describe how one might work with \mathcal{Z} -stability in terms of its standard building blocks. Recall that, for $n, m \geq 2$, the dimension drop algebras are

$$(2.7) \quad \mathcal{Z}_{n,m} := \{f \in C([0,1], M_n \otimes M_m) \mid f(0) \in M_n \otimes 1_{M_m}, f(1) \in 1_{M_n} \otimes M_m\}.$$

Such an algebra is called a prime dimension drop algebra when n and m are coprime. The Jiang–Su algebra \mathcal{Z} is the unique separable simple C^* -algebra with unique tracial state which is an inductive limit of prime dimension drop algebras with unital connecting maps [32] (in fact, the dimension drop algebras can be chosen to have the form $\mathcal{Z}_{n,n+1}$). It is KK -equivalent to \mathbb{C} and \mathcal{Z} -stability is a often necessary condition for K -theoretic classification.

By [52, Proposition 5.1] (or [54, Proposition 2.1] for our desired formulation), $\mathcal{Z}_{n,n+1}$ is the universal C^* -algebra generated by elements c_1, \dots, c_n and s such that:

- $c_1 \geq 0$;
- $c_i c_j^* = \delta_{ij} c_1^2$;
- $s^* s + \sum_{i=1}^n c_i^* c_i = 1$;
- $c_1 s = s$.

If there are uniformly tracially large (in the sense of [65, Definition 2.2]) order zero¹ c.p.c. maps $M_n \rightarrow A_{\omega} \cap A'$, these give rise to elements $c_1, \dots, c_n \in A_{\omega} \cap A'$ with $c_1 \geq 0$ and $c_i c_j^* = \delta_{ij} c_1^2$, along with certain tracial information. If A has strict comparison, Matui and Sato used this tracial information to show that A has property (SI) [39], from which one can get an element $s \in A_{\omega} \cap A'$ such that $s^* s + \sum_{i=1}^n c_i^* c_i = 1$ and $c_1 s = s$. This gives a $*$ -homomorphism $\mathcal{Z}_{n,n+1} \rightarrow A_{\omega} \cap A'$, which if can be done for each $n \in \mathbb{N}$, is enough to conclude that $\mathcal{Z} \hookrightarrow A_{\omega} \cap A'$ unitaly and hence $A \simeq A \otimes \mathcal{Z}$.

¹Order zero meaning orthogonality preserving: $\phi : A \rightarrow B$ is c.p.c. order zero if it is c.p.c. and $\phi(a)\phi(b) = 0$ whenever $ab = 0$.

(see [67, 71]). In fact, it suffices to show that $\mathcal{Z}_{2,3} \hookrightarrow A_\omega \cap A'$ (or $\mathcal{Z}_{n,n+1}$ for some $n \geq 2$) (see [52, Theorem 3.4(ii)] and [56, Theorem 5.15]).

3 Approximately central approximate embeddings

Here, we formalize some results on approximate embeddings. When $B \subseteq A$ is a unital inclusion of separable C^* -algebras, this will yield local characterizations of nuclear subalgebras of $B_\omega \cap A'$, as defined in (2.6). Recall that we write u.c.p. or c.p.c. to mean that a map is unital and completely positive or completely positive and contractive, respectively.

Definition 3.1 Let $B \subseteq A$ be a unital inclusion of C^* -algebras, and let D be a unital, simple, nuclear C^* -algebra. Let $\mathcal{F} \subseteq D, \mathcal{G} \subseteq A$ be finite sets and $\varepsilon > 0$. We say that a u.c.p. map $\phi : D \rightarrow B$ is an $(\mathcal{F}, \varepsilon)$ -approximate embedding if:

$$(1) \quad \phi(cd) \approx_\varepsilon \phi(c)\phi(d) \text{ for all } c, d \in \mathcal{F}.$$

If ϕ additionally satisfies

$$2. \quad [\phi(c), a] \approx_\varepsilon 0 \text{ for all } c \in \mathcal{F} \text{ and } a \in \mathcal{G},$$

then we say that ϕ is an $(\mathcal{F}, \varepsilon, \mathcal{G})$ -approximately central approximate embedding.

We will usually write that ϕ is a $(\mathcal{F}, \varepsilon)$ -embedding or $(\mathcal{F}, \varepsilon, \mathcal{G})$ -embedding to mean that ϕ is an $(\mathcal{F}, \varepsilon)$ -approximate embedding or $(\mathcal{F}, \varepsilon, \mathcal{G})$ -approximately central approximate embedding, respectively.

Remark 3.1 One can make a similar definition to the above if D is not simple or nuclear (or even unital). The aim is to discuss subalgebras of $B_\omega \cap A'$, and if $D \hookrightarrow B_\omega \cap A'$ is nuclear, then one can use the Choi–Effros lifting theorem [10, Theorem 3.10] (see also [6, Theorem C.3]) to lift the embedding to a sequence of u.c.p. maps which are approximately isometric, approximately multiplicative, and approximately commute with finite subsets of A . If D is simple, the approximate isometry condition follows for free since the embedding $D \hookrightarrow B_\omega \cap A'$ must be isometric.

If we loosen the simple and nuclear assumptions on D , we can still speak of bounded linear maps $\phi : D \rightarrow B$ (no longer necessarily u.c.p.) which are approximately isometric, approximately multiplicative, approximately adjoint-preserving, and approximately commute with a finite prescribed subset of A . This will allow one to discuss general subalgebras of $B_\omega \cap A'$. As we will only be interested in strongly self-absorbing subalgebras of $B_\omega \cap A'$, which are unital, separable, simple, and nuclear [66, Section 1.6], we restrict ourselves to u.c.p. maps from a unital, simple, nuclear C^* -algebras which are approximately multiplicative and approximately commute with finite subsets of A .

Most of the work in this section can be done without assumptions of simplicity and nuclearity.

Lemma 3.2 Suppose that A, B, D are unital C^* -algebras with B separable and D simple, separable and nuclear. Suppose that $B \subseteq A$ is a unital inclusion and let $S \subseteq A$ be a separable subset. There are $(\mathcal{F}, \varepsilon, \mathcal{G})$ -approximately central approximate embeddings $D \rightarrow B$ for all $\mathcal{F} \subseteq D, \mathcal{G} \subseteq S$ and $\varepsilon > 0$ if and only if there is a unital embedding $D \hookrightarrow B_\omega \cap S'$.

Proof Let $(F_n)_{n \in \mathbb{N}}$ be an increasing sequence of finite subsets of D with dense union, and let $(G_n)_{n \in \mathbb{N}}$ be an increasing sequence of finite subsets of S with dense union. Let $\phi_n : D \rightarrow B$ be $(F_n, \frac{1}{n}, G_n)$ -approximately central approximate embeddings. Let $\pi : \ell^\infty(B) \rightarrow B_\omega$ denote the quotient map and set

$$(3.1) \quad \psi := \pi \circ ((\phi_n)_{n \in \mathbb{N}}) : D \rightarrow B_\omega,$$

which is a unital embedding such that $[\psi(d), a] = 0$ for all $d \in D$ and $a \in S$.

Conversely, suppose that $\psi : D \rightarrow B_\omega \cap S'$ is a unital embedding, $\mathcal{F} \subseteq D$, $\mathcal{G} \subseteq S$ are finite and $\varepsilon > 0$. By the Choi–Effros lifting theorem, there is a u.c.p. lift $\tilde{\psi} = (\tilde{\psi}_n)_{n \in \mathbb{N}} : D \rightarrow \ell^\infty(B)$ such that:

- $\|\tilde{\psi}_n(cd) - \tilde{\psi}_n(c)\tilde{\psi}_n(d)\| \rightarrow^{n \rightarrow \omega} 0$,
- $\|[\tilde{\psi}_n(d), a]\| \rightarrow^{n \rightarrow \omega} 0$

for all $c, d \in D$ and $a \in A$. Take n large enough and set $\phi = \psi_n$, so that ϕ will be a $(\mathcal{F}, \varepsilon, \mathcal{G})$ -approximately central approximate embedding. ■

Corollary 3.3 Let A, B, D be unital C^* -algebras with B, D separable, simple, and nuclear and $B \subseteq A$ be a unital inclusion. Suppose that there are unital embeddings $\phi : D \rightarrow B_\omega$ and $\psi : B \rightarrow A_\omega$. Then there is a unital embedding $\xi : D \hookrightarrow A_\omega$. If $S \subseteq A_\omega$ is a separable subset with $\psi(B) \subseteq A_\omega \cap S'$, then ξ can be chosen with $\xi(D) \subseteq A_\omega \cap S'$.

Proof Let $\mathcal{F} \subseteq D$ be finite and $\varepsilon > 0$. Let $L := \max\{\max_{d \in \mathcal{F}} \|d\|, 1\}$. By the above lemma, there is an $(\mathcal{F}, \frac{\varepsilon}{2L})$ -approximate embedding $\phi : D \rightarrow B$, so let $\mathcal{F}' = \phi(\mathcal{F})$. Now there is an $(\mathcal{F}', \frac{\varepsilon}{2L})$ -approximate embedding $\psi : B \rightarrow A$. An easy calculation shows that $\psi \circ \phi : D \rightarrow A$ is an approximate $(\mathcal{F}, \varepsilon)$ -embedding.

Appending the condition that $\psi : B \rightarrow A_\omega \cap S'$, then, for any finite subset $\mathcal{G} \subseteq S$, we can take $\psi : B \rightarrow A$ to be a $(\mathcal{F}', \frac{\varepsilon}{2L}, \mathcal{G})$ -approximately central approximate embedding. This gives that $\psi \circ \phi : D \rightarrow A$ is a $(\mathcal{F}, \varepsilon, \mathcal{G})$ -approximately central approximate embedding. ■

Corollary 3.4 Let D be a C^* -algebra and $B \subseteq A$ be a unital inclusion of separable C^* -algebras such that B and D are unital, separable, simple, and nuclear. Suppose that there is an embedding $\pi : A \hookrightarrow A_\omega \cap A'$ with $\pi(B) \subseteq B_\omega \cap A'$. If $D \hookrightarrow B_\omega$ unitaly, then $D \hookrightarrow B_\omega \cap A'$ unitaly.

Proof As $D \hookrightarrow B_\omega$ and $B \hookrightarrow B_\omega \cap A' \subseteq A_\omega \cap A'$, the above yields $D \hookrightarrow B_\omega \cap A'$. ■

The following is useful for discussing \mathcal{D} -stability for some inclusions of fixed point subalgebras by certain automorphisms on UHF algebras. In particular, the following will work for automorphisms on UHF algebras of product-type, as well as tensor permutations (of finite tensor powers of UHF algebras).

Corollary 3.5 Let $A = \bigotimes_{\mathbb{N}} B$ be an infinite tensor product of a unital, separable, nuclear C^* -algebra B , and let D be unital, separable, simple, and nuclear. Let $\lambda \in \text{End}(A)$ be the Bernoulli shift $\lambda(a) = 1 \otimes a$. If $\sigma \in \text{Aut}(A)$ is such that $\lambda \circ \sigma = \sigma \circ \lambda$, and $D \hookrightarrow (A^\sigma)_\omega$ unitaly, then $D \hookrightarrow (A^\sigma)_\omega \cap A'$ unitaly.

Proof Note that $\pi = (\lambda^n)$ induces an embedding $A \hookrightarrow A_\omega \cap A'$. We just need to show that $\pi(A^\sigma) \subseteq (A^\sigma)_\omega \cap A'$, which is true since $\lambda^n \circ \sigma = \sigma \circ \lambda^n$ for all n by hypothesis. The result now follows from the above. ■

We note that if we have approximately central approximate embeddings $D \rightarrow B \subseteq A$, then we can also find approximately central approximate embeddings $D \rightarrow u^*Bu \subseteq A$ for any $u \in U(A)$. In the separable setting, this just means $D \hookrightarrow B_\omega \cap A'$ implies that $D \hookrightarrow u^*B_\omega u \cap A'$ for any $u \in U(A)$.

Lemma 3.6 *Let $B \subseteq A$ be a unital inclusion of C^* -algebras, and let D be a unital, separable, simple, nuclear C^* -algebra. Let $u \in U(A)$. If there are $(\mathcal{F}, \varepsilon, \mathcal{G})$ -approximately central approximate embeddings $D \rightarrow B$ for all $\mathcal{F} \subseteq D, \mathcal{G} \subseteq A$ finite subsets and $\varepsilon > 0$, then there are $(\mathcal{F}, \varepsilon, \mathcal{G})$ -approximately central approximate embeddings $D \rightarrow u^*Bu \subseteq A$ for all $\mathcal{F}, \varepsilon, \mathcal{G}$.*

Proof Let $\mathcal{F} \subseteq D, \mathcal{G} \subseteq A$ be finite and $\varepsilon > 0$. Let $L = \max\{1, \max_{d \in \mathcal{F}} \|d\|\}$ and $\phi : D \rightarrow B$ be a $(\mathcal{F}, \frac{\varepsilon}{3L}, \mathcal{G} \cup \{u\})$ -approximately central approximate embedding. Then $\psi = \text{Ad}_u \circ \phi : D \rightarrow u^*Bu$ will be an $(\mathcal{F}, \varepsilon, \mathcal{G})$ -embedding. ■

We can also discuss existence of approximately central approximate embeddings in inductive limits (with injective connecting maps). This is an adaptation of [67, Proposition 2.2] to our setting.

Proposition 3.7 *Suppose that we have increasing sequences $(B_n)_{n \in \mathbb{N}}$ and $(A_n)_{n \in \mathbb{N}}$ of C^* -algebras such that $B_n \subseteq A_n$ are unital inclusions. If $B = \bigcup_n B_n, A = \bigcup_n A_n$, and $D = \bigcup_n D_n$, where $(D_n)_{n \in \mathbb{N}}$ is an increasing sequence of unital, separable, simple, nuclear C^* -algebras and there are $(\mathcal{F}, \varepsilon, \mathcal{G})$ -embeddings $D_n \rightarrow B_n \subseteq A_n$ whenever $n \in \mathbb{N}, \mathcal{F} \subseteq D_n, \mathcal{G} \subseteq A_n$ are finite and $\varepsilon > 0$, then there are $(\mathcal{F}, \varepsilon, \mathcal{G})$ -embeddings $D \rightarrow B \subseteq A$ for all $\mathcal{F} \subseteq D, \mathcal{G} \subseteq A$ finite and $\varepsilon > 0$.*

Proof Let $\mathcal{F} \subseteq D$ and $\mathcal{G} \subseteq A$ be finite sets and $\varepsilon > 0$. Let

$$(3.2) \quad L := \max\{1, \max_{d \in \mathcal{F}} \|d\|, \max_{a \in \mathcal{G}} \|a\|\}$$

and set $\delta := \frac{\varepsilon}{6L+5}$. Without loss of generality, assume that $\varepsilon < 1$. Label $\mathcal{F} = \{d_1, \dots, d_p\}$ and $\mathcal{G} = \{a_1, \dots, a_q\}$ and find N large enough so that there are $d'_1, \dots, d'_p \in D_N$ and $a'_1, \dots, a'_q \in A_N$ with $d'_i \approx_\delta d_i, i = 1, \dots, p$, and $a'_j \approx_\delta a_j, j = 1, \dots, q$. Let $\mathcal{F}' := \{d'_1, \dots, d'_p\}, \mathcal{G}' := \{a'_1, \dots, a'_q\}$ and let $\phi : D_N \rightarrow B_N \subseteq A_N$ be an $(\mathcal{F}', \delta, \mathcal{G}')$ -embedding. As D_N is nuclear, there are $k \in \mathbb{N}$ and u.c.p. maps $\rho : D_N \rightarrow M_k$ and $\eta : M_k \rightarrow B_N$ such that $\eta \circ \rho(d'_i) \approx_\delta \phi(d'_i)$ and $\eta \circ \rho(d'_i d'_j) \approx_\delta \phi(d'_i d'_j)$. By Arveson's extension theorem (see [6, Section 1.6]), we can extend ρ to a u.c.p. map $\tilde{\rho} : D \rightarrow M_k$ and let $\psi := \eta \circ \tilde{\rho} : D \rightarrow B_N$. As $B_N \subseteq B$, we can think of ψ as a map $\psi : D \rightarrow B$. Now for $i, j = 1, \dots, p$, we have

$$(3.3) \quad \begin{aligned} \psi(d_i d_j) &\approx_{(2L+1)\delta} \psi(d'_i d'_j) \\ &= \eta \circ \rho(d'_i d'_j) \\ &\approx_\delta \phi(d'_i d'_j) \\ &\approx_\delta \phi(d'_i) \phi(d'_j) \\ &\approx_{2L\delta} \eta \circ \rho(d'_i) \eta \circ \rho(d'_j) \\ &= \psi(d'_i) \psi(d'_j) \\ &\approx_{(2L+1)\delta} \psi(d_i) \psi(d_j). \end{aligned}$$

Thus $\psi(d_i d_j) \approx_{(4+6L)\delta} \psi(d_i) \psi(d_j)$, and as $(4 + 6L)\delta \leq (6L + 5)\delta = \varepsilon$, this implies that $\psi(d_i d_j) \approx_\varepsilon \psi(d_i) \psi(d_j)$. For approximate commutation with \mathcal{G} , we make use of the following two approximations: for a, a', a'', b, b' elements in a C^* -algebra,

$$(3.4) \quad \begin{aligned} \|[a, b]\| &\leq (\|a\| + \|a'\|)\|b - b'\| + (\|b\| + \|b'\|)\|a - a'\| + \|[a', b']\|, \\ \|[a', b']\| &\leq 2\|b'\|\|a' - a''\| + \|[a'', b']\|. \end{aligned}$$

Note that for $a = \psi(d_i)$, $a' = \psi(d'_i)$, $a'' = \phi(d'_i)$, $b = a_j$, $b' = a'_j$, we have that $\|a\|, \|b\| \leq L + 1$ and $\|a'\|, \|a''\|, \|b'\| \leq L$. Therefore, from the above two inequalities, we get

$$(3.5) \quad \begin{aligned} \|\psi(d_i), a_j\| &\leq 2L\|\psi(d_i) - \psi(d'_i)\| + 2(L + 1)\|a_j - a'_j\| + \|\psi(d'_i), a_j\|; \\ \|\psi(d'_i), a'_j\| &\leq 2(L + 1)\|\psi(d'_i) - \phi(d'_i)\| + \|\phi(d'_i), a'_j\| \end{aligned}$$

whenever $i = 1, \dots, p$, $j = 1, \dots, q$. Using these approximations, we have

$$(3.6) \quad \begin{aligned} \|\psi(d_i), a_j\| &\leq 2L\|\psi(d_i) - \psi(d'_i)\| + 2(L + 1)\|a_j - a'_j\| + \|\psi(d'_i), a_j\| \\ &< (4L + 2)\delta + \|\psi(d'_i), a_j\| \\ &\leq (4L + 2)\delta + 2(L + 1)\|\psi(d'_i) - \phi(d'_i)\| + \|\phi(d'_i), a'_j\| \\ &< (4L + 2)\delta + 2(L + 1)\delta + \delta \\ &= (6L + 5)\delta = \varepsilon. \end{aligned}$$

■

The following will be useful to show that there are many \mathcal{D} -stable embeddings.

Lemma 3.8 *Let $\phi : B_0 \simeq B_1$ and $\psi : A_0 \simeq A_1$ be $*$ -isomorphisms between unital C^* -algebras, and let D be a unital, simple, nuclear C^* -algebra. Suppose that there is a unital $*$ -homomorphism $\eta : B_1 \hookrightarrow A_1$ such that there are $(\mathcal{F}, \varepsilon, \mathcal{G})$ -embeddings $D \rightarrow \eta(B_1) \subseteq A_1$ for all finite subsets $\mathcal{F} \subseteq D$, $\mathcal{G} \subseteq A_1$ and $\varepsilon > 0$. Let $\sigma = \psi^{-1} \circ \eta \circ \phi : B_0 \rightarrow A_0$. Then there are $(\mathcal{F}, \varepsilon, \mathcal{G})$ -embeddings $D \rightarrow \sigma(B_0) \subseteq A_0$ for all $\mathcal{F} \subseteq D$, $\mathcal{G} \subseteq A_0$ finite and $\varepsilon > 0$.*

Proof The diagram

$$(3.7) \quad \begin{array}{ccc} A_0 & \xrightarrow{\psi} & A_1 \\ \sigma \uparrow & & \uparrow \eta \\ B_0 & \xrightarrow{\phi} & B_1 \end{array}$$

commutes and so if $\mathcal{F} \subseteq D$, $\mathcal{G} \subseteq A_0$ are finite, $\varepsilon > 0$ and $\xi : D \rightarrow \eta(B_1) \subseteq A_1$ is an $(\mathcal{F}, \varepsilon, \psi(\mathcal{G}))$ -embedding, then $\psi^{-1} \circ \xi : D \rightarrow \psi^{-1}(\eta(B_1)) \subseteq \psi^{-1}(A_1) = A_0$ is an $(\mathcal{F}, \varepsilon, \mathcal{G})$ -embedding. Moreover, from

$$(3.8) \quad \psi^{-1}(\eta(B_1)) = \psi^{-1}(\eta(\phi(B_0))) = \sigma(B_0),$$

it is clear that $\psi^{-1} \circ \xi$ is an $(\mathcal{F}, \varepsilon, \mathcal{G})$ -embedding $D \rightarrow \sigma(B_0) \subseteq A_0$. ■

4 Relative intertwining and \mathcal{D} -stable embeddings

4.1 Relative intertwining

It is well known that a strongly self-absorbing C^* -algebra \mathcal{D} embeds unitaly into the central sequence algebra $(\mathcal{M}(A))_\omega \cap A'$ of a separable C^* -algebra A if and only if $A \simeq A \otimes \mathcal{D}$, where $\mathcal{M}(A)$ is the multiplier algebra of A (see, for example, [49, Theorem 7.2.2(i)]). We alter the proof to keep track of a subalgebra in order to show that for a unital inclusion $B \subseteq A$ of separable C^* -algebras, $\mathcal{D} \hookrightarrow B_\omega \cap A'$ unitaly if and only if there is an isomorphism $\Phi : A \rightarrow A \otimes \mathcal{D}$ which is approximately unitarily equivalent to the first factor embedding and satisfies $\Phi(B) = B \otimes \mathcal{D}$. This was initially done for (irreducible) inclusions of II_1 factors in [2] and commented on in [31] for \mathcal{D} being $M_{n^\infty}, \mathcal{O}_2, \mathcal{O}_\infty$. The proof we alter is Elliott's intertwining argument, which can be found as a combination of Propositions 2.3.5 and 7.2.1 and Theorem 7.2.2 of [49].

Proposition 4.1 (Relative intertwining) *Let A, B, C be unital, separable C^* -algebras, and let $\phi : A \hookrightarrow C, \theta : B \rightarrow A, \psi : B \rightarrow C$ be unital $*$ -homomorphisms such that $\phi \circ \theta(B) \subseteq \psi(B)$. Suppose there is a sequence $(u_n)_{n \in \mathbb{N}}$ of unitaries in $\psi(B)_\omega \cap \phi(A)'$ such that:*

- $\text{dist}(v_n^* c v_n, \phi(A)_\omega) \rightarrow 0$ for all $c \in C$;
- $\text{dist}(v_n^* \psi(b) v_n, \phi \circ \theta(B)_\omega) \rightarrow 0$ for all $b \in B$.

Then ϕ is approximately unitarily equivalent to an isomorphism $\Phi : A \simeq C$ such that $\Phi \circ \theta(B) = \psi(B)$.

Proof Apply the below proposition with $B_m := B, \theta_m := \theta, \psi_m := \psi$ for all $m \in \mathbb{N}$. ■

Proposition 4.2 (Countable relative intertwining) *Let A, B_m, C be unital, separable C^* -algebras, $m \in \mathbb{N}$, and $\phi : A \hookrightarrow C, \theta_m : B_m \rightarrow A, \psi_m : B_m \rightarrow C$ be such that $\phi \circ \theta_m(B_m) \subseteq \psi_m(B_m)$ and $\psi_1(B_1) \subseteq \psi_m(B_m)$. Suppose there is a sequence $(v_n)_{n \in \mathbb{N}} \subseteq \psi_1(B_1)_\omega \cap \phi(A)'$ of unitaries such that:*

- $\text{dist}(v_n^* c v_n, \phi(A)_\omega) \rightarrow 0$ for all $c \in C$;
- $\text{dist}(v_n^* \psi_m(b) v_n, \phi \circ \theta_m(B_m)_\omega) \rightarrow 0$ for all $b \in B_m$.

Then ϕ is approximately unitarily equivalent to an isomorphism $\Phi : A \simeq C$ such that $\Phi \circ \theta_m(B_m) = \psi_m(B_m)$ for all $m \in \mathbb{N}$.

Proof We show that if there are unitaries $(v_n)_{n \in \mathbb{N}} \subseteq \psi_1(B_1)$ satisfying:

- $[v_n, \phi(a)] \rightarrow 0$ for all $a \in A$;
- $\text{dist}(v_n^* c v_n, \phi(A)) \rightarrow 0$ for all $c \in C$;
- $\text{dist}(v_n^* \psi_m(b) v_n, \phi \circ \theta_m(B_m)) \rightarrow 0$ for all $b \in B_m$,

then the conclusion holds. Such unitaries can be found using Kirchberg's ε -test (Lemma 2.1).

Let $(a_n)_{n \in \mathbb{N}}, (b_n^{(m)})_{n \in \mathbb{N}}, (c_n)_{n \in \mathbb{N}}$ be dense sequences of A, B_m, C , respectively, $m \in \mathbb{N}$. We can inductively choose v_n , forming a subsequence $(v_n)_{n \in \mathbb{N}}$ of the unitaries above (after re-indexing, we are still calling them v_n), such that there are $a_{j_n} \in A, b_{j_n}^{(m)} \in B_m$ with:

- $v_n^* \dots v_1^* c_j v_1 \dots v_n \approx_{\frac{1}{n}} \phi(a_{jn})$;
- $v_n^* \dots v_1^* \psi(b_j^{(m)}) v_1 \dots v_n \approx_{\frac{1}{n}} \phi \circ \theta_m(b_{jn}^{(m)})$;
- $[v_n, \phi(a_j)] \approx_{\frac{1}{2^n}} 0$;
- $[v_n, \phi(a_{jl})] \approx_{\frac{1}{2^n}} 0$;
- $[v_n, \phi \circ \theta_m(b_j^{(m)})] \approx_{\frac{1}{2^n}} 0$;
- $[v_n, \phi \circ \theta_m(b_{jl}^{(m)})] \approx_{\frac{1}{2^n}} 0$,

where $j, m = 1, \dots, n$ and $l = 1, \dots, n-1$. Define, for $a \in \{a_n \mid n \in \mathbb{N}\}$,

$$(4.1) \quad \Phi(a) = \lim_n v_1 \dots v_n \phi(a) v_n^* \dots v_1^*$$

which extends to a $*$ -isomorphism $\Phi : A \simeq C$, as in [49, Proposition 2.3.5]. The proof also yields the following useful approximation:

$$(4.2) \quad \Phi \circ \theta_m(b_{jn}^{(m)}) \approx_{\frac{1}{2^n}} v_1 \dots v_n \phi \circ \theta_m(b_{jn}^{(m)}) v_n^* \dots v_1^*$$

for appropriate $n \geq j, m$.

We now need to check that $\Phi \circ \theta_m(B_m) = \psi_m(B_m)$. Approximate

$$(4.3) \quad \psi_m(b_j^{(m)}) \approx_{\frac{1}{n}} v_1 \dots v_n \phi \circ \theta_m(b_{jn}^{(m)}) v_n^* \dots v_1^* \approx_{\frac{1}{2^n}} \Phi \circ \theta_m(b_{jn}^{(m)}).$$

As $n \in \mathbb{N}$ can be made arbitrarily large, this yields $\psi_m(B_m) \subseteq \overline{\Phi \circ \theta_m(B_m)} = \Phi \circ \theta_m(B_m)$. On the other hand, for any $\varepsilon > 0$ and $b \in B_m$, we can find n such that

$$(4.4) \quad \Phi \circ \theta_m(b) \approx_\varepsilon v_1 \dots v_n \phi \circ \theta_m(b) v_n^* \dots v_1^* \in \psi_m(B_m)$$

since $v_i \in \psi_1(B_1) \subseteq \psi_m(B_m)$ and $\phi \circ \theta_m(B_m) \subseteq \psi_m(B_m)$. Hence $\Phi \circ \theta_m(B_m) \subseteq \overline{\psi_m(B_m)} = \psi_m(B_m)$. ■

4.2 \mathcal{D} -stable embeddings

Definition 4.1 Let $\iota : B \hookrightarrow A$ be an embedding and \mathcal{D} be strongly self-absorbing. We say that ι is \mathcal{D} -stable (or \mathcal{D} -absorbing) if there exists an isomorphism $\Phi : A \simeq A \otimes \mathcal{D}$ such that $\Phi \circ \iota(B) = \iota(B) \otimes \mathcal{D}$.

We will mostly have interest in the case where ι corresponds to the inclusion map and $B \subseteq A$ is a subalgebra. In this form, we will say $B \subseteq A$ is \mathcal{D} -stable (or \mathcal{D} -absorbing). Clearly, ι being \mathcal{D} -stable is the same as $\iota(B) \subseteq A$ being \mathcal{D} -stable. We note that we can define the above for any $*$ -homomorphism. Namely, a $*$ -homomorphism $\phi : B \rightarrow A$ is \mathcal{D} -stable if $\phi(B) \subseteq A$ is.

Lemma 4.3 If $\iota : B \hookrightarrow A$ is an embedding, then $\iota \otimes id_{\mathcal{D}} : B \otimes \mathcal{D} \hookrightarrow A \otimes \mathcal{D}$ is \mathcal{D} -stable.

Proof Let $\phi : D \simeq D \otimes \mathcal{D}$ be an isomorphism. Then

$$(4.5) \quad \Phi := id_A \otimes \phi : A \otimes \mathcal{D} \rightarrow A \otimes \mathcal{D} \otimes \mathcal{D}$$

is an isomorphism with

$$(4.6) \quad \Phi(\iota \otimes id_{\mathcal{D}}(B \otimes \mathcal{D})) = (\iota \otimes id_{\mathcal{D}}(B \otimes \mathcal{D})) \otimes \mathcal{D}. \quad \blacksquare$$

We note that this is a strengthening of the notion of \mathcal{D} -stability for C^* -algebras because if $\iota := \text{id}_A : A \rightarrow A$, then ι is \mathcal{D} -stable if and only if A is \mathcal{D} -stable. This condition is different from the notion of \mathcal{O}_2 or \mathcal{O}_∞ -absorbing morphisms discussed in [4, 21, 22] – they require sequences from a larger algebra to commute with a smaller algebra, while we require sequences from a smaller algebra to commute with the larger algebra. In the former, neither of the algebras are required to be \mathcal{D} -stable, while the latter necessitates both to be \mathcal{D} -stable.

The following adapts [49, Theorem 7.2.2].

Theorem 4.4 *Suppose that $B \subseteq A$ is a unital inclusion of separable C^* -algebras. If \mathcal{D} is strongly self-absorbing, then $B \subseteq A$ is \mathcal{D} -stable if and only if there is a unital inclusion $\mathcal{D} \hookrightarrow B_\omega \cap A'$.*

Proof Let $\phi : A \rightarrow A \otimes \mathcal{D}$ be the first factor embedding $\phi(a) := a \otimes 1_{\mathcal{D}}$. First, suppose that $\xi : \mathcal{D} \hookrightarrow B_\omega \cap A' \simeq (B \otimes 1_{\mathcal{D}})_\omega \cap (A \otimes 1_{\mathcal{D}})'$ is an embedding (so that $\phi(a)\xi(d) \in \phi(A)_\omega$ and $\phi(b)\xi(d) \in \phi(B)_\omega$). Let $\eta : \mathcal{D} \hookrightarrow (B \otimes \mathcal{D})_\omega \cap (A \otimes 1_{\mathcal{D}})'$ be given by $\eta(d) := (1 \otimes d)_n$ and notice that ξ, η have commuting ranges. As all endomorphisms of \mathcal{D} are approximately unitarily equivalent by [66, Corollary 1.12], let $(v_n)_{n \in \mathbb{N}} \subseteq C^*(\xi(\mathcal{D}), \eta(\mathcal{D})) \simeq \mathcal{D} \otimes \mathcal{D}$ be such that $v_n^* \eta(d) v_n \rightarrow \xi(d)$ for $d \in \mathcal{D}$. For $b \in B$ and $d \in \mathcal{D}$, we have

$$\begin{aligned} v_n^*(b \otimes d)v_n &= v_n^*(b \otimes 1_{\mathcal{D}})(1_A \otimes d)v_n \\ (4.7) \qquad &= v_n^*\phi(b)\eta(d)v_n \\ &= \phi(b)v_n^*\eta(d)v_n \\ &\rightarrow \phi(b)\xi(d) \in \phi(B)_\omega. \end{aligned}$$

Moreover, the same argument shows that, for $a \in A$, we have

$$(4.8) \qquad v_n^*(a \otimes d)v_n \rightarrow \phi(a)\xi(d) \in \phi(A)_\omega.$$

Now $(v_n)_{n \in \mathbb{N}}$ satisfy the hypothesis of Proposition 4.1 with $C := A \otimes \mathcal{D}$, ϕ being the first factor embedding, $\theta : B \rightarrow A$ being the inclusion and $\psi : B \simeq B \otimes \mathcal{D} \subseteq A \otimes \mathcal{D} = C$ (where this isomorphism exists since if $\mathcal{D} \hookrightarrow B_\omega \cap A'$, then clearly $\mathcal{D} \hookrightarrow B_\omega \cap B'$). From this, we see that ϕ is approximately unitarily equivalent to an isomorphism $\Phi : A \simeq A \otimes \mathcal{D}$ such that $\Phi(B) = B \otimes \mathcal{D}$.

Conversely, if $B \subseteq A$ is \mathcal{D} -stable, let $\Phi : A \simeq A \otimes \mathcal{D}$ be an isomorphism such that $\Phi(B) = B \otimes \mathcal{D}$. By [66, Proposition 1.10(iv)], we can identify $\mathcal{D} \simeq \mathcal{D}^{\otimes \infty}$ and take $\xi : \mathcal{D} \hookrightarrow B_\omega \cap A'$ to be given by

$$(4.9) \qquad \xi(d) = (\Phi^{-1}(1_A \otimes 1_{\mathcal{D}}^{\otimes n-1} \otimes d \otimes 1_{\mathcal{D}}^{\otimes \infty}))_n.$$

■

Corollary 4.5 *Let $\iota : B \hookrightarrow A$ be a unital embedding between separable C^* -algebras. If \mathcal{D} is strongly self-absorbing and ι is \mathcal{D} -stable, then for every intermediate unital C^* -algebra C with $\iota(B) \subseteq C \subseteq A$, we have that $\iota(B) \subseteq C$ and $C \subseteq A$ are \mathcal{D} -stable. In particular, $C \simeq C \otimes \mathcal{D}$ for all such C .*

Proof We have

$$(4.10) \qquad \mathcal{D} \hookrightarrow B_\omega \cap A' \subseteq B_\omega \cap C'$$

and

$$(4.11) \quad \mathcal{D} \hookrightarrow B_\omega \cap A' \subseteq C_\omega \cap A'.$$

Now apply Theorem 4.4. ■

It is not, however, the case that any isomorphism $\Phi : A \simeq A \otimes \mathcal{D}$ with $\Phi(B) = B \otimes \mathcal{D}$ maps C to $C \otimes \mathcal{D}$.

Example 4.6 Let \mathcal{D} be strongly self-absorbing and consider

$$(4.12) \quad \begin{aligned} A &:= \mathcal{D} \otimes \mathcal{D} \otimes \mathcal{D}, \\ C_1 &:= \mathcal{D} \otimes 1_{\mathcal{D}} \otimes \mathcal{D}, \\ C_2 &:= 1_{\mathcal{D}} \otimes \mathcal{D} \otimes \mathcal{D}, \\ B &:= 1_{\mathcal{D}} \otimes 1_{\mathcal{D}} \otimes \mathcal{D}. \end{aligned}$$

If $f : \mathcal{D} \otimes \mathcal{D} \rightarrow \mathcal{D} \otimes \mathcal{D}$ is the tensor flip and $\phi : \mathcal{D} \simeq \mathcal{D} \otimes \mathcal{D}$ is an isomorphism, let

$$(4.13) \quad \Phi := f \otimes \phi : A \simeq A \otimes \mathcal{D}$$

which satisfies $\Phi(B) = B \otimes \mathcal{D}$ (in particular, $B \subseteq A$ is \mathcal{D} -stable). However,

$$(4.14) \quad \Phi(C_1) = C_2 \otimes \mathcal{D} \text{ and } \Phi(C_2) = C_1 \otimes \mathcal{D}.$$

In fact, the above example can be generalized to show that for any \mathcal{D} -stable inclusion $B \subseteq A$, there are an isomorphism $\Phi : A \simeq A \otimes \mathcal{D}$ such that $\Phi(B) = B \otimes \mathcal{D}$ and an intermediate algebra $B \subseteq C \subseteq A$ with $\Phi(C) \neq C \otimes \mathcal{D}$ (obviously, we may still have that $\Phi(C) \simeq C \otimes \mathcal{D}$, but equality may not happen).

Corollary 4.7 Let $B \subseteq A$ be a \mathcal{D} -stable inclusion. There exist a C^* -algebra C with $B \subseteq C \subseteq A$ and an isomorphism $\Phi : A \simeq A \otimes \mathcal{D}$ such that $\Phi(B) = B \otimes \mathcal{D}$ but $\Phi(C) \neq C \otimes \mathcal{D}$.

Proof We first claim that if $B \subseteq A$ is \mathcal{D} -stable, then we can identify $B \subseteq A$ with $B \otimes 1_{\mathcal{D}} \subseteq A \otimes \mathcal{D}$. If $\Psi : A \simeq A \otimes \mathcal{D}$ is such that $\Psi(B) = B \otimes \mathcal{D}$ and $f : \mathcal{D} \otimes \mathcal{D} \simeq \mathcal{D} \otimes \mathcal{D}$ is the tensor flip, we have

$$(4.15) \quad \Xi := (\text{id}_A \otimes f) \circ (\Psi \otimes \text{id}_{\mathcal{D}}) : A \otimes \mathcal{D} \simeq A \otimes \mathcal{D} \otimes \mathcal{D}$$

is such that $\Xi(B \otimes 1_{\mathcal{D}}) = B \otimes 1_{\mathcal{D}} \otimes \mathcal{D}$. This proves the claim.

Now, by applying the claim twice, we can identify $B \subseteq A$ with the inclusion

$$(4.16) \quad B \otimes 1_{\mathcal{D}} \otimes 1_{\mathcal{D}} \otimes \mathcal{D} \subseteq A \otimes \mathcal{D} \otimes \mathcal{D} \otimes \mathcal{D}.$$

If $\phi : \mathcal{D} \simeq \mathcal{D} \otimes \mathcal{D}$ is any $*$ -isomorphism,

$$(4.17) \quad \Phi := \text{id}_A \otimes f \otimes \phi : A \otimes \mathcal{D} \otimes \mathcal{D} \otimes \mathcal{D} \simeq A \otimes \mathcal{D} \otimes \mathcal{D} \otimes \mathcal{D}$$

is such that

$$(4.18) \quad \Phi(B \otimes 1_{\mathcal{D}} \otimes 1_{\mathcal{D}} \otimes \mathcal{D}) = B \otimes 1_{\mathcal{D}} \otimes 1_{\mathcal{D}} \otimes \mathcal{D} \otimes \mathcal{D}.$$

Taking C_1 and C_2 as in Example 4.6, we have that

$$(4.19) \quad \Phi(B \otimes C_1) = B \otimes C_2 \otimes \mathcal{D} \text{ and } \Phi(B \otimes C_2) = B \otimes C_1 \otimes \mathcal{D}. \quad \blacksquare$$

However, we can always realize \mathcal{D} -stability for countably many intermediate C^* -algebras at once using *some* isomorphism $A \simeq A \otimes \mathcal{D}$.

Theorem 4.8 *Suppose that $B_1 \subseteq B_m \subseteq A$ are unital inclusions of separable C^* -algebras (note that we are not asking for (B_m) to form a chain). If \mathcal{D} is strongly self-absorbing and $\mathcal{D} \hookrightarrow (B_1)_\omega \cap A'$ unitaly, there exists an isomorphism $\Phi : A \simeq A \otimes \mathcal{D}$ such that $\Phi(B_m) = B_m \otimes \mathcal{D}$ for all $m \in \mathbb{N}$.*

Proof This is essentially the same proof as Theorem 4.4, except we use the countable relative intertwining (Proposition 4.2) in place of Proposition 4.1. Let ξ, η be as before, and let $(v_n)_{n \in \mathbb{N}} \subseteq C^*(\xi(\mathcal{D}), \eta(\mathcal{D})) \simeq \mathcal{D} \otimes \mathcal{D}$ be such that $v_n^* \eta(d) v_n \rightarrow \xi(d)$ for $d \in \mathcal{D}$.

- If $a \in A, d \in \mathcal{D}, v_n^*(a \otimes d)v_n \rightarrow \phi(a)\xi(d) \in \phi(A)_\omega$;
- if $b \in B_m, v_n^*(b \otimes d)v_n \rightarrow \phi(b)\xi(d) \in \phi(B_m)_\omega$.

Now with $\phi : A \rightarrow A \otimes \mathcal{D}$, the first factor embedding, $\theta_m : B_m \rightarrow A$ the inclusion maps, and $\psi_m : B_m \simeq B_m \otimes \mathcal{D}$ (these exist since $\mathcal{D} \hookrightarrow (B_1)_\omega \cap A'$ implies that $\mathcal{D} \hookrightarrow (B_m)_\omega \cap B'_m$), our unitaries satisfy the hypothesis of Proposition 4.2 and therefore ϕ is approximately unitarily equivalent to a $*$ -isomorphism $\Phi : A \simeq A \otimes \mathcal{D}$ such that $\Phi(B_m) = B_m \otimes \mathcal{D}$ for all m . ■

The above works since norm ultrapowers have the property that unitaries lift to sequences of unitaries.² Tracial ultrapowers of II_1 von Neumann algebras also have this property.³ Consequently, if we work with the 2-norm $\|x\|_2 = \tau(x^*x)^{\frac{1}{2}}$, where τ is the unique trace on a II_1 factor, all of the above arguments with the C^* -norm replaced by $\|\cdot\|_2$ will allow us to recover Bisch's result [2, Theorem 3.1], provided we have the appropriate separability conditions.

Theorem 4.9 *Let $\mathcal{N} \subseteq \mathcal{M}$ be an inclusion of II_1 factors with separable preduals. Then $\mathcal{R} \hookrightarrow \mathcal{N}^\omega \cap \mathcal{M}'$ if and only if there exists an isomorphism $\Phi : \mathcal{M} \rightarrow \mathcal{M} \overline{\otimes} \mathcal{R}$ such that $\Phi(\mathcal{N}) = \mathcal{N} \overline{\otimes} \mathcal{R}$.*

4.3 Existence of \mathcal{D} -stable embeddings

We move to discuss the existence of \mathcal{D} -stable embeddings. First, we show that each unital embedding of unital, separable \mathcal{D} -stable C^* -algebras is approximately unitarily equivalent to a \mathcal{D} -stable embedding. From this, it will follow that there are many \mathcal{D} -stable embeddings.

Lemma 4.10 *Let \mathcal{D} be strongly self-absorbing. If $\iota : B \hookrightarrow A$ is a unital, \mathcal{D} -stable inclusion of separable C^* -algebras and $u \in U(A)$, then $Ad_u \circ \iota : B \hookrightarrow A$ is \mathcal{D} -stable.*

Proof Apply Lemma 3.6. ■

²If $u = (u_n)_{n \in \mathbb{N}} \in A_\omega$ is unitary, then $\{n \in \mathbb{N} \mid \|u_n^* u_n - 1\|, \|u_n u_n^* - 1\| < 1\} \in \omega$. If n is in the set, replace u_n with the unitary part of its polar decomposition, and replace u_n with 1 otherwise.

³The tracial ultrapower of a II_1 von Neumann algebra is again a II_1 von Neumann algebra. Therefore, if $u \in \mathcal{M}^\omega$ is unitary, it is of the form e^{ia} for some $a = a^* \in \mathcal{M}^\omega$. Lift a to a sequence $(a_n)_{n \in \mathbb{N}}$ of self-adjoints in \mathcal{M} and note that $u = (e^{ia_n})$, so that u has a unitary lift.

Proposition 4.11 *Let \mathcal{D} be strongly self-absorbing, A, B be unital separable \mathcal{D} -stable C^* -algebras, and let $\iota : B \hookrightarrow A$ be a unital embedding. Then ι is approximately unitarily equivalent to a unital \mathcal{D} -stable embedding $B \hookrightarrow A$.*

Proof As A, B are \mathcal{D} -stable, there are isomorphisms

$$(4.20) \quad \phi : B \simeq B \otimes \mathcal{D} \text{ and } \psi : A \simeq A \otimes \mathcal{D},$$

which are approximately unitarily equivalent to the first factor embeddings $b \mapsto b \otimes 1_{\mathcal{D}}, b \in B$ and $a \mapsto a \otimes 1_{\mathcal{D}}, a \in A$, respectively. As $\iota \otimes \text{id}_{\mathcal{D}} : B \otimes \mathcal{D} \hookrightarrow A \otimes \mathcal{D}$ is \mathcal{D} -stable by Lemma 4.3,

$$(4.21) \quad \sigma := \psi^{-1} \circ (\iota \otimes \text{id}_{\mathcal{D}}) \circ \phi : B \hookrightarrow A$$

is \mathcal{D} -stable by Lemma 3.8. Now we show that σ is approximately unitarily equivalent to ι . Let $\mathcal{F} \subseteq B$ be finite and $\varepsilon > 0$. Let $u \in U(B \otimes \mathcal{D})$ be such that $u^*(b \otimes 1_{\mathcal{D}})u \approx_{\frac{\varepsilon}{2}} \phi(b)$ for $b \in \mathcal{F}$ and $v \in U(A \otimes \mathcal{D})$ be such that $v^*(\iota(b) \otimes 1_{\mathcal{D}})v \approx_{\frac{\varepsilon}{2}} \psi \circ \iota(b)$ for $b \in \mathcal{F}$. Set $w = \psi^{-1}(\iota \otimes \text{id}_{\mathcal{D}}(u))^* \psi^{-1}(v) \in U(A)$. Then for $b \in \mathcal{F}$,

$$(4.22) \quad \begin{aligned} w^* \sigma(b) w &= \psi^{-1}(v)^* \psi^{-1}(\iota \otimes \text{id}_{\mathcal{D}}(u \phi(b) u^*)) \psi^{-1}(v) \\ &\approx_{\frac{\varepsilon}{2}} \psi^{-1}(v)^* \psi^{-1}(\iota \otimes \text{id}_{\mathcal{D}}(b \otimes 1_{\mathcal{D}})) \psi^{-1}(v) \\ &= \psi^{-1}(v)^* \psi^{-1}(\iota(b) \otimes 1_{\mathcal{D}}) \psi^{-1}(v) \\ &\approx_{\frac{\varepsilon}{2}} \psi^{-1}(\psi(\iota(b))) \\ &= \iota(b). \end{aligned}$$

Corollary 4.12 *Let \mathcal{D} be strongly self-absorbing. The set of unital \mathcal{D} -stable embeddings $B \hookrightarrow A$ of unital, separable, \mathcal{D} -stable C^* -algebras is point-norm dense in the set of unital embeddings $B \hookrightarrow A$.*

Proof Every embedding is approximately unitarily equivalent to a \mathcal{D} -stable embedding. As \mathcal{D} -stability of an embedding is preserved if one composes with Ad_u , it follows that every embedding is the point-norm limit of \mathcal{D} -stable embeddings. ■

Remark 4.13 We note that it is not actually necessary that ι is an embedding. If $\pi : B \rightarrow A$ is any unital $*$ -homomorphism between unital, separable, \mathcal{D} -stable C^* -algebras, then π is approximately unitarily equivalent to a $*$ -homomorphism $\pi' : B \rightarrow A$ such that $\pi'(B) \subseteq A$ is \mathcal{D} -stable. Consequently, the set of unital $*$ -homomorphisms $\pi : B \rightarrow A$ with $\pi(B) \subseteq A$ being \mathcal{D} -stable is in fact dense in the set of unital $*$ -homomorphisms $B \rightarrow A$.

Later on, there will be some examples of non- \mathcal{D} -stable embeddings between \mathcal{D} -stable C^* -algebras. Consequently, despite the fact \mathcal{D} -stable embeddings are point-norm dense, the set of unital \mathcal{D} -stable embeddings need not coincide with the set of all unital embeddings $B \hookrightarrow A$. Another clear consequence is that despite \mathcal{D} -stability of an embedding being closed under conjugation by a unitary, it is not true that it is preserved under approximate unitary equivalence (in fact, the examples in question show that \mathcal{D} -stability is not even preserved under asymptotic unitary equivalence). We finish with a corollary about embeddings into the Cuntz algebra \mathcal{O}_2 [15].

Corollary 4.14 *Let B be a unital, separable, exact \mathcal{D} -stable C^* -algebra, where \mathcal{D} is strongly self-absorbing. Then there is a \mathcal{D} -stable embedding $B \hookrightarrow \mathcal{O}_2$.*

Proof As \mathcal{D} is unital, simple, separable, and nuclear by [66, Section 1.6], $\mathcal{O}_2 \simeq \mathcal{O}_2 \otimes \mathcal{D}$ and $B \hookrightarrow \mathcal{O}_2$ unittally by Theorems 3.7 and 2.8 of [36], respectively. The above results then yield a \mathcal{D} -stable embedding $B \hookrightarrow \mathcal{O}_2$. ■

We include this last result about the classification of morphisms via functors.

Theorem 4.15 *Let \mathcal{D} be strongly self-absorbing, and let F be a functor from a class of unital, separable, \mathcal{D} -stable C^* -algebras satisfying the following.*

- (E) *If there exists a morphism $\Phi : F(B) \rightarrow F(A)$, then there exists a unital $*$ -homomorphism $\phi : B \rightarrow A$ such that $F(\phi) = \Phi$.*
- (U) *If $\phi, \psi : B \rightarrow A$ are unital $*$ -homomorphisms which are approximately unitarily equivalent, then*

$$(4.23) \quad F(\phi) = F(\psi).$$

Then whenever there is a morphism $\Phi : F(B) \rightarrow F(A)$, there exists a unital $$ -homomorphism $\phi : B \rightarrow A$ such that $F(\phi) = \Phi$ and $\phi(B) \subseteq A$ is \mathcal{D} -stable. Moreover, ϕ is unique up to approximate unitary equivalence.*

Proof By the existence (E), there exists a $*$ -homomorphism $\phi : B \rightarrow A$. Now by Proposition 4.11 (Remark 4.13 allows us to work with general $*$ -homomorphisms), there exists a $*$ -homomorphism $\phi' : B \rightarrow A$ which is approximately unitarily equivalent to ϕ and $\phi'(B) \subseteq A$ is \mathcal{D} -stable. Uniqueness (U) gives that this is unique up to approximate unitary equivalence. ■

4.4 Permanence properties

We now discuss some permanence properties.

Lemma 4.16 *Let \mathcal{D} be strongly self-absorbing. Suppose that $\iota_j : B_j \hookrightarrow A_j$, $j = 1, 2$ are \mathcal{D} -stable inclusions. Then $\iota_1 \oplus \iota_2 : B_1 \oplus B_2 \hookrightarrow A_1 \oplus A_2$ is \mathcal{D} -stable.*

Proof Let $\Phi_j : A_j \simeq A_j \otimes \mathcal{D}$ be isomorphisms such that $\Phi_j \circ \iota_j(B_j) = \iota_j(B_j) \otimes \mathcal{D}$ and consider

$$(4.24) \quad \Phi : A_1 \oplus A_2 \simeq (A_1 \oplus A_2) \otimes \mathcal{D}$$

given by the composition

$$(4.25) \quad A_1 \oplus A_2 \xrightarrow{\Phi_1 \oplus \Phi_2} (A_1 \otimes \mathcal{D}) \oplus (A_2 \otimes \mathcal{D}) \xrightarrow{\simeq} (A_1 \oplus A_2) \otimes \mathcal{D},$$

where the last isomorphism follows from (finite) distributivity of the min-tensor. Then we see that

$$(4.26) \quad \Phi(\iota_1(B_1) \oplus \iota_2(B_2)) = (\iota_1(B_1) \oplus \iota_2(B_2)) \otimes \mathcal{D}. \quad \blacksquare$$

Lemma 4.17 *Let \mathcal{D} be strongly self-absorbing. Suppose that $\iota_j : B_j \hookrightarrow A_j$, $j = 1, 2$ are inclusions and that at least one of ι_1 or ι_2 is \mathcal{D} -stable. Then $\iota_1 \otimes \iota_2 : B_1 \otimes B_2 \hookrightarrow A_1 \otimes A_2$ is \mathcal{D} -stable.*

Proof We prove this if ι_2 is \mathcal{D} -stable, and a symmetric argument will yield the result if ι_1 is. Let $\Phi_2 : A_2 \simeq A_2 \otimes \mathcal{D}$ be such that $\Phi_2 \circ \iota_2(B_2) = \iota(B_2) \otimes \mathcal{D}$. Taking

$$(4.27) \quad \Phi := \text{id}_{A_1} \otimes \Phi_2 : A_1 \otimes A_2 \simeq A_1 \otimes A_2 \otimes \mathcal{D},$$

we have that

$$(4.28) \quad \Phi(\iota_1(B_1) \otimes \iota_2(B_2)) = \iota_1(B_1) \otimes \iota_2(B_2) \otimes \mathcal{D}. \quad \blacksquare$$

Proposition 4.18 *Let \mathcal{D} be strongly self-absorbing. Suppose that we have increasing sequences of unital separable C^* -algebras $(B_n)_{n \in \mathbb{N}}$ and $(A_n)_{n \in \mathbb{N}}$ such that $B_n \subseteq A_n$ unitaly. Let $B = \overline{\cup_n B_n}$ and $A = \overline{\cup_n A_n}$. If $B_n \subseteq A_n$ is \mathcal{D} -stable for all n , then $B \subseteq A$ is \mathcal{D} -stable.*

Proof This follows from Proposition 3.7, together with Lemma 3.2 and Theorem 4.4. \blacksquare

Lastly, we discuss unital inclusions $B \subseteq A$ of $C(X)$ algebras, where X is a compact Hausdorff space. We show that if X has finite covering dimension, then such an inclusion is \mathcal{D} -stable if and only if the inclusion $B_x \subseteq A_x$ along each fiber is \mathcal{D} -stable.

Lemma 4.19 *Let \mathcal{D} be strongly self-absorbing. Suppose that $B_i \subseteq A_i$ are unital inclusions, for $i = 1, 2$, and $\psi : A_1 \rightarrow A_2$ is a surjective $*$ -homomorphism such that $\psi(B_1) = B_2$. If $B_1 \subseteq A_1$ is \mathcal{D} -stable, then so is $B_2 \subseteq A_2$.*

Proof We note that ψ induces a $*$ -homomorphism

$$(4.29) \quad \tilde{\psi} : (B_1)_\omega \cap A'_1 \rightarrow (B_2)_\omega \cap A'_2$$

and consequently if $\xi : \mathcal{D} \hookrightarrow (B_1)_\omega \cap A'_1$, we have a unital $*$ -homomorphism

$$(4.30) \quad \eta := \tilde{\psi} \circ \xi : \mathcal{D} \rightarrow (B_2)_\omega \cap A'_2.$$

The homomorphism η is automatically injective since \mathcal{D} is simple. \blacksquare

Rephrasing the above in terms of commutative diagrams, it says that if we have a commutative diagram

$$(4.31) \quad \begin{array}{ccc} A_1 & \longrightarrow & A_2 \\ \uparrow & & \uparrow \\ B_1 & \longrightarrow & B_2 \end{array}$$

where the left inclusion is \mathcal{D} -stable, then the right inclusion is \mathcal{D} -stable as well.

Now we consider many of the results discussed in [27, Section 4], except for inclusions of C^* -algebras.

Definition 4.2 Let X be a compact Hausdorff space. A $C(X)$ -algebra is a C^* -algebra A endowed with a unital $*$ -homomorphism $C(X) \rightarrow \mathcal{Z}(\mathcal{M}(A))$, where $\mathcal{Z}(\mathcal{M}(A))$ is the center of the multiplier algebra $\mathcal{M}(A)$ of A .

If $Y \subseteq X$ is a closed subset, we set $I_Y := C_0(X \setminus Y)A$, which is a closed two-sided ideal in A . We denote $A_Y := A/I_Y$ and the quotient map $A \rightarrow A_Y$ by π_Y . For an element $a \in A$, we write $a_Y := \pi_Y(a)$ and if Y consists of a single point x , we write A_x, I_x, π_x and a_x . We say that A_x is the fiber of A at x . We note that $A_X = A$.

If $B \subseteq A$ is a unital inclusion and $\theta_A : C(X) \rightarrow A$, $\theta_B : C(X) \rightarrow B$ are morphisms which witness A and B as $C(X)$ -algebras, respectively, we say that $B \subseteq A$ is an inclusion of $C(X)$ -algebras if

$$(4.32) \quad \begin{array}{ccc} B & \hookrightarrow & A \\ \theta_B \uparrow & \nearrow \theta_A & \\ C(X) & & \end{array}$$

commutes. Note that $\theta_B(C(X)) \subseteq \mathcal{Z}(A)$, and when discussing an inclusion of fibers $B_Y \subseteq A_Y$ we are considering $B_Y := \pi_Y^A(B) \subseteq \pi_Y^A(A) =: A_Y$, where $\pi_Y^A : A \rightarrow A_Y$ is the associated quotient map.

Remark 4.20 (Upper semi-continuity) In [27, Section 1.3], it was pointed out that the norm on a $C(X)$ -algebra A is upper semi-continuous. This means that, fixing some $a \in A$, the function $x \mapsto \|a_x\|$ from X to \mathbb{R} is upper semi-continuous (as it is the infimum of a family of continuous functions), and consequently the set $\{x \in X \mid \|a_x\| < \varepsilon\} \subseteq X$ is open for all $a \in A$ and $\varepsilon > 0$.

We note that Lemma 4.19 gives that if $B \subseteq A$ is \mathcal{D} -stable and $Y \subseteq X$ is closed, then $B_Y \subseteq A_Y$ is automatically \mathcal{D} -stable as well since we have the commuting diagram

$$(4.33) \quad \begin{array}{ccc} A & \xrightarrow{\pi_Y} & A_Y \\ \uparrow & & \uparrow \\ B & \xrightarrow{\pi_Y|_B} & B_Y. \end{array}$$

The converse needs a bit of work. This is the embedding analog of the beginning of [27, Section 4]. We discuss how the proofs can be adapted and often omit approximations that were otherwise done there. We want a version of [27, Lemma 4.5], which is a result about *gluing* c.c.p. maps together along fibers. In our setting, we are only interested in u.c.p. maps, and we want to show that if we *glue* two u.c.p. maps together whose images are contained in some $C(X)$ -subalgebra B , then the *glued* map also has image contained in B . We borrow their Definition 4.2.

Definition 4.3 Let A be a unital $C(X)$ -algebra, for a compact Hausdorff space X , and let D be a unital C^* -algebra. Let $\phi : D \rightarrow A$ be a u.c.p. map and $Y \subseteq X$ a closed subset. If $\mathcal{F} \subseteq D$, $\mathcal{G} \subseteq A$ are finite and $\varepsilon > 0$, we say that ϕ is $(\mathcal{F}, \varepsilon, \mathcal{G})$ -good for Y if:

- (1) $([\phi(d), a])_Y \approx_\varepsilon 0$ and
- (2) $\phi(dd')_Y \approx_\varepsilon \phi(d)_Y \phi(d')_Y$

whenever $d, d' \in \mathcal{F}$ and $a \in \mathcal{G}$. If $X = [0, 1]$, $Y \subseteq X$ is a closed interval, $\mathcal{F}' \supseteq \mathcal{F}$ is another finite set and $0 < \varepsilon' < \varepsilon$, we say that ϕ is $(\mathcal{F}, \varepsilon, \mathcal{G}; \mathcal{F}', \varepsilon')$ -good for Y if ϕ is $(\mathcal{F}, \varepsilon, \mathcal{G})$ -good for Y and there exists some closed neighborhood V of the endpoints of Y such that ϕ is $(\mathcal{F}', \varepsilon', \mathcal{G})$ -good for V .

First, we need a lemma that follows as a consequence of \mathcal{D} -stability. It is the embedding analog of [27, Proposition 4.1].

Lemma 4.21 *Let \mathcal{D} be strongly self-absorbing, and $B \subseteq A$ be a unital, \mathcal{D} -stable inclusion of separable C^* -algebras. Then for any $\mathcal{G} \subseteq A$ finite and $\varepsilon > 0$, there exist unital $*$ -homomorphisms $\kappa : A \rightarrow A$ and $\mu : \mathcal{D} \rightarrow B$ such that:*

- (1) $\kappa(B) \subseteq B$,
- (2) $[\kappa(A), \mu(\mathcal{D})] = 0$,
- (3) $\kappa(a) \approx_\varepsilon a$ for all $a \in \mathcal{G}$.

Proof The proof is essentially the same as the proof of (a) \Rightarrow (c) in [27, Proposition 4.1]. As $B \subseteq A$ is \mathcal{D} -stable, let us identify $B \subseteq A$ with $B \otimes \mathcal{D} \subseteq A \otimes \mathcal{D}$. As \mathcal{D} is strongly self-absorbing, [66, Theorem 2.3] gives a sequence $(\phi_n)_{n \in \mathbb{N}}$ of $*$ -homomorphisms $\phi_n : \mathcal{D} \otimes \mathcal{D} \rightarrow \mathcal{D}$ such that

$$(4.34) \quad \phi_n(d \otimes 1_{\mathcal{D}}) \rightarrow d \text{ for all } d \in \mathcal{D}.$$

Define $\kappa_n : A \otimes \mathcal{D} \rightarrow A \otimes \mathcal{D}$ by

$$(4.35) \quad \kappa_n := (\text{id}_A \otimes \phi) \circ (\text{id}_A \otimes \text{id}_{\mathcal{D}} \otimes 1_{\mathcal{D}}),$$

and $\mu_n : \mathcal{D} \rightarrow B \otimes \mathcal{D}$ by

$$(4.36) \quad \mu_n := (\text{id}_B \otimes \phi_n) \circ (1_A \otimes 1_{\mathcal{D}} \otimes \text{id}_{\mathcal{D}}).$$

Then taking n large enough and letting κ and μ be κ_n and μ_n , respectively, it is clear that $\kappa(B \otimes \mathcal{D}) \subseteq B \otimes \mathcal{D}$, $[\kappa(A), \mu(\mathcal{D})] = 0$ and that $\kappa(a) \approx_\varepsilon a$ whenever a is in some prescribed finite subset $\mathcal{G} \subseteq A$ and $\varepsilon > 0$ is some prescribed error. ■

Lemma 4.22 *Let \mathcal{D} be strongly self-absorbing and A be a unital, separable $C([0, 1])$ -algebra. Suppose $\mathcal{F} \subseteq \mathcal{D}$, $\mathcal{G} \subseteq A$ are finite self-adjoint subsets of contractions with $1_{\mathcal{D}} \in \mathcal{F}$. Suppose that we have points $0 \leq r < s < t \leq 1$ and two u.c.p. maps $\rho, \sigma : \mathcal{D} \rightarrow A$ which are $(\mathcal{F}, \varepsilon, \mathcal{G})$ -good for $[r, s]$, $[s, t]$, respectively. Suppose that A_s is \mathcal{D} -stable.*

Then there are u.c.p. maps $\rho', \sigma' : \mathcal{D} \rightarrow A$ which are $(\mathcal{F}, \varepsilon, \mathcal{G})$ -good for $[r, s]$, $[s, t]$, respectively, and u.c.p. maps $\nu_{\rho'}, \nu_{\sigma'} : \mathcal{D} \rightarrow A$, $\mu_{\rho'}, \mu_{\sigma'} : \mathcal{D} \otimes \mathcal{D} \rightarrow A$ such that $\nu_{\rho'}, \nu_{\sigma'}$ are $(\mathcal{F}, 3\varepsilon, \mathcal{G})$ -good for some interval $I \subseteq (r, t)$ containing s in its interior, and such that for any $a \in \mathcal{G}$, $d, d' \in \mathcal{F}$, we have:

- (1) $([\rho'(d), \nu_{\rho'}(d')])_I \approx_{2\varepsilon} 0$,
- (2) $([\sigma'(d), \nu_{\sigma'}(d')])_I \approx_{2\varepsilon} 0$,
- (3) $\rho'(d)_I \nu_{\rho'}(d')_I \approx_\varepsilon \mu_{\rho'}(d \otimes d')_I$,
- (4) $\sigma'(d)_I \nu_{\sigma'}(d')_I \approx_\varepsilon \mu_{\sigma'}(d \otimes d')_I$,
- (5) $\nu_{\rho'}(d)_I \approx_{2\varepsilon} \nu_{\sigma'}(d)_I$.

If ρ, σ are $(\mathcal{F}, \varepsilon, \mathcal{G}; \mathcal{F}', \varepsilon)$ -good for $[r, s]$, $[s, t]$, respectively, for some finite $\mathcal{F}' \supseteq \mathcal{F}$ set of contractions and for some $0 < \varepsilon' < \varepsilon$, then we can arrange so that $\rho', \sigma', \nu_{\rho'}, \nu_{\sigma'}$ are $(\mathcal{F}', 3\varepsilon', \mathcal{G})$ -good for the interval I , and that the above five conditions hold with ε' in place of ε and \mathcal{F}' in place of \mathcal{F} .

Moreover, if $B \subseteq A$ is a unital inclusion of $C([0, 1])$ -algebras such that $\rho(\mathcal{D}) \subseteq B$, $\sigma(\mathcal{D}) \subseteq B$ and $B_s \subseteq A_s$ is \mathcal{D} -stable, then the images of all $\rho', \sigma', \mu_{\rho'}, \mu_{\sigma'}$ are contained in B (as are the images of $\nu_{\rho'}$ and $\nu_{\sigma'}$).

Proof This is [27, Lemma 4.4], except we have replaced c.c.p. maps with u.c.p. maps. One can easily check that the resulting maps are u.c.p. maps.

As for the “moreover” part, which is the only addition besides the unitality, we outline the definitions of these maps to show that the images of $\rho', \sigma', \mu_{\rho'}, \mu_{\sigma'}$ are contained in B . As $B_s \subseteq A_s$ is \mathcal{D} -stable, we can find $\kappa : A_s \rightarrow A_s$ and $\mu : \mathcal{D} \rightarrow B_s$ as in Lemma 4.21, where $\kappa(a_s) \approx a_s$ for an appropriate error whenever $a \in \mathcal{G}$. We use Choi–Effros to find u.c.p. lifts $\tilde{\rho}, \tilde{\sigma} : \mathcal{D} \rightarrow B$ for the maps $\kappa \circ \pi_s \circ \rho$ and $\kappa \circ \pi_s \circ \sigma$, respectively (note that $\kappa \circ \pi_s \circ \rho$ and $\kappa \circ \pi_s \circ \sigma$ lie in B_s , which is a $*$ -homomorphic image of B). One then defines piece-wise linear functions $f, g : [0, 1] \rightarrow [0, 1]$ which attain both values 0 and 1 at the end points (their definition is not important to show the “moreover” part). Then ρ', σ' are defined as

$$(4.37) \quad \rho'(d) := (1 - f) \cdot \rho(d) + f \cdot \tilde{\rho}(d) \text{ and } \sigma'(d) := (1 - g) \cdot \sigma(d) + g \cdot \tilde{\sigma}(d).$$

Clearly, ρ', σ' take values in B as $\rho, \tilde{\rho}, \sigma, \tilde{\sigma}$ all do and $(1 - f), f, (1 - g), g$ are in B . Now we define u.c.p. maps $\tilde{\mu}_{\rho'}, \tilde{\mu}_{\sigma'} : \mathcal{D} \otimes \mathcal{D} \rightarrow B_s$ by

$$(4.38) \quad \tilde{\mu}_{\rho'}(d \otimes d') := \rho'(d)_s \mu(d') \text{ and } \tilde{\mu}_{\sigma'}(d \otimes d') := \sigma'(d)_s \mu(d').$$

Now by Choi–Effros, we can take u.c.p. lifts $\mu_{\rho'}$ and $\mu_{\sigma'}$ of $\tilde{\mu}_{\rho'}$ and $\tilde{\mu}_{\sigma'}$, respectively. As the images of $\tilde{\mu}_{\rho'}$ and $\tilde{\mu}_{\sigma'}$ lie in B_s , the images of $\mu_{\rho'}$ and $\mu_{\sigma'}$ will lie in B . ■

Lemma 4.23 *Let A be a unital, separable $C([0, 1])$ -algebra. Suppose $\mathcal{F} \subseteq \mathcal{D}, \mathcal{G} \subseteq A$ are finite self-adjoint subsets with $1_{\mathcal{D}} \in \mathcal{F}$ and $\varepsilon > 0$. There exists $0 < \varepsilon' < \varepsilon$ and a finite subset $\mathcal{F}' \supseteq \mathcal{F}$ such that if $\rho, \sigma : \mathcal{D} \rightarrow A$ are u.p.c. maps and $0 \leq r < s < t \leq 1$ are points such that ρ is $(\mathcal{F}, \varepsilon, \mathcal{G}; \mathcal{F}', \varepsilon')$ -good for $[r, s]$, σ is $(\mathcal{F}, \varepsilon, \mathcal{G}; \mathcal{F}', \varepsilon')$ -good for $[s, t]$ and A_s is \mathcal{D} -stable, then there is a u.c.p. map $\psi : \mathcal{D} \rightarrow A$ which is $(\mathcal{F}, \varepsilon, \mathcal{G}; \mathcal{F}', \varepsilon')$ -good for $[r, t]$.*

Moreover, if $B \subseteq A$ is a unital inclusion of $C([0, 1])$ -algebras such that $\rho(\mathcal{D}) \subseteq B$, $\sigma(\mathcal{D}) \subseteq B$ and $B_s \subseteq A_s$ is \mathcal{D} -stable, then $\psi(\mathcal{D}) \subseteq B$.

Proof The first part is [27, Lemma 4.5], except we have replaced c.c.p. maps with u.c.p. maps. One has to check that the resulting ψ is unital, but this follows easily if ρ and σ are.

We outline the construction of ψ to show unitality, as it will also be useful to show the “moreover” part, which is the only real addition. Let $u \in C([0, 1], \mathcal{D} \otimes \mathcal{D})$ be a path of unitaries such that $u_0 = 1_{\mathcal{D} \otimes \mathcal{D}}$ and

$$(4.39) \quad u_1(d \otimes 1_{\mathcal{D}})u_1^* \approx_{\frac{\varepsilon}{4}} 1_{\mathcal{D}} \otimes d.$$

We replace ρ, σ with ρ', σ' as in the above lemma and this yields u.c.p. maps μ_{ρ}, μ_{σ} satisfying the hypotheses above for some interval $I \subseteq (r, t)$ with s in its interior. Define

$$(4.40) \quad \phi_{\rho}, \phi_{\sigma} : C([0, 1]) \otimes \mathcal{D} \otimes \mathcal{D} \rightarrow A$$

by

$$(4.41) \quad \begin{aligned} \phi_{\rho}(f \otimes d \otimes d') &:= f \cdot \mu_{\rho}(d \otimes d'), \\ \phi_{\sigma}(f \otimes d \otimes d') &:= f \cdot \mu_{\sigma}(d \otimes d'). \end{aligned}$$

Note that these maps are unital. Take nonzero piece-wise linear functions

$$(4.42) \quad h_1, h_2, h_3, h_4 : [0, 1] \rightarrow [0, 1]$$

which sum to 1 (their specific form does not matter to show unitality of ψ nor the “moreover” part) and $g_\rho, g_\sigma : [0, 1] \rightarrow [0, 1]$ which sum to 1 (again, their specific form does not matter to show unitality of ψ nor the “moreover” part). Define unitaries $u_\rho, u_\sigma \in C([0, 1]) \otimes \mathcal{D} \otimes \mathcal{D} \simeq C([0, 1], \mathcal{D} \otimes \mathcal{D})$ by

$$(4.43) \quad u_{\rho x} := u_{g_\rho(x)} \text{ and } u_{\sigma x} := u_{g_\sigma(x)}.$$

Now define $\zeta_\rho, \zeta_\sigma : \mathcal{D} \rightarrow A$ by

$$(4.44) \quad \begin{aligned} \zeta_\rho(d) &:= \phi_\rho(u_\rho(1_{C([0,1])} \otimes d \otimes 1_{\mathcal{D}})u_\rho^*), \\ \zeta_\sigma(d) &:= \phi_\sigma(u_\sigma(1_{C([0,1])} \otimes d \otimes 1_{\mathcal{D}})u_\sigma^*), \end{aligned}$$

which are clearly unital. Finally, the map $\psi : \mathcal{D} \rightarrow A$ is defined by

$$(4.45) \quad \psi(d) := h_1 \cdot \rho(d) + h_2 \cdot \zeta_\rho(d) + h_3 \cdot \zeta_\sigma(d) + h_4 \cdot \sigma(d).$$

Clearly, ψ is unital.

Now for the “moreover” part. If $\rho(\mathcal{D}) \subseteq B$ and $\sigma(\mathcal{D}) \subseteq B$, clearly the first and fourth terms in the definition of ψ will lie in B . So it suffices to show that $\zeta_\rho(\mathcal{D}) \subseteq B$ and $\zeta_\sigma(\mathcal{D}) \subseteq B$, and for this it suffices to show that $\mu_\rho(\mathcal{D} \otimes \mathcal{D}) \subseteq B$ and $\mu_\sigma(\mathcal{D} \otimes \mathcal{D}) \subseteq B$ (since h_1, h_2, h_3, h_4 all lie in B). But this follows from the “moreover” part of the previous lemma. ■

With this, we get the analog of [28, Theorem 4.6], the proof being essentially the same as well, except we insist that the our u.c.p. maps commute with a prescribed finite subset of A .

Proposition 4.24 *Let \mathcal{D} be strongly self-absorbing, and X be a compact Hausdorff space with finite covering dimension. Suppose that $B \subseteq A$ is a unital inclusion of $C(X)$ -algebras. Then $B_x \subseteq A_x$ is \mathcal{D} -stable for all $x \in X$ if and only if $B \subseteq A$ is \mathcal{D} -stable.*

Proof As previously mentioned, if $B \subseteq A$ is \mathcal{D} -stable, then $B_x \subseteq A_x$ is \mathcal{D} -stable for all x .

For the converse, the proof is essentially the same as [27, Theorem 4.6]. Using the arguments there, one can simplify to the case where we can argue this for $C([0, 1])$ -algebras (by using [29, Theorem V.3], which says that a compact space of dimension $\leq n$ is homeomorphic to a subset of $[0, 1]^{2n+1}$, and then working component-wise). Now for $\mathcal{F} \subseteq \mathcal{D}, \mathcal{G} \subseteq A$ and $\varepsilon > 0$, let $\mathcal{G}_x := \{a_x \mid a \in \mathcal{G}\}$. Without loss of generality suppose that $\mathcal{F}^* = \mathcal{F}, \mathcal{G}^* = \mathcal{G}$ and that $1_{\mathcal{D}} \in \mathcal{F}$. Let $\mathcal{F}', \varepsilon'$ be as in Lemma 4.23.

By \mathcal{D} -stability of the inclusion $B_x \subseteq A_x$ there are u.c.p. $(\mathcal{F}', \varepsilon', \mathcal{G}_x)$ -embeddings $\psi_x : \mathcal{D} \rightarrow B_x \subseteq A_x$ which lift by Choi–Effros to u.c.p. maps $\psi'_x : \mathcal{D} \rightarrow B$. The norm is upper semi-continuous (Remark 4.20), and this yields intervals $I_x \subseteq [0, 1]$ such that ψ'_x is $(\mathcal{F}', \varepsilon', \mathcal{G})$ -good for $\overline{I_x}$. Note that ψ'_x being $(\mathcal{F}', \varepsilon', \mathcal{G})$ -good for the whole of I_x implies that it is $(\mathcal{F}, \varepsilon, \mathcal{G}; \mathcal{F}', \varepsilon')$ -good for $\overline{I_x}$. Compactness then allows us to split the interval as

$$(4.46) \quad 0 = t_0 < t_1 < \cdots < t_n = 1$$

and to take $\psi_i : \mathcal{D} \rightarrow B$ u.c.p. which are $(\mathcal{F}, \varepsilon, \mathcal{G}; \mathcal{F}', \varepsilon')$ -good for $[t_{i-1}, t_i]$ for $i = 1, \dots, n$ ($\psi_i = \psi'_x$ for some $x \in [0, 1]$). Now by repeatedly using the gluing lemma

(Lemma 4.23) to glue these maps together, we can find a u.c.p. map $\psi : \mathcal{D} \rightarrow B$ which is an $(\mathcal{F}, \varepsilon, \mathcal{G})$ -embedding. ■

5 Crossed products

In this section, we discuss how inclusions coming from noncommutative dynamics fit into the framework of tensorially absorbing inclusions. We will briefly discuss group actions $G \curvearrowright^\alpha A$ with Rokhlin properties and consider the inclusion of a C^* -algebra in its crossed product $A \subseteq A \rtimes_\alpha G$, as well as the inclusion of the fixed point subalgebra of the action in the C^* -algebra $A^\alpha \subseteq A$. We then discuss diagonal inclusions associated with certain group actions.

This first result says that if we have an isomorphism $A \simeq A \otimes \mathcal{D}$ which is G -equivariant with respect to an action point-wise fixing the right tensor factor, up to a 1-cocycle, then the corresponding inclusion $A \subseteq A \rtimes_{r,\alpha} G$ is \mathcal{D} -stable. Recall that if $\beta : G \curvearrowright B$ is an action of a countable discrete group on a unital C^* -algebra B , then a β -1-cocycle is a map $u : G \rightarrow U(B)$ satisfying the cocycle identity:

$$(5.1) \quad u_{gh} = u_g \beta_g(u_h).$$

If $(A, \alpha), (B, \beta)$ are G - C^* -algebras, we say that they are cocycle conjugate, denoted $(A, \alpha) \simeq_{c.c.} (B, \beta)$, if there are an isomorphism $\phi : A \simeq B$ and a β -1-cocycle $u : G \rightarrow U(B)$ such that

$$(5.2) \quad \begin{array}{ccc} A & \xrightarrow{\phi} & B \\ \alpha_g \downarrow & & \downarrow \text{Ad}(u_g) \circ \beta_g \\ A & \xrightarrow{\phi} & B \end{array}$$

commutes for all $g \in G$. Conjugacy is usually too strong a notion of equivalence, whereas cocycle conjugacy has allowed for quite deep classification of automorphisms. For example, this notion has been used for classifying automorphisms of von Neumann factors [11, 12, 14, 33, 57].

Proposition 5.1 *Let $G \curvearrowright^\alpha A$ be an action of a countable discrete group on a unital separable C^* -algebra. Suppose that $\alpha \simeq_{c.c.} \alpha \otimes id_{\mathcal{D}}$. That is, there is an $\alpha \otimes id_{\mathcal{D}}$ -1-cocycle $u : G \rightarrow U(A \otimes \mathcal{D})$ and an isomorphism $\Phi : A \simeq A \otimes \mathcal{D}$ such that*

$$(5.3) \quad \begin{array}{ccc} A & \xrightarrow{\Phi} & A \otimes \mathcal{D} \\ \alpha_g \downarrow & & \downarrow \text{Ad}(u_g) \circ (\alpha_g \otimes id_{\mathcal{D}}) \\ A & \xrightarrow{\Phi} & A \otimes \mathcal{D} \end{array}$$

commutes for all $g \in G$. Then $A \subseteq A \rtimes_{r,\alpha} G$ is \mathcal{D} -stable.

Proof Let $\psi : \mathcal{D} \simeq \mathcal{D}^{\otimes \infty}$, and let $\phi_n : \mathcal{D} \rightarrow \mathcal{D}^{\otimes \infty}$ be the n th factor embedding:

$$(5.4) \quad \phi_n(d) := 1_{\mathcal{D}}^{\otimes n-1} \otimes d \otimes 1_{\mathcal{D}}^{\otimes \infty}.$$

We claim that $\xi(d) := (\Phi^{-1}(1_A \otimes \psi^{-1} \circ \phi_n(d)))_n : \mathcal{D} \rightarrow A_\omega$ is an embedding such that $\xi(\mathcal{D}) \subseteq A_\omega \cap A'$ and $(\alpha_g)_\omega \circ \xi = \xi$ for all $g \in G$ – that is, ξ is an embedding

$\mathcal{D} \hookrightarrow A_\omega \cap (A \rtimes_{r,\alpha} G)'$. The first part of the claim is obvious, so we prove the second. We have

$$\begin{aligned}
 & \|\alpha_g(\Phi^{-1}(1_A \otimes \psi^{-1}(\phi_n(d)))) - \Phi^{-1}(1_A \otimes \psi^{-1}(\phi_n(d)))\| \\
 &= \|\Phi \circ \alpha_g(\Phi^{-1}(1_A \otimes \psi^{-1}(\phi_n(d)))) - \Phi(\Phi^{-1}(1_A \otimes \psi^{-1}(\phi_n(d))))\| \\
 &= \|\text{Ad}(u_g) \circ (\alpha_g \otimes \text{id}_{\mathcal{D}})(1_A \otimes \psi^{-1}(\phi_n(d))) - 1_A \otimes \psi^{-1}(\phi_n(d))\| \\
 &= \|\text{Ad}(u_g)(1_A \otimes \psi^{-1}(\phi_n(d))) - 1_A \otimes \psi^{-1}(\phi_n(d))\| \\
 &= \|u_g^*(1_A \otimes \psi^{-1}(\phi_n(d)))u_g - 1_A \otimes \psi^{-1}(\phi_n(d))\| \\
 (5.5) \quad & \rightarrow 0
 \end{aligned}$$

since $(1_A \otimes \psi^{-1}(\phi_n(d)))_n$ is asymptotically central in $A \otimes \mathcal{D}$. \blacksquare

Actions satisfying the hypotheses of Proposition 5.1 are said to be *equivariantly \mathcal{D} -absorbing, up to cocycle conjugacy*. These actions are fairly common and there are a wide range of positive results (see, for example, [58, 59]).

The next lemma of note is the following.

Lemma 5.2 *Suppose that $G \curvearrowright^\alpha A$ is an action of a finite group on a unital separable C^* -algebra A such that $A \subseteq A \rtimes_\alpha G$ is \mathcal{D} -stable. Then $A^\alpha \subseteq A \rtimes_\alpha G$ is \mathcal{D} -stable. In particular, if $A \subseteq A \rtimes_\alpha G$ is \mathcal{D} -stable, then $C \simeq C \otimes \mathcal{D}$ whenever $A^\alpha \subseteq C \subseteq A \rtimes_\alpha \mathcal{D}$.*

Proof For an element $(x_n)_{n \in \mathbb{N}} \in A_\omega \cap (A \rtimes_\alpha G)'$, an easy averaging argument shows that

$$(5.6) \quad (x_n)_{n \in \mathbb{N}} = \left(\frac{1}{|G|} \sum_{g \in G} \alpha_g(x_n) \right)_{n \in \mathbb{N}}$$

in A_ω , and the right is clearly point-wise fixed by α_g for all $g \in G$. So $A_\omega \cap (A \rtimes_\alpha G)'$ is actually equal to $(A^\alpha)_\omega \cap (A \rtimes_\alpha G)'$, and the existence of a unital embedding of \mathcal{D} in $A_\omega \cap (A \rtimes_\alpha G)'$ is in fact equivalent to the existence of a unital embedding of \mathcal{D} into $(A^\alpha)_\omega \cap (A \rtimes_\alpha G)'$. The result follows. \blacksquare

The Galois correspondence of Izumi [30] yields the following.

Theorem 5.3 *Let A be a unital, simple, separable C^* -algebra, and let $G \curvearrowright^\alpha A$ be an action of a finite group by outer automorphisms. If $A \subseteq A \rtimes_\alpha \mathcal{D}$ is \mathcal{D} -stable, then there exists an isomorphism $\Phi: A \rtimes_\alpha G \simeq (A \rtimes_\alpha G) \otimes \mathcal{D}$ such that whenever C is a unital C^* -algebra satisfying either:*

- (1) $A^\alpha \subseteq C \subseteq A$ or
- (2) $A \subseteq C \subseteq A \rtimes_\alpha G$,

we have $\Phi(C) = C \otimes \mathcal{D}$.

Proof Applying [30, Corollary 6.6] gives the following two correspondences:

- (1) there is a one-to-one correspondence between subgroups of G with intermediate C^* -algebras $A^\alpha \subseteq C \subseteq A$ given by

$$(5.7) \quad H \leftrightarrow A^{\alpha_H};$$

- (2) there is a one-to-one correspondence between subgroups of G and intermediate C^* -algebras $A \subseteq C \subseteq A \rtimes_\alpha G$ given by

$$(5.8) \quad H \leftrightarrow A \rtimes_{\alpha|_H} H.$$

In particular, there are only finitely many C^* -algebras C between either $A^\alpha \subseteq A$ or $A \subseteq A \rtimes_\alpha G$. As all such lie between the \mathcal{D} -stable inclusion $A^\alpha \subseteq A \rtimes_\alpha G$, Theorem 4.8 yields the desired isomorphism. ■

5.1 (Tracial) Rokhlin properties

Here, we will restrict ourselves to finite groups for simplicity, although many results hold more generally (see [23, 26, 28]).

Definition 5.1 Let A be a unital, separable C^* -algebra. We say that a finite group action $G \curvearrowright^\alpha A$ has the Rokhlin property if there are pairwise orthogonal projections $(p_g)_{g \in G} \subseteq A_\omega \cap A'$ summing to 1_{A_ω} such that $(\alpha_g)_\omega(p_h) = p_{gh}$ for $g, h \in G$.

Proposition 5.4 Let A be a unital, separable \mathcal{D} -stable C^* -algebra. If $G \curvearrowright^\alpha A$ is an action of a finite group with the Rokhlin property, then $A^\alpha \subseteq A \rtimes_\alpha G$ is \mathcal{D} -stable.

Proof This follows from [28, Theorem 3.3], together with Lemma 5.2. ■

Definition 5.2 Let A be a unital, separable C^* -algebra. We say that a finite group action $G \curvearrowright^\alpha A$ has the weak tracial Rokhlin property if for all $\mathcal{F} \subseteq A$ finite, $\varepsilon > 0$ and $0 \neq a \in A_+$, there are pairwise orthogonal normalized positive contractions $(e_g)_{g \in G} \subseteq A$ such that:

- (1) $1 - \sum_g e_g \lesssim a$;⁴
- (2) $[e_g, x] \approx_\varepsilon 0$ for all $x \in \mathcal{F}$, $g \in G$;
- (3) $\alpha_g(e_h) \approx_\varepsilon e_{gh}$ for all $g, h \in G$.

It is easy to see that a Rokhlin action is outer, since the central projections must commute with any unitary. The fact that weak tracial Rokhlin actions are outer is [26, Proposition 5.3].

Proposition 5.5 Let A be a unital, simple, separable, nuclear, \mathbb{Z} -stable C^* -algebra. If $G \curvearrowright^\alpha A$ is an action of a finite group with the weak tracial Rokhlin property, then $A^\alpha \subseteq A \rtimes_\alpha G$ is \mathbb{Z} -stable.

Proof Let $k \in \mathbb{N}$. By [26, Theorem 5.6] $A \rtimes_\alpha G$ is tracially \mathbb{Z} -absorbing, meaning there are tracially large (in the sense of [65]) c.p.c. order zero maps $\phi: M_k \rightarrow (A \rtimes_\alpha G)_\omega \cap (A \rtimes_\alpha G)'$, which can be chosen to be c.p.c. order zero maps $\phi: M_k \rightarrow A_\omega \cap (A \rtimes_\alpha G)'$ by the proof of [26, Lemma 5.5]. These tracially large c.p.c. order zero maps yield sequences of positive contractions $c_1 = (c_{1n}), \dots, c_k = (c_{kn}) \in A_\omega \cap (A \rtimes_\alpha G)'$ such that if $(e_n)_{n \in \mathbb{N}} = e := 1 - \sum_i c_i^* c_i$, we have

$$(5.9) \quad \lim_{n \rightarrow \infty} \max_{\tau \in T(A)} \tau(e_n) = 0, \inf_m \lim_{n \rightarrow \infty} \min_{\tau \in T(A)} \tau(c_{1n}^m) > 0$$

⁴For two positive elements x, y in a C^* -algebra, we write $x \lesssim y$ to mean that x is Cuntz-subequivalent to y . That is, there are $(r_n)_{n \in \mathbb{N}}$ in the C^* -algebra such that $r_n^* y r_n \rightarrow x$ (see [26, Section 2]).

and $c_i c_j^* = \delta_{ij} c_1^2$. By [23, Proposition 4.11] (which is much more general, applicable to all countable amenable groups), $A \subseteq A \rtimes_\alpha G$ has equivariant property (SI) since A has property (SI).⁵ Consequently, there exists $s \in A_\omega \cap (A \rtimes_\alpha G)'$ such that $s^* s = 1 - \sum_i c_i^* c_i$ and $c_1 s = s$. Altogether,

- $c_1 \geq 0$;
- $c_i c_j^* = \delta_{ij} c_1^2$;
- $s^* s + \sum_i c_i^* c_i = 1$;
- $c_1 s = s$.

As mentioned in the proof of (iv) \Rightarrow (i) of [39], $\mathcal{Z}_{n,n+1}$ is the universal C^* -algebra generated by $n+1$ elements satisfying the above four relations (see [52, Proposition 5.1] and [54, Proposition 2.1]), and consequently we have a unital $*$ -homomorphism $\mathcal{Z}_{n,n+1} \rightarrow A_\omega \cap (A \rtimes_\alpha G)'$. Therefore $\mathcal{Z} \hookrightarrow A_\omega \cap (A \rtimes_\alpha G)'$, giving that the desired inclusion is \mathcal{Z} -stable by Lemma 5.2. ■

Corollary 5.6 *Let A be a unital, simple, separable, nuclear, \mathcal{Z} -stable C^* -algebra, and let $G \curvearrowright^\alpha A$ be an action of a finite group with the weak tracial Rokhlin property. There exists an isomorphism $\Phi : A \rtimes_\alpha G \simeq (A \rtimes_\alpha G) \otimes \mathcal{Z}$ such that whenever C is a unital C^* -algebra satisfying either:*

- (1) $A^\alpha \subseteq C \subseteq A$ or
 - (2) $A \subseteq C \subseteq A \rtimes_\alpha G$,
- we have $\Phi(C) = C \otimes \mathcal{Z}$.*

Proof This results from combining Proposition 5.5 together with Theorem 5.3, making note that this is an outer action. ■

5.2 The diagonal inclusion associated with a group action

In the von Neumann setting, a certain diagonal inclusion associated with several automorphisms was considered in [7, 34, 43], and they play a role in subfactor theory. Here, we consider a unital C^* -algebraic inclusion of the same form.

Definition 5.3 Let A be a C^* -algebra, $\alpha_1, \dots, \alpha_n \in \text{Aut}(A)$. The diagonal inclusion associated with $\alpha_1, \dots, \alpha_n$ is

$$(5.10) \quad B(\alpha_1, \dots, \alpha_n) = \left\{ \bigoplus_{i=1}^n \alpha_i(a) \mid a \in A \right\} \subseteq M_n(A).$$

If $G \curvearrowright^\alpha A$ is an action of a finite group, we write

$$(5.11) \quad B(\alpha) = \left\{ \bigoplus_{g \in G} \alpha_g(a) \mid a \in A \right\} \subseteq M_{|G|}(A).$$

We note that a diagonal $B(\alpha) \subseteq M_{|G|}(A)$ is unique up to unitary conjugation (by permutation unitaries). As \mathcal{D} -stability of an inclusion is preserved under unitary conjugation, there is no ambiguity in speaking of \mathcal{D} -stability of the inclusion $B(\alpha) \subseteq M_{|G|}(A)$.

⁵A unital, separable, simple, nuclear, \mathcal{Z} -stable C^* -algebra has property (SI) as in [39].

Proposition 5.7 *Let $G \curvearrowright^\alpha A$ be an action of a countable discrete group on a unital, separable C^* -algebra. If $G = \langle g_1, \dots, g_n \rangle$, then $A \subseteq A \rtimes_\alpha G$ is \mathcal{D} -stable if and only if*

$$(5.12) \quad B(\text{id}_A, \alpha_{g_1}, \dots, \alpha_{g_n}) \subseteq M_{n+1}(A)$$

is \mathcal{D} -stable.

Proof First, suppose that $A \subseteq A \rtimes_\alpha G$ is \mathcal{D} -stable. Let $\mathcal{F} \subseteq \mathcal{D}$, $\mathcal{G} \subseteq M_{n+1}(A)$ be finite and $\varepsilon > 0$. Let $\mathcal{G}' \subseteq A$ be the set of matrix coefficients of elements of \mathcal{G} , together with the identity of A , and let $L := \max\{1, \max_{a \in \mathcal{G}'} \|a\|\}$. Relabel $\text{id}_A, \alpha_{g_1}, \dots, \alpha_{g_n}$ as $\alpha_1, \dots, \alpha_{n+1}$. Let

$$(5.13) \quad \delta := \frac{\varepsilon}{(4L+1)(n+1)^2},$$

and let $\psi : \mathcal{D} \rightarrow A$ be a u.c.p. $(\mathcal{F}, \delta, \mathcal{G}' \cup \{u_{g_i}\}_{i=1}^n)$ -embedding, where (u_g) are the implementing unitaries for α . Let $\phi : \mathcal{D} \rightarrow B(\alpha) \subseteq M_{|G|}(A)$ be given by

$$(5.14) \quad \phi(d) := \bigoplus_{i=1}^{n+1} (\alpha_i \circ \psi)(d).$$

Clearly, ϕ will be (\mathcal{F}, δ) -multiplicative since each component is the composition of a $*$ -homomorphism (which are contractive) with a map which is (\mathcal{F}, δ) -multiplicative. Now for $d \in \mathcal{F}$ and $a = (a_{ij}) \in \mathcal{G}$, we have

$$\begin{aligned} \|\phi(d), (a_{ij})\| &\leq \sum_{i,j=1}^{n+1} \|\alpha_i(\psi(d))a_{ij} - a_{ij}\alpha_j(\psi(d))\| \\ &\leq \sum_{i,j=1}^{n+1} \|\alpha_i(\psi(d))a_{ij} - \psi(d)a_{ij}\| \\ (5.15) \quad &+ \|\psi(d)a_{ij} - a_{ij}\psi(d)\| + \|a_{ij}\psi(d) - a_{ij}\alpha_j(\psi(d))\| \\ &\leq \sum_{i,j=1}^{n+1} \|a_{ij}\| (\|\alpha_i(\psi(d)) - \psi(d)\| + \|\psi(d) - \alpha_j(\psi(d))\|) \\ &+ \|\psi(d), a_{i,j}\| \\ &< (n+1)^2(2L(\delta + \delta) + \delta) \\ &= (n+1)^2(4L+1)\delta = \varepsilon. \end{aligned}$$

Conversely, if the associated diagonal inclusion is \mathcal{D} -stable, we note that if $(x_k) \subseteq B(\text{id}_A, \alpha_{g_1}, \dots, \alpha_{g_n})$ is central for $M_{n+1}(A)$, writing

$$(5.16) \quad x_k = \bigoplus_{i=1}^{n+1} \alpha_i(a_k)$$

yields that $(a_k) \subseteq A$ is central for A and is asymptotically fixed by α_{g_i} , $i = 1, \dots, n$. In particular, if $\mathcal{D} \hookrightarrow B(\text{id}_A, \alpha_{g_1}, \dots, \alpha_{g_n})_\omega \cap (M_{n+1}(A))'$, then $\mathcal{D} \hookrightarrow A_\omega \cap (A \rtimes_\alpha G)'$. ■

Corollary 5.8 *Let $G \curvearrowright^\alpha A$ be an action of a finite group on a unital, separable C^* -algebra. Then $A \subseteq A \rtimes_\alpha G$ is \mathcal{D} -stable if and only if*

$$(5.17) \quad B(\alpha) \subseteq M_{|G|}(A)$$

is \mathcal{D} -stable.

6 Examples

6.1 Non-examples

We first start with some non-examples. Villadsen's C^* -algebras with perforation will be useful (see [68] for good exposition). Let $\mathcal{Q} = \bigotimes_n M_n$ denote the universal UHF C^* -algebra.

Theorem 6.1 ([64, 69]) *There exists a unital, simple, separable, nuclear C^* -algebra C satisfying the UCT such that $C \not\subseteq C \otimes \mathbb{Z}$ and C contains the universal UHF algebra unittally. Moreover, C is tracial and can be chosen to be AH with*

$$(6.1) \quad (K_0(C), K_0(C)^+, [1]_0, K_1(C)) = (\mathbb{Q}, \mathbb{Q}_+, 1, 0).$$

Corollary 6.2 *There exists an embedding $\mathcal{Q} \hookrightarrow \mathcal{Q}$ which is not \mathbb{Z} -stable. In particular, it is not \mathcal{Q} -stable.*

Proof Let C be as above. Note that $\mathcal{Q} \subseteq C$ so we must find an embedding $C \hookrightarrow \mathcal{Q}$. As C is unital, separable, exact, satisfies the UCT and has a faithful amenable trace (it has traces, and every such trace will be faithful and amenable since C is nuclear and simple) and there is clearly a morphism between K_0 -groups, [55, Theorem D] gives an embedding $C \hookrightarrow \mathcal{Q}$. Consequently, there is an embedding

$$(6.2) \quad \mathcal{Q} \hookrightarrow C \hookrightarrow \mathcal{Q}$$

which is not \mathcal{Q} -stable since there is an intermediate C^* -algebra C with $C \not\subseteq C \otimes \mathbb{Z}$. ■

Corollary 6.3 *There is an embedding $\mathbb{Z} \hookrightarrow \mathcal{Q}$ which is not \mathbb{Z} -stable.*

Proof Take C as above and take the chain of embeddings (noting that \mathcal{Q} is \mathbb{Z} -stable)

$$(6.3) \quad \mathbb{Z} \hookrightarrow \mathcal{Q} \otimes \mathbb{Z} \simeq \mathcal{Q} \hookrightarrow C \hookrightarrow \mathcal{Q}.$$

■

Corollary 6.4 *There is an embedding $\mathbb{Z} \hookrightarrow \mathcal{O}_2$ which is not \mathbb{Z} -stable.*

Proof Just take the same embedding as above together with an embedding $\mathcal{Q} \hookrightarrow \mathcal{O}_2$. ■

Remark 6.5 All $*$ -homomorphisms between strongly self-absorbing C^* -algebras are approximately unitarily equivalent by [66, Corollary 1.12], or even asymptotically unitarily equivalent by [16, Theorem 2.2]. Therefore, \mathcal{D} -stability is not closed under these equivalences (nor homotopy, see [16, Corollary 3.1]).

The only method we have used to show that an inclusion is not \mathcal{D} -stable is by finding an intermediate algebra which is not \mathcal{D} -stable. There are plenty of examples of stably finite C^* -algebras with perforation or higher-stable rank (in particular, non- \mathcal{L} -stable C^* -algebras [50]) [19, 27, 37, 60, 62–64, 68–70]. This gives rise to the following two questions.

- (1) Is there a unital inclusion $B \subseteq A$ of separable C^* -algebras such that whenever C is such that $B \subseteq C \subseteq A$, we have $C \simeq C \otimes \mathcal{D}$ but $B \subseteq A$ is not \mathcal{D} -stable? Is \mathcal{D} -stability equivalent to every intermediate C^* -algebra being \mathcal{D} -stable?
- (2) To get non-examples, we use stably finite C^* -algebras with perforation in between sufficiently regular C^* -algebras. Is there a way to do this for purely infinite C^* -algebras, or is finiteness the only obstruction? Thus we can ask: if \mathcal{D} is a purely infinite strongly self-absorbing C^* -algebra, is every embedding of \mathcal{D} into itself \mathcal{D} -stable? More specifically, if $B \subseteq A$ is a unital inclusion of simple, separable, purely infinite C^* -algebras, is the inclusion \mathcal{O}_∞ -stable?

Our third question asks if we can get non-examples arising from dynamical systems.

- (3) Is there a unital, separable \mathcal{D} -stable C^* -algebra and a (finite) group action $G \curvearrowright^\alpha A$ such that $A \rtimes_\alpha G$ is \mathcal{D} -stable, but the inclusion is not? One would need $A \rtimes_\alpha G$ to be \mathcal{D} -stable for non-dynamical reasons.

6.2 Cyclically permuting tensor powers

Here, we give a dynamical example to illustrate the discussion in Section 5. In particular, we can look at a consequence of Corollary 3.5.

Example 6.6 Let $p, q \in \mathbb{N}$ be coprime and consider the q th tensor power of the UHF algebra $A = M_{p^\infty}^{\otimes q}$. Let us examine the action $\mathbb{Z}_q \curvearrowright^\sigma A$ given by cyclically permuting the tensors:

$$(6.4) \quad \sigma(a_1 \otimes \cdots \otimes a_q) = a_2 \otimes \cdots \otimes a_q \otimes a_1.$$

One can prove directly or use [26] or [1] in order to conclude that this action has the weak tracial Rokhlin property, or that this action is \mathcal{L} -equivariantly absorbing, and consequently that $A^\sigma \subseteq A \rtimes_\sigma \mathbb{Z}_q$ is \mathcal{L} -stable.

Alternatively, one can use techniques similar to [24], [25], or [28] in order to compute the K -theory of the fixed point algebra A^σ to be

$$(6.5) \quad K_0((M_{p^\infty}^{\otimes q})^\sigma) \simeq \lim_{\rightarrow} \left(\mathbb{Z}^q, \begin{pmatrix} p + \frac{p^q - p}{q} & \frac{p^q - p}{q} & \cdots & \frac{p^q - p}{q} \\ \frac{p^q - p}{q} & p + \frac{p^q - p}{q} & \cdots & \frac{p^q - p}{q} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{p^q - p}{q} & \frac{p^q - p}{q} & \cdots & p + \frac{p^q - p}{q} \end{pmatrix} \right),$$

from which one can show that $K_0(A^\sigma)$ is p -divisible. Then using the fact that $K_0(A^\sigma)$ is p -divisible and A^σ is AF, it follows that A^σ is M_{p^∞} -stable. Using Corollary 3.5, we then see that $M_{p^\infty} \hookrightarrow (A^\sigma)_\omega \cap A'$. In particular, we have that $A^\sigma \subseteq A \rtimes_\sigma \mathbb{Z}_q$ is M_{p^∞} -stable (since clearly if this embedding is fixed by \mathbb{Z}_q , it will commute with the implementing unitaries as well).

Example 6.7 Following up on the previous example, if we consider the embedding

$$(6.6) \quad B := \left\{ \begin{pmatrix} x & & & \\ & \sigma(x) & & \\ & & \ddots & \\ & & & \sigma^{q-1}(x) \end{pmatrix} \mid x \in M_{p^\infty}^{\otimes q} \right\} \subseteq M_q(M_{p^\infty}^{\otimes q}) := A,$$

then $B \subseteq A$ is M_{p^∞} -stable by Proposition 5.7.

6.3 The canonical inclusion of the CAR algebra in \mathcal{O}_2

Example 6.8 Let $\mathcal{O}_2 = C^*(s_1, s_2)$ be the Cuntz algebra generated by two isometries [15], and consider the inclusion

$$(6.7) \quad M_{2^\infty} \simeq \overline{\text{span}}\{s_\mu s_\nu^* \mid |\mu| = |\nu|\} \subseteq \mathcal{O}_2,$$

where for a word $\mu = \{i_1, \dots, i_p\} \in \{1, 2\}^p$, $s_\mu = s_{i_1} \dots s_{i_p}$. This copy of the CAR algebra is precisely the fixed point subalgebra of the gauge action (see [45]). Consider the endomorphism $\lambda : \mathcal{O}_2 \rightarrow \mathcal{O}_2$ given by

$$(6.8) \quad \lambda(x) := s_1 x s_1^* + s_2 x s_2^*.$$

We note that a sequence $(x_n)_{n \in \mathbb{N}}$ is ω -asymptotically central for \mathcal{O}_2 if and only if it is ω -asymptotically fixed by λ . Indeed, if $(x_n)_{n \in \mathbb{N}}$ is central, then $\|\lambda(x_n) - x_n\| \xrightarrow{n \rightarrow \omega} 0$ since $[x_n, s_i] \rightarrow 0$ for $i = 1, 2$. On the other hand, if $(x_n)_{n \in \mathbb{N}}$ is asymptotically fixed by λ , then the inequalities

$$(6.9) \quad \begin{aligned} \|s_i x_n - x_n s_i\| &= \|s_1 x_n s_1^* s_i + s_2 x_n s_2^* s_i - x_n s_i\| \leq \|\lambda(x_n) - x_n\| \|s_i\| \\ \|s_i^* x_n - x_n s_i^*\| &= \|s_i^* x_n - s_i^* s_1 s_1^* - s_i^* s_2 s_2^* s_i\| \leq \|s_i^*\| \|\lambda(x_n) - x_n\| \end{aligned}$$

imply that $(x_n)_{n \in \mathbb{N}}$ is asymptotically central.

We note that $\lambda|_{M_{2^\infty}}$ is the forward tensor shift if we identify $M_{2^\infty} = \bigotimes_{\mathbb{N}} M_2$ (see, for example, [17, Section V.4]). Now [5] gives an embedding $\xi : M_2 \hookrightarrow (M_{2^\infty})_\omega$ such that $\lambda_\omega \circ \xi = \xi$. In particular, $M_{2^\infty} \hookrightarrow (M_{2^\infty})_\omega \cap \mathcal{O}'_2$ so that this inclusion is M_{2^∞} -stable.

Thinking of \mathcal{O}_2 as the semigroup crossed product $\mathcal{O}_2 \simeq M_{2^\infty} \rtimes_{\lambda} \mathbb{N}$ (see [48, 51]), any intermediate C^* -algebra is automatically CAR stable. Consequently, each intermediate subalgebra $M_{2^\infty} \rtimes d\mathbb{N} = C^*(M_{2^\infty}, s_1^d)$ is M_{2^∞} -stable. We can do this all concurrently.

Corollary 6.9 *There exists an isomorphism $\Phi : \mathcal{O}_2 \simeq \mathcal{O}_2 \otimes M_{2^\infty}$ such that*

$$(6.10) \quad \Phi(C^*(M_{2^\infty}, s_1^d)) \simeq C^*(M_{2^\infty}, s_1^d) \otimes M_{2^\infty}$$

for all $d \in \mathbb{N}$. The same holds if we replace M_{2^∞} by \mathbb{Z} .

Now let us play with some diagonal inclusions associated with powers of the Bernoulli shift λ on \mathcal{O}_2 above. This will be similar to what was discussed in Section 5.2, except we allow endomorphisms.

Example 6.10 Consider, for $n \in \mathbb{N}$, the diagonal inclusion

$$(6.11) \quad B_n := \left\{ \begin{pmatrix} x & & & \\ & \lambda(x) & & \\ & & \ddots & \\ & & & \lambda^{n-1}(x) \end{pmatrix} \mid x \in \mathcal{O}_2 \right\} \subseteq M_n(\mathcal{O}_2) =: A_n.$$

Note that both A_n and B_n are isomorphic to \mathcal{O}_2 , and in fact, this gives a nontrivial inclusion of \mathcal{O}_2 into itself which is \mathcal{O}_2 -stable. This is \mathcal{O}_2 -stable since a sequence is asymptotically fixed by λ if and only if it asymptotically commutes with the algebra. A similar argument to that of Proposition 5.7 will yield that this inclusion is \mathcal{O}_2 -stable.

One can even restrict the diagonal to elements of the CAR algebra $M_{2^\infty} \subseteq \mathcal{O}_2$ sitting as the fixed point subalgebra of the gauge action as above.

Example 6.11 Consider

$$(6.12) \quad B_n^{(2)} := \left\{ \begin{pmatrix} x & & & \\ & \lambda(x) & & \\ & & \ddots & \\ & & & \lambda^{n-1}(x) \end{pmatrix} \mid x \in M_{2^\infty} \right\} \subseteq M_n(\mathcal{O}_2) = A_n.$$

This is M_{2^∞} -stable for the same reasons as above. This gives another inclusion $M_{2^\infty} \simeq B_n^{(2)} \subseteq M_n(\mathcal{O}_2) \simeq \mathcal{O}_2$ which is CAR-stable.

Acknowledgments Many thanks to my Ph.D. supervisors Thierry Giordano and Aaron Tikuisis for many helpful discussions, as well as to Eusebio Gardella for helpful comments.

References

- [1] M. Amini, N. Golestani, S. Jamali, and N. Christopher Phillips, *Group actions on simple tracially \mathcal{Z} -absorbing C^* -algebras*. Preprint, 2022. [arXiv:2204.03615](https://arxiv.org/abs/2204.03615)
- [2] D. H. Bisch, *On the existence of central sequences in subfactors*. Trans. Amer. Math. Soc. 321(1990), no. 1, 117–128. MR 1005075
- [3] D. H. Bisch, *Central sequences in subfactors. II*. Proc. Amer. Math. Soc. 121(1994), no. 3, 725–731. MR 1209417
- [4] J. Bosa, J. Gabe, A. Sims, and S. White, *The nuclear dimension of \mathcal{O}_∞ -stable C^* -algebras*. Adv. Math. 401(2022), Article no. 108250, 51 pp. MR 4392219
- [5] O. Bratteli, E. Størmer, A. Kishimoto, and M. Rørdam, *The crossed product of a UHF algebra by a shift*. Ergodic Theory Dynam. Systems 13(1993), no. 4, 615–626. MR 1257025
- [6] N. P. Brown and N. Ozawa, *C^* -algebras and finite-dimensional approximations*, Graduate Studies in Mathematics, 88, American Mathematical Society, Providence, RI, 2008. MR 2391387
- [7] R. D. Burstein, *Commuting square subfactors and central sequences*. Internat. J. Math. 21(2010), no. 1, 117–131. MR 2642989
- [8] J. Cameron and R. R. Smith, *A Galois correspondence for reduced crossed products of simple C^* -algebras by discrete groups*. Canad. J. Math. 71(2019), no. 5, 1103–1125. MR 4010423
- [9] J. R. Carrión, J. Gabe, C. Schafhauser, A. Tikuisis, and S. White, *Classifying $*$ -homomorphisms I: Unital simple nuclear C^* -algebras*. Preprint, 2023. [arXiv:2307.06480](https://arxiv.org/abs/2307.06480)
- [10] M. D. Choi and E. G. Effros, *The completely positive lifting problem for C^* -algebras*. Ann. of Math. (2) 104(1976), no. 3, 585–609. MR 417795
- [11] A. Connes, *Classification of automorphisms of hyperfinite factors of type II_1 and II_∞ and application to type III factors*. Bull. Amer. Math. Soc. 81(1975), no. 6, 1090–1092. MR 388117
- [12] A. Connes, *Outer conjugacy classes of automorphisms of factors*. Ann. Sci. École Norm. Sup. (4) 8(1975), no. 3, 383–419. MR 394228

- [13] A. Connes, *Classification of injective factors cases II_1 , II_∞ , III_λ , $\lambda \neq 1$* . Ann. of Math. (2) 104(1976), no. 1, 73–115. MR 454659
- [14] A. Connes, *Periodic automorphisms of the hyperfinite factor of type II_1* . Acta Sci. Math. (Szeged) 39(1977), nos. 1–2, 39–66. MR 448101
- [15] J. Cuntz, *Simple C^* -algebras generated by isometries*. Comm. Math. Phys. 57(1977), no. 2, 173–185. MR 467330
- [16] M. Dadarlat and W. Winter, *On the KK-theory of strongly self-absorbing C^* -algebras*. Math. Scand. 104(2009), no. 1, 95–107. MR 2498373
- [17] K. R. Davidson, *C^* -algebras by example*, Fields Institute Monographs, 6, American Mathematical Society, Providence, RI, 1996. MR 1402012
- [18] S. Echterhoff and M. Rørdam, *Inclusions of C^* -algebras arising from fixed-point algebras*. Groups Geom. Dyn. 18 (2024), no. 1, pp. 127–145
- [19] G. A. Elliott and J. Villadsen, *Perforated ordered K_0 -groups*. Canad. J. Math. 52(2000), no. 6, 1164–1191. MR 1794301
- [20] D. Enders, A. Schemaitat, and A. Tikuisis, *Corrigendum to “K-theoretic characterization of C^* -algebras with approximately inner flip”*. Preprint, 2023. [arXiv:2303.11106](https://arxiv.org/abs/2303.11106)
- [21] J. Gabe, *Classification of \mathcal{O}_∞ -stable C^* -algebras*. Mem. Amer. Math. Soc. 293 (2024), no. 1461, v+115 pp.
- [22] J. Gabe, *A new proof of Kirchberg’s ${}_2$ -stable classification*. J. Reine Angew. Math. 761(2020), 247–289. MR 4080250
- [23] E. Gardella and I. Hirshberg, *Strongly outer actions of amenable groups on \mathbb{Z} -stable C^* -algebras*. Preprint, 2018. [arXiv:1811.00447](https://arxiv.org/abs/1811.00447)
- [24] D. Handelman and W. Rossmann, *Product type actions of finite and compact groups*. Indiana Univ. Math. J. 33(1984), no. 4, 479–509. MR 749311
- [25] D. Handelman and W. Rossmann, *Actions of compact groups on AF C^* -algebras*. Illinois J. Math. 29(1985), no. 1, 51–95. MR 769758
- [26] I. Hirshberg and J. Orovitz, *Tracially \mathbb{Z} -absorbing C^* -algebras*. J. Funct. Anal. 265(2013), no. 5, 765–785. MR 3063095
- [27] I. Hirshberg, M. Rørdam, and W. Winter, *${}_0(X)$ -algebras, stability and strongly self-absorbing C^* -algebras*. Math. Ann. 339(2007), no. 3, 695–732. MR 2336064
- [28] I. Hirshberg and W. Winter, *Rokhlin actions and self-absorbing C^* -algebras*. Pacific J. Math. 233(2007), no. 1, 125–143. MR 2366371
- [29] W. Hurewicz and H. Wallman, *Dimension theory*, Princeton Mathematical Series, 4, Princeton University Press, Princeton, NJ, 1941. MR 0006493
- [30] M. Izumi, *Inclusions of simple C^* -algebras*. J. Reine Angew. Math. 547(2002), 97–138.
- [31] M. Izumi, *Finite group actions on C^* -algebras with the Rohlin property*. I. Duke Math. J. 122(2004), no. 2, 233–280. MR 2053753
- [32] X. Jiang and H. Su, *On a simple unital projectionless C^* -algebra*. Amer. J. Math. 121(1999), no. 2, 359–413. MR 1680321
- [33] Y. Kawahigashi, C. E. Sutherland, and M. Takesaki, *The structure of the automorphism group of an injective factor and the cocycle conjugacy of discrete abelian group actions*. Acta Math. 169(1992), nos. 1–2, 105–130. MR 1179014
- [34] K. Kawamuro, *Central sequence subfactors and double commutant properties*. Internat. J. Math. 10(1999), no. 1, 53–77. MR 1678538
- [35] E. Kirchberg, *Central sequences in C^* -algebras and strongly purely infinite algebras*. In: Operator Algebras: The Abel Symposium 2004, vol. 1, Springer, Berlin, 2006, pp. 175–231. MR 2265050
- [36] E. Kirchberg and N. C. Phillips, *Embedding of exact C^* -algebras in the Cuntz algebra \mathcal{O}_2* . J. Reine Angew. Math. 525(2000), 17–53. MR 1780426
- [37] F. de Lacerda Mortari, *Tracial state spaces of higher stable rank simple C^* -algebras*, ProQuest LLC, Ann Arbor, MI, 2009, Thesis (Ph.D.)—University of Toronto (Canada). MR 2753146
- [38] H. H. Lee and H. Osaka, *On permanence of regularity properties*. J. Topol. Anal. (2023).
- [39] H. Matui and Y. Sato, *Strict comparison and \mathbb{Z} -absorption of nuclear C^* -algebras*. Acta Math. 209(2012), no. 1, 179–196. MR 2979512
- [40] D. McDuff, *Uncountably many II_1 factors*. Ann. of Math. (2) 90(1969), 372–377. MR 259625
- [41] F. J. Murray and J. von Neumann, *On rings of operators IV*. Ann. of Math. (2) 44(1943), 716–808. MR 9096
- [42] H. Osaka and T. Teruya, *The Jiang–Su absorption for inclusions of unital C^* -algebras*. Canad. J. Math. 70(2018), no. 2, 400–425. MR 3759005
- [43] S. Popa, *Sousfacteurs, actions des groupes et cohomologie*. C. R. Acad. Sci. Paris Sér. I Math. 309(1989), no. 12, 771–776. MR 1054961

- [44] S. Popa, *On the relative Dixmier property for inclusions of C^* -algebras*. J. Funct. Anal. 171(2000), no. 1, 139–154. MR 1742862
- [45] I. Raeburn, *Graph algebras*, CBMS Regional Conference Series in Mathematics, 103, Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2005. MR 2135030
- [46] M. Rørdam, *On the structure of simple C^* -algebras tensored with a UHF-algebra*. J. Funct. Anal. 100(1991), no. 1, 1–17. MR 1124289
- [47] M. Rørdam, *On the structure of simple C^* -algebras tensored with a UHF-algebra. II*. J. Funct. Anal. 107(1992), no. 2, 255–269. MR 1172023
- [48] M. Rørdam, *Classification of certain infinite simple C^* -algebras*. J. Funct. Anal. 131(1995), no. 2, 415–458. MR 1345038
- [49] M. Rørdam, *Classification of nuclear, simple C^* -algebras*. In: J. Cuntz and V. Jones (eds.), *Classification of nuclear C^* -algebras. Entropy in operator algebras*, Encyclopaedia Mathematical Sciences, 126, Springer, Berlin, 2002, pp. 1–145. MR 1878882
- [50] M. Rørdam, *The stable and the real rank of \mathbb{Z} -absorbing C^* -algebras*. Internat. J. Math. 15(2004), no. 10, 1065–1084. MR 2106263
- [51] M. Rørdam, *Irreducible inclusions of simple C^* -algebras*. Enseign. Math. 69(2023), nos. 3–4, 275–314. MR 4599249
- [52] M. Rørdam and W. Winter, *The Jiang–Su algebra revisited*. J. Reine Angew. Math. 642(2010), 129–155. MR 2658184
- [53] J. Rosenberg and C. Schochet, *The Künneth theorem and the universal coefficient theorem for Kasparov’s generalized K -functor*. Duke Math. J. 55(1987), no. 2, 431–474. MR 894590
- [54] Y. Sato, *The Rohlin property for automorphisms of the Jiang–Su algebra*. J. Funct. Anal. 259(2010), no. 2, 453–476. MR 2644109
- [55] C. Schafhauser, *Subalgebras of simple AF-algebras*. Ann. of Math. (2) 192(2020), no. 2, 309–352. MR 4151079
- [56] A. Schemaitat, *The Jiang–Su algebra is strongly self-absorbing revisited*. J. Funct. Anal. 282(2022), no. 6, Article no. 109347, 39 pp. MR 4360358
- [57] C. E. Sutherland and M. Takesaki, *Actions of discrete amenable groups on injective factors of type III_λ , $\lambda \neq 1$* . Pacific J. Math. 137(1989), no. 2, 405–444. MR 990219
- [58] Y. Suzuki, *Equivariant \mathcal{O}_2 -absorption theorem for exact groups*. Compos. Math. 157(2021), no. 7, 1492–1506. MR 4275465
- [59] G. Szabó, *Equivariant Kirchberg–Phillips-type absorption for amenable group actions*. Comm. Math. Phys. 361(2018), no. 3, 1115–1154. MR 3830263
- [60] A. Tikuisis, *Regularity for stably projectionless, simple C^* -algebras*. J. Funct. Anal. 263(2012), no. 5, 1382–1407. MR 2943734
- [61] A. Tikuisis, *K -theoretic characterization of C^* -algebras with approximately inner flip*. Int. Math. Res. Not. IMRN 2016(2016), no. 18, 5670–5694. MR 3567256
- [62] A. S. Toms, *On the independence of K -theory and stable rank for simple C^* -algebras*. J. Reine Angew. Math. 578(2005), 185–199. MR 2113894
- [63] A. S. Toms, *An infinite family of non-isomorphic C^* -algebras with identical K -theory*. Trans. Amer. Math. Soc. 360(2008), no. 10, 5343–5354. MR 2415076
- [64] A. S. Toms, *On the classification problem for nuclear C^* -algebras*. Ann. of Math. (2) 167(2008), no. 3, 1029–1044. MR 2415391
- [65] A. S. Toms, S. White, and W. Winter, *\mathbb{Z} -stability and finite-dimensional tracial boundaries*. Int. Math. Res. Not. IMRN 2015(2015), no. 10, 2702–2727. MR 3352253
- [66] A. S. Toms and W. Winter, *Strongly self-absorbing C^* -algebras*. Trans. Amer. Math. Soc. 359(2007), no. 8, 3999–4029. MR 2302521
- [67] A. S. Toms and W. Winter, *\mathbb{Z} -stable ASH algebras*. Canad. J. Math. 60(2008), no. 3, 703–720. MR 2414961
- [68] A. S. Toms and W. Winter, *The Elliott conjecture for Villadsen algebras of the first type*. J. Funct. Anal. 256(2009), no. 5, 1311–1340. MR 2490221
- [69] J. Villadsen, *Simple C^* -algebras with perforation*. J. Funct. Anal. 154(1998), no. 1, 110–116. MR 1616504
- [70] J. Villadsen, *On the stable rank of simple C^* -algebras*. J. Amer. Math. Soc. 12(1999), no. 4, 1091–1102. MR 1691013
- [71] W. Winter, *Strongly self-absorbing C^* -algebras are \mathbb{Z} -stable*. J. Noncommut. Geom. 5(2011), no. 2, 253–264. MR 2784504

Department of Pure Mathematics, University of Waterloo, Waterloo, ON N2L 3G1, Canada

e-mail: psarkowi@uwaterloo.ca