

The Journal of Laryngology and Otology

EDITED BY
G. H. BATEMAN

ASSISTANT EDITOR
LIONEL TAYLOR

Contents

Hereditary conductive deafness due to incus-stapes abnormalities and associated with pinna deformity	T. J. WILMOT	469 ✓
A survey of hearing by evoked response audiometry in a group of normally hearing school-children	H. A. BEAGLEY and SUSAN E. KELLOGG	481 ✓
Familial congenital moderate neural hearing loss	BRUCE W. KONIGSMARK, MARVIN C. MENGEL and HARRIET HASKINS	495 ✓
Ageing of the otosclerotic focus	GEORGE KELEMEN	507 ✓
Experiences with SAL audiometry	GEORGE RÉVÉSZ	515 ✓
Clinical records—		
The Papousek cradle: a device for measuring babies' head movement responses to auditory stimulation	R. J. BENCH and K. P. MURPHY	521 ✓
A case of primary localized amyloidosis in both tonsils	H. ELLERMANN ERIKSEN	525 ✓
Osteomyelitis of maxilla with extrusion of teeth in the floor of the nose requiring extraction	S. R. REGE, K. L. SHAH and P. T. MARFATIA	533 ✓
Ectopic supernumerary tooth in the nasal cavity	G. S. KOHLI and P. L. VERMA	537 ✓
An unusual foreign body in the nose	CHANDER MALHOTRA, M. M. L. ARORA and Y. N. MEHRA	539 ✓
General notes		

Headley Brothers Ltd

Ashford, Kent

The Journal of Laryngology and Otology

(FOUNDED in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY

G. H. BATEMAN

ASSISTANT EDITOR

LIONEL TAYLOR

1. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the understanding that they are contributed to this *Journal* solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgement must be made.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and sent at the same time as original photographs, etc.

The Harvard system of recording references should be used, e.g. GREEN, C., and BROWN, D. (1951), *Journal of Laryngology and Otology*, 65, 33. The titles of all Journals should be given without abbreviation.

It is most important that authors should verify personally the accuracy of every reference before submitting a paper for publication.

3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.

4. Illustration blocks will normally be held by the Printers for three years, after which they will be destroyed. Any author is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to HEADLEY BROTHERS LTD., The Invicta Press, Ashford, Kent. Coloured illustrations will be charged to authors, unless a special grant is authorized by the Editor.

5. Orders for reprints must be sent when returning galley proofs, and for this purpose special forms are supplied.

6. Editorial communications may be addressed to THE EDITOR, *Journal of Laryngology*, c/o HEADLEY BROTHERS LTD., The Invicta Press, Ashford, Kent.

7. The annual subscription is £8 (eight pounds sterling) post free; U.S.A. \$23 post free, and payable in advance.

8. Single copies of current or back numbers (when available) will be on sale at £1 (including postage) each.


9. All subscriptions, advertising and business communications should be sent to the publishers, or subscription agents.

HEADLEY BROTHERS LTD.

The Invicta Press, ASHFORD, Kent.

© *Journal of Laryngology and Otology Ltd.*, 1970

Thought reader!!



With the complex tests that are being done today clinical audiometry can become altogether too complicated.

By switching only one knob, the model 103 Clinical Audiometer will do the thinking for you, and enable you to select up to 18 different functions:

- Pure Tone Air Conduction
- Pure Tone Bone Conduction
- Narrow Band Masking
- Wide Band (White Noise) Masking
- Alternative Binaural Loudness Balance
- Simultaneous Binaural Loudness Balance
- Speech, Live Voice
- Speech, Recorded
- Speech with Masking
- SISI Test
- Lüscher DL Test
- Audiometer Weber Test
- SAL Test
- Rainville Test
- Shifting Voice
- Stenger Test
- Modified Stenger Test
- Tone Decay Test

The model 103 is functional engineering with the operator in mind. We believe the audiometer should do the thinking ... and this one does just that.


To: Amplivox Hearing Conservation Ltd.,
Beresford Avenue, Wembley, Middlesex.
Tel: 01-902 8991

Name _____

Address _____

- Please send demonstrator
 Please send leaflet

JL05



PETERS CLINIC AUDIOMETERS AND BÉKÉSY
ARE THE IDEAL TOOLS FOR MORE ADVANCED AUDIOMETRY

The Audiometers offer :

- Two continuously variable attenuators accurate to 1 db.
- SISI Test (built-in) and automatic alternate loudness balance switching.
- Maximum attenuation of 120 db for loudness discomfort level tests.
- Narrow band SYNCHRONOUS masking (a new semi-automatic masking technique).
- Full speech test facilities with white noise masking.
- Plug-in circuit boards and operator output checks.
- Conversion to a true BÉKÉSY AUDIOMETER at will.
- Extra features not listed are offered by the AP/6 compared to the AP/5 AUDIOMETER.

write for full details to:

ALFRED PETERS AND SONS LTD
GELL STREET SHEFFIELD 3

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

What clinicians say about ORBENIN

Clinicians have frequently testified to the success of Orbenin in the treatment of Gram-positive infections. World Literature now contains over 400 references to Orbenin.

Pneumonia Lung Abscess Tonsillitis Pharyngitis Acute bronchitis

Septicaemia

Boils Carbuncles Infected dermatoses

Osteomyelitis Osteitis

Post-operative wound infections Burns Skin graft protection

Acute endocarditis

Staphylococcal enterocolitis Staphylococcal urinary tract infections
Staphylococcal meningitis

Dosage

Adults: 500mg (2 capsules) four times daily, by mouth. 250mg (1 vial) by intramuscular injection, four times daily.

Children: Under 2 years—quarter adult dose. 2-10 years—half adult dose.

In severe infections the dosage may be increased.

Contra-indications

Orbenin should not be given to patients with a penicillin allergy or administered by subconjunctival injection.

Side-effects

As with other penicillins.

Additional information is available on request.



Orbenin (Regd.) (cloxacillin sodium BP) is a product of British research at **Beecham Research Laboratories** Brentford, England, originators of the new penicillins.



A personal announcement by Stanley Ingram

(formerly of Ingrams Hearing Aids)



I am now an independent hearing aid dispenser with no link whatsoever with any manufacturer and can offer a very wide range of aids, and quite impartial advice, to each patient.

I shall be pleased to see personally each patient referred and will undertake to supply a detailed report to the otologist both at the start and the conclusion of the trial period.

My rooms are centrally located in Marylebone Road, a few doors from Baker Street station. Domiciliary visits can be arranged if preferred.

STANLEY INGRAM 146 Marylebone Road, London NW1
Telephone 01-486 3638

Please mention *The Journal of Laryngology and Otology* when replying to advertisements



A Product from the Sea for E.N.T. Surgeons

*calgitex

Calgitex calcium alginate E.N.T. Gauze is made from highly purified specially processed calcium alginate, derived from seaweed. This product is used extensively by E.N.T. surgeons throughout the world, it has a slow absorption rate, and can be easily removed from the nose without effecting or damaging the tissue.

In the field of aural surgery Calgitex is highly suitable for packing procedures due to its solubility, soft texture, and discouragement of haemorrhage.

Packed in vials containing one metre of ribbon gauze.
Obtainable from your usual Surgical or Pharmaceutical Supplier


MEDICAL ALGINATES LTD.

Wadsworth Road, Perivale, Middlesex. Tel. 01-997 4441

A member of the Optrex Group

*Reg'd Trade Mark

Please mention *The Journal of Laryngology and Otology* when replying to advertisements



5 drops
to remove
ear wax
easily
rapidly
safely

CERUMOL[®]


Active ingredients per 100 ml.
p-dichlorobenzene B.P.C. 2 G., Benzocaine B.P. 2.7 G.,
Chlorbutol B.P. 5 G., Ol. Terebinth 10 ml.

In convenient dropper vial for surgery
use. Also in 2 oz. Hospital pack.

 **LABORATORIES FOR APPLIED BIOLOGY LTD**
91 Amhurst Park London N 16
Tel: STAmford Hill 2252

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

Tegretol® controls seizures without sedation *and* improves behaviour and mood



Geigy

Tegretol™ is available as white, scored, compressed tablets containing 5-Carbamoyl-5H-dibenz (b, f) azepine (Carbamazepine) 200 mg.

Detailed literature describing any Geigy product will be supplied on request.

Geigy (U.K.) Limited
Pharmaceuticals Division,
Macclesfield, Cheshire.

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

THE LARYNGOSCOPE

A Monthly Journal
devoted to the disease of
EAR, NOSE AND THROAT

*Official organ for the American Laryngological
Rhinological and Otological Society*

Price \$22.00 per year Canada \$23.00 per year
Foreign \$23.00 per year

ESTABLISHED 1896

THEODORE E. WALSH, M.D.
EDITOR

222 PINE LAKE ROAD, COLLINSVILLE,
ILLINOIS 62234


For advertisement space in this Journal

apply to

HEADLEY BROTHERS LTD.

Ashford, Kent

Please mention *The Journal of Laryngology and Otology* when replying to advertisements



the drop that soothes and clears infected ears

soothes by reducing inflammation with hydrocortisone
clears by rapidly destroying virtually all bacteria commonly
found in ear infections

*
OTOSPORIN drops contain polymyxin B sulphate, neomycin sulphate
and hydrocortisone in a bland suspension designed to facilitate penetration.
Supplied in dropper bottles of 5 ml. Full information is available on request.

*Trade Mark


OTOSPORIN



Burroughs Wellcome & Co.
(The Wellcome Foundation Ltd.) Dartford, Kent

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

New for the infected ear OTOSEPTIL eardrops



Containing a broad spectrum antibiotic and possessing antifungal properties

Candida and aspergillus are found in about 20% of cases of otitis externa.

Otoseptil ear drops destroy both pathogenic bacteria and fungi commonly found in ear infections.

Otoseptil ear drops contain:

Neomycin Undecylenate – antibacterial and antifungal.

Tyrosin – active against gram-positive cocci.
(Containing gramicidin and tyrocin).

Hydrocortisone alcohol – anti-inflammatory.

Ethylene oxide polyoxypropylene glycol condensate –
to ensure rapid tissue contact.

Otoseptil ear drops control infection, relieve pain, erythema and itching.

Neomycin Undecylenate was researched by H R Napp Ltd.

Presentation : 8 ml bottle with separate dropper.

Samples and full information available to the medical profession upon request.
H R Napp Ltd Watford WD2 7RA England



A new product by Nepp—Developed by International Research Group. Dedicated to Physician and Patient for over 40 years.