

A NEW CHARACTERIZATION OF THE BRUHAT DECOMPOSITION

YOSHIFUMI KATO

§ 0. Introduction

By an algebraic homogeneous space, we mean the factor space $X = G/P$, where G is a simply-connected, complex, semi-simple Lie group and P is a parabolic subgroup of G . Many typical manifolds such as the projective spaces and the Grassmann varieties belong to this class of manifolds. For instance, the Grassmann variety $G(k, n)$ can be expressed as $SL(n+1, C)/P$, where P is a maximal parabolic subgroup of $SL(n+1, C)$ leaving a suitable $k+1$ dimensional subspace invariant. In this paper, we devote ourselves to study the Bruhat decomposition of an algebraic homogeneous space $X = G/P$. For that purpose, in the first place, we construct a holomorphic vector field V_H on X for an element H of the positive Weyl chamber as follows

$$(V_H f)(\bar{g}) = \lim_{\varepsilon \rightarrow 0} \frac{f(\exp(\varepsilon H)\bar{g}) - f(\bar{g})}{\varepsilon},$$

where $\bar{g} \in X$ and f is a local function around \bar{g} . Then the vector field V_H vanishes only at the isolated points and in fact at the quotient set $W^1 = W/W_1$, which is naturally embedded into $X = G/P$ as the set of all fixed points of the left T action. Here W and W_1 are the Weyl groups associated to G and P respectively. And there is a nice coordinate neighborhood $(w\bar{N}^*, \phi^{-1} \circ \psi^{-1} \circ w^{-1})$ around each $w \in W^1$ such that;

1) The set $w\bar{N}^*$ is a T invariant Zariski open set. Let w_0 be the element of W whose length $\ell(w_0)$ is maximal among all. Then the set $w_0\bar{N}^*$ coincides with the cell of maximal dimension appeared in the Bruhat decomposition of X .

2) The set $w\bar{N}^*$ is mapped by $\phi^{-1} \circ \psi^{-1} \circ w^{-1}$ onto the Lie algebra \mathfrak{n}^* of the Lie group N^* and hence we can write as

Received November 26, 1979.

$$\phi^{-1} \circ \psi^{-1} \circ \iota v^{-1}(\iota v \bar{n}^*) = \sum_{\alpha \in \mathcal{A}(\mathfrak{n}^*)} z_\alpha(\iota v \bar{n}^*) X_\alpha.$$

3) With respect to the local coordinate $\{z_\alpha(\iota v \bar{n}^*)\}_{\alpha \in \mathcal{A}(\mathfrak{n}^*)}$, the vector field V_H can be written explicitly as follows

$$V_H = \sum_{\alpha \in \mathcal{A}(\mathfrak{n}^*)} (\iota v \alpha)(H) z_\alpha \frac{\partial}{\partial z_\alpha}.$$

From the above expression of V_H on $\iota v \bar{N}^*$, any flow $P(t)$, parametrized by $t \in \mathbb{C}$, of V_H on $\iota v \bar{N}^*$ can be written as $z_\alpha(P(t)) = c_\alpha e^{(\iota v \alpha)(H)t}$, $c_\alpha \in \mathbb{C}$, for $\alpha \in \mathcal{A}(\mathfrak{n}^*)$.

Every algebraic homogeneous space X has the distinguished cellular decomposition $X = \bigcup_{\mathfrak{w} \in W^1} X_{\mathfrak{w}}$ parametrized by W^1 , which is called the Bruhat one of X . We recall the definition precisely in section 2. The new point of view in this note is to understand that any Bruhat cell $X_{\mathfrak{w}}$, $\mathfrak{w} \in W^1$, consists of some flows $P(t)$ of V_H . We embed the space X into a certain projective space \mathbf{P}^N and construct a vector field V_H^A on \mathbf{P}^N whose restriction to X coincides with the vector field V_H . And we indentify the flow $P(t)$ as that of V_H^A lying entirely on X and investigate the behavior of it. As a result we obtain a rule to seek for the limit points $\lim_{t \rightarrow \pm\infty} P(t)$. And we show that to compactify $X_{\mathfrak{w}}$ to $\bar{X}_{\mathfrak{w}}$, it is enough to attach such limit points $\lim_{t \rightarrow \pm\infty} P(t)$ successively. The set $\bar{X}_{\mathfrak{w}}$ is a subvariety of X and is called the generalized Schubert variety. The family of subvarieties $\bar{X}_{\mathfrak{w}}$, $\mathfrak{w} \in W^1$, has a lot of important meanings for the geometry of $X = G/P$. For instance, the homology classes of $\{X_{\mathfrak{w}}\}_{\mathfrak{w} \in W^1}$ form the free \mathbb{Z} -basis of the integral homology group $H_*(X, \mathbb{Z})$ of X . In case $X = \text{Gr}(k, n)$, the classical Schubert calculus is to study the ring structure of $H_*(X, \mathbb{Z})$ by using the basis.

Further we give the following results in Theorem 4.9 and Theorem 4.12.

- 1) We give a new geometrical meaning to the reflections S_α , $\alpha \in \mathcal{A}$.
- 2) We obtain a simple necessary sufficient condition to determine the Bruhat ordering \leq on W^1 .

3) From 1) and 2), we clarify what Bruhat cells of lower dimension should be attached to the boundary of $X_{\mathfrak{w}}$ of the appointed direction to compactify $X_{\mathfrak{w}}$.

It seems interesting to study the singularities of $\bar{X}_{\mathfrak{w}}$ as a sequel of this note. And we also wish to clarify the relations between our results and the representation theory of nilpotent Lie algebras.

This paper is divided into four parts.

1. Semisimple Lie algebras.
2. Bruhat decomposition of G/P .
3. Vector field V_H .
4. Proof of our theorems.

The author wishes to express his thanks to Profs. H. Morikawa, K. Aomoto and H. Umemura for their valuable suggestions.

§1. Semisimple Lie algebras

First we fix the terminology and recall the fundamental facts of semisimple Lie algebras.

Let \mathfrak{g} be a complex semisimple Lie algebra. We choose a real Lie subalgebra \mathfrak{t} of \mathfrak{g} which satisfies;

- 1) $\mathfrak{g} = \mathfrak{t} + i\mathfrak{t}$ is a direct sum,
- 2) the Killing form $(,)$ is negative definite on \mathfrak{t} .

We fix the above chosen algebra \mathfrak{t} later and call it a compact form of \mathfrak{g} . Let $\mathfrak{q} = i\mathfrak{t}$. Then \mathfrak{q} is a real subspace of \mathfrak{g} and the Killing form $(,)$ is positive definite there. We introduce a $*$ -operation, which is conjugate linear, as follows

$$(u + iv)^* = u - iv, \quad u, v \in \mathfrak{q}.$$

From the definition, $*^2$ is the identity operation of \mathfrak{g} . We denote

$$\{X, Y\} = (X, Y^*), \quad X, Y \in \mathfrak{g},$$

then the form $\{, \}$ becomes a positive definite hermitian form on \mathfrak{g} . For any vector subspace α of \mathfrak{g} , we set

$$\begin{aligned} \alpha^0 &= \{X \in \mathfrak{g} \mid (X, Y) = 0 \text{ for any } Y \in \alpha\} \\ &= \{X \in \mathfrak{g} \mid \{X, Y^*\} = 0 \text{ for any } Y \in \alpha\}. \end{aligned}$$

Let \mathfrak{b} be a Borel subalgebra of \mathfrak{g} and fix it once and for all. Put $\mathfrak{h} = \mathfrak{b} \cap \mathfrak{b}^*$. Then \mathfrak{h} is a Cartan subalgebra of \mathfrak{g} . The rank l of \mathfrak{g} is, by definition, the dimension of \mathfrak{h} . Let \mathfrak{h}_R be the real part of \mathfrak{h} with respect to the $*$ -operation. We denote by \mathfrak{h}_R^* the dual vector space of \mathfrak{h}_R . The root system Δ of \mathfrak{h} in \mathfrak{g} is contained in \mathfrak{h}_R^* . The set Δ is divided into two classes, positive roots Δ_+ and negative roots Δ_- with respect to \mathfrak{b} ;

- 1) $\Delta = \Delta_+ \cup \Delta_-$, disjoint union,
- 2) $\Delta_- = -\Delta_+$.
- 3) if $\alpha, \beta \in \Delta_+$ (respectively Δ_-) and $\alpha + \beta \in \Delta$ then $\alpha + \beta \in \Delta_+(\Delta_-)$.

Each connected component of the set

$$\{H \in \mathfrak{h}_R \mid \alpha(H) \neq 0 \text{ for } \alpha \in \Delta_+\}$$

is called a Weyl chamber. The following set

$$(1.1) \quad \mathfrak{h}_R^+ = \{H \in \mathfrak{h}_R \mid \alpha(H) > 0 \text{ for } \alpha \in \Delta_+\}$$

is, by definition, the positive Weyl chamber. Then conversely for any $H \in \mathfrak{h}_R^+$, we have

$$\Delta_+ = \{\alpha \in \Delta \mid \alpha(H) > 0\}.$$

We denote by $\Pi = \{\alpha_1, \dots, \alpha_l\}$ the set of all simple roots with respect to Δ_+ and take the set as the basis of $\hat{\mathfrak{h}}_R$. We can write any root $\phi \in \Delta$ as $\phi = \sum n_{\alpha_i}(\phi)\alpha_i$ where $n_{\alpha_i}(\phi)$ are non negative or non positive integers according to $\phi \in \Delta_+$ or Δ_- .

We decompose the algebra \mathfrak{g} into root spaces;

$$(1.2) \quad \mathfrak{g} = \mathfrak{h} + \sum_{\alpha \in \Delta_+} \mathfrak{g}_\alpha + \sum_{\beta \in \Delta_-} \mathfrak{g}_\beta,$$

where

$$\mathfrak{h} = \mathfrak{h} + \sum_{\alpha \in \Delta_+} \mathfrak{g}_\alpha.$$

For any $\alpha \in \Delta$, $\dim \mathfrak{g}_\alpha = 1$ and from the definition of \mathfrak{g}_α , it follows

$$\text{ad}(H)(X) = [H, X] = \alpha(H)X, \quad X \in \mathfrak{g}_\alpha, \quad H \in \mathfrak{h}.$$

We can choose the elements $H_\alpha \in \mathfrak{h}$, $X_\alpha \in \mathfrak{g}_\alpha$, $X_{-\alpha} \in \mathfrak{g}_{-\alpha}$ for $\alpha \in \Delta_+$ in the way

$$\begin{aligned} (H, H_\alpha) &= \alpha(H), & H \in \mathfrak{h}, \\ (X_\alpha, X_\beta) &= \begin{cases} 1 & \text{if } \alpha = -\beta \\ 0 & \text{if } \alpha \neq -\beta, \end{cases} \end{aligned}$$

and consequently

$$[X_\alpha, X_{-\alpha}] = H_\alpha.$$

Since the elements H_α , $\alpha \in \Delta_+$, are uniquely determined if we put

$$(1.3) \quad (\alpha, \beta) = (H_\alpha, H_\beta), \quad \alpha, \beta \in \Delta_+,$$

we can define a positive definite symmetric bilinear form $(,)$ on $\hat{\mathfrak{h}}_R$.

Let $[\mathfrak{p}]$ denote the set of all parabolic subalgebras which contain \mathfrak{h} . For each $\mathfrak{p} \in [\mathfrak{p}]$, we know the following decomposition of \mathfrak{g} . See [6].

THEOREM 1.1. *Let $\mathfrak{p} \in [\mathfrak{p}]$. We put $\mathfrak{n} = \mathfrak{p}^0$. Then \mathfrak{n} is the maximal*

nilpotent ideal of \mathfrak{p} and also the set of all nilpotent elements in the radical of \mathfrak{p} . If we put $\mathfrak{g}_1 = \mathfrak{p} \cap \mathfrak{p}^*$ then \mathfrak{g} is decomposed into;

$$(1.4) \quad \mathfrak{g} = \mathfrak{n}^* + \mathfrak{g}_1 + \mathfrak{n} ,$$

where

$$\mathfrak{p} = \mathfrak{g}_1 + \mathfrak{n} .$$

Moreover \mathfrak{g}_1 lies in the normalizers of both \mathfrak{n} and \mathfrak{n}^* , i.e., $[\mathfrak{g}_1, \mathfrak{n}] \subset \mathfrak{n}$ and $[\mathfrak{g}_1, \mathfrak{n}^*] \subset \mathfrak{n}^*$.

For any vector space α which is invariant by the adjoint representation of \mathfrak{h} , we define the subset $\Delta(\alpha) \subset \Delta$ as follows

$$(1.5) \quad \Delta(\alpha) = \left\{ \hat{\mathfrak{h}}_R \ni \alpha \neq 0 \left| \begin{array}{l} [H, X] = \alpha(H)X \text{ for some } 0 \neq X \in \alpha \\ \text{and any } H \in \mathfrak{h} \end{array} \right. \right\} .$$

Since we have $\mathfrak{p} \supset \mathfrak{b} \supset \mathfrak{h}$, we can see \mathfrak{p} as soon as we know $\Delta(\mathfrak{p})$ or in fact $\Delta(\mathfrak{p}) \cap \Delta_-$. The following theorem is also in [6].

- THEOREM 1.2.** 1) *There exists a one-to-one mapping, $\mathfrak{p} \rightarrow \Pi(\mathfrak{p})$, from $[\mathfrak{P}]$ onto the set of all subset of Π .*
 2) *The subset $\Pi(\mathfrak{p})$ of Π satisfies*

$$\Delta(\mathfrak{p}) \cap \Delta_- = \{ \phi \in \Delta_- \mid n_{\alpha_i}(\phi) = 0 \text{ for all } \alpha_i \in \Pi(\mathfrak{p}) \} .$$

Since there are l elements in Π , the number of $[\mathfrak{P}]$ is exactly 2^l . The Lie algebras $\mathfrak{g}_1, \mathfrak{n}$ and \mathfrak{n}^* are all invariant by the adjoint action of \mathfrak{h} . And we have;

$$(1.6) \quad \Delta(\mathfrak{g}_1) = \{ \phi \in \Delta \mid n_{\alpha_i}(\phi) = 0 \text{ for } \alpha_i \in \Pi(\mathfrak{p}) \} ,$$

$$(1.7) \quad \Delta(\mathfrak{n}) = \{ \phi \in \Delta_+ \mid n_{\alpha_i}(\phi) > 0 \text{ for } \alpha_i \in \Pi(\mathfrak{p}) \} ,$$

$$(1.8) \quad \Delta(\mathfrak{n}^*) = -\Delta(\mathfrak{n}) .$$

§ 2. Bruhat decomposition of G/P

We use the notations in § 1., i.e., \mathfrak{g} denotes a semisimple Lie algebra, \mathfrak{b} a Borel subalgebra, $\mathfrak{h} \subset \mathfrak{b}$ a Cartan subalgebra, \dots , etc.

Let G be a simply-connected Lie group whose Lie algebra is \mathfrak{g} . Let B be the Borel subgroup of G whose Lie algebra is \mathfrak{b} . Let T be the maximal torus with Lie algebra \mathfrak{h} . We denote by P the parabolic subgroup corresponding to $\mathfrak{p} \in [\mathfrak{P}]$. Then we have $G \supset P \supset B \supset T$. We call the factor space $X = G/P$ an algebraic homogeneous space. The space X

is a compact kähler manifold and in fact is a projective algebraic manifold. We recall how to embed the space $X = G/P$ into a projective space in section 4.

Let G_1, N and N^* be the Lie subgroups of G whose Lie algebras are $\mathfrak{g}_1, \mathfrak{n}$ and \mathfrak{n}^* respectively. Then the group P is the semidirect product of G_1 and N , i.e.;

$$(2.1) \quad P = G_1 \cdot N,$$

$$(2.2) \quad G_1 \cap N = \{I\},$$

$$(2.3) \quad N \text{ is normal in } P.$$

And the groups P and N^* satisfy

$$(2.4) \quad P \cap N^* = \{I\}.$$

See [11].

We denote by $N(T)$ the normalizer of T in G . We call the group $W = N(T)/T$ the Weyl group of G with respect to T . We define a subgroup W_1 of W and the quotient W^1 in the following

$$\begin{aligned} W_1 &= N(T) \cap P/T, \\ W^1 &= W/W_1 = N(T)/N(T) \cap P. \end{aligned}$$

The group $N(T)$ acts on T, \mathfrak{h} and \mathcal{A} and the formulae

$$w \cdot \exp H \cdot w^{-1} = \exp(\text{Ad}(w)H), \quad (\text{Ad}(w)^*\alpha)(H) = \alpha(\text{Ad}(w)^{-1}(H)),$$

for $w \in N(T), H \in \mathfrak{h}, \alpha \in \mathcal{A}$, are valid. But the actions of $w \in T$ are all trivial so we can regard as the group $W = N(T)/T$ acts on T, \mathfrak{h} and \mathcal{A} . For simplicity, we use the same letter w for $w, \text{Ad}(w)$ and $\text{Ad}(w)^*$. The action of W on \mathcal{A} can be naturally extended on $\hat{\mathfrak{h}}_R$ and then any element of W acts as an orthogonal transformation with respect to the norm $(,)$ introduced in section 1. For any $\alpha \in \mathcal{A}$, we define the reflection S_α ;

$$(2.5) \quad S_\alpha(\phi) = \phi - \frac{2(\alpha, \phi)}{(\alpha, \alpha)}\alpha, \quad \phi \in \hat{\mathfrak{h}}_R,$$

then S_α belongs to W . Further the Weyl group W is generated by the reflections $S_{\alpha_i}, \alpha_i \in \Pi$, and the subgroup W_1 is generated by the reflections $S_{\alpha_j}, \alpha_j \in \Pi - \Pi(p)$. The length $\ell(w)$ of $w \in W$ is the number of reflections appeared in the reduced expressions of w by using the reflections $S_{\alpha_i}, \alpha_i \in \Pi$.

Remark 2.1. By virtue of (1.8), the group W_1 keeps the set $\mathcal{A}(n^*)$ invariant.

Remark 2.2. We fix some representatives $\{w_1, \dots, w_k\}$ of W^1 in $N(T)$. Here $k = \#W^1$. And we sometimes equate the representatives with those of $W^1 = W/W_1$ in W and further with the set W^1 itself.

Every semisimple Lie group G admits the following double coset decomposition and there are three ways to describe;

$$(2.6) \quad G = \bigcup_{w \in W^1} NwP = \bigcup_{w \in W^1} MwP = \bigcup_{w \in W^1} PwP,$$

where M is the maximal nilpotent Lie group of B and w runs over all the above chosen representatives. Let $X_w = NwP/P$ be the image of NwP in X . Then X_w is a cell and the dimension of X_w is represented in terms of the length $\ell(w)$ of w . Confer [1] or Proposition 4.1 in this paper. From (2.6) X_w can be considered as the N, M or P orbit of \bar{w} in X . Then the space X is decomposed into;

$$(2.7) \quad X = \bigcup_{w \in W^1} X_w$$

and we call this decomposition the Bruhat one of X . The topological closure \bar{X}_w of X_w , that is also the Zariski closure, is a subvariety of X and is called a generalized Schubert variety. As a fundamental character, the homology classes of $\{\bar{X}_w\}_{w \in W^1}$ form the free \mathbf{Z} -basis of the integral homology group $H_*(X, \mathbf{Z})$ of X .

In the set W^1 , we can introduce a partial ordering \leq , called the Bruhat ordering;

$$(2.8) \quad w_1 \leq w_2 \text{ if and only if } \bar{X}_{w_1} \subset \bar{X}_{w_2}$$

for $w_1, w_2 \in W^1$. By using this ordering, we can write;

$$(2.9) \quad \bar{X}_w = \bigcup_{w' \leq w} X_{w'}.$$

Namely, to take the closure \bar{X}_w of X_w , it suffices that we attach suitable Bruhat cells of lower dimension to the boundary of X_w successively. Though this ordering is relatively easily defined, it is not easy to judge the order of arbitrarily given two elements $w_1, w_2 \in W^1$. Hence we wish to restate the ordering in another easily understandable words. And further we wish to make the geometrical meanings of the ordering clear. Roughly speaking, for that purpose there exist two ways until now, firstly

the way to compare the reduced expressions of $w \in W^1$ by using the reflections S_{α_i} , $\alpha_i \in \Pi$, and secondly the way to compare the inclusion relation among Verma modules $M(w\lambda)$ associated to $w \in W^1$. Here λ is the highest weight which determines the embedding of X into the projective space $P^N(V)$. See [3].

But as will be shown in section 4, we take a different approach. We first show that any Bruhat cell can be written as the union of suitable flows $P(t)$ of the vector field V_H on X . The vector field V_H is defined in the next section. We investigate the behavior of the flows and show what Bruhat cells of lower dimension should be attached to the boundary of the Bruhat cell X_w of the appointed direction for the compactification of X_w . We give a simple necessary-sufficient condition to determine the ordering \leq in Theorem 4.12 and ties the two ways stated above in some sense.

§3. Vector field V_H

We first prove the following proposition.

PROPOSITION 3.1. *Let us act the maximal torus T on $X = G/P$. Then the quotient set $W^1 = W/W_1 = N(T)/N(T) \cap P$ is naturally realized as the set of all T fixed points in X .*

Proof. An element $\bar{g} \in X$ is fixed by the action of T if and only if $g^{-1}Tg \subset P$ where g is a representative of \bar{g} in G . The group $g^{-1}Tg$ is also a maximal torus of G contained in P and hence we have

$$g^{-1}Tg = pTp^{-1}$$

for some $p \in P$. See [9]. This means $gp \in N(T)$. Hence \bar{g} defines a coset \widetilde{gp} in W^1 . If \bar{g} and \bar{g}' define the same coset in W^1 , we can write $gp = g'p'p''$ for some $p, p' \in P$ and $p'' \in N(T) \cap P$. So we have $\bar{g} = \bar{g}'$ in X . If we take any element $w \in N(T)$, the coset corresponding to w is $\bar{w} \in W^1$. So the mapping is onto. The proof is completed.

Let us consider the following diagram

$$(3.1) \quad \begin{array}{ccccc} & G & G/P & G/P & \\ & \cup & \cup & \cup & \\ n^* & \xrightarrow{\phi} N^* & \xrightarrow{\psi} \bar{N}^* & \xrightarrow{w} w\bar{N}^* & \\ \psi & \psi & \psi & \psi & \\ Z & \longrightarrow \exp Z & \longrightarrow \exp \bar{Z} & \longrightarrow w \exp \bar{Z} & . \end{array}$$

We take the elements X_α , $\alpha \in \mathcal{A}(\mathfrak{n}^*)$, chosen in section 1, as the basis of \mathfrak{n}^* and write an element Z of \mathfrak{n}^* as $Z = \sum_{\alpha \in \mathcal{A}(\mathfrak{n}^*)} z_\alpha X_\alpha$. Since the Lie algebra \mathfrak{n}^* is nilpotent, we have $\log(\exp Z) = Z$ and hence the mapping ϕ is one-to-one and onto. Since $N^* \cap P = \{I\}$, the mapping ψ is also one-to-one. See (2.4). The left multiplication by \mathfrak{w} is clearly one-to-one. Hence we can take the pair $(\mathfrak{w}\bar{N}^*, \phi^{-1} \circ \psi^{-1} \circ \mathfrak{w}^{-1})$ as a coordinate neighborhood around $\mathfrak{w} \in W^1$ and then $\{z_\alpha(\mathfrak{w}\bar{n}^*)\}_{\alpha \in \mathcal{A}(\mathfrak{n}^*)}$ becomes the local coordinate.

Remark 3.2. Let \mathfrak{w}_0 be the element of W whose length $\ell(\mathfrak{w}_0)$ is maximal among all. Then since $\mathfrak{w}_0^{-1}N\mathfrak{w}_0 = N^*$, $\mathfrak{w}_0\bar{N}^* = N\mathfrak{w}_0P/P = X_{\mathfrak{w}_0}$. So the set $\mathfrak{w}_0\bar{N}^*$ is the Bruhat cell of maximal dimension and is a Zariski open set.

We sum up our assertions.

THEOREM 3.3. *The quotient set $W^1 = W/W_1$ can be naturally embedded into $X = G/P$ as the set of all T -fixed points and the pair $(\mathfrak{w}\bar{N}^*, \phi^{-1} \circ \psi^{-1} \circ \mathfrak{w}^{-1})$ is a coordinate neighborhood around $\mathfrak{w} \in W^1$. The sets $\mathfrak{w}\bar{N}^*$, $\mathfrak{w} \in W^1$, are all T invariant Zariski open sets. In fact if we multiply $\exp H \in T$ from the left side on $\mathfrak{w}\bar{N}^*$, the local coordinate $\{z_\alpha(\mathfrak{w}\bar{n}^*)\}_{\alpha \in \mathcal{A}(\mathfrak{n}^*)}$ changes to $\{e^{(\mathfrak{w}\alpha)(H)} \cdot z_\alpha(\mathfrak{w}\bar{n}^*)\}_{\alpha \in \mathcal{A}(\mathfrak{n}^*)}$. Further the space X is completely covered with the family of the open sets $\{\mathfrak{w}\bar{N}^*\}_{\mathfrak{w} \in W^1}$, i.e., $X = \bigcup_{\mathfrak{w} \in W^1} \mathfrak{w}\bar{N}^*$.*

Proof. The first and second sentences have been proved. Since for $\exp Z \in N^*$ we have

$$\begin{aligned} \exp H \cdot \mathfrak{w} \exp Z \cdot P &= \mathfrak{w}\mathfrak{w}^{-1} \exp H \cdot \mathfrak{w} \exp Z \cdot \mathfrak{w}^{-1} \exp(-H)\mathfrak{w} \cdot P \\ &= \mathfrak{w} \exp(\mathfrak{w}^{-1}(H)) \exp Z \exp(-\mathfrak{w}^{-1}(H)) \cdot P \\ &= \mathfrak{w} \exp(\text{Ad}(\exp(\mathfrak{w}^{-1}(H)))Z) \cdot P \\ &= \mathfrak{w} \exp(\text{Exp}(\text{ad}(\mathfrak{w}^{-1}(H)))Z) \cdot P \end{aligned}$$

and

$$\text{Exp}(\text{ad}(\mathfrak{w}^{-1}(H))) \cdot Z \in \mathfrak{n}^*,$$

so

$$\begin{aligned} (\phi^{-1} \circ \psi^{-1} \circ \mathfrak{w}^{-1})(\exp H \cdot \mathfrak{w} \overline{\exp Z}) &= \log(\mathfrak{w}^{-1}\mathfrak{w} \exp(\text{Exp}(\text{ad}(\mathfrak{w}^{-1}(H))) \cdot Z)) \\ &= \text{Exp}(\text{ad}(\mathfrak{w}^{-1}(H))) \cdot Z. \end{aligned}$$

If we write $Z = \sum_{\alpha \in \mathcal{A}(\mathfrak{n}^*)} z_\alpha X_\alpha$, we have

$$\begin{aligned} \text{ad}(\mathfrak{w}^{-1}(H)) \cdot Z &= [\mathfrak{w}^{-1}(H), Z] = [\mathfrak{w}^{-1}(H), \sum_{\alpha \in \mathcal{J}(\mathfrak{n}^*)} z_\alpha X_\alpha] \\ &= \sum_{\alpha \in \mathcal{J}(\mathfrak{n}^*)} z_\alpha [\mathfrak{w}^{-1}(H), X_\alpha] \\ &= \sum_{\alpha \in \mathcal{J}(\mathfrak{n}^*)} \alpha(\mathfrak{w}^{-1}(H)) z_\alpha X_\alpha \\ &= \sum_{\alpha \in \mathcal{J}(\mathfrak{n}^*)} (\mathfrak{w}\alpha)(H) z_\alpha X_\alpha \end{aligned}$$

and hence

$$\text{Exp}(\text{ad}(\mathfrak{w}^{-1}(H))) \cdot Z = \sum_{\alpha \in \mathcal{J}(\mathfrak{n}^*)} e^{(\mathfrak{w}\alpha)(H)} z_\alpha X_\alpha .$$

To prove $X = \bigcup_{\mathfrak{w} \in W^1} \mathfrak{w}\bar{N}^*$, we need the following fact. See [8].

FACT. If Y is a compact kähler manifold and $H^1(Y, \mathbb{C}) = 0$, then a complex connected solvable Lie group S acting holomorphically on Y always has a fixed point inside any analytic subvariety that S leaves invariant.

The space X satisfies these assumptions and we can take T as S . Let us apply this fact to our case. Since $\mathfrak{w}\bar{N}^*$ is a T invariant Zariski open set, the complement $X' = X - \bigcup_{\mathfrak{w} \in W^1} \mathfrak{w}\bar{N}^*$ becomes a T invariant subvariety. Hence if X' is not empty, it must have a fixed point of T . But the set W^1 , which is the set of all T fixed points, is of course outside X' . This is a contradiction.

For any $Y \in \mathfrak{g}$, we can define a holomorphic vector field V_Y by operating Y infinitesimally on X ;

$$(3.2) \quad (V_Y f)(\bar{g}) = \lim_{\varepsilon \rightarrow 0} \frac{f(\exp(\varepsilon Y)\bar{g}) - f(\bar{g})}{\varepsilon} ,$$

where $\bar{g} \in X$ and f is a local function around \bar{g} . In particular for an element $H \in \mathfrak{h}$, V_H can be written as follows

$$(3.3) \quad V_H = \sum_{\alpha \in \mathcal{J}(\mathfrak{n}^*)} (\mathfrak{w}\alpha)(H) z_\alpha \frac{\partial}{\partial z_\alpha}$$

on $\mathfrak{w}\bar{N}^*$ by using the local coordinate $\{z_\alpha(\mathfrak{w}\bar{n}^*)\}_{\alpha \in \mathcal{J}(\mathfrak{n}^*)}$. This claim is obvious from Theorem 3.3. If we choose H from the Weyl chambers, we have $0 \neq (\mathfrak{w}\alpha)(H) \in \mathbb{R}$, for $\mathfrak{w} \in W^1$, $\alpha \in \mathcal{J}(\mathfrak{n}^*)$. Hence the set of all points where V_H vanishes agrees with W^1 and V_H vanishes with first order there. Hereafter we further assume that H belongs to the positive Weyl chamber $\mathfrak{h}_{\mathbb{R}}^+$. Then for $\alpha \in \mathcal{A}$, we note that $\alpha \in \mathcal{A}_+$ if and only if $\alpha(H) > 0$.

From the expression (3.3) of V_H , if we use the local coordinate $\{z_\alpha(w\bar{N}^*)\}_{\alpha \in \mathcal{A}(n^*)}$ on $w\bar{N}^*$, any flow $P(t)$, parametrized by $t \in \mathbb{C}$, of V_H can be written;

$$(3.4) \quad z_\alpha(P(t)) = c_\alpha^w e^{(w\alpha)(H)t}, \quad c_\alpha^w \in \mathbb{C}, \quad \text{for } \alpha \in \mathcal{A}(n^*).$$

Let us introduce some notations.

$$\begin{aligned} \mathcal{A}_+(n^*, w) &= \{\alpha \in \mathcal{A}(n^*) \mid w\alpha \in \mathcal{A}_+\}, \\ \mathcal{A}_-(n^*, w) &= \{\alpha \in \mathcal{A}(n^*) \mid w\alpha \in \mathcal{A}_-\}, \\ \mathcal{A}(n^*, w, P(t)) &= \{\alpha \in \mathcal{A}(n^*) \mid c_\alpha^w \neq 0\}, \\ \mathcal{A}_+(n^*, w, P(t)) &= \mathcal{A}_+(n^*, w) \cap \mathcal{A}(n^*, w, P(t)) \\ &= \{\alpha \in \mathcal{A}(n^*) \mid c_\alpha^w \neq 0, w\alpha \in \mathcal{A}_+\}, \\ \mathcal{A}_-(n^*, w, P(t)) &= \mathcal{A}_-(n^*, w) \cap \mathcal{A}(n^*, w, P(t)), \end{aligned}$$

where $P(t)$ is the flow of V_H on $w\bar{N}^*$ as above.

We divide the flow $P(t)$ of V_H on $w\bar{N}^*$ into three types;

- 1) $\mathcal{A}_-(n^*, w, P(t)) = \emptyset$, i.e., a flow starting from w .
- 2) $\mathcal{A}_+(n^*, w, P(t)) = \emptyset$, i.e., a flow arriving at w .
- 3) the others.




Fig. 1

For instance, let us choose a flow $P(t)$ of type 2). Then if we fix the

imaginary part of t and approach the real part of t to $+\infty$, we obtain $\lim_{t \rightarrow +\infty} P(t) = w$.

§ 4. Proof of our theorems

Let V_H be the same as in section 3. Then since $\Delta(n^*) \subset \Delta_-$, we have $\alpha(H) < 0$ for $\alpha \in \Delta(n^*)$. We further assume that H satisfies

$$(w\alpha_i)(H) \neq (w\alpha_j)(H) \quad \text{if } \alpha_i \neq \alpha_j \text{ in } \Delta(n^*)$$

for each $w \in W^1$.

We denote

$$(4.1) \quad W^1(p) = \{w \in W^1 \mid \text{the number of } \Delta_+(n^*, w) \text{ is } p\}$$

and

$$(4.2) \quad X'_w = \{w\bar{n}^* \in w\bar{N}^* \mid z_\beta(w\bar{n}^*) = 0 \text{ for } \beta \in \Delta_-(n^*, w)\}.$$




Fig. 2. $p = 2$.

From the definition of $\Delta_+(n^*, w)$, the set X'_w can be considered as the union of all flows of V_H which start from w . And if $w \in W^1(p)$, X'_w is biholomorphic to C^p .

PROPOSITION 4.1. *The space X is decomposed into the disjoint union of the cells X'_w , $w \in W^1$. And this decomposition is identical to the Bruhat decomposition of X .*

Proof. We choose $x \in X$. If x is fixed by the infinitesimal action of H , it belongs to W^1 . Hence $x = w \in X'_w$ for some $w \in W^1$. Conversely if x is not fixed, there is the unique flow $P(t)$ of V_H which passes through x and the starting point $\lim_{t \rightarrow -\infty} P(t)$ is fixed by the infinitesimal action of H . We give the precise proof of the existence of the limit points $\lim_{t \rightarrow \pm\infty} P(t)$ in Remark 4.6 later. If we denote $w = \lim_{t \rightarrow -\infty} P(t)$, x belongs to X'_w . To prove the equivalence of two decompositions, it suffices to show $X_w \supset X'_w$ for each $w \in W^1$. We use the second description of the Bruhat

cell, i.e., $X_{\mathfrak{w}} = M\mathfrak{w}P/P = M\bar{\mathfrak{w}}$. See (2.6). Since M is the maximal nilpotent Lie group contained in B , the Lie algebra \mathfrak{m} of M is written as $\mathfrak{m} = \sum_{\alpha \in \mathcal{J}_+} \mathfrak{g}_\alpha$. Then we have

$$\begin{aligned} X_{\mathfrak{w}} \cap \mathfrak{w}\bar{N}^* &= M\bar{\mathfrak{w}} \cap \mathfrak{w}\bar{N}^* \\ &= \mathfrak{w}\mathfrak{w}^{-1}M\bar{\mathfrak{w}} \cap \mathfrak{w}\bar{N}^* \supset \overline{\mathfrak{w}(\mathfrak{w}^{-1}M\mathfrak{w})} \cap \bar{N}^* . \end{aligned}$$

Since the Lie algebra of $\mathfrak{w}^{-1}M\mathfrak{w} \cap N^*$ is

$$\begin{aligned} \left(\mathfrak{w}^{-1} \sum_{\beta \in \mathcal{J}_+} \mathfrak{g}_\beta \mathfrak{w} \right) \cap \left(\sum_{\alpha \in \mathcal{J}(N^*)} \mathfrak{g}_\alpha \right) &= \left(\sum_{\beta \in \mathcal{J}_+} \mathfrak{g}_{\mathfrak{w}^{-1}\beta} \right) \cap \left(\sum_{\alpha \in \mathcal{J}(N^*)} \mathfrak{g}_\alpha \right) \\ &= \sum_{\substack{\alpha \in \mathcal{J}(N^*) \\ \mathfrak{w}\alpha \in \mathcal{J}_+}} \mathfrak{g}_\alpha , \end{aligned}$$

$\overline{\mathfrak{w}(\mathfrak{w}^{-1}M\mathfrak{w})} \cap \bar{N}^*$ coincides with $X'_{\mathfrak{w}}$. Hence we have $X_{\mathfrak{w}} \supset X'_{\mathfrak{w}}$ and complete the proof.

To investigate the behavior of the flow $P(t)$ of V_H , let us embed the space X into a certain projective space. See [10]. Let λ be an integral dominant weight such that $(\lambda, \alpha_i) = 0$ for $\alpha_i \in \Pi - \Pi(\mathfrak{p})$ and $(\lambda, \alpha_j) > 0$ for $\alpha_j \in \Pi(\mathfrak{p})$. We denote by (ϕ, \mathcal{V}) the irreducible representation of G with the highest weight λ and by $\langle \lambda \rangle$ the set of all weights of the representation. Here we count the weights with multiplicities. So even if the letters of weights are different as γ and δ , it does not necessarily mean that they are different as weights. The Weyl group W acts on the space $\hat{\mathfrak{h}}_R$ as the orthogonal transformations and the subgroup W_λ is realized as the isotropy subgroup of W at λ ;

$$W_\lambda = \{w \in W \mid w\lambda = \lambda\} .$$

We call such a weight as $w\lambda$, $w \in W$, extremal and denote

$$(4.3) \quad \langle \lambda \rangle^e = \{w\lambda \mid w \in W\} \subset \langle \lambda \rangle .$$

By virtue of the irreducibility of the representation, the multiplicity of $w\lambda$ is one. There is a one-to-one correspondence;

$$(4.4) \quad \begin{array}{ccc} W^1 & \longrightarrow & \langle \lambda \rangle^e \\ \cup & & \cup \\ \mathfrak{w} & \longrightarrow & \mathfrak{w}\lambda . \end{array}$$

We decompose the vector space \mathcal{V} into the weight spaces and write in the following;

$$(4.5) \quad \mathcal{V} = \sum_{\gamma \in \langle \lambda \rangle} \mathcal{V}_\gamma .$$

From the remark stated above, $\dim V_\gamma = 1$ for all $\gamma \in \langle \Lambda \rangle$. We choose a nonzero element $Y_\gamma \in V_\gamma$ as the base of V_γ . By virtue of the choice of Λ , the space V_Λ of the highest weight is invariant by P and conversely any element of G which keeps V_Λ invariant belongs to P . Hence the restriction $(\phi|_P, V_\Lambda)$ of the representation (ϕ, V) is a representation of P of one dimension. We write $\ell(p) = \phi|_P(p)$, $p \in P$. Then $d\ell|_{\mathfrak{p}} = \Lambda$. The representation (ℓ, V_Λ) induces a homogeneous line bundle L on $X = G/P$ and the space V is concretely realized as the vector space $H^0(X, L)$ of all global sections of L . Every section $\psi \in H^0(X, L)$ can be considered as a function on G satisfying

$$\psi(g) = \ell(p)\psi(gp)$$

for $g \in G$ and $p \in P$. The representation ϕ of G on $V = H^0(X, L)$ is as follows

$$(\phi(g) \cdot \psi)(g') = \psi(g^{-1}g')$$

for $\psi \in H^0(X, L)$ and $g, g' \in G$.

For simplicity, we omit the letter ϕ later and write in the way

$$\phi(g) \cdot Y = g \cdot Y$$

for $g \in G, Y \in V$. We offer the embedding theorem. See [10].

THEOREM 4.2. *Let ι be the projection of G onto the G -orbit of $Y_\Lambda \in V_\Lambda$. Since P leaves the space V_Λ invariant, the mapping ι can be projected to $\bar{\iota}$;*

$$(4.6) \quad \begin{array}{ccc} G & \xrightarrow{\quad \phi \quad} & V \\ \downarrow \iota & \searrow \cong & \downarrow \\ X = G/P & \xrightarrow{\quad \bar{\iota} \quad} & P^N(V) = V/\sim \end{array} \quad \begin{array}{ccc} & \xrightarrow{\quad g \quad} & g \cdot Y_\Lambda \\ & \searrow \cong & \downarrow \\ & \xrightarrow{\quad \bar{g} \quad} & \bar{g} \cdot Y_\Lambda \end{array}$$

Then all the mappings above are G -equivariant. And the horizontal mapping $\bar{\iota}$ embeds the space X into $P^N(V)$.

By considering the mapping

$$\begin{array}{ccc} V & \longrightarrow & \mathbb{C}^{N+1} \\ \psi & & \psi \\ \sum_{\gamma \in \langle \Lambda \rangle} x_\gamma Y_\gamma & \longrightarrow & (x_\gamma)_{\gamma \in \langle \Lambda \rangle}, \end{array}$$

we obtain the homogeneous coordinate $[x_\gamma]_{\gamma \in \langle \Lambda \rangle}$ of $P^N(V)$. We abbreviate \overline{Y}_δ to δ . As usual, we define affine open sets

$$U_\gamma = \{[x_\delta] \in P^N(V) \mid x_\gamma \neq 0\}, \quad \gamma \in \langle \Lambda \rangle,$$

and denote by $y_\delta = x_\delta/x_\gamma$, $\delta \in \langle \Lambda \rangle - \gamma$, the inhomogeneous coordinate on U_γ .

If we denote

$$S = \{[x_\gamma] \in P^N(V) \mid [x_\gamma] \text{ is fixed by the left action of } T\}$$

then it is decomposed into

$$S = \bigcup_{\gamma \in \langle \Lambda \rangle} S_\gamma, \quad \text{disjoint union,}$$

where

$$S_\gamma = \{[x_\delta] \in P^N(V) \mid x_\delta = 0 \text{ for any } \delta \in \langle \Lambda \rangle \text{ such that } \delta \neq \gamma \text{ as weights}\}.$$

For an element $w \in N(T)$, we have

$$\begin{aligned} twY_A &= tw^{-1}twY_A \\ &= w\ell(w^{-1}tw)Y_A \\ &= (w\ell)(t)wY_A, \quad \text{for } t \in T, \end{aligned}$$

so wY_A belongs to V_{wA} . Hence the projected element $\bar{w} \in W^1 = N(T)/N(T) \cap P \hookrightarrow X$ is mapped to $\overline{wY_A} = \overline{Y_{wA}} = wA \in \langle \Lambda \rangle^e$ by $\bar{\iota}$. In other words, the correspondence (4.4) coincides with $\bar{\iota} \mid W^1$. The embedded manifold $\bar{\iota}(X)$ intersects with S only at the points $\langle \Lambda \rangle^e$. Because if $\bar{\iota}(X)$ has a common point x with S outside $\langle \Lambda \rangle^e$, the inverse image $\bar{\iota}^{-1}(x) \in X$ must be fixed by the T action. But this case can not occur.

EXAMPLE 4.3. Let $X = G(k, n)$ be the Grassmann variety. We can write X as follows

$$X = SL(n+1, \mathbf{C})/P,$$

where

$$P = \left\{ A \in SL(n+1, \mathbf{C}) \mid A = \begin{pmatrix} A_1 & A_2 \\ 0 & A_3 \end{pmatrix}, A_1: (k+1) \times (k+1) \text{ matrix} \right\}.$$

The manifold X has the well known extension of vector bundles;

$$0 \longrightarrow U \longrightarrow X \times E \longrightarrow Q \longrightarrow 0$$

where U is the universal bundle and Q is the quotient bundle. The bundle U is induced by the representation

$$\begin{array}{ccc} \phi: P & \longrightarrow & GL(k + 1, \mathbb{C}) \\ \psi & & \psi \\ A & \longrightarrow & A_1. \end{array}$$

In this case, we can take $L = \det U^{-1} = \det Q$ and $V = H^0(X, L)$ and then we obtain the usual Plücker embedding;

$$X \hookrightarrow P^N(V),$$

where $N = \binom{n + 1}{k + 1} - 1$. Since the Weyl group W of $SL(n + 1, \mathbb{C})$ is isomorphic to \mathfrak{S}_{n+1} , the permutation group of $n + 1$ elements, and the subgroup W_1 is isomorphic to $\mathfrak{S}_{k+1} \times \mathfrak{S}_{n-k}$, we have

$$\begin{aligned} \text{the number of } W^1 &= \frac{(n + 1)!}{(k + 1)!(n - k)!} = \binom{n + 1}{k + 1} \\ &= \dim V. \end{aligned}$$

Hence $\langle \lambda \rangle^e = \langle \lambda \rangle$, i.e., the weights of (ϕ, V) are all extremal. If we take as the basis of V , we obtain the usual Plücker coordinate $[x_{\mathfrak{m}\lambda}]_{\mathfrak{m}\lambda \in \langle \lambda \rangle^e}$ of $P^N(V)$.

We impose one more condition on the selection of H ;
 $\delta(H) \neq \gamma(H)$ for $\gamma, \delta \in \langle \lambda \rangle$ such that $\gamma \neq \delta$ as the elements of $\hat{\mathfrak{h}}_{\mathbb{R}}$.

Remark 4.4. We note that the conditions imposed on the choice of H are all open.

Let us lift the T action on X to $P^N(V)$ and construct a vector field V_H^A . Since

$$\begin{aligned} &\lim_{\varepsilon \rightarrow 0} \frac{\exp(\varepsilon H) \cdot (\sum_{\gamma \in \langle \lambda \rangle} x_{\gamma} Y_{\gamma}) - (\sum_{\gamma \in \langle \lambda \rangle} x_{\gamma} Y_{\gamma})}{\varepsilon} \\ &= \lim_{\varepsilon \rightarrow 0} \frac{(I + \varepsilon H/1! + (\varepsilon H)^2/2! + \dots) \cdot (\sum_{\gamma \in \langle \lambda \rangle} x_{\gamma} Y_{\gamma}) - (\sum_{\gamma \in \langle \lambda \rangle} x_{\gamma} Y_{\gamma})}{\varepsilon} \\ &= \sum_{\gamma \in \langle \lambda \rangle} \gamma(H) x_{\gamma} Y_{\gamma}, \end{aligned}$$

we have a vector field \tilde{V}_H^A on V such that

$$\tilde{V}_H^A = \sum_{\gamma \in \langle \lambda \rangle} \gamma(H) x_{\gamma} \frac{\partial}{\partial x_{\gamma}}.$$

Because the coefficients of $\partial/\partial x_{\gamma}$, $\gamma \in \langle \lambda \rangle$, in \tilde{V}_H^A are all linear functions, the vector field \tilde{V}_H^A can be descended to $P^N(V)$. By using the inhomoge-

neous coordinate $\{y_\delta\}_{\delta \in \langle \Lambda \rangle - \gamma}$ on U_γ , we can write the descended vector field V_H^A as follows

$$(4.7) \quad V_H^A = \sum_{\delta \in \langle \Lambda \rangle - \gamma} (\delta(H) - \gamma(H))y_\delta \frac{\partial}{\partial y_\delta} = \sum_{\delta \in \langle \Lambda \rangle - \gamma} (\delta - \gamma)(H)y_\delta \frac{\partial}{\partial y_\delta}.$$

In general, there are multiplicities in weights so the zero set of V_H^A is not necessarily isolated and in fact coincides with S . Every flow $P(t)$ of V_H^A on U_γ can be expressed as follows

$$(4.8) \quad y_\delta(P(t)) = c_\delta e^{(\delta - \gamma)(H)t}, \quad c_\delta \in \mathbb{C}, \quad \text{for } \delta \in \langle \Lambda \rangle - \gamma.$$

PROPOSITION 4.5. 1) For any flow $P(t)$ of V_H^A , the limit points $\lim_{t \rightarrow \pm\infty} P(t)$ exist and belong to the zero set S of V_H^A .

2) There is a flow of V_H^A which starts from γ and arrives at δ if and only if it is satisfied that $(\delta - \gamma)(H) > 0$.

Proof. To prove 1), we can assume that a part of the flow $P(t)$ lies on U_γ for some $\gamma \in \langle \Lambda \rangle$. Then $P(t)$ is written as (4.8) there. If the maximal value m of $\{(\delta - \gamma)(H) \mid c_\delta \neq 0\}$ is positive, we set

$$\begin{aligned} M &= \{\delta \in \langle \Lambda \rangle \mid (\delta - \gamma)(H) = m, c_\delta \neq 0\} \\ &= \{\delta_1, \dots, \delta_k\}. \end{aligned}$$

Then by using the homogeneous coordinate $[x_\tau]_{\tau \in \langle \Lambda \rangle}$, we have

$$x_\tau \left(\lim_{t \rightarrow +\infty} P(t) \right) = \begin{cases} c_{\delta_i} & \text{if } \tau = \delta_i \in M \\ 0 & \text{if } \tau \notin M. \end{cases}$$

From Remark 4.4., we have $\delta_1 - \gamma = \delta_2 - \gamma = \dots = \delta_k - \gamma$ and hence $\delta_1 = \dots = \delta_k$ as an element of $\hat{\mathfrak{h}}_R$. So the limit point $\lim_{t \rightarrow +\infty} P(t)$ exists and belongs to $S_{\delta_1} \subset S$. In case $m \leq 0$, the limit point $\lim_{t \rightarrow +\infty} P(t)$ also exists and belongs to S_γ . Similarly we can prove the existence of $\lim_{t \rightarrow +\infty} P(t)$. If we consider the flow $P_\tau^\delta(t)$ such that

$$y_\tau(P_\tau^\delta(t)) = \begin{cases} e^{(\delta - \gamma)(H)t} & \text{if } \tau = \delta \\ 0 & \text{if } \tau \neq \delta, \end{cases}$$

for $\tau \in \langle \Lambda \rangle - \gamma$, we can prove 2). We omit the detailed proof.

Remark 4.6. If we embed the space X into $P^N(V)$, we can consider that the vector field V_H is the restriction of V_H^A to X . Then the flow $P(t)$ of V_H is that of V_H^A lying entirely on X . Hence the limit points $\lim_{t \rightarrow \pm\infty} P(t)$ always exist and belong to $\langle \Lambda \rangle^\varepsilon$.

In the following discussions, we consider the space X as the T invariant submanifold of $P^N(V)$. Let us compare the expressions of V_H^A and V_H around $\mathfrak{w}A = i(\mathfrak{w})$. By using the coordinate $\{y_\delta\}_{\delta \in \langle A \rangle - \mathfrak{w}A}$, we have

$$(4.9) \quad V_H^A = \sum_{\delta \in \langle A \rangle - \mathfrak{w}A} (\delta - \mathfrak{w}A)(H)y_\delta \frac{\partial}{\partial y_\delta} .$$

On the other hand, since the space X is nonsingular around $\mathfrak{w}A = i(\mathfrak{w})$, we add some local functions $\{u_i\}_{i \in I}$ to $\{z_\alpha\}_{\alpha \in \mathcal{A}(n^*)}$ and make them a local coordinate in some neighborhood U of $\mathfrak{w}A = i(\mathfrak{w})$. We can assume that the functions $\{u_i\}_{i \in I}$ satisfy;

- 1) $X \cap U = \{x \in U \mid u_i(x) = 0, i \in I\}$
- 2) With respect to the local coordinate $\{z_\alpha, u_i\}_{\alpha \in \mathcal{A}(n^*), i \in I}$, it is satisfied

$$(4.10) \quad V_H^A = \sum_{\alpha \in \mathcal{A}(n^*)} \{(\mathfrak{w}\alpha)(H)z_\alpha + f_\alpha(z_\beta; u_j)\} \frac{\partial}{\partial z_\alpha} + \sum_{i \in I} \{\gamma_i u_i + g_i(z_\beta; u_j)\} \frac{\partial}{\partial u_i} ,$$

where $f_\alpha(z_\beta; u_j)$ and $g_i(z_\beta; u_j)$ are the functions of higher terms and $f_\alpha(z_\beta; 0) = g_i(z_\beta; 0) = 0, \alpha \in \mathcal{A}(n^*), i \in I$. If we compare (4.9) and (4.10), we have

$$\{(\delta - \mathfrak{w}A)(H)\}_{\delta \in \langle A \rangle - \mathfrak{w}A} = \{(\mathfrak{w}\alpha)(H); \gamma_i\}_{\alpha \in \mathcal{A}(n^*)} , \quad i \in I .$$

Let us consider a flow $P_\alpha^{\mathfrak{w}}(t)$ with special direction of V_H on $\mathfrak{w}\bar{N}^*$ such that

$$(4.11) \quad z_\beta(P_\alpha^{\mathfrak{w}}(t)) = \begin{cases} c_\alpha^{\mathfrak{w}} e^{(\mathfrak{w}\alpha)(H)t} & \text{if } \beta = \alpha \\ 0 & \text{if } \beta \neq \alpha , \end{cases}$$

where $C \ni c_\alpha^{\mathfrak{w}} \neq 0$. In other words, $\mathcal{A}(n^*, \mathfrak{w}, P_\alpha^{\mathfrak{w}}(t)) = \alpha$. If $\mathfrak{w}\alpha \in \mathcal{A}_+$ then $P_\alpha^{\mathfrak{w}}(t)$ starts from \mathfrak{w} . Conversely if $\mathfrak{w}\alpha \in \mathcal{A}_-$, $P_\alpha^{\mathfrak{w}}(t)$ arrives at \mathfrak{w} .

PROPOSITION 4.7. *Let $P_\alpha^{\mathfrak{w}}(t)$ be the flow of V_H on $\mathfrak{w}\bar{N}^*$ as above. We assume $\mathfrak{w}\alpha \in \mathcal{A}_+$ and put $\mathfrak{w}' = \lim_{t \rightarrow +\infty} P_\alpha^{\mathfrak{w}}(t) \in W^1$. If we write the flow by using the local coordinate $\{z_\beta(\mathfrak{w}'\bar{n}^*)\}_{\beta \in \mathcal{A}(n^*)}$ around \mathfrak{w}' , it follows*

$$z_\beta(P_\alpha^{\mathfrak{w}'}(t)) = \begin{cases} c_\alpha^{\mathfrak{w}'} e^{(\mathfrak{w}'\beta)(H)t} & \text{if } \mathfrak{w}\alpha = -\mathfrak{w}'\beta \\ 0 & \text{otherwise .} \end{cases}$$

Namely the flows $P_\alpha^{\mathfrak{w}}(t)$ and $P_\beta^{\mathfrak{w}'}(t)$ agree with each other on $\mathfrak{w}\bar{N}^ \cap \mathfrak{w}'\bar{N}^*$.*

Proof. To avoid confusion, we write the local coordinate $\{z_\beta(\mathfrak{w}\bar{n}^*)\}_{\beta \in \mathcal{A}(n^*)}$ as $\{z_\beta\}_{\beta \in \mathcal{A}(n^*)}$ and the local coordinate $\{z_\beta(\mathfrak{w}'\bar{n}^*)\}_{\beta \in \mathcal{A}(n^*)}$ as $\{z'_\beta\}_{\beta \in \mathcal{A}(n^*)}$. We choose a point $P_\alpha^{\mathfrak{w}}(t_0), 0 \neq t_0 \in C$, on the flow and put $z_\beta(P_\alpha^{\mathfrak{w}}(t_0)) = z_{\beta,0}, \beta \in \mathcal{A}(n^*)$, and $z'_\beta(P_\alpha^{\mathfrak{w}}(t_0)) = z'_{\beta,0}, \beta \in \mathcal{A}(n^*)$. Then both $\{z_\beta - z_{\beta,0}\}_{\beta \in \mathcal{A}(n^*)}$ and $\{z'_\beta -$

$z'_{\beta,0}$ } $_{\beta \in \Delta(n^*)}$ are local coordinates around $P_\alpha^w(t_0)$. If we compare the expressions of the flow $P_\alpha^w(t)$ by using the two local coordinate, we have

$$\begin{aligned} z'_\beta(P_\alpha^w(t)) - z'_{\beta,0} &= \sum_{n=0}^\infty d_{\beta,n}(z_\alpha(P_\alpha^w(t)) - z_{\alpha,0})^n \\ &= \sum_{n=0}^\infty d_{\beta,n}(c_\alpha^w e^{(w\alpha)(H)t} - z_{\alpha,0})^n \\ &= \sum_{n=0}^\infty d'_{\beta,n} e^{n(w\alpha)(H)t} \end{aligned}$$

and hence

$$(4.12) \quad z'_\beta(P_\alpha^w(t)) = \sum_{n=0}^\infty d''_{\beta,n} e^{n(w\alpha)(H)t} \quad \text{for } \beta \in \Delta(n^*).$$

On the other hand, any flow $P(t)$ of V_H on $w'\bar{N}^*$ must be written as follows

$$(4.13) \quad z'_\beta(P(t)) = c_\beta^{w'} e^{(w'\beta)(H)t}, \quad c_\beta^{w'} \in \mathbb{C}, \quad \text{for } \beta \in \Delta(n^*).$$

Hence if we compare the coefficients of t in (4.12) and (4.13), we have

$$c_\beta^{w'}(w'\beta)(H) = \left(\sum_{n=0}^\infty n d''_{\beta,n} \right) (w\alpha)(H).$$

From Remark 4.4., this means

$$c_\beta^{w'}(w'\beta) = \left(\sum_{n=0}^\infty n d''_{\beta,n} \right) (w\alpha)$$

as an element of $\hat{\mathfrak{h}}_R$. We note that if $\alpha \in \Delta$ and $c\alpha \in \Delta$ then $c = \pm 1$. So if $c_\beta^{w'} \neq 0$ then $(\sum_{n=0}^\infty n d''_{\beta,n})/c_\beta^{w'} = \pm 1$ and $w'\beta = \pm w\alpha$. Further since the flow $P_\alpha^w(t)$ arrives at w' , we have $w'\beta = -w\alpha$ and $\beta = -w'^{-1}w\alpha$. We complete the proof.

Remark 4.8. From the above proposition, it is easily shown that the flow $P_\alpha^w(t)$ does not lie on any open set $w''\bar{N}^*$, $w'' \in W^1$, except $w\bar{N}^*$ and $w'\bar{N}^*$.

THEOREM 4.9. *Let $P_\alpha^w(t)$ be the flow as (4.11) and assume $w\alpha \in \Delta_+$, i.e., $\lim_{t \rightarrow -\infty} P_\alpha^w(t) = w$. We consider the space X as the T invariant submanifold of $\mathbb{P}^N(V)$ and identify the point $w \in W^1$ and $w\Lambda \in \langle \Lambda \rangle^e$. Then the following three conditions are all equivalent;*

- 1) $\lim_{t \rightarrow +\infty} P_\alpha^w(t) = w'\Lambda \in \langle \Lambda \rangle^e$,
- 2) $w'\Lambda - w\Lambda = -g \cdot w\alpha \in \hat{\mathfrak{h}}_R$,

where

$$g = \frac{2(\Lambda, \alpha)}{(\alpha, \alpha)},$$

3) $w^{-1}S_{w\alpha}w' \in W_1$.

Proof. If we change the local coordinate from $\{z_\alpha; u_i\}_{\alpha \in \Lambda(n^*), i \in I}$ to $\{y_\delta\}_{\delta \in \langle \Lambda \rangle - w\Lambda}$ around $i(w) = w\Lambda$ then we have the convergence power series

$$\begin{aligned} (4.14) \quad y_\delta(P_\alpha^w(t)) &= \sum_{n=0}^{\infty} d_{\delta,n} (c_\alpha^w e^{(w\alpha)(H)t})^n \\ &= \sum_{n=0}^{\infty} d'_{\delta,n} e^{n(w\alpha)(H)t} \end{aligned}$$

for $\delta \in \langle \Lambda \rangle - w\Lambda$. But any flow $P(t)$ of V_H^A on $U_{w\Lambda}$ must be written as (4.8). So if we compare the coefficients of t in (4.8) and (4.14), we have

$$b_\delta(\delta - w\Lambda)(H) = \left(\sum_{n=0}^{\infty} n d'_{\delta,n} \right) (w\alpha)(H).$$

And from Remark 4.4, we have

$$b_\delta(\delta - w\Lambda) = \left(\sum_{n=0}^{\infty} n d'_{\delta,n} \right) (w\alpha)$$

as an element of $\hat{\mathfrak{h}}_R$.

The limit point $\lim_{t \rightarrow +\infty} P_\alpha^w(t)$ is determined by comparing the magnitude of the values $\{(\delta - w\Lambda)(H) \mid b_\delta \neq 0\}$. And from Remark 4.6, there must exist at least one element $w'\Lambda \in \langle \Lambda \rangle^e - w\Lambda$ such that $b_{w'\Lambda} \neq 0$. Then we have

$$b_{w'\Lambda}(w'\Lambda - w\Lambda) = \left(\sum_{n=0}^{\infty} n d'_{w'\Lambda,n} \right) (w\alpha) \in \hat{\mathfrak{h}}_R$$

and hence

$$w'\Lambda - w\Lambda = c w\alpha,$$

where

$$c = \left(\sum_{n=0}^{\infty} n d'_{w'\Lambda,n} \right) / b_{w'\Lambda}.$$

Fortunately this condition determines the value c uniquely as

$$c = -g = -2 \frac{(\Lambda, \alpha)}{(\alpha, \alpha)}$$

and consequently the extremal weight $w'\Lambda$ is also uniquely determined. If

$$c \neq -g = -2 \frac{(A, \alpha)}{(\alpha, \alpha)},$$

we have three distinct extremal weights $A, w^{-1}w'A = A + c\alpha$ and $S_\alpha A = A - g\alpha$ lying on a straight line. But since the elements of the Weyl group W act on $\hat{\mathfrak{h}}_R$ as orthogonal transformations with respect to the norm $(,)$ determined by the Killing form, the points $A, w^{-1}w'A$ and $S_\alpha A$ also lie on the hypersphere of radius $|A| = (A, A)^{1/2}$. This is a contradiction. Hence we have

$$c = -g = -2 \frac{(A, \alpha)}{(\alpha, \alpha)}$$

and then $w^{-1}w'A = A - g\alpha = S_\alpha A$. Since W_1 is the isotropy subgroup of W at A , we have $S_\alpha w^{-1}w' = w^{-1}wS_\alpha w^{-1}w' = w^{-1}S_{w\alpha}w' \in W_1$. We complete the proof.

Remark 4.10. In the case $X = G/B$, we have $W_1 = \{1\}$. So the condition 3) means $S_{w\alpha}w' = w$.

Remark 4.11. If we change the representatives w, w' of $\bar{w}, \bar{w}' \in W^1 = W/W_1$ to w_1, w'_1 respectively, where $w_1, w'_1 \in W_1$, we have

$$w^{-1}S_{w\alpha}w' = w_1(w_1w_1)^{-1}S_{(w_1w_1)(w_1^{-1}\alpha)}(w'_1w'_1)w_1'^{-1} \in W_1$$

and hence

$$(w_1w_1)^{-1}S_{(w_1w_1)(w_1^{-1}\alpha)}(w'_1w'_1) \in W_1.$$

We note that from Remark 2.1., $w_1^{-1}\alpha$ belongs to $\Delta(n^*)$.

THEOREM 4.12. *For an element $w' \in W^1$, the following are equivalent:*

- 1) $\bar{X}_{w'}$ is contained in \bar{X}_w , i.e., $w' \leq w$,
- 2) w' belongs to \bar{X}_w ,
- 3) *There exist two series $w = w_0, w_1, \dots, w_k = w' \in W^1$ and $\alpha_0, \alpha_1, \dots, \alpha_{k-1} \in \Delta(n^*)$ such that*

$$w_i^{-1}S_{w_i\alpha_i}w_{i+1} \in W_1 \quad \text{and} \quad w_i\alpha_i \in \Delta_+, \quad 0 \leq i \leq k-1.$$

Proof. 1) \Rightarrow 2) is obvious. 2) \Rightarrow 1). If w' belongs to \bar{X}_w , since \bar{X}_w is N invariant, $X_{w'} = N\bar{w}' \subset \bar{X}_w$. Hence we have $\bar{X}_{w'} \subset \bar{X}_w$. 3) \Rightarrow 2). By virtue of Theorem 4.9., we have

$$w_1 = \lim_{t \rightarrow +\infty} P_{\alpha_0}^{w_0}(t) \in \bar{X}_{w_0} \quad \text{and} \quad w_2 = \lim_{t \rightarrow +\infty} P_{\alpha_1}^{w_1}(t) \in \bar{X}_{w_1} \subset \bar{X}_{w_0}.$$

Repeating this procedure we can show $w_k = w' \in \bar{X}_w$. Let us prove the converse 2) \Rightarrow 3) by induction on the dimension of \bar{X}_w . If $\dim \bar{X}_w = 0$, it is obvious. Hence we assume that if $\dim \bar{X}_w = n - 1$, the assertion has been already proved. Put $\dim \bar{X}_w = n$ and choose an element $w' \in W^1$ which belongs to \bar{X}_w . If $\dim \bar{X}_{w'} \leq n - 1$, we select $\bar{X}_{w''}$ such that $\bar{X}_{w'} \subset \bar{X}_{w''} \subset \bar{X}_w$ and $\dim \bar{X}_{w''} = n - 1$. From the induction there exist two series $w'' = w_0, w_1, \dots, w_{k-1} = w' \in W^1$ and $\alpha_0, \alpha_1, \dots, \alpha_{k-2} \in \mathcal{A}(n^*)$ such that

$$w_i^{-1} S_{w_i \alpha_i} w_{i+1} \in W_1 \quad \text{and} \quad w_i \alpha_i \in \mathcal{A}_+, \quad 0 \leq i \leq k - 2.$$

So it is sufficient to show that if $\dim \bar{X}_{w'} = n - 1$ and $\bar{X}_{w'} \subset \bar{X}_w$, there exists an element $\alpha \in \mathcal{A}(n^*)$ such that

$$w'^{-1} S_{w' \alpha} w' \in W_1 \quad \text{and} \quad w' \alpha \in \mathcal{A}_+.$$

We denote

$$U_{w'} = \{x \in w' \bar{N}^* \mid z_\alpha(x) = 0 \text{ for any } \alpha \in \mathcal{A}_+(n^*, w')\}.$$

Namely, $U_{w'}$ is the union of all flows of V_H which arrives at w' . Then $U_{w'}$ is invariant by the action of T and $\dim U_{w'} = \dim X - \dim \bar{X}_{w'} = \dim n^* - (n - 1)$. From the definition of $U_{w'}$, we have $w' = X_{w'} \cap U_{w'} \subseteq \bar{X}_w \cap U_{w'}$ and $\dim \bar{X}_w \cap U_{w'} = 1$. So there exists an element x such that $w' \neq x \in \bar{X}_w \cap U_{w'}$. By virtue of T invariantness of the both sets \bar{X}_w and $U_{w'}$, the flow $P(t)$ of V_H which passes through x lies entirely on $\bar{X}_w \cap U_{w'}$. And from the definition of $U_{w'}$, $\lim_{t \rightarrow +\infty} P(t) = w'$. If $\lim_{t \rightarrow -\infty} P(t) = w'' \neq w$ then we have $\bar{X}_{w''} \subseteq \bar{X}_{w'} \subseteq \bar{X}_w$. This contradicts to our assumption. So $\lim_{t \rightarrow -\infty} P(t) = w$. The flow $P(t)$ starts from w so $\mathcal{A}_-(n^*, w, P(t)) = \emptyset$. We denote

$$T \cdot P = \{\exp(H') \cdot P(t) \in X \mid H' \in \mathfrak{h}, t \in \mathbb{C}\}.$$

From the T invariantness of \bar{X}_w and $U_{w'}$, the set $T \cdot P$ is contained in $\bar{X}_w \cap U_{w'}$. If the number of $\mathcal{A}_+(n^*, w, P(t))$ is greater than two, from Theorem 3.3, $\dim T \cdot P \geq 2$. This is a contradiction. Hence there exists an element $\alpha \in \mathcal{A}(n^*)$ such that $\alpha = \mathcal{A}_+(n^*, w, P(t))$ and then the flow $P(t)$ coincides with the distinguished flow $P_\alpha(t)$. By virtue of Theorem 4.9., we complete the proof.

Remark 4.13. From Remark 4.11 the condition 3) is independent of the choice of the representatives of W^1 in W .

EXAMPLE 4.14. Let $P^3 = SL(4, \mathbb{C})/P$, where

$$P = \left\{ \left(\begin{array}{cc|c} *_1 & *_2 & \\ \hline 0 & & \\ 0 & *_3 & \\ 0 & & \end{array} \right) \in SL(4, \mathbf{C}) \right\}.$$

The Weyl group W of $SL(4, \mathbf{C})$ is isomorphic to \mathfrak{S}_4 and the subgroup W_1 is isomorphic to $\{1\} \times \mathfrak{S}_3$. Hence we have $W^1 = \{w_0, w_1, w_2, w_3\}$. We can assume $w_i \in W^1(i)$, $0 \leq i \leq 3$. We obtain the following figure.




Fig. 3

REFERENCES

- [1] I. N. Bernstein, I. M. Gel'fand and S. I. Gel'fand, Schubert cells and cohomology of the spaces G/P , *Russ. Math. Surveys*, **28** (1975), 1–26.
- [2] ———, Differential operators on the base affine space and a study of \mathfrak{g} -modules, *Lie groups and their representations*, Halsted Press (1975), 21–64.
- [3] J. Dixmier, *Enveloping algebras*, North-Holland Mathematical Library, Vol. 14.
- [4] V. V. Deodhar, Some characterizations of Bruhat ordering on a Coxeter group and determination of the relative Möbius function, *Inventiones Math.*, **39** (1977), 187–198.
- [5] Y. Kato, On the vector fields on an algebraic homogeneous space, preprint.
- [6] B. Kostant, Lie algebra cohomology and generalized Borel-Weil theorem, *Ann. Math.*, **74**(2) (1961), 329–387.
- [7] ———, Lie algebra cohomology and generalized Schubert cells, *Ann. Math.*, **77**(1) (1963), 72–144.
- [8] A. J. Sommese, Holomorphic vector fields on compact kaehler manifolds, *Math. Ann.*, **210** (1974), 75–82.
- [9] V. S. Varadarajan, *Lie groups, Lie algebras and their representations*, Prentice-Hall, Inc.
- [10] N. Wallach, *Harmonic analysis on homogeneous spaces*, Marcel Dekker, Inc.
- [11] G. Warner, *Harmonic analysis on semisimple Lie groups I*, Springer Verlag.

Nagoya University