

# Quantitative Spectral Analyses of CSPNs of Early [WC]-Type

L. Koesterke and W.-R. Hamann

Institut für Theoretische Physik und Astrophysik der Universität Potsdam

## Introduction

Among the Central Stars of Planetary Nebulae (CSPN) there are several stars which show Wolf-Rayet-type spectra resembling those of Pop. I Wolf-Rayet (WR) stars of the carbon sequence (WC). Due to progress in computer technology and new solution techniques it became possible ten years ago to calculate models which account for the very complex physical conditions in Wolf-Rayet atmospheres (Hillier 1989, Hamann et al. 1992). These models have been successfully applied to the vast majority of Pop. I WR stars in the Galaxy and the Magellanic Clouds (Hamann et al. 1995) and, in the last three years, to an increasing sample of CSPNs of [WC] type (Koesterke & Hamann 1996, Leuenhagen & Hamann 1996, Leuenhagen this meeting). Here we present the analyses of ten CSPNs of early [WC] type, i.e. from [WC 2] to [WC 4].

## Model atmospheres and observations

The model atmospheres are based on the so-called *Standard Model* for Wolf-Rayet stars, assuming homogeneity and stationarity of a spherically symmetric outflow. The consistent solution of the equations of the radiation transfer in the co-moving frame and equations of statistical equilibrium in non-LTE is achieved by the Accelerated Lambda Iteration (ALI). Up to 200 non-LTE levels are accounted for. Each model is specified by the basic parameters  $T_*$ ,  $R_*$ ,  $\dot{M}$ ,  $v_\infty$  and the chemical abundances  $\beta_{\text{He}}$ ,  $\beta_{\text{C}}$  and  $\beta_{\text{O}}$ . Optical spectra of the program stars have been collected at the DSAZ (Calar Alto, Spain) and at ESO (La Silla, Chile). Three spectra have been kindly provided by M. Peña. IUE spectra were retrieved from the IUEDAC.

## Results and discussion


While  $R_t = R_* \left( \frac{v_\infty}{2500 \text{ km/s}} / \frac{\dot{M}}{10^{-4} M_\odot/\text{yr}} \right)^{2/3}$  can be determined spectroscopically,  $\dot{M}$  and  $R_*$  scale with the adopted distance which is not known but inferred from the assumption of  $L = 5000 L_\odot$  as a typical value for CSPNs.

The determination of nitrogen remains uncertain for fainter stars for which no useful IUE observation in the short wavelength range (covering N V 1240 Å) is available. The program stars (diamonds) are shown in the figure together with late-type [WC] stars (squares) and Weak Emission Line Stars (WELS, circles). The [WCE]-stars build up a homogeneous group between 125 and 150 kK and transformed radii between 2.0 and 6.3  $R_\odot$ . It is remarkable that the stellar parameters and abundances are not correlated with the spectral subtype, although there is a weak trend to lower terminal velocities for [WC 4] stars.

Usually it is supposed that hydrogen-deficient post-AGB stars are formed within the “born-again scenario”. In this scenario the stars are thrown back to the AGB when they suffer a late thermal pulse after having arrived at the White Dwarf cooling sequence. The hydrogen-rich layers are completely ejected and the bare hydrogen-deficient core becomes visible. Together with analyses of [WCL] (Leuenhagen & Hamann 1996) and PG-1159

### III. Central Stars

Name	HCLS	Type	$m_B$ [mag]	$T_*$ [kK]	$R_t$ [ $R_\odot$ ]	$R_*$ [ $R_\odot$ ]	$\dot{M}$ [ $M_\odot/\text{yr}$ ]	$v_\infty$ [km/s]	$\beta_C$	$\beta_O$	$\beta_N$
									mass fraction		
NGC 5189	[WR 13]	[WC 2]	14.9	135.0	5.0	0.13	-6.30	3000	0.16	0.08	0.003
PB 6		[WC 2]	17.4	140.0	4.5	0.12	-6.28	3000	0.24	0.14	0.003
NGC 2452	[WR 10]	[WC 2]	17.7	141.0	4.0	0.12	-6.20	3000	0.15	0.07	0:
NGC 2867	[WR 12]	[WC 2]	16.6	141.0	4.0	0.12	-6.24	1800	0.25	0.09	0:
NGC 6905	[WR 39]	[WC 2-3]	16.3	141.0	3.4	0.12	-6.32	1800	0.25	0.15	0
NGC 7026	[WR 40]	[WC 3]	15.3	130.5	6.3	0.14	-6.34	3500	0.20	0.10	0:
He 2-55		[WC 3]	17.6	128.0	5.2	0.15	-6.25	3000	0.15	0.06	0:
Sand 3		[WC 3]	(14.2V)	140.0	3.0	0.12	-6.15	2200	0.26	0.12	0.005
IC 1747	[WR 3]	[WC 4]	15.8	126.0	6.3	0.15	-6.58	1800	0.25	0.06	0
NGC 1501	[WR 6]	[WC 4]	15.2	134.5	3.5	0.13	-6.28	1800	0.35	0.15	0
NGC 6369		[WC 4]	17.0	150.0	1.8	0.11	-6.15	1200	0.30	0.14	+



stars (Dreizler et al. 1995), the results presented here give constraints concerning the post-AGB stages of low- and intermediate mass stars. Unfortunately detailed evolutionary tracks are not available so far but, hopefully, in the near future a detailed comparison between the analyses and the predictions of evolutionary calculations become practicable.

#### References

- Hamann W.-R., Leuenhagen U., Koesterke L., Wessolowski U. 1992, *A&A* 255, 200  
 Hamann W.-R., Koesterke L., Wessolowski U. 1995, *A&A* 299, 151  
 van der Hucht K.A., Conti P.S., Lundström I., Stenholm B. 181, *Space Sci. Rev.* 28, 227  
 Hillier D.J. 1989, *ApJ* 347, 392  
 Koesterke L., Hamann W.-R. 1996, *A&A* in press  
 Leuenhagen U., Hamann W.-R. 1996, *A&A* 312, 167  
 Dreizler S., Werner K., Heber U. 1995, in "White Dwarfs", Lecture Notes in Physics, D. Koester & K. Werner (eds), Springer-Verlag, p. 160